



## Diversity of staphylococcal species in food producing animals in Spain, with detection of PVL-positive MRSA ST8 (USA300)



Olouwafemi Mistourath Mama<sup>a</sup>, Elena Gómez-Sanz<sup>b,c</sup>, Laura Ruiz-Ripa<sup>a</sup>, Paula Gómez<sup>a</sup>, Carmen Torres<sup>a,\*</sup>

<sup>a</sup> Área de Bioquímica y Biología Molecular, Universidad de La Rioja, Logroño, Spain

<sup>b</sup> Laboratory of Food Microbiology, Institute of Food, Nutrition and Health, ETH Zurich, Zurich, Switzerland

<sup>c</sup> Área de Microbiología Molecular, Centro de Investigación Biomédica de La Rioja (CIBIR), Logroño, Spain

### ARTICLE INFO

#### Keywords:

*Staphylococcus* spp  
MRSA USA300  
PVL  
IEC  
Virulence genes  
Food producing animals

### ABSTRACT

This work aimed to determine the prevalence, diversity, antibiotic-resistance phenotype/genotype and virulence factors in staphylococci of farm-animals. Nasal samples of 117 farm-animals (calve: 72; lamb: 37; goat: 8) were collected from one slaughterhouse in La Rioja/Spain and cultured for staphylococci and methicillin-resistant *Staphylococcus* (MRS) recovery. Identification was performed by MALDI-TOF. Antimicrobial resistance phenotype/genotype was determined by susceptibility testing and specific PCRs. Molecular typing (*spa*-typing, multilocus-sequence-typing, *agr*-typing, SCC*mec*), and detection of 12 virulence genes and human Immune-evasive-cluster (IEC) genes were performed by PCR/sequencing in *S. aureus*. Two marker genes of arginine catabolic mobile element (ACME) were determined by PCR (USA300-MRSA detection). Staphylococci were identified in 50%, 54% and 21% of goat, lamb and calve samples, respectively. Among the 13 *S. aureus* isolates recovered, 11 were susceptible to all antimicrobials tested, and two were multidrug-resistant-MRSA [beta-lactams (*bla<sub>Z</sub>*, *mecA*), macrolides [(*msr(A)*/*msr(B)*) and fluoroquinolones]. The MSSA harboured either *tst* or enterotoxin genes, while the MRSA harboured the *lukF/lukS-PV* genes. Five sequence-types were detected. The two MRSA strains (from lamb and goat) were typed as t5173/ST8/*agr-1*/SCC*mec-IVa*/ACME-positive, corresponding to USA300 clone, and were IEC-B-positive. Among the 47 coagulase-negative staphylococci (CoNS), six species were identified, predominating *S. simulans* (n = 25) and *S. sciuri* (n = 11). Fifty-three percent of CoNS showed resistance to at least one antimicrobial agent (six multidrug-resistant strains), and the following resistance phenotypes/genotypes were detected: streptomycin [27.6%; *ant(6)-Ia*, *str*], tetracycline [23.4%; *tet(M)*, *tet(L)*, *tet(K)*], clindamycin [19.1%; *lnu(A)*, *vgaA*], erythromycin [10.6%; *erm(C)*, *msr(A)*/*msr(B)*], chloramphenicol (8.5%; *fexA*), tobramycin (6.4%), penicillin-cefoxitin (4.3%; *bla<sub>Z</sub>*, *mecA*), and SXT (2.1%). The detection of the MRSA-USA300 lineage in food animals is worrisome and should be further monitored.

### 1. Introduction

Staphylococcal species are divided in two groups: coagulase-positive staphylococci (CoPS, as *S. aureus*, *S. pseudintermedius* or *S. intermedius*, among others), and coagulase-negative staphylococci (CoNS, as *S. sciuri*, *S. epidermidis*, or *S. saprophyticus*, among others). They generally colonize the skin and mucous membranes of humans and animals (mammals and birds), but are also opportunistic pathogens (Kluytmans, 2010). In humans, staphylococci may be responsible for either skin and soft tissue infections, pneumonia or food intoxication. In livestock, especially among bovines, they cause mastitis, among others (Pantosti, 2012). Staphylococci are also common contaminants of animal-derived

foods, such as raw meats or milk-derived products (Lozano et al., 2009; Kluytmans, 2010; Becker et al., 2014). In that context, livestock is a source of staphylococci that can be transmitted to humans, highlighting the need to study and characterize the species from food producing animals.

Methicillin resistant *S. aureus* (MRSA) of the lineage CC398 is a colonizer of pigs, humans exposed to pig farming and other farm species. MRSA CC398 is also implicated in farm-related infections, and it is therefore designated as livestock associated (LA) MRSA (Benito et al., 2014b; Pantosti, 2012). Furthermore, other *S. aureus* clonal lineages such as CC130, CC599, CC59, CC9, CC133 or CC425 are animal-adapted lineages, causing infections in animals and zoonosis in humans

\* Corresponding author at: Área de Bioquímica y Biología Molecular, Universidad de La Rioja, Madre de Dios 51, 26006, Logroño, Spain.  
E-mail address: [carmen.torres@unirioja.es](mailto:carmen.torres@unirioja.es) (C. Torres).

(Gharsa et al., 2012; Cuny et al., 2013; Ben Said et al., 2017). MRSA is of great concern for public health, partly due to its pathogenic potential. *S. aureus* disposes a large variety of virulence factors, such as the staphylococcal enterotoxins, toxic shock syndrome toxin (TSST-1), or Pantone-Valentine leucocidin (PVL), among others (Jans et al., 2017). A very pathogenic and invasive MRSA clone globally distributed, named USA300, initially emerged in the USA since 1999 in very crowded-communities, but was then spread to the general population, becoming an important cause of skin and soft-tissue infections (SSTIs) (Glaser et al., 2016; Planet, 2017). MRSA USA300 was associated to the community [Community-Associated (CA)], and combined the following molecular characteristics: multilocus sequence type 8 (ST8), staphylococcal cassette chromosome *mec* type IVa (SCC*mec*IVa) encoding *mecA*, presence of PVL, and presence of the arginine catabolic mobile element (ACME) in 99.9% of the strains (Diep et al., 2008; Planet, 2017). MRSA USA300 clone has also been described as CA-MRSA in few occasions in some European countries, such as Spain, Switzerland or France, while closely related strains have been found in Latin America (Diep et al., 2008; Gómez-Sanz et al., 2013b; Vindel et al., 2014; Jung et al., 2016; Planet, 2017). Moreover, MRSA USA300 clone was detected in wild boar meat in Germany, and from retail meat in the USA (Kraushaar and Fetsch, 2014; Ge et al., 2017).

Since farm animals are known to be reservoirs for *Staphylococcus* spp., the objective of this work was to determine the prevalence and diversity of staphylococcal species in food producing animals such as calves, lamb and goat, and to study their antimicrobial resistance phenotype/genotype as well as the content in genes encoding virulence factors.

## 2. Material and methods

### 2.1. *Staphylococcus* spp. isolation and identification

Nasal samples of 117 farm animals (calve, 72; lamb, 37; goat, 8) were collected from one slaughterhouse in La Rioja, Spain, during the period of June-September 2009. The calves came from 4 farms, the lambs from 2 farms and the goats from one farm. Samples were inoculated into mannitol-salt-agar and oxacillin-resistance-screening-agar-base for staphylococci and methicillin-resistant *Staphylococcus* (MRS) recovery, respectively. After incubation at 37 °C for 24 h, up to five colonies with different staphylococcal morphology were submitted to identification using the matrix-assisted laser desorption/ionization time of flight mass spectrometry (MALDI-TOF, Bruker). For identification of *S. pseudintermedius*, digestion of *pta* gene amplicon with *Mbo*I endonuclease was performed by PCR plus restriction fragment length polymorphism (RFLP) (Bannoehr et al., 2009). All staphylococci with different species (or *spa*-type in case of *S. aureus*) and/or different phenotype of antimicrobial resistance per sample were further characterized.

### 2.2. Molecular typing and virulence study

All coagulase-positive staphylococcal (CoPS) strains were typed. In the case of *S. aureus*, *spa*-, *agr*- and SCC*mec*-typing were performed (Benito et al., 2014a). The presence of the *agrD* gene was tested in *S. pseudintermedius* isolates (Bannoehr et al., 2007). Multilocus sequence typing (MLST) was performed for both species. For this purpose, PCR and partial sequencing of seven housekeeping genes for *S. aureus* and *S. pseudintermedius* ([www.pubmlst.org](http://www.pubmlst.org)) were performed to determine the sequence type (ST) and the clonal complex (CC), as previously described (Enright et al., 2000; Bannoehr et al., 2007; Solyman et al., 2013).

The presence of the genes encoding Pantone Valentine Leucocidin (*lukF/lukS-PV*), exfoliative toxins (*eta* and *etb*), toxic shock syndrome toxin (*tst*) and enterotoxins genes (*sea*, *see*, *seg*, *sei*, *sem*, *sen*, *seo*, *seu*) was studied by PCR for *S. aureus* (Benito et al., 2014a; Hwang et al., 2007).

The gene *scn*, belonging to the human Immune Evasive Cluster (IEC), was studied for *S. aureus* isolates, and the IEC type was determined for *scn*-positive isolates (Benito et al., 2014b). In addition, the virulence genes for the *S. pseudintermedius* leucocidin (*lukS/F-I*), exfoliative toxins (*siet*, *expA*, *expB*) and enterotoxin (*sec<sub>canine</sub>*) were tested for *S. pseudintermedius* isolates (Gharsa et al., 2015).

Specific PCRs were performed to determine the presence of the ACME in the CC8 strains, using the primer pairs AIPS.27 + AIPS.28 and AIPS.45 + AIPS.46 for *arcA*, and *opp3* genes, respectively (Diep et al., 2008). The PCR program used in this study consists of an initial denaturation for 7 min at 94 °C, followed by 30 cycles of denaturation for 1 min at 94 °C, annealing for 1 min at either 55 °C (*arcA*) or 61 °C (*opp3*), and an extension for 1 min at 72 °C, and a final extension for 10 min at 72 °C.

### 2.3. Antibiotic susceptibility testing and antibiotic resistance gene detection

For all the isolates (CoPS and CoNS), susceptibility to penicillin, cefoxitin (oxacillin in the case of *S. pseudintermedius*), gentamicin, tobramycin, tetracycline, chloramphenicol, erythromycin, clindamycin, ciprofloxacin, linezolid, and trimethoprim/sulfamethoxazole (SXT) was analysed by disk-diffusion method (CLSI, 2018). In addition, streptomycin susceptibility was also tested (CASFM, 2018). The detection of the following antimicrobial resistance genes was performed by PCR, according to the phenotype of resistance: beta-lactams (*blaZ*, *mecA*, and *mecC*), tetracycline [*tet(K)*, *tet(M)*, and *tet(L)*], aminoglycosides [*aac* (6')-*aph*(2''), and *ant*(4')-Ia], macrolides-lincosamides [*erm(A)*, *erm(B)*, *erm(C)*, *erm(T)*, *msr(A)*/*msr(B)*, *lnu(A)*, *lnu(B)*, and *vgaA*], streptomycin [*str*, and *ant*(6)-Ia] trimethoprim (*dfrA*, *dfrD*, *dfrG*, and *dfrK*) and phenicols (*cat<sub>pc194</sub>*, *cat<sub>pc221</sub>*, *cat<sub>pc223</sub>*, *cfr*, *fexA*, *optrA*, and *poxtA*) (Sutcliffe et al., 1996; Kehrenberg and Schwarz, 2005; Schnellmann et al., 2006; Gómez-Sanz et al., 2013b; Liu et al., 2013; Benito et al., 2014b; Wang et al., 2015; Antonelli et al., 2018).

## 3. Results

### 3.1. Detection of staphylococcal species

*Staphylococcus* spp. (CoPS and CoNS) were detected in 50%, 54% and 21% of goat, lamb and calve samples, respectively. Twelve animals harboured CoPS isolates, and 14 isolates were obtained (one isolate/animal, except for two animals in which two *S. aureus* of different *spa*-types were obtained). In addition, 33 animals harboured CoNS and 47 isolates were recovered (1–5 isolates/sample, with different species or antibiotic resistance phenotypes). Six of the tested animals harboured both CoPS and CoNS isolates.

Thirteen of the 14 CoPS isolates were identified as *S. aureus*, and one isolate as *S. pseudintermedius* (lamb origin). *S. aureus* isolates were recovered from lamb (n = 11; 29.7%), goat (n = 1; 12.5%) and calve samples (n = 1; 1.4%) (Table 1). The 47 CoNS isolates (33/117 animals) belonged to six distinct species, with predominance of *S. simulans* (n = 25) and *S. sciuri* (n = 11), followed by *S. cohnii* (n = 5), *S. chromogenes* (n = 4), *S. epidermidis* (n = 1) and *S. hominis* (n = 1) (Table 2).

### 3.2. Coagulase-positive *Staphylococcus*: molecular typing, antimicrobial resistance and virulence

Among the 13 *S. aureus* isolates, six *spa*-types were detected, including one novel, registered as t1754, and five STs, two of them being especially relevant: ST8-CC8 (n = 2) and ST133/CC133 (n = 7) (Table 1). The *S. aureus* isolates were all typed *agr*-I, except for one *agr*-IV. Most of the strains (n = 11, 84.6%) showed susceptibility to all the antimicrobials tested, while only two (15.4%) showed resistance to some antimicrobials. The nine isolates of the clonal complexes CC133 and CC522 were included among the antimicrobial susceptible strains; they all harboured the virulence gene *tst* and were isolated from lamb

**Table 1**  
Phenotypic and genotypic characteristics of coagulase-positive *Staphylococcus* isolates from different origins.

Origin	Species	Number of isolates	ST-CC-SCCmec	spa-type (N° of isolates)	agr	Antimicrobial resistance			Virulence genes	scn gene	IEC type	
						Phenotype <sup>c</sup>	Genotype	Genotype				
Lamb	<i>S. aureus</i>	7	ST133-CC133	t17754 <sup>b</sup> (4)	I	Susceptible	–	–	tst	–	–	
		2	ST522-CC522	t2678 (3)	I	Susceptible	–	–	tst	–	–	
	1	ST18-CC8-IV <sup>a</sup>	t1534	I	Susceptible	blaZ-mecA-msr(A)/msr(B)	–	–	lukF/lukS-PV	+	B	
	1	ST2329	t5173	I	PEN-FOX-ERY-CIP	–	–	–	–	–	–	
Goat	<i>S. pseudintermedius</i>	1	ST932 <sup>b</sup>	t8765	IV	Susceptible	–	–	sei-sem-sei-seu	–	–	
		1	ST18-CC8-IV <sup>a</sup>	–	II	TET	tet(M)	–	–	lukF/S-I-siet-expA-expB	–	–
Calve	<i>S. aureus</i>	1	ST45-CC45	t5173	I	PEN-FOX-ERY-CIP	blaZ-mecA-msr(A)/msr(B)	–	–	–	–	B
		1	–	t505	I	Susceptible	–	–	–	–	–	–

<sup>a</sup> strain positive in the ACME specific PCR.

<sup>b</sup> new spa-type or new ST.

<sup>c</sup> PEN: penicillin; FOX: cefoxitin; TET: tetracycline, ERY: erythromycin, CIP: ciprofloxacin.

samples. The two other antimicrobial susceptible isolates (ST45 and ST2329) hosted enterotoxin genes (*seg*, *sei*, *sem*, *sen*, *seo*, *seu*), while only one was positive for the *scn* gene (IEC type-B) (Table 1). The remaining two isolates (recovered from lamb and goat samples) showed resistance to cefoxitin (*mecA*), penicillin (*blaZ*), erythromycin [*msr(A)*/*msr(B)*], and ciprofloxacin, and harboured the gene encoding PVL, as well as the *scn* gene (IEC type B). Both MRSA strains were typed as t5173/ST8/CC8/*agr*-I/SCCmecIVa, thus, considered as the USA300 clone (Table 1). Furthermore, they were positive for the ACME marker genes (*arcA* and *opp3* genes).

The *S. pseudintermedius* strain, recovered from a lamb sample, was resistant to tetracycline [carrying the *tet(M)* gene] and harboured the following virulence genes: *lukS/F-I*, *siet*, *expA* and *expB*. This strain was typed as *agrD*-II and presented a new MLST allelic combination (*ack*:1; *cpn60*:9; *fdh*:2; *pta*:1; *purA*:8; *sar*:1; *tuf*:1), rendering a new sequence type registered as ST932. The ST932 is a double locus variant of the known ST309.

### 3.3. Coagulase-negative *Staphylococcus*: antimicrobial resistance phenotype and genotype

The CoNS isolates presented diverse resistance phenotypes and genotypes, which are summarized in Table 2. Twenty-five isolates (53%) were susceptible to all the tested antimicrobial agents while twenty-two (47%) were resistant to at least one of them. Among the resistant isolates, six strains of the species *S. simulans* (3/25), *S. sciuri* (2/11) and *S. cohnii* (1/5) presented a multidrug resistance phenotype (including at least three antimicrobial families). The following resistance rates were detected among the CoNS isolates: streptomycin (27.6%), tetracycline (23.4%), clindamycin (19.1%), erythromycin (10.6%), chloramphenicol (8.5%), tobramycin (6.4%), penicillin (4.3%), cefoxitin (4.3%), and SXT (2.1%). Streptomycin resistance was mediated by *str* gene for most strains, and *ant(6)-Ia* gene in one case. Resistance to tetracycline was encoded by either *tet(K)* or *tet(L)* genes in the strains from calve, and by *tet(K)* alone or associated to either *tet(L)* or *tet(M)* in the isolates from lamb (Table 2). *lnu(A)* gene was detected in 66.6% of the clindamycin-resistant strains, and *vgaA* gene in 22.2% of them. Resistance to erythromycin was mediated by either *erm(C)* or *msr(A)*/*msr(B)*. All the chloramphenicol-resistant strains harboured the *fxeA* gene but were negative for the *oprA*, *poxtA* and *cfr* genes. The penicillin and cefoxitin resistant isolates carried *blaZ* and *mecA* genes, respectively, and SXT-resistant isolates harboured the *dfrK* gene.

## 4. Discussion

Staphylococcal species are opportunistic pathogens and commensals for humans and animals (Pantosti, 2012). Animal-to-human transmission of bacteria may occur by either contact with the animals or consumption of animal-derived products (Kluytmans, 2010), which can contribute to the spread of resistance genes and pathogenicity factors in different ecosystems. In that context, this work aimed to isolate and characterize staphylococcal strains from food producing animals.

The prevalence rate of *S. aureus* for lamb (29.7%) was higher than the rates for goat (12.5%) and calve samples (1.4%). A similar pattern, but with higher percentages, was observed in a recent similar study performed in Greece (Papadopoulos et al., 2018). Furthermore, one isolate of *S. pseudintermedius* was detected in a lamb sample, although this staphylococcal species is generally associated to dogs (Gómez-Sanz et al., 2013a). This species has been formerly identified in equine and bovine at low rates (Rubin et al., 2011), as well as in wildlife (Mama et al., 2019a).

Molecular typing revealed CC133 as the most frequent clonal complex among *S. aureus* recovered (7/13), always as methicillin-susceptible *S. aureus* (MSSA) isolates. Lineage CC133 is particularly frequent among MSSA from small ruminants such as sheep and goat (Eriksson et al., 2013; Ben Said et al., 2017), as well as from donkeys

**Table 2**  
Coagulase negative *Staphylococcus* of different origins, and antimicrobial resistance phenotypes and genotypes.

Origin (N° of positive animals)	Species (N° of isolates)	Antibiotic resistance phenotype <sup>a</sup> (N° of isolates)	Genes detected (n° of isolates)
Lamb (16)	<i>S. simulans</i> (19)	STR (2)	<i>str</i> (2)
		STR-TET (3)	<i>tet</i> (K) (3) - <i>tet</i> (L) (1) - <i>str</i> (3)
		TET-SXT (1)	<i>tet</i> (K) - <i>tet</i> (L) - <i>dfrK</i>
		STR-TET-CLOR (1)	<i>tet</i> (K) - <i>tet</i> (M) - <i>ant</i> (6)-Ia - <i>fexA</i>
		ERY-CLIN-STR-TET-CLOR (1)	<i>tet</i> (K) - <i>tet</i> (M) - <i>erm</i> (C) - <i>fexA</i>
	<i>S. sciuri</i> (4)	Susceptible (11)	
		CLIN (1)	<i>lnu</i> (A)
		CLIN-STR (1)	<i>lnu</i> (A) - <i>str</i>
	<i>S. cohnii</i> (4)	FOX-STR-TET (1)	<i>tet</i> (K) - <i>tet</i> (L) - <i>mecA</i> - <i>str</i>
		Susceptible (3)	
<i>S. chromogenes</i> (2)	Susceptible (2)		
	<i>S. epidermidis</i> (1)	PEN	<i>blaZ</i>
Calve (14)	<i>S. simulans</i> (3)	TET (1)	<i>tet</i> (K)
		ERY-CLIN-STR-CLOR (1)	<i>fexA</i> - <i>str</i>
		Susceptible (1)	
	<i>S. sciuri</i> (7)	CLIN-TOB (1)	<i>lnu</i> (A) (1)
		TOB-TET (1)	<i>tet</i> (K)
		CLIN-TET (1)	<i>tet</i> (K) - <i>lnu</i> (A)
		FOX-STR-TOB-TET (1)	<i>tet</i> (L) - <i>mecA</i>
		CLIN-STR-TOB-TET (1)	<i>tet</i> (K) - <i>lnu</i> (A)
		Susceptible (2)	
	<i>S. cohnii</i> (1)	Susceptible (1)	
		<i>S. chromogenes</i> (2)	ERY-CLIN-CLOR (1)
	<i>S. hominis</i> (1)	ERY-CLIN-STR (1)	<i>erm</i> (C) - <i>vgaA</i> - <i>lnu</i> (A)
		PEN-ERY	<i>blaZ</i> - <i>msr</i> (A)/ <i>msr</i> (B)
Goat (3)	<i>S. simulans</i> (3)	Susceptible (3)	

<sup>a</sup> PEN: penicillin; FOX: cefoxitin; TET: tetracycline; CLIN: clindamycin; ERY: erythromycin; TOB: tobramycin; CLOR: chloramphenicol; SXT: trimethoprim/sulfamethoxazole.

(Gharsa et al., 2012). In addition, *S. aureus* CC133 has also been found in wildlife in Spain (Mama et al., 2019a). The clonal complex CC522, commonly observed among ruminants and animal-derived products (Gharsa et al., 2012; Ben Said et al., 2017), was also present among our isolates. All strains belonging to the CC133 and CC522 lineages were obtained from lamb samples and harboured the *tst* gene related to the toxic shock syndrome. This was not surprising as such gene was highly frequent among sheep strains of the same genetic lineages (Ben Said et al., 2017). On the other hand, the presence of a calve strain MSSA/ST45 harbouring the *egc*-like operon genes (*seg-sei-sem-sen-seo-seu*) is relevant. This hospital and/or community-associated clone was detected before in healthy humans (Gómez-Sanz et al., 2013b), and in meat samples in Spain (Benito et al., 2014a). In the last case, the strains were typed as IEC-B, as in our work, suggesting a human origin.

The strains ST8 (n = 2), isolated from lamb and goat and typed as t5173/*agrI*/SCC*mecIVa*, were PVL-positive and ACME-positive, thus designated as clone USA300. The clone USA300 is a CA-MRSA that emerged mainly in the USA, representing an important cause of SSTIs (Diep et al., 2008; Glaser et al., 2016; Planet, 2017). It was also sporadically found in some European countries, while a similar variant was found in Latin America (Vindel et al., 2014; Jung et al., 2016; Planet, 2017). According to some authors, the CA-MRSA dominant in the USA is the USA300 North American (USA300-NA) MRSA clone, which descended from USA500-like MSSA strains by acquisition of various mobile genetic elements (MGEs) (Glaser et al., 2016). The acquired MGEs include: a) the SCC*mecIVa* element (with *mecA* gene); b) *S. aureus* pathogenicity island 5 containing *sek* and *seq* genes; c) phage phiSA2 carrying the Pantone-Valentine leukocidin genes; and d) ACME type I, which are assumed to confer enhance pathogenicity and fitness on USA300-NA MRSA strains (Diep et al., 2008; Glaser et al., 2016; Planet, 2017). In addition, plasmid analysis showed the presence of the plasmid p18805-p03 encoding diverse antibiotic resistance genes [aminoglycosides, *ant*(6)-Ia and *aph*(3')-III; beta-lactams, *blaZ*; macrolides *mph*(C); macrolides-lincosamides-streptogramin B, *msr*(A)] (Glaser et al., 2016). Our strains could correspond to the USA300-NA clone, and the

multidrug resistance and the presence of PVL genes, suggest the acquisition of MGEs. Furthermore, the presence of IEC system (type B) confirms the potential adaptation to humans. Unexpectedly, that clone has been found, as in this work, in animals, such as an infected dog in France (Haenni et al., 2012), wild boar meat in Germany (Kraushaar and Fetsch, 2014), and retail meat in the USA (Ge et al., 2017). In those cases, the USA300 strains detected were suspected to be transmitted by humans (hypothetically the dog owner or food handlers). These data are important in the context of the One Health perspective, as it reflects how the dissemination of bacteria with their mobilizable resistance and virulence genes content can occur between the clinic, community, animals and food. Additionally, the presence of *S. pseudintermedius* in a lamb sample may suggest a contact between the animal and dogs or dog owners.

Six species of CoNS were identified among the tested animals (28% were positive): *S. simulans*, *S. sciuri*, *S. cohnii*, *S. chromogenes*, *S. epidermidis* and *S. hominis*. Those species seem to be recurrent in cattle samples (De Visscher et al., 2014; Cuny et al., 2017). CoNS could be a reservoir of clinically relevant resistance genes that could be transferred to *S. aureus* isolates (Becker et al., 2014). Half of the CoNS strains were resistant to at least one tested antimicrobial agent. The resistance phenotypes observed among our isolates were not unexpected since beta-lactams, tetracyclines, lincosamides and sulphonamides are antimicrobial families very frequently used in the veterinary sector (food producing animals and pets) (Prestinaci et al., 2015). In addition, the phenotypic and genotypic data were in accordance with a previous study on animal-derived food (Chajęcka-Wierzychowska et al., 2015). Alternatively, the *fexA* gene, though uncommon, seems to become increasingly present in CoNS isolated from animals (Mama et al., 2019b). It was first identified in a bovine *S. lentus* isolate (Kehrenberg and Schwarz, 2004), and was later found in other staphylococcal species such as *S. sciuri*, *S. simulans* and *S. aureus* from food producing animals, equidae and wild boar (Kehrenberg and Schwarz, 2006; Mama et al., 2019b). In this study, that gene was detected in *S. simulans* and *S. chromogenes* isolates. Recently, the *fexA* gene has been detected

associated with the *cfr* and *oprA* linezolid resistance genes in the same plasmid (Li et al., 2016), although those genes were absent in our *fexA* positive strains.

## 5. Conclusion

Farm animals are reservoir for *Staphylococcus* spp., including CoPS and CoNS carrying relevant resistance genes [*mecA*, *lnu(A)*, *vgaA*, *fexA*], and important virulence factors (*lukF/lukS-PV*, *tst*, enterotoxins). The presence of the CA-MRSA-USA300/ACME+/IEC-B in farm animals as well as the CA-MSSA/ST45/IEC-B is a subject of concern, which opens the question about the origin of the isolates and the transference of virulent staphylococci between different ecosystems.

## Acknowledgments

This work was supported by project SAF2016-76571-R from the Agencia Española de Investigación (AEI) and FEDER. OMM has a predoctoral fellowship of Mujeres por África-Universidad de La Rioja (Spain). LRR has a predoctoral FPI fellowship of the Universidad de La Rioja (Spain).

## References

- Antonelli, A., D'Andrea, M.M., Brenciani, A., Galeotti, C.L., Morroni, G., Pollini, S., Varaldo, P.E., Rossolini, G.M., 2018. Characterization of *poxtA*, a novel phenicol-oxazolidinone-tetracycline resistance gene from an MRSA of clinical origin. *J. Antimicrob. Chemother.* 73, 1763–1769. <https://doi.org/10.1093/jac/dky088>.
- Bannoehr, J., Ben Zakour, N.L., Waller, A.S., Guardabassi, L., Thoday, K.L., Van Den Broek, A.H., Fitzgerald, J.R., 2007. Population genetic structure of the *Staphylococcus intermedius* group: insights into *agr* diversification and the emergence of methicillin-resistant strains. *J. Bacteriol.* 189, 8685–8692. <https://doi.org/10.5021/ad.2011.23.3.338>.
- Bannoehr, J., Franco, A., Iurescia, M., Battisti, A., Fitzgerald, J.R., 2009. Molecular diagnostic identification of *Staphylococcus pseudintermedius*. *J. Clin. Microbiol.* 47, 469–471. <https://doi.org/10.1128/JCM.01915-08>.
- Becker, K., Heilmann, C., Peters, G., 2014. Coagulase-negative staphylococci. *Clin. Microbiol. Rev.* 27, 870–926. <https://doi.org/10.1128/CMR.00109-13>.
- Ben Said, M., Abbassi, M.S., Gómez, P., Ruiz-Ripa, L., Sghaier, S., El Fekih, O., Hassen, A., Torres, C., 2017. Genetic characterization of *Staphylococcus aureus* isolated from nasal samples of healthy ewes in Tunisia. High prevalence of CC130 and CC522 lineages. *Comp. Immunol. Microbiol. Infect. Dis.* 51, 37–40. <https://doi.org/10.1016/j.cimid.2017.03.002>.
- Benito, D., Gómez, P., Lozano, C., Estepa, V., Gómez-Sanz, E., Zarazaga, M., Torres, C., 2014a. Genetic lineages, antimicrobial resistance, and virulence in *Staphylococcus aureus* of meat samples in Spain: analysis of immune evasion cluster (IEC) genes. *Foodborne Pathog. Dis.* 11, 354–356. <https://doi.org/10.1089/fpd.2013.1689>.
- Benito, D., Lozano, C., Rezusta, A., Ferrer, I., Vasquez, M.A., Ceballos, S., Zarazaga, M., Revillo, M.J., Torres, C., 2014b. Characterization of tetracycline and methicillin resistant *Staphylococcus aureus* strains in a Spanish hospital: is livestock-contact a risk factor in infections caused by MRSA CC398? *Int. J. Med. Microbiol.* 304, 1226–1232. <https://doi.org/10.1016/j.ijmm.2014.09.004>.
- CASFM, 2018. Comité De l'antibiogramme De La Société Française De Microbiologie. Recommendations vétérinaires. <https://www.sfm-microbiologie.org/>.
- Chajęcka-Wierzchowska, W., Zadernowska, A., Nalepa, B., Sierpińska, M., Laniewska-Trokenheim, L., 2015. Coagulase-negative staphylococci (CoNS) isolated from ready-to-eat food of animal origin - phenotypic and genotypic antibiotic resistance. *Food Microbiol.* 46, 222–226. <https://doi.org/10.1016/j.fm.2014.08.001>.
- CLSI, 2018. Performance Standards for Antimicrobial Susceptibility Testing, 28<sup>th</sup> ed. CLSI supplement M100, clinical laboratory standards institute, Wayne, PA. <https://clsi.org/>.
- Cuny, C., Köck, R., Witte, W., 2013. Livestock associated MRSA (LA-MRSA) and its relevance for humans in Germany. *Int. J. Med. Microbiol.* 303, 331–337. <https://doi.org/10.1016/j.ijmm.2013.02.010>.
- Cuny, C., Arnold, P., Hermes, J., Eckmanns, T., Mehraj, J., Schoenfelder, S., Ziebuhr, W., Zhao, Q., Wang, Y., Feßler, A.T., Krause, G., Schwarz, S., Witte, W., 2017. Occurrence of *cfr*-mediated multiresistance in staphylococci from veal calves and pigs, from humans at the corresponding farms, and from veterinarians and their family members. *Vet. Microbiol.* 200, 88–94. <https://doi.org/10.1016/j.vetmic.2016.04.002>.
- De Visscher, A., Supré, K., Haesebrouck, F., Zadoks, R.N., Piessens, V., Van Coillie, E., Piepers, S., De Vliegher, S., 2014. Further evidence for the existence of environmental and host-associated species of coagulase-negative staphylococci in dairy cattle. *Vet. Microbiol.* 172, 466–474. <https://doi.org/10.1016/j.vetmic.2014.06.011>.
- Diep, B.A., Stone, G.G., Basuino, L., Graber, C.J., Miller, A., Etages, S., des Jones, A., Palazzolo-Ballance, A.M., Perdreau-Remington, F., Sensabaugh, G.F., DeLeo, F.R., Chambers, H.F., 2008. The arginine catabolic mobile element and staphylococcal chromosomal cassette *mec* linkage: convergence of virulence and resistance in the USA300 clone of methicillin-resistant *Staphylococcus aureus*. *J. Infect. Dis.* 197, 1523–1530. <https://doi.org/10.1086/587907>.
- Enright, M.C., Day, N.P.J., Davies, C.E., Peacock, S.J., Spratt, B.G., 2000. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J. Clin. Microbiol.* 38, 1008–1015. <https://doi.org/10.1128/jcm.01546-06>.
- Eriksson, J., Espinosa-Gongora, C., Stamphøj, I., Larsen, A.R., Guardabassi, L., 2013. Carriage frequency, diversity and methicillin resistance of *Staphylococcus aureus* in Danish small ruminants. *Vet. Microbiol.* 163, 110–115. <https://doi.org/10.1016/j.vetmic.2012.12.006>.
- Ge, B., Mukherjee, S., Hsu, C.H., Davis, J.A., Tran, T.T.T., Yang, Q., Abbott, J.W., Ayers, S.L., Young, S.R., Crarey, E.T., Womack, N.A., Zhao, S., McDermott, P.F., 2017. MRSA and multidrug-resistant *Staphylococcus aureus* in U.S. retail meats, 2010–2011. *Food Microbiol.* 62, 289–297. <https://doi.org/10.1016/j.fm.2016.10.029>.
- Gharsa, H., Ben Sallem, R., Ben Slama, K., Gómez-Sanz, E., Lozano, C., Jouini, A., Klibi, N., Zarazaga, M., Boudabous, A., Torres, C., 2012. High diversity of genetic lineages and virulence genes in nasal *Staphylococcus aureus* isolates from donkeys destined to food consumption in Tunisia with predominance of the ruminant associated CC133 lineage. *BMC Vet. Res.* 8, 203. <https://doi.org/10.1186/1746-6148-8-203>.
- Gharsa, H., Slama, K., Ben Gómez-Sanz, E., Gómez, P., Klibi, N., Zarazaga, M., Boudabous, A., Torres, C., 2015. Characterisation of nasal *Staphylococcus delphini* and *Staphylococcus pseudintermedius* isolates from healthy donkeys in Tunisia. *Equine Vet. J.* 47, 463–466. <https://doi.org/10.1111/evj.12305>.
- Glaser, P., Martins-Simões, P., Villain, A., Barbier, M., Tristan, A., Bouchier, C., 2016. Demography and intercontinental spread of the USA300 community-acquired methicillin-resistant *Staphylococcus aureus* lineage. *mBio* 7, e02183-15. <https://doi.org/10.1128/mBio.02183-15>.
- Gómez-Sanz, E., Torres, C., Ceballos, S., Lozano, C., Zarazaga, M., 2013a. Clonal dynamics of nasal *Staphylococcus aureus* and *Staphylococcus pseudintermedius* in dog-owning household members. Detection of MSSA ST398. *PLoS One* 8. <https://doi.org/10.1371/journal.pone.0069337>.
- Gómez-Sanz, E., Torres, C., Lozano, C., Zarazaga, M., 2013b. High diversity of *Staphylococcus aureus* and *Staphylococcus pseudintermedius* lineages and toxigenic traits in healthy pet-owning household members. Underestimating normal household contact? *Comp. Immunol. Microbiol. Infect. Dis.* 36, 83–94. <https://doi.org/10.1016/j.cimid.2012.10.001>.
- Haenni, M., Saras, E., Châtre, P., Médaille, C., Bes, M., Madec, J.Y., Laurent, F., 2012. A USA300 variant and other human-related methicillin-resistant *Staphylococcus aureus* strains infecting cats and dogs in France. *J. Antimicrob. Chemother.* 67, 326–329. <https://doi.org/10.1093/jac/dkr499>.
- Hwang, S.Y., Kim, S.H., Jang, E.J., Kwon, N.H., Park, Y.K., Koo, H.C., Jung, W.K., Kim, J.M., Park, Y.H., 2007. Novel multiplex PCR for the detection of the *Staphylococcus aureus* superantigen and its application to raw meat isolates in Korea. *Int. J. Food Microbiol.* 117, 99–105. <https://doi.org/10.1016/j.ijfoodmicro.2007.02.013>.
- Jans, C., Merz, A., Johler, S., Younan, M., Tanner, S.A., Kaindi, D.W.M., Wangoh, J., Bonfoh, B., Meile, L., Tasara, T., 2017. East and West African milk products are reservoirs for human and livestock-associated *Staphylococcus aureus*. *Food Microbiol.* 65, 64–73. <https://doi.org/10.1016/j.fm.2017.01.017>.
- Jung, J., Song, E.H., Park, S.Y., Lee, S., Park, S., Sung, H., Kim, M., 2016. Emergence of Pantone-Valentine leucocidin-positive ST8-methicillin-resistant *Staphylococcus aureus* (USA300 clone) in Korea causing healthcare-associated and hospital-acquired bacteraemia. *Eur. J. Clin. Microbiol. Infect. Dis.* 35, 1323–1329. <https://doi.org/10.1007/s10096-016-2668-y>.
- Kehrenberg, C., Schwarz, S., 2004. *fexA*, a novel *Staphylococcus lentus* gene encoding resistance to florfenicol and chloramphenicol. *Antimicrob. Agents Chemother.* 48, 615–618. <https://doi.org/10.1128/AAC.48.2.615>.
- Kehrenberg, C., Schwarz, S., 2005. Florfenicol-chloramphenicol exporter gene *fexA* is part of the novel transposon Tn558. *Antimicrob. Agents Chemother.* 49, 813–815. <https://doi.org/10.1128/AAC.49.2.813-815.2005>.
- Kehrenberg, C., Schwarz, S., 2006. Distribution of florfenicol resistance genes *fexA* and *cfr* among chloramphenicol-resistant *Staphylococcus* isolates. *Antimicrob. Agents Chemother.* 50, 1156–1163. <https://doi.org/10.1128/AAC.50.4.1156-1163.2006>.
- Kluytmans, J.A.J.W., 2010. Methicillin-resistant *Staphylococcus aureus* in food products: cause for concern or case for complacency? *Clin. Microbiol. Infect.* 16, 11–15. <https://doi.org/10.1111/j.1469-0691.2009.03110.x>.
- Kraushaar, B., Fetsch, A., 2014. First description of PVL-positive methicillin-resistant *Staphylococcus aureus* (MRSA) in wild boar meat. *Int. J. Food Microbiol.* 186, 68–73. <https://doi.org/10.1016/j.ijfoodmicro.2014.06.018>.
- Li, D., Wang, Y., Schwarz, S., Cai, J., Fan, R., Li, J., Feßler, A.T., Zhang, R., Wu, C., Shen, J., 2016. Co-location of the oxazolidinone resistance genes *oprA* and *cfr* on a multiresistance plasmid from *Staphylococcus sciuri*. *J. Antimicrob. Chemother.* 71, 1474–1478. <https://doi.org/10.1093/jac/dkw040>.
- Liu, Y., Wang, Y., Schwarz, S., Li, Y., Shen, Z., Zhang, Q., Wu, C., Shen, J., 2013. Transferable multiresistance plasmids carrying *cfr* in *Enterococcus* spp. from swine and farm environment. *Antimicrob. Agents Chemother.* 57, 42–48. <https://doi.org/10.1128/AAC.01605-12>.
- Lozano, C., López, M., Gómez-Sanz, E., Ruiz-Larrea, F., Torres, C., Zarazaga, M., 2009. Detection of methicillin-resistant *Staphylococcus aureus* ST398 in food samples of animal origin in Spain. *J. Antimicrob. Chemother.* 64, 1325–1346. <https://doi.org/10.1093/JAC/DKP378>.
- Mama, O.M., Ruiz-Ripa, L., Fernández-Fernández, R., González-Barrio, D., Ruiz-Fons, J.F., Torres, C., 2019a. High frequency of coagulase-positive staphylococci carriage in healthy wild boar with detection of MRSA of lineage ST398-t011. *FEMS Microbiol. Lett.* <https://doi.org/10.1093/femsle/fny292>.
- Mama, O.M., Ruiz-Ripa, L., Lozano, C., González-Barrio, D., Ruiz-Fons, J.F., Torres, C., 2019b. High diversity of coagulase negative staphylococci species in wild boars, with low antimicrobial resistance rates but detection of unusual genes. *Comp. Immunol.*

- Microbiol. Infect. Dis. 64, 125–129. <https://doi.org/10.1016/j.cimid.2019.03.006>.
- Pantosti, A., 2012. Methicillin-resistant *Staphylococcus aureus* associated with animals and its relevance to human health. *Front. Microbiol.* 3, 1–12. <https://doi.org/10.3389/fmicb.2012.00127>.
- Papadopoulos, P., Papadopoulos, T., Angelidis, A.S., Boukouvala, E., Zdragas, A., Papa, A., Hadjichristodoulou, C., Sergelidis, D., 2018. Prevalence of *Staphylococcus aureus* and of methicillin-resistant *S. aureus* (MRSA) along the production chain of dairy products in north-western Greece. *Food Microbiol.* 69, 43–50. <https://doi.org/10.1016/j.fm.2017.07.016>.
- Planet, P.J., 2017. Life after USA300: the rise and fall of a superbug. *J. Infect. Dis.* 215, S71–S77. <https://doi.org/10.1093/infdis/jiw444>.
- Prestinaci, F., Pezzotti, P., Pantosti, A., 2015. Antimicrobial resistance: a global multifaceted phenomenon. *Pathog. Glob. Health* 109, 309–318. <https://doi.org/10.1179/2047773215Y.0000000030>.
- Schnellmann, C., Gerber, V., Rossano, A., Jaquier, V., Panchaud, Y., Doherr, M.G., Thomann, A., Straub, R., Perreten, V., 2006. Presence of new *mecA* and *mph(C)* variants conferring antibiotic resistance in *Staphylococcus* spp. isolated from the skin of horses before and after clinic admission. *J. Clin. Microbiol.* 44, 4444–4454. <https://doi.org/10.1128/JCM.00868-06>.
- Solyman, S.M., Black, C.C., Duim, B., Perreten, V., Van Duijkeren, E., Wagenaar, J.A., Eberlein, L.C., Sadeghi, L.N., Videla, R., Bemis, D.A., Kania, S.A., 2013. Multilocus sequence typing for characterization of *Staphylococcus pseudintermedius*. *J. Clin. Microbiol.* 51, 306–310. <https://doi.org/10.1128/JCM.02421-12>.
- Sutcliffe, J., Grebe, T., Tait-Kamradt, A., Wondrack, L., 1996. Detection of erythromycin-resistant determinants by PCR. *Antimicrob. Agents Chemother.* 40, 2562–2566. <https://doi.org/10.1097/00005537-200307000-00001>.
- Vindel, A., Trincado, P., Cuevas, O., Ballesteros, C., Bouza, E., Cercenado, E., 2014. Molecular epidemiology of community-associated methicillin-resistant *Staphylococcus aureus* in Spain: 2004–12. *J. Antimicrob. Chemother.* 69, 2913–2919. <https://doi.org/10.1093/jac/dku232>.
- Wang, Y., Lv, Y., Cai, J., Schwarz, S., Cui, L., Hu, Z., Zhang, R., Li, J., Zhao, Q., He, T., Wang, D., Wang, Z., Shen, Y., Li, Y., Fessler, A.T., Wu, C., Yu, H., Deng, X., Xia, X., Shen, J., 2015. A novel gene, *optrA*, that confers transferable resistance to oxazolidinones and phenicols and its presence in *Enterococcus faecalis* and *Enterococcus faecium* of human and animal origin. *J. Antimicrob. Chemother.* 70, 2182–2190. <https://doi.org/10.1093/jac/dkv116>.