



# Predominance of *bla*<sub>CTX-M-65</sub> and *bla*<sub>CTX-M-55</sub> in extended-spectrum $\beta$ -lactamase-producing *Escherichia coli* from raw retail chicken in South Korea

Hyeeun Park<sup>a</sup>, Jinshil Kim<sup>a</sup>, Sangryeol Ryu<sup>a,\*</sup>, Byeonghwa Jeon<sup>a,b,\*</sup>

<sup>a</sup> Department of Food and Animal Biotechnology, Department of Agricultural Biotechnology, Research Institute for Agriculture and Life Sciences, and Center for Food and Bioconvergence, Seoul National University, Seoul 08826, South Korea

<sup>b</sup> School of Public Health, University of Alberta, Edmonton, Alberta, Canada



## ARTICLE INFO

### Article history:

Received 30 October 2018

Received in revised form 19 December 2018

Accepted 6 January 2019

Available online 15 January 2019

### Keywords:

Antimicrobial resistance

Extended-spectrum  $\beta$ -lactamase

ESBL

CTX-M

*Escherichia coli*

Chicken

## ABSTRACT

**Objectives:** Extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* (ESBL-EC) are a serious public health concern worldwide. The aim of this study was to characterise ESBL-EC isolated from raw retail chicken in South Korea.

**Methods:** The antimicrobial resistance, phylogenetic group and virulence gene prevalence of 67 ESBL-EC isolated from retail chicken in South Korea were investigated.

**Results:** All of the isolates possessed *bla*<sub>CTX-M</sub> genes, predominantly *bla*<sub>CTX-M-65</sub> (52.2%) and *bla*<sub>CTX-M-55</sub> (25.4%), and three isolates harboured both *bla*<sub>CTX-M-65</sub> and *bla*<sub>CTX-M-55</sub>. More than one-half of the ESBL-EC strains also carried *bla*<sub>TEM</sub>. Antimicrobial susceptibility testing revealed that 98.5% of the strains were multidrug-resistant (MDR). Phylogenetic analysis showed that group A was predominant (56.7%), followed by B1 (19.4%), E (8.9%), B2 (6.0%) and D (6.0%). Virulence genes associated with extraintestinal pathogenic *E. coli* (ExPEC) were frequently detected in isolates of phylogenetic groups B1, B2, D and E. **Conclusion:** The results in this study demonstrate that retail chicken in South Korea is highly contaminated with MDR ESBL-EC and may serve as a reservoir for transmitting ExPEC to humans.

© 2019 The Authors. Published by Elsevier Ltd on behalf of International Society for Chemotherapy of Infection and Cancer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Extended-spectrum  $\beta$ -lactamases (ESBLs) are a group of enzymes that hydrolyse most  $\beta$ -lactams, including penicillins, cephalosporins and monobactams, but not carbapenems [1]. Since ESBL genes are typically encoded on mobile genetic elements, usually plasmids, these genes may be easily disseminated [2]. The mobility of ESBL genes has resulted in a rapid increase in the prevalence of ESBL-producing Enterobacteriaceae in food-producing and companion animals, environmental samples (e.g. wastewater) and even food [3]. ESBL-producing *Escherichia coli* (ESBL-EC) are frequently isolated from chicken meat [4–8]. In some cases, the high prevalence of ESBL-EC hampers the isolation of other fastidious bacteria (e.g. *Campylobacter*) from chicken as it outgrows during the enrichment step using cephalosporins as a selective supplement [9].

Pathogenic *E. coli* are a major cause of not only enteric diseases but also extraintestinal infections such as urinary tract infections [10]. Although extraintestinal pathogenic *E. coli* (ExPEC) cause infections outside the intestines, ExPEC first colonise the gastrointestinal tract and are transmitted primarily by the consumption of food, particularly chicken [11]. Commensal *E. coli* isolates usually harbour no or only a very few virulence genes; however, ExPEC possess a broad range of virulence genes involved in bacterial adhesion, iron acquisition and serum survival as well as toxins associated with extraintestinal disease [12]. ExPEC commonly possess large, transmissible, multidrug resistance plasmids encoding ESBLs [13], suggesting that chicken could be a source both for ExPEC and ESBL-EC.

A number of studies have shown that retail chicken is significantly involved in transmitting ESBL-EC and ExPEC to humans [4,13,14]. Despite its public health importance, the prevalence of ESBL-EC on retail chicken in South Korea has been reported in only a single study that characterised only a limited number ( $n=6$ ) of ESBL-EC from retail chicken [15]. Furthermore, there is no study regarding ESBL-EC from retail chicken, although they may have the potential to cause

\* Corresponding authors.

E-mail addresses: [sangryu@snu.ac.kr](mailto:sangryu@snu.ac.kr) (S. Ryu), [bjeon@snu.ac.kr](mailto:bjeon@snu.ac.kr), [bjeon@ualberta.ca](mailto:bjeon@ualberta.ca) (B. Jeon).

**Table 2**Antimicrobial resistance patterns of extended-spectrum  $\beta$ -lactamase (ESBL)-producing *Escherichia coli* isolates ( $n=67$ ) according to ESBL gene type.

Antimicrobial agent	Resistance breakpoint ( $\mu\text{g}/\text{mL}$ )	No. (%) of resistant strains			
		CTX-M ( $n=29$ )	CTX-M+TEM ( $n=32$ )	CTX-M+TEM+OXA ( $n=6$ )	Total ( $n=67$ )
CIP	$\geq 4$	13 (44.8)	21 (65.6)	5 (83.3)	39 (58.2)
TET	$\geq 16$	26 (89.7)	23 (71.9)	6 (100)	55 (82.1)
CHL	$\geq 32$	19 (65.5)	25 (78.1)	6 (100)	50 (74.6)
KAN	$\geq 64$	6 (20.7)	8 (25.0)	4 (66.7)	18 (26.9)
GEN	$\geq 16$	12 (41.4)	17 (53.1)	3 (50.0)	32 (47.8)
STR	$\geq 64$	25 (86.2)	24 (75.0)	6 (100)	55 (82.1)
COL	$\geq 8$	2 (6.9)	0	0	2 (3.0)
AMP	$\geq 32$	29 (100)	32 (100)	6 (100)	67 (100)
CEF	$\geq 32$	29 (100)	32 (100)	6 (100)	67 (100)
CRO	$\geq 4$	29 (100)	32 (100)	6 (100)	67 (100)
CTX	$\geq 4$	29 (100)	32 (100)	6 (100)	67 (100)

CIP, ciprofloxacin; TET, tetracycline; CHL, chloramphenicol; KAN, kanamycin; GEN, gentamicin; STR, streptomycin; COL, colistin; AMP, ampicillin; CEF, cefalotin, CRO, ceftriaxone; CTX, cefotaxime.

extraintestinal infections in humans. To fill this important knowledge gap, in this study the antimicrobial resistance and virulence gene prevalence of ESBL-EC isolated from retail raw chicken in South Korea were characterised.

## 2. Materials and methods

### 2.1. Collection of extended-spectrum $\beta$ -lactamase-producing *E. coli* from raw retail chicken in South Korea

A total of 67 ESBL-EC were isolated from 40 retail raw whole chicken samples from 28 companies in six different provinces of South Korea in our previous study (submitted). *E. coli* were grown on MacConkey agar and were confirmed by 16S rRNA sequencing (Macrogen, Seoul, South Korea). To confirm the ESBL phenotype, the *E. coli* isolates were subjected to the modified ESBL confirmatory test of the Clinical and Laboratory Standards Institute (CLSI), which uses antimicrobial disks of cefotaxime and ceftazidime with or without the ESBL inhibitor clavulanic acid, and boric acid and ethylene diamine tetra-acetic acid (EDTA) to inhibit AmpC  $\beta$ -lactamases and carbapenemases, respectively [16]. *E. coli* ATCC 25922, a CLSI quality control strain, was used as a negative control. The presence of ESBL genes ( $bla_{SHV}$ ,  $bla_{TEM}$ ,  $bla_{CTX-M}$  and  $bla_{OXA}$ ) was determined by PCR using previously described primers [17]. PCR amplicons were sequenced and the translated amino acid sequences were used to determine the ESBL gene type [18–20]. The *E. coli* isolates were routinely cultured on Luria–Bertani medium.

### 2.2. Antimicrobial susceptibility testing

Antimicrobial susceptibility of the isolates was determined by the broth dilution method using a total of 11 antibiotics, including ampicillin, cefalotin, tetracycline, chloramphenicol, kanamycin, gentamicin, streptomycin, colistin, ceftriaxone, cefotaxime and ciprofloxacin. Minimum inhibitory concentrations (MICs) were determined according to CLSI guidelines and previous reports [21–23]. *E. coli* ATCC 25922 was used as a quality control strain according to the CLSI protocol.

### 2.3. Phylogenetic analysis of extended-spectrum $\beta$ -lactamase-producing *E. coli*

The phylogenetic group of the ESBL-EC isolates was determined using a quadruplex PCR-based method amplifying *chuA*, *yjaA*, DNA fragment *TspE4C2* and *arpA* [24]. *E. coli* strains MG1655 and ATCC 25922 were used as controls for phylogenetic groups A and B2, respectively.

### 2.4. Random amplified polymorphic DNA (RAPD) analysis

RAPD analysis was used as a PCR-based DNA fingerprinting method to analyse clonal similarity of the isolates. DNA extracted from the 67 *E. coli* isolates was subjected to PCR using previously reported primers [25] and the PCR results were analysed using BioNumerics v.7 software (Applied Maths, Sint-Martens-Latem, Belgium).

### 2.5. Detection of virulence genes associated with pathogenic *E. coli*

PCR was performed to analyse the presence of virulence genes associated with five major intestinal pathogenic *E. coli* groups, including Shiga toxin-producing *E. coli* (STEC) (*stx1* and *stx2* encoding Shiga toxins 1 and 2, *hlyA* encoding enterohaemolysin and *espP* encoding serine protease), enteropathogenic *E. coli* (EPEC) (*eaeA* encoding intimin), enterotoxigenic *E. coli* (ETEC) (*st* and *lt* encoding heat-stable and heat-labile enterotoxins) and enteroaggregative *E. coli* (EAEC) (*aggR* encoding a transcription regulator for aggregative adherence fimbria I) and ExPEC [26,27]. The tested virulence genes related to ExPEC included extraintestinal *E. coli* attachment factors (*fimH* and *iha*), iron uptake factors (*aer*, *irp2* and *iutA*), iron transporter (*feoB*), increased serum survival protein (*iss*) and heat-stable enterotoxin (*astA*) [28–30]. *E. coli* ATCC 35150 was used as a positive control for STEC and EPEC. *E. coli* NCCT 14039 was a positive control for EAEC. *E. coli* MG1655 was used as a negative control for the tested virulence genes except for *fimH* and *feoB*. *E. coli* strains ATCC 43888 and O169 were used as positive controls for *iha* and *astA*, respectively. *E. coli* strains MG1655 and ATCC 25922

**Table 1**Distribution of extended-spectrum  $\beta$ -lactamase (ESBL) types in ESBL-producing *Escherichia coli* isolates ( $n=67$ ) from raw retail chicken in South Korea.

ESBL group	ESBL type	No. of strains
CTX-M group 1	CTX-M-55	2
CTX-M group 1, TEM	CTX-M-55, TEM-1	6
	CTX-M-55, TEM-116	6
	CTX-M-15, TEM-1	1
	CTX-M-15, TEM-135	1
CTX-M group 9	CTX-M-65	17
	CTX-M-14	7
	CTX-M-27	1
CTX-M group 9, TEM	CTX-M-65, TEM-1	3
	CTX-M-65, TEM-116	7
	CTX-M-14, TEM-1	6
	CTX-M-14, TEM-116	2
	Miscellaneous	CTX-M-55, CTX-M-65
	CTX-M-65, OXA-1, TEM-1	5
	CTX-M-55, CTX-M-65, OXA-1, TEM-1	1

**Table 3**

Phylogenetic group of extended-spectrum  $\beta$ -lactamase (ESBL)-producing *Escherichia coli* isolates ( $n=67$ ) from raw retail chicken in South Korea.

Phylogenetic group	No. (%) of strains
A	38 (56.7)
B1	13 (19.4)
B2	4 (6.0)
D	4 (6.0)
E	6 (9.0)
F	1 (1.5)
Unknown	1 (1.5)

were used as positive and negative controls, respectively, for *fimH* and *feoB*.

### 2.6. Statistical analysis

The statistical significance of the distribution of virulence genes was analysed by  $\chi^2$  test using GraphPad Prism software v.5 (GraphPad Software Inc., La Jolla, CA).

## 3. Results

### 3.1. Distribution of extended-spectrum $\beta$ -lactamase genes in *E. coli* from raw retail chicken

All of the ESBL-EC strains from retail poultry possessed *bla*<sub>CTX-M</sub> (Table 1), suggesting that CTX-M is the predominant ESBL type in *E. coli* from retail chicken in South Korea. More than one-half (56.7%; 38/67) of the *bla*<sub>CTX-M</sub>-positive strains also carried *bla*<sub>TEM</sub> and/or *bla*<sub>OXA</sub>, whereas *bla*<sub>SHV</sub> was not detected (Table 1). The dominant CTX-M types included CTX-M-65 (52.2%; 35/67) in the CTX-M group 9 and CTX-M-55 (25.4%; 17/67) in the CTX-M group 1; three strains harboured both *bla*<sub>CTX-M-65</sub> and *bla*<sub>CTX-M-55</sub> (Table 1). The *bla*<sub>OXA-1</sub> gene was detected in six (9.0%) of the 67 ESBL-EC strains, and the *bla*<sub>OXA-1</sub>-positive strains also harboured *bla*<sub>TEM-1</sub> and *bla*<sub>CTX-M-65</sub> and/or *bla*<sub>CTX-M-55</sub> (Table 1).

### 3.2. Antimicrobial susceptibility of extended-spectrum $\beta$ -lactamase-producing *E. coli* from raw retail chicken

Strains harbouring all of the *bla*<sub>CTX-M</sub>, *bla*<sub>TEM</sub> and *bla*<sub>OXA</sub> genes were highly resistant to all tested antibiotics except for colistin (Table 2). Antimicrobial susceptibility testing showed that 98.5% (66/67) of the tested ESBL-EC isolates were resistant to at least three antibiotic classes tested in this study, suggesting that ESBL-EC from retail chicken in South Korea is highly multidrug-resistant (MDR).

### 3.3. Phylogenetic group analysis of extended-spectrum $\beta$ -lactamase-producing *E. coli* from raw retail chicken

The predominant phylogenetic groups of the ESBL-EC isolates from retail chicken were group A (56.7%) and group B1 (19.4%). Four *E. coli* isolates each belonged to groups B2 and D, respectively. Eight strains were classified as minor group, of which six belonged to group E, one to group F and one strain was unknown (Table 3). Phylogenetic analysis using RAPD-PCR showed that ESBL-EC isolates belonging to the same phylogenetic group tended to form the same cluster (Supplementary Fig. S1).

### 3.4. Prevalence of virulence genes in extended-spectrum $\beta$ -lactamase-producing *E. coli* from raw retail chicken

The prevalence of virulence genes representing the five major pathogenic groups of *E. coli*, including ETEC, EPEC, EAEC, STEC and ExPEC, was examined. The *espP* genes was detected and in one strain in group B1 and the *eaeA* gene was detected five strains in group A, whereas toxin genes were not detected. Interestingly, all 67 ESBL-EC isolates carried at least one ExPEC-related virulence gene. Compared with group A, the prevalence of ExPEC-related virulence genes was more frequent in groups B1, B2, D and E (Table 4), and those harbouring at least six ExPEC-related virulence genes belonged to phylogenetic groups B1, B2, D and E with statistical significance (Table 5).

**Table 4**

Prevalence of virulence genes of extended-spectrum  $\beta$ -lactamase (ESBL)-producing *Escherichia coli* isolates according to phylogenetic group.

Pathogenetic group	Virulence gene	No. (%) of strains							
		Group A (n=38)	Group B1 (n=13)	Group B2 (n=4)	Group D (n=4)	Group E (n=6)	Group F (n=1)	Unknown (n=1)	Total (n=67)
ExPEC	<i>astA</i>	3 (7.9)	5 (38.5)*	1 (25.0)	4 (100)***	3 (50.0)**	0	0	16 (23.9)
	<i>iss</i>	27 (71.1)	9 (69.2)	3 (75.0)	2 (50.0)	6 (100)	1 (100)	1 (100)	49 (73.1)
	<i>fimH</i>	21 (55.3)	13 (100)**	4 (100)	4 (100)	6 (100)*	1 (100)	1 (100)	50 (74.6)
	<i>aer</i>	22 (57.9)	10 (76.9)	4 (100)	4 (100)	6 (100)*	1 (100)	1 (100)	48 (71.6)
	<i>irp2</i>	3 (7.9)	7 (53.8)***	3 (75.0)***	1 (25.0)	0	0	0	14 (20.9)
	<i>iha</i>	1 (2.6)	3 (23.1)*	0	0	0	0	0	4 (6.0)
	<i>iutA</i>	21 (55.3)	10 (76.9)	4 (100)	4 (100)	6 (100)*	1 (100)	1 (100)	47 (70.1)
	<i>feoB</i>	38 (100)	13 (100)	4 (100)	4 (100)	6 (100)	1 (100)	1 (100)	67 (100)
	STEC	<i>stx1</i>	0	0	0	0	0	0	1 (100)
<i>stx2</i>		0	0	0	0	0	0	0	0
<i>hlyA</i>		0	0	0	0	0	0	0	0
<i>espP</i>		0	1 (7.7)	0	0	0	0	0	1 (1.5)
EPEC	<i>eaeA</i>	5 (13.2)	0	0	0	0	0	0	5 (7.5)
ETEC	<i>st</i>	0	0	0	0	0	0	0	0
	<i>lt</i>	0	0	0	0	0	0	0	0
EAEC	<i>aggR</i>	0	0	0	0	0	0	0	0

ExPEC, extraintestinal pathogenic *E. coli*; STEC, Shiga toxin-producing *E. coli*; EPEC, enteropathogenic *E. coli*; ETEC, enterotoxigenic *E. coli*; EAEC, enteroaggregative *E. coli*.

\*  $P < 0.05$ , distribution in indicated group versus group A.

\*\*  $P < 0.01$ , distribution in indicated group versus group A.

\*\*\*  $P < 0.001$ , distribution in indicated group versus group A.

**Table 5**  
Distribution of extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* isolates in phylogenetic groups according to the number of virulence genes.

Phylogenetic group	No. (%) of virulence genes						
	7	6	5	4	3	2	1
Group A (n=38)	0	0	7 (18.4)	15 (39.5)	11 (28.9)	3 (7.9)	2 (5.3)
Group B1 (n=13)	3 (23.1)**	5 (38.5)***	2 (15.4)	0**	3 (23.1)	0	0
Group B2 (n=4)	1 (2.5)**	1 (2.5)**	2 (50)	0	0	0	0
Group D (n=4)	1 (2.5)**	1 (2.5)**	2 (50)	0	0	0	0
Group E (n=6)	0	3 (50)***	3 (50)	0	0	0	0
Group F (n=1)	0	0	1 (100)*	0	0	0	0
Unknown (n=1)	0	0	1 (100)*	0	0	0	0

\*  $P < 0.05$ , distribution in indicated group versus group A.

\*\*  $P < 0.01$ , distribution in indicated group versus group A.

\*\*\*  $P < 0.001$ , distribution in indicated group versus group A.

#### 4. Discussion

ESBL-EC are frequently isolated from poultry [6], and retail chicken is considered as an important vehicle transmitting ESBL-EC to humans [5]. In this study, ESBL-EC isolates from retail chicken in South Korea were extensively characterised. Consistent with the global expansion of CTX-M ESBLs [31], CTX-M was the predominant ESBL type in *E. coli* strains from retail chicken in South Korea. In this study, *bla*<sub>CTX-M-65</sub> (52.2%), *bla*<sub>CTX-M-55</sub> (25.4%) and *bla*<sub>CTX-M-14</sub> (22.4%) were the most common ESBL genes. Similarly, *bla*<sub>CTX-M-55</sub>, *bla*<sub>CTX-M-65</sub> and *bla*<sub>CTX-M-14</sub> were commonly detected in ESBL-EC from chicken in China [32]. In Japan, *bla*<sub>CTX-M-2</sub>, *bla*<sub>TEM</sub> and *bla*<sub>CTX-M-1</sub> were present in 45%, 36% and 34% of ESBL-EC isolates from domestic retail chicken meat samples, respectively [7]. In the Netherlands, among 87 strains of ESBL-EC from chicken meat, 69% harboured *bla*<sub>CTX-M-1</sub> [33]. Similarly, 65.4% of chicken meat samples in the UK were contaminated with ESBL-EC, and *bla*<sub>CTX-M-1</sub> was predominant (82.7%) [34]. Based on the findings in this and other studies, ESBL-EC are highly prevalent on retail chicken in most countries, however the dominant ESBL gene types are different depending on the geographic region.

All of the ESBL-EC isolates from retail chicken in the current study were resistant to multiple drugs belonging to different classes, such as ampicillin, tetracycline, chloramphenicol and streptomycin (Table 2). A high prevalence of MDR ESBL-EC has been reported previously. In China, 96.9% of ESBL-EC isolated from chickens were resistant to at least three different antimicrobial classes [35]. MDR strains are highly distributed in clinical as well as chicken samples. For instance, all of the ESBL-EC isolated from hospitals in India were MDR [36]. In the current study, two strains were resistant to colistin, an antibiotic of last-resort (Table 2). Further investigation found that the strains were positive for *mcr-1* (data not shown), the plasmid-encoded gene conferring resistance to colistin [37]. This observation may be explained by the fact that ESBL genes are usually encoded on transmissible plasmids harbouring multiple resistance genes [38]. Strains carrying three different ESBL genes (i.e. *bla*<sub>CTX-M</sub>, *bla*<sub>TEM</sub> and *bla*<sub>OXA</sub>) exhibited increased antimicrobial resistance compared with those harbouring fewer ESBL genes, although this was not statistically significant owing to the small number of strains (n=6) (Table 2). Presumably, the co-existence of different ESBL genes may result from the co-presence of multiple resistance plasmids in the same strain. For instance, it has been demonstrated that plasmid-mediated quinolone resistance genes are frequently detected in ESBL-EC isolates from poultry [39].

Virulent ExPEC usually belong to phylogenetic groups B2 and D [40,41]. However, in the current study ESBL-EC isolated from retail

chicken mainly belonged to groups A (56.7%) and B1 (19.4%), and only four strains were assigned to each of groups B2 and D (Table 3). Interestingly, the number of strains in group E, which is classified as a minor group, was larger than those belonging to groups B2 and D (Table 3). *E. coli* O157:H7 EDL933 is the best-known member belonging to group E [24]. Consistently, a previous study in Jeonnam Province in South Korea showed that the majority of *E. coli* isolates from chicken were limited to phylogenetic group A, followed by group B1, and no isolate belonged to group B2 [42]. A similar pattern of distribution was also found among isolates from chicken carcasses in China [43]. In contrast, *E. coli* isolates from chicken meat had the greatest percentage of group B1 strains (44%), followed by groups A (28%) and D (23%) in the Netherlands [33]. Among *E. coli* isolates from chicken meat in the USA, groups D, B2, A and B1 were dominant in that order [44]. Thus, the distribution of phylogenetic groups may be affected by their geographical region. However, the prevalence of strains belonging to group E in the current study was similar to a previous report from the Netherlands [33]. Whereas the isolates harboured no or only very few virulence factors related to gastrointestinal infection, various virulence factors associated with ExPEC were frequently detected in ESBL-EC isolates, particularly in those belonging to groups B1, B2, D and E, but not to group A (Tables 4 and 5). These results suggest that ESBL-EC isolates in groups B1, B2, D and E from retail chicken may potentially be implicated in extraintestinal infections in humans in South Korea.

The findings of this study demonstrate that raw retail chicken in South Korea could be a major source of disseminating ESBL-EC with great potential to cause extraintestinal infections. Since ESBL-EC from chicken are often MDR, further investigation is required to control and reduce the contamination of chicken meat by ESBL-EC to protect public health.

#### Funding

This research was supported by a grant [16162MFDS029] from the Ministry of Food and Drug Safety, South Korea. HP and JK were supported by the BK21 Plus Program of the Department of Agricultural Biotechnology, Seoul National University, Seoul, South Korea.

#### Competing interests

None declared.

#### Ethical approval

Not required.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found in the online version, at <https://doi.org/10.1016/j.jgar.2019.01.005>.

## References

- [1] Rawat D, Nair D. Extended-spectrum  $\beta$ -lactamases in Gram negative bacteria. *J Glob Infect Dis* 2010;2:263–74.
- [2] Pitout JD, Nordmann P, Laupland KB, Poirel L. Emergence of Enterobacteriaceae producing extended-spectrum  $\beta$ -lactamases (ESBLs) in the community. *J Antimicrob Chemother* 2005;56:52–9.
- [3] Iovleva A, Bonomo RA. The ecology of extended-spectrum  $\beta$ -lactamases (ESBLs) in the developed world. *J Travel Med* 2017;24(Suppl. 1):S44–51.
- [4] Ewers C, Antão E-M, Diehl I, Philipp H-C, Wieler LH. Intestine and environment of the chicken as reservoirs for extraintestinal pathogenic *Escherichia coli* strains with zoonotic potential. *Appl Environ Microbiol* 2009;75:184–92.
- [5] Overdeest I, Willemsen I, Rijnsburger M, Eustace A, Xu L, Hawkey P, et al. Extended-spectrum  $\beta$ -lactamase genes of *Escherichia coli* in chicken meat and humans, the Netherlands. *Emerg Infect Dis* 2011;17:1216–22.
- [6] Olsen RH, Bisgaard M, Löhren U, Robineau B, Christensen H. Extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* isolated from poultry: a review of current problems, illustrated with some laboratory findings. *Avian Pathol* 2014;43:199–208.
- [7] Nahar A, Awasthi SP, Hatanaka N, Okuno K, Hoang PH, Hassan J, et al. Prevalence and characteristics of extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* in domestic and imported chicken meats in Japan. *J Vet Med Sci* 2018;80:510–7.
- [8] Yuan L, Liu J-H, Hu G-Z, Pan Y-S, Liu Z-M, Mo J, et al. Molecular characterization of extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* isolates from chickens in Henan Province, China. *J Med Microbiol* 2009;58:1449–53.
- [9] Hazeleger WC, Jacobs-Reitsma WF, den Besten HM. Quantification of growth of *Campylobacter* and extended spectrum  $\beta$ -lactamase producing bacteria sheds light on black box of enrichment procedures. *Front Microbiol* 2016;7:1430.
- [10] Smith JL, Fratamico PM, Gunther NW. Extraintestinal pathogenic *Escherichia coli*. *Foodborne Pathog Dis* 2007;4:134–63.
- [11] Nordstrom L, Liu CM, Price LB. Foodborne urinary tract infections: a new paradigm for antimicrobial-resistant foodborne illness. *Front Microbiol* 2013;4:29.
- [12] Russo TA, Johnson JR. Proposal for a new inclusive designation for extraintestinal pathogenic isolates of *Escherichia coli*: ExPEC. *J Infect Dis* 2000;181:1753–4.
- [13] Johnson TJ, Logue CM, Johnson JR, Kuskowski MA, Sherwood JS, Barnes HJ, et al. Associations between multidrug resistance, plasmid content, and virulence potential among extraintestinal pathogenic and commensal *Escherichia coli* from humans and poultry. *Foodborne Pathog Dis* 2012;9:37–46.
- [14] Manges AR, Smith SP, Lau BJ, Nuval CJ, Eisenberg JN, Dietrich PS, et al. Retail meat consumption and the acquisition of antimicrobial resistant *Escherichia coli* causing urinary tract infections: a case-control study. *Foodborne Pathog Dis* 2007;4:419–31.
- [15] Jo S-J, Woo G-J. Molecular characterization of plasmids encoding CTX-M  $\beta$ -lactamases and their associated addiction systems circulating among *Escherichia coli* from retail chickens, chicken farms, and slaughterhouses in Korea. *J Microbiol Biotechnol* 2016;26:270–6.
- [16] Poulou A, Grivakou E, Vriou G, Koumaki V, Pittaras T, Pournaras S, et al. Modified CLSI extended-spectrum  $\beta$ -lactamase (ESBL) confirmatory test for phenotypic detection of ESBLs among Enterobacteriaceae producing various  $\beta$ -lactamases. *J Clin Microbiol* 2014;52:1483–9.
- [17] Fang H, Ataker F, Hedin G, Dornbusch K. Molecular epidemiology of extended-spectrum  $\beta$ -lactamases among *Escherichia coli* isolates collected in a Swedish hospital and its associated health care facilities from 2001 to 2006. *J Clin Microbiol* 2008;46:707–12.
- [18] Costa D, Poeta P, Sáenz Y, Vinué L, Rojo-Bezares B, Jouini A, et al. Detection of *Escherichia coli* harbouring extended-spectrum  $\beta$ -lactamases of the CTX-M, TEM and SHV classes in faecal samples of wild animals in Portugal. *J Antimicrob Chemother* 2006;58:1311–2.
- [19] Kim J, Lim Y-M, Rheem I, Lee Y, Lee J-C, Seol S-Y, et al. CTX-M and SHV-12  $\beta$ -lactamases are the most common extended-spectrum enzymes in clinical isolates of *Escherichia coli* and *Klebsiella pneumoniae* collected from 3 university hospitals within Korea. *FEMS Microbiol Lett* 2005;245:93–8.
- [20] Olesen I, Hasman H, Møller Aarestrup F. Prevalence of  $\beta$ -lactamases among ampicillin-resistant *Escherichia coli* and *Salmonella* isolated from food animals in Denmark. *Microb Drug Resist* 2004;10:334–40.
- [21] Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing. 27th ed. CLSI supplement M100. Wayne, PA: CLSI; 2017.
- [22] Bell SM, Gatus BJ, Pham JN. Antibiotic susceptibility testing by the CDS method. A concise laboratory manual. Sydney, NSW: Australia Arthur Productions Pty Ltd.; 1999.
- [23] Sader HS, Rhomberg PR, Flamm RK, Jones RN. Use of a surfactant (polysorbate 80) to improve MIC susceptibility testing results for polymyxin B and colistin. *Diagn Microbiol Infect Dis* 2012;74:412–4.
- [24] Clermont O, Christenson JK, Denamur E, Gordon DM. The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environ Microbiol Rep* 2013;5:58–65.
- [25] Madico G, Akopyants NS, Berg DE. Arbitrarily primed PCR DNA fingerprinting of *Escherichia coli* O157: H7 strains by using templates from boiled cultures. *J Clin Microbiol* 1995;33:1534–6.
- [26] Paton AW, Paton JC. Detection and characterization of Shiga toxin-producing *Escherichia coli* by using multiplex PCR assays for *stx1*, *stx2*, *eaeA*, enterohemorrhagic *E. coli* *hlyA*, *rfbO111*, and *rfbO157*. *J Clin Microbiol* 1998;36:598–602.
- [27] Cerna JF, Nataro JP, Estrada-García T. Multiplex PCR for detection of three plasmid-borne genes of enteroaggregative *Escherichia coli* strains. *J Clin Microbiol* 2003;41:2138–40.
- [28] Lee J, Subhadra B, Son YJ, Kim D, Park H, Kim J, et al. Phylogenetic group distributions, virulence factors and antimicrobial resistance properties of uropathogenic *Escherichia coli* strains isolated from patients with urinary tract infections in South Korea. *Lett Appl Microbiol* 2016;62:84–90.
- [29] Jakobsen L, Spangholm DJ, Pedersen K, Jensen LB, Emborg H-D, Agersø Y, et al. Broiler chickens, broiler chicken meat, pigs and pork as sources of ExPEC related virulence genes and resistance in *Escherichia coli* isolates from community-dwelling humans and UTI patients. *Int J Food Microbiol* 2010;142:264–72.
- [30] Dezfulian H, Batisson I, Fairbrother JM, Lau PC, Nassar A, Szatmari G, et al. Presence and characterization of extraintestinal pathogenic *Escherichia coli* virulence genes in F165-positive *E. coli* strains isolated from diseased calves and pigs. *J Clin Microbiol* 2003;41:1375–85.
- [31] Cantón R, González-Alba JM, Galán JC. CTX-M enzymes: origin and diffusion. *Front Microbiol* 2012;3:110.
- [32] Rao L, Lv L, Zeng Z, Chen S, He D, Chen X, et al. Increasing prevalence of extended-spectrum cephalosporin-resistant *Escherichia coli* in food animals and the diversity of CTX-M genotypes during 2003–2012. *Vet Microbiol* 2014;172:534–41.
- [33] Kluytmans JA, Overdeest IT, Willemsen I, Kluytmans-Van Den Bergh MF, Van Der Zwaluw K, Heck M, et al. Extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* from retail chicken meat and humans: comparison of strains, plasmids, resistance genes, and virulence factors. *Clin Infect Dis* 2012;56:478–87.
- [34] Randall L, Lodge M, Elviss N, Lemma F, Hopkins K, Teale C, et al. Evaluation of meat, fruit and vegetables from retail stores in five United Kingdom regions as sources of extended-spectrum  $\beta$ -lactamase (ESBL)-producing and carbapenem-resistant *Escherichia coli*. *Int J Food Microbiol* 2017;241:283–90.
- [35] Tong P, Sun Y, Ji X, Du X, Guo X, Liu J, et al. Characterization of antimicrobial resistance and extended-spectrum  $\beta$ -lactamase genes in *Escherichia coli* isolated from chickens. *Foodborne Pathog Dis* 2015;12:345–52.
- [36] Sharma M, Pathak S, Srivastava P. Prevalence and antibiogram of extended spectrum  $\beta$ -lactamase (ESBL) producing Gram negative bacilli and further molecular characterization of ESBL producing *Escherichia coli* and *Klebsiella* spp. *J Clin Diagn Res* 2013;7:2173–7.
- [37] Liu Y-Y, Wang Y, Walsh TR, Yi L-X, Zhang R, Spencer J, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* 2016;16:161–8.
- [38] Paterson D. Recommendation for treatment of severe infections caused by Enterobacteriaceae producing extended-spectrum  $\beta$ -lactamases (ESBLs). *Clin Microbiol Infect* 2000;6:460–3.
- [39] Xia L-N, Tao X-Q, Shen J-Z, Dai L, Wang Y, Chen X, et al. A survey of  $\beta$ -lactamase and 16S rRNA methylase genes among fluoroquinolone-resistant *Escherichia coli* isolates and their horizontal transmission in Shandong, China. *Foodborne Pathog Dis* 2011;8:1241–8.
- [40] Cortés P, Blanc V, Mora A, Dahbi G, Blanco JE, Blanco M, et al. Isolation and characterization of potentially pathogenic antimicrobial-resistant *Escherichia coli* strains from chicken and pig farms in Spain. *Appl Environ Microbiol* 2010;76:2799–805.
- [41] Picard B, Garcia JS, Gouriou S, Duriez P, Brahimi N, Bingen E, et al. The link between phylogeny and virulence in *Escherichia coli* extraintestinal infection. *Infect Immun* 1999;67:546–53.
- [42] Unno T, Han D, Jang J, Lee S-N, Ko G, Choi HY, et al. Absence of *Escherichia coli* phylogenetic group B2 strains in humans and domesticated animals from Jeonnam Province, Republic of Korea. *Appl Environ Microbiol* 2009;75:5659–66.
- [43] Xu X, Cui S, Zhang F, Luo Y, Gu Y, Yang B, et al. Prevalence and characterization of cefotaxime and ciprofloxacin co-resistant *Escherichia coli* isolates in retail chicken carcasses and ground pork, China. *Microb Drug Resist* 2014;20:73–81.
- [44] Mitchell NM, Johnson JR, Johnston B, Curtiss R, Mellata M. Zoonotic potential of *Escherichia coli* isolates from retail chicken meat products and eggs. *Appl Environ Microbiol* 2015;81:1177–87.