



## Review article

## Bridging the gap: Mechanisms of plasticity and repair after pediatric TBI

Naomi S. Sta Maria<sup>a,\*</sup>, Saman Sargolzaei<sup>b</sup>, Mayumi L. Prins<sup>b,g</sup>, Emily L. Dennis<sup>c</sup>,  
Robert F. Asarnow<sup>d</sup>, David A. Hovda<sup>b,e</sup>, Neil G. Harris<sup>b,f</sup>, Christopher C. Giza<sup>b,g,h</sup>

<sup>a</sup> Department of Physiology and Neuroscience, Zilkha Neurogenetic Institute, University of Southern California, 1501 San Pablo Street, ZNI115, Los Angeles, CA 90033, United States of America

<sup>b</sup> UCLA Brain Injury Research Center, Department of Neurosurgery, University of California at Los Angeles, Box 956901, 300 Stein Plaza, Ste 562, 5th Floor, Los Angeles, CA 90095-6901, United States of America

<sup>c</sup> Brigham and Women's Hospital/Harvard University and Department of Psychology, Stanford University, 1249 Boylston Street, Boston, MA 02215, United States of America

<sup>d</sup> Department of Psychiatry and Biobehavioral Sciences, University of California at Los Angeles, Box 951759, 760 Westwood Plaza, 48-240C Semel Institute, Los Angeles, CA 90095-1759, United States of America

<sup>e</sup> Department of Medical and Molecular Pharmacology, University of California at Los Angeles, Box 956901, 300 Stein Plaza, Ste 562 & Semel 18-228A, Los Angeles, CA 90095-6901, United States of America

<sup>f</sup> Intellectual Development and Disabilities Research Center, David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA, United States of America

<sup>g</sup> Steve Tisch BrainSPORT Program, University of California at Los Angeles, Los Angeles, CA, United States of America

<sup>h</sup> Division of Pediatric Neurology, Mattel Children's Hospital – UCLA, Los Angeles, CA, United States of America

## ARTICLE INFO

## Keywords:

Traumatic brain injury  
Plasticity  
Development  
Pediatric TBI

## ABSTRACT

Traumatic brain injury is the leading cause of death and disability in the United States, and may be associated with long lasting impairments into adulthood. The multitude of ongoing neurobiological processes that occur during brain maturation confer both considerable vulnerability to TBI but may also provide adaptability and potential for recovery. This review will examine and synthesize our current understanding of developmental neurobiology in the context of pediatric TBI. Delineating this biology will facilitate more targeted initial care, mechanism-based therapeutic interventions and better long-term prognostication and follow-up.

## 1. Introduction: The problem of developmental plasticity and brain injury

Traumatic injury to the developing brain has been called the most complex injury to the most complex organ at the most complex time (Kenzie et al., 2017; Marklund and Hillered, 2011; Wheble and Menon, 2016). Traumatic Brain Injury (TBI) is the leading cause of death and acquired disability in children and is responsible for approximately

640,000 emergency department visits, 18,000 hospitalizations and 1500 deaths in the U.S. alone (Schuchat et al., 2016; Taylor et al., 2017). A significant number of these (325,000 in the U.S.) occur in settings of sports and recreation (Coronado et al., 2015). Emergency department (ED) estimates of pediatric TBI certainly underestimate the public health burden, as a recent study reported only 12% of pediatric mild TBI patients presented to the health care system through the ED, with 82% initially visiting primary care providers and 5% presenting to

**Abbreviations:** AMPAR,  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole-propionate receptor; BDNF, brain-derived neurotrophic factor; BOLD, blood-oxygen-level-dependent signal; cAMP, cyclic adenosine monophosphate; CAMK, calcium-calmodulin protein kinase; CBF, cerebral blood flow; CCI, cortical contusion injury; CHI, closed head injury; CMRg, cerebral metabolic rates of glucose; CPEB, cytoplasmic polyadenylation element-binding protein; CREB, cAMP response element binding protein; DCS, *D*-cycloserine; ED, emergency department; E, embryonic; EE, enriched environment; ERK, extracellular signal regulated kinase; fc, functional connectivity; FPI, fluid percussion injury; fMRI, functional magnetic resonance imaging; GABA,  $\gamma$ -aminobutyric acid; GAD, glutamic acid decarboxylase; GluN, glutamate NMDAR subunit; IHHT, interhemispheric transit time; JNK, c-Jun N terminal kinase; LTD, long-term depression; LTP, long-term potentiation; MRI, magnetic resonance imaging; MRS, magnetic resonance spectroscopy; MWM, morris water maze; NMDAR, *N*-methyl-*D*-aspartate receptor; NOR, novel object recognition; NSC, neural stem cell; OL, oligodendrocyte; OPC, oligodendrocyte precursor cell; P, postnatal; pHMRI, pharmacological MRI; PID, post-injury day; PSD, post synaptic density; rCBV, relative cerebral blood volume; TBI, traumatic brain injury; WM, white matter

\* Corresponding author.

E-mail addresses: [nstamari@usc.edu](mailto:nstamari@usc.edu) (N.S. Sta Maria), [ssargolzaei@mednet.ucla.edu](mailto:ssargolzaei@mednet.ucla.edu) (S. Sargolzaei), [mprins@mednet.ucla.edu](mailto:mprins@mednet.ucla.edu) (M.L. Prins), [edennis@bwh.harvard.edu](mailto:edennis@bwh.harvard.edu) (E.L. Dennis), [rasarnow@mednet.ucla.edu](mailto:rasarnow@mednet.ucla.edu) (R.F. Asarnow), [dhovda@mednet.ucla.edu](mailto:dhovda@mednet.ucla.edu) (D.A. Hovda), [ngharris@ucla.edu](mailto:ngharris@ucla.edu) (N.G. Harris), [cgiza@mednet.ucla.edu](mailto:cgiza@mednet.ucla.edu) (C.C. Giza).

<https://doi.org/10.1016/j.expneurol.2019.04.016>

Received 25 December 2018; Received in revised form 9 March 2019; Accepted 25 April 2019

Available online 02 May 2019

0014-4886/© 2019 Elsevier Inc. All rights reserved.

specialty providers (Arbogast et al., 2016). Worldwide, the incidence of pediatric TBI is quite variable, ranging from 47 to 280/100,000, with peaks in the very young (0–2 yrs) and in adolescence (15–18 yrs) (Dewan et al., 2016). Falls, motor vehicle collisions, nonaccidental trauma and sports/recreation were contributing mechanisms.

While more severe pediatric TBI may be associated with immediate and persistent deficits (Hyder et al., 2007), this injury can also alter developmental trajectories, with some impairments becoming more evident later. Injury severity is an important predictor for long-term outcome. Anderson and colleagues showed that children with severe TBI (Glasgow Coma Score 3–8) recorded persistent depressed intellectual abilities up to 10 years postinjury (Anderson et al., 2012). Age at injury is the strongest and consistent predictor of neurocognitive outcome and behavior problems. Children sustaining TBI, whether diffuse or focal, before 3 years of age recorded poorer neurocognitive outcomes (Anderson et al., 2012, 2014; Karver et al., 2012, 2014). Anderson et al. further demonstrates there is no severity-related differences in recovery rates in children (2–7 yrs. old) who experience TBI. There is an initial protracted period of disrupted development in young children following early life TBI, but the brain shows evidence of recovery. Age-standardized intellectual quotient (IQ) scores are stable in injured children from 30 months to 10 years post-insult (Anderson et al., 2012). This suggests that intervention may be effective even after many years post TBI. Injury severity, however, is also a predictor for behavior problems, specifically ADHD symptoms and anxiety. Younger children who sustain a severe injury exhibit higher levels of symptoms over time (Karver et al., 2012, 2014).

The myriad of ongoing neurobiological processes that occur during brain maturation confer both considerable vulnerability to TBI but may also provide adaptability and potential for recovery. Changes in metabolism, axonal outgrowth, synaptogenesis, synaptic pruning and myelination are major ongoing biological processes occurring in the young brain. This review will examine and synthesize our current understanding of developmental neurobiology in the context of pediatric TBI. Delineating this biology will facilitate more targeted initial care, mechanism-based therapeutic interventions and better long-term prognosis and follow-up.

## 2. Mechanisms of normal development: Plug it in and turn it on

It is important to understand normal cerebral development as a context for comprehending injury effects, as well as for developing age-appropriate treatments. The rat model has been widely used to investigate the effects of TBI on developmental plasticity and recovery (Xiong et al., 2013). While there is no single timeline that links rodent development to human, the timing of critical developmental processes such as neurogenesis, cell migration, dendritic growth, and synaptic pruning can be used to compare across species (Kolb et al., 2000; Semple et al., 2013). Fig. 1 shows the timeline of cerebral development in rats and a comparable timeline in humans (Kolb et al., 2000; Semple et al., 2013). Kolb and colleagues describe that cells that eventually form the cortex are generated between embryonic days 12 (E12) and E21 (Kolb et al., 2000). Further, rats are born on day 22 (postnatal day 0 – P0), with neurogenesis already completed, and that these newly-generated cells migrate to appropriate locations, from the E12 to P7-10 and begin to differentiate (Kolb et al., 2000; Pressler and Auvin, 2013). Cell differentiation is almost complete by eye opening (P15), but neuronal maturation continues and peaks for another 2 to 3 weeks then declines.


It is important to note that the rat cerebrum is less developed at birth relative to the human brain, and that the first week of life is approximately equivalent to the third trimester in humans. Maximum dendritic growth occurs around P14 in rats, which corresponds to around 8 months in humans; whereas synaptogenesis peaks around P35 in rats and beginning at 1 year of age to 5–6 years in humans, depending on the brain region (Kolb et al., 2000; Pressler and Auvin,

2013; Stiles and Jernigan, 2010). Injury at various time points during these developmental stages has been shown to result in different functional outcomes by disrupting the trajectory of neural network formation and maturation (Kolb and Cioe, 2003; Kolb and Tomie, 1988).

Axonal outgrowth is a crucial developmental process that requires tight control to ensure appropriate neural connectivity to specific targets. This outgrowth is directed by extrinsic guidance molecules that are synthesized at a particular place and time to either attract or repulse axonal growth and direct its trajectory (Hirata, 2009). The inability to regenerate axons coincides with the maturation of CNS glial cells, such as astrocytes and oligodendrocytes, and to the production of myelin (Goldberg, 2003). However, the ability of neuronal axons to grow and regenerate may be an intrinsic property. The presence of brain-derived neurotrophic factor (BDNF) regulated enzyme arginase I in neurons overcomes inhibition of axonal regrowth by myelin. (Cai et al., 2002).

Neurons are supported by glia, such as astrocytes, microglia and oligodendrocytes. Astrocytes and microglia in TBI contribute to damage control and scar formation by releasing growth factors, modifying extracellular matrix components or by removal of cellular debris (Burda et al., 2016; Loane and Byrnes, 2010; Scheller et al., 2017). Oligodendrocytes (OLs) provide trophic support and produce the protective myelin that envelops neuronal axons (Scheller et al., 2017). Neural stem cells (NSC) give rise to oligodendrocyte precursor cells (OPCs) that differentiate into pre-mature OLs and then into mature OLs that contact neurons and their axons (Scheller et al., 2017). In development, rodents and humans share similar spatial and temporal patterns. Myelination occurs caudal-to-rostral, early in the brain stem, midbrain and cerebellum and later in the telencephalon, beginning with the occipital cortex and progressing to the prefrontal cortex (van Tilborg et al., 2018). OLs contain glutamate transporters and glutamate receptors (NMDAR, AMPAR, kainate), and thus are particularly sensitive to excitotoxic insults (Saab et al., 2016; Scheller et al., 2017). Significant OL loss leads to impaired electrical signal transduction across neurons and subsequently makes neurons more vulnerable (Scheller et al., 2017). Scheller and colleagues describe a significant increase of OPCs in the lesion area as a general acute response to TBI in 4 week old mice, which is accompanied by cell-specific changes in epigenetic regulation of gene expression and re-modeling of tissue structure (Scheller et al., 2017).

Functional magnetic resonance imaging (fMRI) during rest or during a task can demonstrate changes in blood-oxygen-level-dependent (BOLD) signal presumably linked to neural activity. The temporal coincidence of BOLD signal is used to infer presumed functional connectivity (fc) between brain regions, so that wider regions of coincident signal are taken to indicate co-dependent regions of functionally similar cellular activity, or discrete functional networks within the brain. A growing body of evidence (Gilmore et al., 2018; Hagmann et al., 2010; Huang et al., 2015; Menon, 2013) suggests that the functional and structural formation of the brain networks are inseparable, and that they progressively mature together during the early years after birth into adolescence. The age of 2 years has been repeatedly flagged as a landmark of rapid anatomical volume increase, cortical thickening and surface expansion, and elaboration of new synapses (Gilmore et al., 2018; Huang et al., 2015); after which a rapid increase in regional-specific myelination and axonal diameter triggers progressive white matter (WM) maturation (Hagmann et al., 2010). The structural maturation is characterized by heterogeneous strengthening and pruning of WM tracts that proceeds in a region-dependent way that, in turn, influences fc. Functional connectivity has been shown to be a surrogate of brain development into both healthy and pathologic brain (Menon, 2013). By the age of 2 years, core sub-networks (frontoparietal central executive network, default mode network, and salience network) are fully developed, however further changes in fc occur into adolescence, a phenomenon thought to support more complex cognitive demands. Apart from heterogeneous structural reconfigurations at the connectome level, local properties that influence sub-networks such as



**Fig. 1.** Schematic illustration of brain development in rat and in human. Abbreviation: E = embryonic day; P = postnatal day; mo = month; yr = year. Colored bars indicate main developmental processes. Adapted from Kolb et al. (2000) and Semple et al. (2013). Note that different cortical regions may have differing time profiles.

inhibitory-excitatory synaptic plasticity, neuronal migration, and synaptogenesis, are found to be dominant regulators of fc maturation (Menon, 2013). The analysis of fc data is still not completely standardized but generally falls into three categories for analysis of either static, time-averaged data or dynamic data: seed based analysis, independent component analysis and graph theory algorithms to characterize differences in network architecture at the local and global level of connectivity.

### 2.1. Plasticity

Once “plugged in and turned on,” the brain has the capacity to alter structure and function in response to environmental diversity (Kolb et al., 2011). This is called experience-dependent plasticity. For instance, housing animals in enriched environments (EE) leads to increased cortical thickness, increased dendritic arborization, and enhanced cognition (Fineman et al., 2000; Rosenzweig and Bennett, 1996; Rosenzweig et al., 1962). EE-induced plasticity is age-dependent. The EE-induced increases in dendritic arbors were highest in the weanling compared to the adult and senescent rats. In contrast, while adult and senescent rats showed increased spine density after EE exposure, the weanling rats showed a decrease in spine density (Kolb et al., 2003). There are also sex differences in spine density responses after EE. While adult females showed increases in spine density in the parietal cortex after EE exposure, weanling females showed decreases in parietal spine density (Kolb et al., 2003). There is reported an increase in global methylation in both female and male rats in the medial prefrontal cortex, orbital and insular prefrontal cortex, but a decrease in the hippocampus in males not in females (Kolb and Gibb, 2015). Epigenetic effects of chronic stress exposure is associated with a greater number of changes in gene expression in females (72 genes) compared to male rats (58 genes) in the prefrontal cortex and hippocampus combined (Kolb and Gibb, 2015). Behavioral training and environmental stimulation can also be used to restore learning and memory deficits (Rosenzweig and Bennett, 1996).

### 2.2. Neurotransmission and receptor ontogeny


Glutamate is the major excitatory neurotransmitter in the mammalian central nervous system and is crucial in many forms of synaptic plasticity, such as long-term potentiation (LTP) and long-term depression (LTD), which mediate learning and memory formation (Cull-Candy et al., 2001, 2006). Ionotropic glutamate receptors have been named based on the ligands that activate them: *N*-methyl-D-aspartate (NMDA),  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole-propionate (AMPA), and kainate. The NMDA receptor (NMDAR) and AMPA receptor (AMPA) are integral to normal development and experience-dependent plasticity, whose maturation is inevitably disrupted by TBI. Understanding the structure and function of these receptor systems could elucidate the effects of targeting glutamatergic neurotransmission for treating cognitive impairments that result from injury or disease.

#### 2.2.1. NMDAR

The NMDAR is a cation channel permeable to  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Ca}^{2+}$ . Under resting conditions, the NMDAR ligand binding sites are open and the ion channel pore is blocked by  $\text{Mg}^{2+}$ . Activation requires glutamate binding and post-synaptic depolarization to alleviate the  $\text{Mg}^{2+}$  block and allow full ionic flow.

The NMDAR is a heteromeric assembly containing 4 major subunits. The GluN1 transcript can be spliced to produce 8 distinct isoforms at three sites in the subunit protein, one in the N-terminus and 2 in the C-terminus. Four separate genes make up GluN2 (A-D) and 2 genes produce GluN3 subtypes (A-B). The GluN1 subunit is a necessary constituent for a functional receptor, whereas the GluN2 or GluN3 subunit confer modulatory properties that regulate glutamate sensitivity and channel opening (Benarroch, 2011; Cull-Candy et al., 2001; Ishii, 1993; Low and Wee, 2010). The diversity of NMDARs depends on the RNA splicing of GluN1, the type of NR2 subunit, and the presence or absence of the GluN3 subunit.

While GluN2B is predominant in the early postnatal period, GluN2A begins to be expressed during the second and third postnatal weeks (Scheller et al., 2017) and goes on to exceed GluN2B expression in adulthood. GluN2C and GluN2D are also developmentally regulated, predominantly in subcortical structures (Cull-Candy et al., 2001; Laurie



**Fig. 2.** NMDAR-mediated secondary effectors. Simplified schematic of intracellular NMDAR activated secondary effectors crucial in synaptic plasticity. NMDAR is shown as a heterodimer of GluN1 and GluN2 receptors bound with neurotransmitters glycine and glutamate. Once activated, NMDAR allows  $Ca^{2+}$  to enter the cell.  $Ca^{2+}$  can activate calcium-dependent kinases, such as CAMKII and CAMIV, which in turn can phosphorylate (circled P) downstream targets (e.g. Ras, ERK, and CREB). CREB activation leads to transcription of key target genes, like BDNF, that are important in cell survival and synaptic plasticity.

et al., 1997). GluN3A is only detected in the fetal rodent brain until two weeks postnatally, whereas GluN3B gradually increases over development (Low and Wee, 2010). When GluN3A co-assembles with GluN2A, the GluN3A subunits reduces channel conductance (Cull-Candy et al., 2001). GluN3B protein is widely expressed in the brain and spinal cord (Low and Wee, 2010), and is detected in both neurons and oligodendrocytes.

Experience also induces the GluN2B-to-GluN2A shift. Eye opening and visual stimuli increase GluN2A, but this enhanced GluN2A expression can be delayed or reversed by dark-rearing (Quinlan et al., 1999a; Quinlan et al., 1999b). Nurturing maternal behaviors, such as arch-back nursing and high frequency of licking, correlate with increased GluN2A levels in rats (Liu et al., 2000). GluN2A-containing NMDARs are primarily found in the synapse, whereas GluN2B is expressed both synaptically and extrasynaptically. Activation of synaptic NMDARs has been demonstrated to promote neuroprotection by triggering pro-survival signal pathways that include cAMP response element binding protein (CREB) dependent gene expression (Papadia et al., 2005) and brain-derived neurotrophic factor protein (BDNF) expression. Extrasynaptic NMDA activity triggers cell death pathways, inactivates CREB and blocks BDNF transcription (Hardingham et al., 2002).

Proper regulation of glutamate receptors (NMDAR, AMPAR) is crucial for experience-dependent synaptic activity. Activation of NMDARs increases intracellular  $Ca^{2+}$ , which activates kinases, such as CAMKII, and phosphatases, such as calcineurin. CAMKII-mediated phosphorylation of GluR1 AMPARs promotes their synaptic insertion, resulting in increased AMPAR conductance and LTP. Calcineurin, on the other hand, can dephosphorylate AMPAR and promote receptor internalization, resulting in LTD. An optimal NMDAR:AMPA ratio is typically maintained depending on neuronal activity (Benarroch, 2011; Cull-Candy et al., 2001; Lau and Zukin, 2007; Paoletti and Neyton, 2007).

### 2.2.2. AMPAR

The AMPAR is also a cation channel permeable to  $Ca^{2+}$ ,  $Na^{+}$ , and

$K^{+}$ . The functional properties of AMPARs depend on the subunit composition. Most AMPARs are heteromeric, assembling as a dimer of dimers containing combinations of GluR1-4. AMPARs are found in both neurons and glia as heteromers containing GluR2. GluR1 and GluR2 subunits are highly expressed in the hippocampus and cortex, with only low levels of GluR3 and GluR4 detected in these regions. Mature pyramidal cells primarily express heterotetramers of GluR1 and GluR2 (Isaac et al., 2007).

GluR2-containing AMPARs are impermeable to  $Ca^{2+}$ , suggesting that AMPARs play a crucial role in developmental synaptic function (Cull-Candy et al., 2006; Isaac et al., 2007; Kumar et al., 2002; Pickard et al., 2000). During the first postnatal weeks of neocortical development, during maximal synaptogenesis and increased expression of spiny neurons, GluR1 levels are higher than GluR2, which increases rapidly during the first week after birth. Additionally, developing GABAergic interneurons also express primarily GluR2-lacking AMPARs. Synaptic inactivity increases the expression of  $Ca^{2+}$ -permeable (GluR2-lacking) AMPARs. Conversely, synaptic potentiation increases the proportion of GluR2-containing AMPARs (Cull-Candy et al., 2006).

AMPA receptors mediate fast excitatory neurotransmission and have been implicated in synaptic plasticity, learning and memory (Cull-Candy et al., 2006).  $Ca^{2+}$ -permeable AMPARs play a crucial role in the maintenance of NMDAR-dependent LTP at synapses. In basal conditions, most AMPARs in the synapses contain GluR2 subunits. After induction of LTP, GluR2-lacking AMPARs (primarily GluR1-containing) are transiently inserted into the postsynaptic membrane, which promotes  $Ca^{2+}$  influx and signal propagation. Although  $Ca^{2+}$  entry via AMPARs is modest compared to that through NMDARs, AMPAR-mediated transmission still plays a crucial role in development, neuroplasticity and disease.

### 2.2.3. Second messenger targets and effectors

Intracellular calcium activates protein kinases (calcium-calmodulin protein kinase-CAMK, extracellular signal regulated kinases-ERK) that facilitate memory formation. ERKs are also known as mitogen-activated kinases (MAPK). These kinases, as well as a downstream effector

(CREB), are implicated in long-lasting changes in brain activity (Purves et al., 2004) (Fig. 2).

CAMKII is a calcium activated enzyme that constitutes about 30–40% of the protein in the post-synaptic density (PSD). Once activated, CAMKII autophosphorylates and then translocates to the PSD and directly binds to the NMDAR (Lisman et al., 2002; Purves et al., 2004). Additionally, CAMKII phosphorylates AMPARs at the serine 831 site, which increases the channel conductance and synaptic strength. CAMKII can also facilitate membrane AMPAR integration (Araki et al., 2015). This results in the increase of AMPARs in the synapse to promote LTP (Araki et al., 2015; Lisman et al., 2002) and confers synapse-specific neuroplasticity (Araki et al., 2015).

The ERK/MAPK pathway is involved in differentiation, proliferation and apoptosis in mammalian cells and has been shown to be necessary for associative learning (Atkins et al., 1998; Purves et al., 2004; Roskoski, 2012; Thomas and Huganir, 2004). In the ERK pathway, Ras activates c-Raf, then c-Raf phosphorylates ERK1/2. ERK1/2 is a known regulator of cell death and survival, as well as axonal growth. This pathway is important in spatial learning and hippocampal-dependent memory formation (Roskoski, 2012). On the other hand, MAPK regulation of c-Jun N terminal kinase (JNK) has been shown to mediate cell stress or death. Activated JNK is evident in apoptotic neurons and glia. Elevated levels of JNK precede neuronal cell death after global cerebral ischemia (Raghupathi, 2003). CREB is a ubiquitous transcriptional activator that binds to a site on DNA called the cAMP response element (CRE). CREB can be phosphorylated via the PKA and Ras pathways or via increased intracellular  $Ca^{2+}$ , which triggers its transcriptional activity (Purves et al., 2004). Increased intracellular  $Ca^{2+}$  can also phosphorylate CREB, wherein CRE site is activated. Here CRE is also called the CaRE (calcium response element) site.  $Ca^{2+}$ -mediated phosphorylation of CREB is mainly due to CAMKIV and ERK. The CREB phosphorylation must be long enough in duration for transcription to occur, even if only increases in  $Ca^{2+}$  levels are transient (Purves et al., 2004).

Hippocampal synaptic plasticity can be perturbed by disturbances in these secondary messenger targets and effectors. CAMKII is necessary for LTP induction. Inhibition of ERK activation blocks LTP maintenance in the hippocampus (Atkins, 2011; Atkins et al., 1998, 2009; English and Sweatt, 1997). Inability to activate ERK may prevent CREB-mediated gene expression of brain derived neurotrophic factor (BDNF). Sustained phosphorylation of CREB is involved in the activation of genes, such as BDNF that promote enhanced cognition, neuroprotection and recovery from injury such as BDNF (Finkbeiner, 2000; Griesbach, 2004).

#### 2.2.4. GABAR

$\gamma$ -aminobutyric acid (GABA) is the major inhibitory neurotransmitter in mammals. GABA is presynaptically synthesized from glutamic acid by glutamic acid decarboxylase (GAD) enzymes GAD67 and GAD65, encoded by the Gad1 and Gad2 genes, respectively (Owens and Kriegstein, 2002). Gene knock out of GAD65 produced mice that exhibit seizures, whereas deletion of GAD67 is lethal at birth and associated with 93% reduction of GABA in the cortex (Simeone et al., 2003).

GABA binds to 2 main subtypes of GABA receptors - GABA<sub>A</sub>R and GABA<sub>B</sub>R. GAD mRNA and protein can be detected in the whole rat brain at embryonic day 15 and levels increases rapidly in late fetal life, persisting through adulthood (Simeone et al., 2003). GABA<sub>A</sub>R is a heteropentameric subunit complex that binds various ligands and is selectively permeable to chloride and bicarbonate ions ( $Cl^-$  and  $HCO_3^-$ ). The majority of GABA<sub>A</sub>R contain  $\alpha$ ,  $\beta$ , and  $\gamma 2$  subunits with a 2:2:1 stoichiometry (Oh and Smith, 2019). Activation of GABA<sub>A</sub>R in mature cortical neurons results in membrane hyperpolarization due to inward flow of  $Cl^-$  currents. However, in immature neurons GABA<sub>A</sub>R activation depolarizes the postsynaptic membrane due to an inverted  $Cl^-$ -electrochemical gradient causing outflow of  $Cl^-$  from the

postsynaptic cell. This is due to expression of the  $Na^+ -K^+ -2Cl^-$  co-transporter NKCC1, which causes  $Cl^-$  ions to accumulate in the cell. Bicarbonate leaving the postsynaptic cell through the GABA<sub>A</sub>R also contributes to membrane depolarization (Simeone et al., 2003). In the first post-natal week GABA<sub>A</sub>R depolarization enables the removal of the  $Mg^{2+}$  block in NMDARs and activates voltage-gated calcium channels. This early-depolarization action of GABA contributes to synaptogenesis of both inhibitory and excitatory synapses, due to activities of NKCC1 and  $Ca^{2+}$  influx (Oh and Smith, 2019). In the second postnatal week, the  $K^+ -Cl^-$  cotransporter KCC2 expression significantly increases, KCC2 extrudes  $Cl^-$  from the cell and promotes more mature hyperpolarizing GABA<sub>A</sub>R kinetics. GABA<sub>A</sub>Rs may be synaptically or extrasynaptically located on the postsynaptic cell. Subunits such as  $\alpha 2$ ,  $\alpha 3$ , and  $\alpha 5$  are dominantly expressed during embryonic development. During the first post-natal week, subunit  $\alpha 1$  levels are low and increases significantly, while  $\alpha 2$  decreases. Subunit  $\alpha 5$  begins to be detected from embryonic day 17 into maturity (Simeone et al., 2003). In the rat, the  $\gamma 2$  subtype becomes the dominant  $\gamma$  subunit expressed at later developmental stages, and mRNA and membrane protein for this subtype are expressed in most brain regions (Owens and Kriegstein, 2002). The  $\gamma 2$  subunit is necessary for postsynaptic clustering of GABA<sub>A</sub>Rs and for forming functional synapses (Owens and Kriegstein, 2002). Location and composition of GABA<sub>A</sub>Rs contribute to the functional properties of these receptors (Owens and Kriegstein, 2002).

GABA<sub>B</sub>R is a metabotropic, G-coupled protein receptor that acts through secondary messenger cascades (Simeone et al., 2003). GABA<sub>B</sub>R may be located on pre or post synaptic cells and its activation results in  $K^+$  channel opening. In presynaptic cells  $K^+$  influx leads to reduction in GABA release. Postsynaptic  $K^+$  influx via activation of GABA<sub>B</sub>R leads to greater hyperpolarization (Simeone et al., 2003).

Once mature, GABAergic neurons contribute to two types of neuronal inhibition: phasic and tonic. GABA<sub>A</sub>R subunits  $\alpha 1$  and  $\gamma 2$  are involved in phasic inhibition are located at the synapse (Kharlamov et al., 2011). Phasic inhibition reduces hyper-excitability of the postsynaptic cell (Farrant and Nusser, 2005), and typically results from activation of postsynaptic GABA<sub>A</sub>R after exposure to high concentration of GABA released from the presynaptic terminal. Phasic inhibition of GABA<sub>A</sub>Rs plays an important role in modulation of theta and gamma oscillations. Tonic inhibition results from activation of GABA<sub>A</sub>R receptors on the presynaptic cell or on neighboring synapses from low concentrations of GABA (Farrant and Nusser, 2005). Tonic inhibition regulates postsynaptic depolarization in magnitude and duration (Farrant and Nusser, 2005), and contributes to the timing and synchronicity of excitatory signals (Guerrero et al., 2015). GABA<sub>A</sub>Rs subunits  $\alpha 4$  and  $\alpha 5$  are important for tonic inhibition and are located extrasynaptically (Farrant and Nusser, 2005; Kharlamov et al., 2011).

### 3. Developmental injury responses

#### 3.1. Energy crisis

Cerebral metabolism of glucose shows dynamic changes during normal development and after TBI. Reliance on glucose metabolism increases after weaning and adult cerebral metabolic rates of glucose (CMRg) are achieved in the rat between postnatal days 35–45 (75  $\mu\text{mol}/100\text{ g}/\text{min}$ ; (Prins and Hovda, 2009)) and in humans by 16–18 years of age (26–27  $\mu\text{mol}/100\text{ g}/\text{min}$ ; (Chugani, 1998)). After puberty, developmental sex differences in brain glucose metabolism through glycolysis are regulated by estrogen at multiple sites. Regional differences in CMRg are observed during the estrus cycle (Nehlig et al., 1985). During proestrus when estradiol is highest, CMRg is highest (124  $\pm$  6  $\mu\text{mol}/100\text{ g}/\text{min}$ ) in prefrontal cortex. This is significantly greater than both estrus females (109  $\pm$  4  $\mu\text{mol}/100\text{ g}/\text{min}$ ) and males (114  $\pm$  4  $\mu\text{mol}/100\text{ g}/\text{min}$ ). These changes in CMRg during development are accompanied by corresponding changes in cerebral blood flow (CBF).

While the overarching pattern of the metabolic response to TBI is consistent, there are distinctions related to injury type and severity. Immediately after impact, indiscriminate release of neurotransmitters activates neural systems causing widespread changes in ionic homeostasis across neuronal membranes (Katayama et al., 1990). Sodium/potassium ATPases are activated to pump ions back across membranes, increasing energy demands and increasing cerebral glucose uptake (3–6 h in animal studies and 7–10d in humans after “moderate” injury) (Yoshino et al., 1991). During this acute state (state 1), microdialysis lactate/pyruvate ratios ( $MD_{LPR}$ ) are increased in adult human subjects (Vespa, 2005), (Bentzer et al., 2000), (Kilbaugh et al., 2011). This increase in  $MD_{LPR}$  may be a potential biomarker for changes in glycolysis. The transient increase in cerebral glucose uptake is followed by a prolonged decrease in glucose metabolism (state 2), which has become a hallmark response to TBI (Yoshino et al., 1991), (Bergsneider et al., 1997). During state 2, there is a 9–12% increase in glucose shunting towards the pentose phosphate pathway (Bartnik et al., 2005). Glycolytic processing of glucose requires a constant supply of nicotinamide adenine dinucleotide ( $NAD^+$ ), which is decreased in the cytosol and thereby reduces the carbon supply to the mitochondria (Cosi and Marien, 1998), (Sheline et al., 2000), (Ying et al., 2003), (Prins, 2012). These acute changes lower pyruvate production and decrease ATP (Lee et al., 1999; Deng-Bryant et al., 2011) creating a state of cerebral metabolic crisis. The magnitude and duration of CMRg depression differs with age and injury type (Prins and Hovda, 2009; Thomas et al., 2000). Mild to moderate fluid percussion injury reveals age-related differences in metabolic recovery. Postnatal day P17 rat pups showed CMRg recovery within 3 days compared to 10 days in adults (Thomas et al., 2000; Yoshino et al., 1991). Even after more severe focal injuries, age differences in metabolic recovery are observed. P35 rats with cortical contusion injury (CCI) injury showed CMRg recovery of subcortical structures within 7 days, whereas adults remained significantly depressed for over 10d (Prins and Hovda, 2009). While there are very few human data, some clinical studies have shown hypometabolism similar to that reported in the animal studies (Roberts et al., 1995; Worley et al., 1995). It remains unclear whether young children and adolescents would show faster metabolic recovery than adults with TBI. While milder concussive injuries do not produce overt pathology, CMRg changes are still observed to a lesser magnitude and duration. In mild concussive brain injuries, the adolescent rat brain shows decreases in glucose uptake at 24 h, but recovery in 3 days (Prins et al., 2013). A second concussive injury during this vulnerable time period prolongs CMRg recovery (Prins et al., 2013). Magnetic resonance spectroscopy (1H-MRS) showed decreases in six metabolites (phosphocreatine, *N*-acetylaspartate, and total choline, GABA, lactate, and myoinositol) following a closed head injury (CHI) model of mild TBI in adult female mice but not in adult male mice (Lyons et al., 2018). These metabolite reductions corresponded with significant decrease in mitochondrial bioenergetics (State-III (adenosine triphosphate [ATP] synthesis capacity), State- $V_{C1}$  and State- $V_{C2}$  (complex I and II driven maximal electron transport) levels) from isolated mitochondria in cortex and hippocampus up to 28 days post CHI (Lyons et al., 2018).

### 3.2. White matter/axonal injury mechanisms

Axonal stretch results in mechanoporation that allows ionic flux directly into the axoplasm. Influx of calcium has been demonstrated to result in neurofilament side-arm cleavage and microtubule collapse (Buki and Povlishock, 2006; Pettus and Povlishock, 1996). This disruption of axonal cytoskeletal integrity impairs critical functions such as axonal transport, resulting in accumulation of molecules within the axon and axonal blebbing, demonstrated both experimentally and in humans (Johnson et al., 2013; Tang-Schomer et al., 2012). In vitro studies have also shown that stretch-induced undulations in axons are associated with greater  $Ca^{2+}$  influx (Tang-Schomer et al., 2012). The complex interactions of acute ionic flux, cytoskeletal breakdown,

impaired axonal transport and blebbing and ultimately axonal disconnection or degeneration all contribute to the longitudinal pathobiology of post-TBI axonal damage (Hill et al., 2016).

These cellular mechanisms of traumatic axonal injury are affected by clinically-important modifiers, including sex and myelination. In vitro and in vivo studies comparing male and female axons have demonstrated baseline differences in axon diameter and number of microtubules (female axons being smaller and more flexible with fewer microtubules) (Dolle et al., 2018). This results in greater undulation and  $Ca^{2+}$  flux after stretch injury, suggesting sex may be a biological substrate for vulnerability to TBI.


Multiple studies have demonstrated that white matter is involved in the acute response to developmental TBI, but challenges remain due to differences in age-at-injury, injury models, age-appropriate behaviors and time course. Midline closed head impact in P17 rats causes blood brain barrier disruption and axonal degeneration, with concomitant learning and memory impairments (Huh et al., 2008). More severe, focal TBI using controlled cortical impact models in P17 rats show alterations in electrophysiology, histological damage and ongoing cell death, in conjunction with emerging behavioral deficits (Ajao et al., 2012; Semple et al., 2014). It is important to note that cellular, molecular and behavioral perturbations may evolve weeks or months after developmental brain injury, making longitudinal follow-up and proper developmental controls critical components of translational pediatric TBI studies.

While the existing basic science examining white matter axonal injury is limited at comparable developmental stages and injury models, there is direct evidence that myelination is important for traumatic axonal injury. Myelinated axons are more resistant to experimental TBI. Midline FPI disrupts transcallosal evoked potentials, and the electrophysiological recovery in myelinated fibers occurs gradually over weeks. However, in unmyelinated axons, the electrophysiological recovery is stalled and incomplete (Reeves et al., 2005). In addition to traditional rodent models of pediatric TBI, newer pre-clinical pediatric models have investigated pathophysiology and behavior in rabbits and piglets (Baker et al., 2019; Zhang et al., 2015). While white matter maturation and axonal myelination continues into adulthood (~30 years of age in humans) (Kochunov et al., 2012), it is likely that this biomechanical vulnerability may play a larger role in the immature, less myelinated brain.

Similar to adult TBI, pediatric functional connectivity (fc) data have been reported as both decreases and increases in connectivity, for example (Risen et al., 2015; Stephens et al., 2017), and both directions of change are variably associated with performance on motor tasks (Risen et al., 2015). Enhancement of motor control to enable postural support was associated with greater cognitive fc from prefrontal regions (Diez et al., 2017). Whether this indicates that larger, or more diffuse brain regions are required to maintain normal function, or that there is a failure of circuits to deactivate, as occurs in adults after TBI is not known (Bonnelle et al., 2012). Clearly there is still much to learn about reorganization of functional networks after pediatric TBI, and how it relates to on-going deficits in the face of large-scale changes in network connectivity that occur as a function of normal development. Fig. 3 summarizes some recent clinical studies related to brain connectivity network alterations after pediatric TBI (Diez et al., 2017; Risen et al., 2015; Stephens et al., 2017, 2018; Yuan et al., 2017a,b).

### 3.3. Synaptic dysfunction & altered plasticity

Biomechanical damage to the synaptic cleft and dendritic spines occurs after TBI (Przekwas et al., 2016). Structural proteins (neurexins, neuroligins, SynCAMs, and integrins) and cell adhesion molecules (CAMs) that connect pre and post synaptic membranes are vulnerable to mechanical insults and typically require synaptic calcium to maintain elasticity and binding. Therefore, it is likely that post-TBI perturbations of synaptic calcium affect the integrity of these structural



		(Risen et al. 2015)	(Yuan et al. 2017b)	(Stephens et al. 2018)	(Stephens et al. 2017)	(Diez et al. 2017)	(Yuan et al. 2017a)
Demographic	TBI	5/9	10/12	5/6	6/12	8/6	7/10
	Control	5/9	7/13	6/5	14	15/12	8/3
Age	TBI	11-17	15.45±1.72	12.6-18.7	12-18	13.14±3.25	9.17-18.88
	Control	11-18.1	16.28±1.38	match	13-18	15.04±2.26	13.37-15.9
Injury severity		1/2	1	1/2 (ACRM)	1/2/3 (ACRM)	2/3 (Mayo)	2/3

Fig. 3. Presentation of current standing in clinical studies related to brain connectivity network alterations after pediatric TBI. “Demographic” is presented in the form of reported group size in Female/Male format. “Age” is reported as either range or mean ± standard deviation. Injury severity of the studied group is reported with (1) mild, (2) moderate, (3) severe. The metric on the basis which severity is reported, when provided, are American Congress of Rehabilitation Medicine (ACRM), and Mayo classification system.

synaptic proteins (Przekwas et al., 2016).

Mild and moderate TBI cause dendrite degeneration in the hippocampal dentate gyrus, with little to no significant cell loss or changes in the hippocampal formation in the adult rat brain at PID3 (Gao et al., 2011). Spared granular neurons in the dentate gyrus showed dendrite swelling with beading, which are signatures of injured dendrites. Adult TBI reduced total dendritic density, complexity and excitability of granule cells (Gao et al., 2011). In contrast, pediatric TBI in P17 rats did not result in any reductions in dendritic length or density in hippocampal DG at PID10. However, P17 TBI did reduce CA1 dendritic arborization and complexity (Casella et al., 2014).

### 3.4. TBI effects on NMDAR and AMPAR

Following TBI, activation of NMDARs and AMPARs results in intracellular Ca<sup>2+</sup> flux, which differs as a function of time post-injury, brain region and age. NMDAR activation dramatically increases 15 min after experimental TBI in adult mice (Biegan et al., 2004; Schumann et al., 2008), followed by diminished NMDAR expression (hours to days) in cortex and hippocampus (Biegan et al., 2004). A decrease in NMDAR binding has been reported three hours post-injury in the hippocampus and neocortex (Miller et al., 1990), and reduced protein expression occurs as early as 6 h after TBI (Kumar et al., 2002; Osteen et al., 2004). This downregulation lasts even longer in the developing brain. In P19 rats that sustained a lateral FPI, the GluN2A subunit protein is reduced by 20–40% in the ipsilateral hippocampus during the first post-injury week (Giza et al., 2006; Sta Maria et al., 2017). This molecular response has correlated with reduced electrophysiological activation through hippocampal NMDAR. FPI in young rats showed reduced amplitude of NMDA-mediated excitatory post-synaptic

currents (EPSCs) (down 40–50%) in hippocampal CA1 neurons. Also, application of subunit specific inhibitors isolating GluN2A currents showed that the reduced NMDA current is due primarily to a 50–60% loss of GluN2A-mediated currents (Li et al., 2005).

After FPI in adult rats, there is an increase in Ca<sup>2+</sup>-permeable AMPARs, as well as a decrease in GluR2 levels (Bell et al., 2007, 2009). In a closed head injury model of TBI, GluR1 levels increased acutely in the ipsilateral hippocampus in adult mice (Schumann et al., 2008).

The global changes in calcium signaling following TBI suggest that downstream signaling molecules would also be affected. TBI induces change in CAMKII levels, first with an initial increase followed by a long lasting reduction that is associated with impaired cognition and experience-dependent plasticity (Glazewski et al., 2000; Schwarzbach et al., 2006; Wu et al., 2009, 2011). FPI induces transient increases in phosphorylated CAMKII (Atkins et al., 2006; Folkerts et al., 2007) and total CAMKII in the hippocampus and cortex ipsilateral to the injury site (30 min) (Atkins et al., 2006; Folkerts et al., 2007). Additionally, downstream effectors of CAMKII (GluR1 and cytoplasmic polyadenylation element-binding protein – CPEB) also increase phosphorylation states in synaptic samples post-injury. CPEB regulates the translation of dendritic mRNAs during hippocampal LTP 30 min after FPI and returns to sham levels by 4 h (Atkins et al., 2006). This suggests that the aberrant increase of autophosphorylated CAMKII can result in loss of synapse specificity of neuronal activation and play a role in deficits long term memory formation. In CNS injury, ERKs have a different pathophysiological response. Following TBI, ERK1/2 is activated in injured cortical neurons, as well as non-neuronal cells in cortex, hippocampus and thalamus, as early as 3 h lasting up to 3 days (Raghupathi, 2003). Reduced ERK activation through pre-TBI treatment with a pERK inhibitor decreased cell death, but worsened

**Table 1**  
Summary of TBI-induced glutamate-mediated responses.

Target	Post TBI response
NMDAR	Transient increase in activity (in adult rats), followed by lasting reduction in activation (up to 1 week, in young rats) (Biegon et al., 2004; Giza et al., 2006; Kumar et al., 2002; Li et al., 2005; Miller et al., 1990; Osteen et al., 2004; Schumann et al., 2008)
AMPA	Adult rats: Increase in Ca <sup>2+</sup> -permeable AMPARs (GluR1 increases), and decrease in Ca <sup>2+</sup> -impermeable AMPARs (GluR2 decreases) (Bell et al., 2007, 2009; Schumann et al., 2008)
CAMKII	Transient increase followed by long lasting reduction (up to 1 week, in adult rats) (Atkins et al., 2006; Folkerts et al., 2007; Glazewski et al., 2000; Schwarzbach et al., 2006; Wu et al., 2009, 2011)
ERK	Long lasting reduction in activation (up to 12 weeks, in adult rats) (Raghupathi, 2003)
CREB	Long lasting reduction in phosphorylated CREB (up to 12 weeks, in adult rats) (Atkins et al., 2009; Griesbach, 2004)

cognitive and motor deficits (Dash et al., 2002). Deficits in ERK activation have been shown up to 12 weeks after FPI (Atkins et al., 2009). JNK activation, in contrast, was not observed in injured regions nor correlated with trauma-induced cell death (Raghupathi, 2004). Lastly, one week after FPI in adult rats, phosphorylated CREB was decreased (Griesbach, 2004). This reduction has been shown to persist up to 12 weeks (Atkins et al., 2009). Table 1 briefly summarizes glutamate-mediated responses following TBI.

The post-TBI alteration of NMDARs can then lead to a loss of plasticity in the young rat. Rats that received FPI at P19 followed immediately by rearing in an enriched environment (EE) failed to show EE-induced anatomical (increased cortical thickness, expanded dendritic arborization) and cognitive enhancements (improved spatial learning and memory) when tested as young adults (Fineman et al., 2000; Giza et al., 2005; Ip et al., 2002; Sta Maria et al., 2017). The period of diminished neural plasticity within one week post-FPI coincides with impaired working memory (Reger et al., 2005; Sta Maria et al., 2017), as well as with a critical period of neural responsiveness to EE-rearing (Giza et al., 2005; Sta Maria et al., 2017). Only when EE-rearing exposure was delayed two weeks following FPI in P19 rats did the animals demonstrate EE-induced enhancement in the MWM task acquisition. The delayed EE time point coincided with already normalized hippocampal NMDAR subunit levels, as well as recovery of NOR working memory to sham levels (Reger et al., 2005). However, probe trial performance in the delayed-EE group showed lingering TBI-associated deficits (Giza et al., 2005).

### 3.5. TBI effects on GABA<sub>A</sub>R

Several studies have reported GABA<sub>A</sub>R mediated alterations of inhibition following TBI that may be attributed to alterations in synaptic and extrasynaptic GABA<sub>A</sub>R subunit composition. Reductions in the GABA<sub>A</sub>R  $\gamma$ 2 subunit and increases in the  $\delta$ 1 have been reported 90 days following CCI in adult rats (Kharlamov et al., 2011). These subunit changes translate to reduced phasic and tonic GABAergic inhibition, respectively.  $\gamma$ 2 is a crucial component for a functional GABA<sub>A</sub>R and is important for GABA<sub>A</sub>R assembly at the synapse.  $\gamma$ 2 reductions or genetic deletion of the  $\gamma$ 2 gene are associated in some forms of epilepsy (Kharlamov et al., 2011). Guerriero and colleagues discusses the glutamate and GABA imbalance at early and late time periods pertinent following TBI in their review (Guerriero et al., 2015).

## 4. Recovery and repair

### 4.1. Linking metabolism & plasticity

The brain's metabolic status can directly impact its ability to adapt from the cellular to the functional level. Nutritional status, physical activity and fuel types can influence neuroplasticity (Stranahan and Mattson, 2008). Age also influences cerebral adaptation with the younger brain having much greater plasticity potential after injury (Weihmuller and Bruno, 1989). The dynamic metabolic changes after TBI can significantly impair neuronal function and plasticity (Babikian

et al., 2010). Immediate release of glutamate can inhibit protein synthesis (Ogawa et al., 1992). While global protein synthesis is depressed early after TBI, protein expression changes within specific pathways can be upregulated. Experimental studies with lateral FPI demonstrated increases in mTOR pathway protein expression 30 min to 1 day after TBI (Chen et al., 2007). In addition to changes in protein synthesis, acute global activation of NMDA receptors has been shown to alter neuronal excitability after the injury (Dietrich et al., 1994) for 24 h. During this time barrel field stimulation failed to increase glucose metabolic rates in injured animals, reflecting widespread circuit dysfunction after TBI. In addition to the effects on protein synthesis and activation, age-dependent calcium accumulation has been observed and this directly affects neurotrophin expression. A time course of <sup>45</sup>Ca<sup>2+</sup> changes after fluid percussion injury was examined in P17, P28 and adult rats (Osteen et al., 2001). Increased calcium accumulation was seen in all age groups immediately post injury with earlier recovery in younger animals. The adolescent (P28) and adult age groups did show a delayed accumulation of <sup>45</sup>Ca<sup>2+</sup> between post-injury days 4–14 in the ipsilateral thalamus, which was not observed in the youngest group. Neural activity and calcium both induce BDNF transcription and affect neuroplasticity (Griesbach and Hovda, 2015). Although overall cerebral glucose metabolism in P16–18 male rats recovers 24 h after TBI, specific reductions in [2-13C]glutamate suggests impairments in either neuronal or astrocytic metabolism (Robertson et al., 2013). Age-related plasticity differences have been observed after TBI (Giza et al., 2009; Giza and Prins, 2006). In weanling rats, TBI inhibits the brain's response to EE rearing (Giza et al., 2005), but in the adolescent rat, TBI seems to interfere with the normal pruning responses (Mychasiuk et al., 2015). There remains a dearth of research addressing age-related mechanisms of plasticity and its relationship to metabolism after TBI.

### 4.2. Protecting/healing axonal networks and promoting recovery

An orderly progression of white matter maturation has been described in both rodents and humans (Samorajski and Friede, 1968; Schonbach et al., 1968). This maturation includes differences in glial cell biology, cytokine expression and synthesis and maintenance of myelin. These normal neurodevelopmental changes have important implications on the distinct response to injury seen in the immature brain.

Basic science studies have demonstrated greater vulnerability of unmyelinated fibers to TBI compared to myelinated fibers (Reeves et al., 2005). Evidence of white matter damage in humans has been demonstrated by neuropathology following more severe TBI, but neuropathology following mild TBI is very limited (Blumbergs et al., 1994). Advanced multimodal imaging can provide information about white matter organization after pediatric TBI (Dennis et al., 2018a,b). Research consistently shows poorer white matter organization following a moderate-severe injury (msTBI) in pediatric patients using diffusion tensor imaging – DTI (Dennis et al., 2015a,b; Oni et al., 2010; Wilde et al., 2011), an MRI modality that looks at the extent and directionality of water in gray and white matter in the brain (Basser, 1997). DTI provides metrics such as anisotropy and diffusivity that has been

further correlated with changes in the corpus callosum macrostructure and with neuropsychological scores in children with chronic TBI (Ewing-Cobbs et al., 2006, 2008; Wilde et al., 2011). Fractional anisotropy (FA) and radial diffusivity are DTI metrics commonly accepted as proxies for fiber organization and myelin integrity. Radial diffusivity has been shown to best discriminate between TBI and comparison groups and is sensitive to traumatic axonal injury, and therefore, would be a good surrogate marker to assess extent of TBI and the effects of therapies restoring myelin integrity. FA is the most sensitive marker of TBI induced deficits in several cognitive and motor outcomes (Ewing-Cobbs et al., 2008). A multimodal approach can reveal different components of altered connectivity, including lower FA and higher radial diffusivity from DTI, altered neurometabolite levels (*N*-acetylaspartate – NAA, a marker of neuronal health, and choline, a marker of cellular turnover), and disrupted functional connectivity (rsfMRI) in msTBI subjects.

Callosal mid-sagittal region is often used as an index of developmental change and an indicator of injury severity (Ewing-Cobbs et al., 2006, 2008). Particularly, the callosal posterior midbody area increases in normal developing children, while the DTI metrics FA increases in the posterior midbody and radial diffusivity decreases in the isthmus and splenium with age (Ewing-Cobbs et al., 2006, 2008). Pediatric TBI decreases FA and increases radial diffusivity, in most of the callosal regions (Ewing-Cobbs et al., 2006, 2008; Wilde et al., 2011). Additionally, although the posterior midbody areas were similar between young normal and TBI children (~7–12 years old), the developmental increase in the posterior midbody area continued in older normal children but is arrested in older TBI children (~13–18 years old) (Ewing-Cobbs et al., 2008). This suggest an atrophy or arrest in callosal development and eventual alternate developmental trajectory for children with TBI.

Work from our group revealed delays in electrophysiological conduction across the corpus callosum – the interhemispheric transfer time (IHTT) – in about 50% of pediatric msTBI patients at a post-acute (2–4 month) time point after injury. This slowing in IHTT correlated with decreased FA and worse neurocognitive function (Dennis et al., 2015a; Ellis et al., 2016). Using a multimodal test battery prospectively and longitudinally from 2 to 4 months post-injury out to 14–18 months, pediatric msTBI subjects show ongoing changes. Uninjured controls show gradually increasing FA and cognitive function as brain development continues at its normal pace. The pediatric msTBI subjects with a normal IHTT show changes in white matter orientation measures (FA) and quantitative volumetrics similar to those seen in the typically developing controls. However, the patients with slow IHTT appear to diverge, showing progressively worsening white matter organization and volume changes inconsistent with healthy maturation (Dennis et al., 2017a,b). Examination of neurometabolic markers using magnetic resonance spectroscopy (MRS) post-acutely shows lower NAA and higher choline. Chronically, NAA increased and choline normalized in those with normal IHTT; while NAA and choline were low in those with slow IHTT (Babikian et al., 2018). The slow IHTT group also showed worse long-term neurocognitive outcomes than healthy controls (Ellis et al., 2016). These studies indicate that while about half of pediatric msTBI patients resume a favorable developmental trajectory, the other half show ongoing evidence of functional and structural degeneration (Dennis et al., 2018b).

The mechanisms underlying these differential white matter trajectories are uncertain but may include ongoing metabolic dysfunction, myelin damage, impaired circuit activity and persistent neuroinflammation. Understanding the contributions of these biological mechanisms post-acutely may open a window of intervention to alleviate this progressive neurodegeneration.

### 4.3. Restoring synaptic function and plasticity

#### 4.3.1. Environment


Experience-dependent plasticity can be induced using environmental enrichment (EE). EE provides opportunity for learning and integration of exploratory, physical, and social elements (Bondi et al., 2014). Neural alterations attributed to EE include increased neurogenesis, increased presynaptic (synaptophysin) and post-synaptic (PSD95) markers, cortical thickening, increases in cortical and hippocampal dendritic arborization, and increased cell survival after TBI (Bondi et al., 2014; Fineman et al., 2000; Hoffman et al., 2008b; Ip et al., 2002; Sozda et al., 2010). EE thus promises to be a viable intervention for impairments associated with TBI, stroke, aging or other neurodegenerative diseases (Bondi et al., 2014; Sampedro-Piquero and Begega, 2017).

Experience-dependent plastic changes are time- and region-dependent. When placed in EE for varying intervals of 4, 8, or 16 days, Comeau and colleagues have demonstrated that there is a transient dendritic length increase in Golgi-stained neurons in the prefrontal cortex after 4 days of EE that was not observed after 16 days of EE. In the sensory cortex, increased dendritic arborization seemed evident after all EE intervals (Comeau et al., 2010). In adult mice, EE rearing reversed the TBI induced learning and memory deficits following closed head weight-drop injury model (Schreiber et al., 2014) and repetitive mild TBI (Liu et al., 2017). In adult rats, shortened EE exposure (6 h per day) for 18 days was demonstrated to be more beneficial in MWM acquisition, whereas continuous EE showed significantly improved recall during the probe trial (de Witt et al., 2011). Delayed EE exposure following TBI in the young (Giza et al., 2005) and adult rats (Hoffman et al., 2008b) showed shortened latency to find the hidden platform in the Morris water maze. EE rearing for 6 weeks following early life (P21) closed head injury significantly reduced alcohol consumption and reward in female mice, as well as normalized relative BDNF gene expression to sham levels, and significantly reduced axonal degeneration (Weil et al., 2016).

Experience-dependent changes accumulate and interact, which is also known as metaplasticity. When methylphenidate and amphetamine were administered to weanling and adult rats, respectively, then exposed to EE, the drug blocked EE-induced dendritic arborization. Although the drugs themselves did not show any disturbance in physiology or morphology, the interaction between the pharmacology and EE resulted in impaired experience-dependent plasticity (Kolb et al., 2003). In experimental TBI in adult rats, EE exposure itself has shown expected enhancements in neuroanatomy and cognition. In several cases, only combined effects of EE and drug treatment or task-specific neurobehavioral experience have demonstrated functional improvement after injury (Hoffman et al., 2008a). Motor ability in TBI adult rats and acquisition of spatial learning in the MWM significantly improved when animals were also exposed to EE during the behavioral tests (Matter et al., 2011). Additionally, subtherapeutic EE experience (2–4 h per day for 19 days) can be used as a rehabilitation technique when combined with acetylcholinesterase inhibitor (galantamine) following CCI in adult rats (de la Tremblaye et al., 2017). Epigenetic studies show EE increases DNA methylation levels and reduces hydroxymethylation levels in senescence accelerated mouse P8 (SAMP8) (Grinan-Ferre et al., 2016), with differential expression in mouse dorsal and ventral hippocampus (Zhang et al., 2018). EE further reduces gene expression that drives oxidative stress and aging-induced inflammation promoting cognitive enhancements and neuroprotection (Grinan-Ferre et al., 2016).

#### 4.3.2. Pharmacology

To protect the brain from acute glutamate-induced excitotoxicity, NMDAR inhibitors have been used as a therapeutic intervention for TBI. Although this approach worked in many experimental TBI models (Han et al., 2009; Rao et al., 2001; Schumann et al., 2008), it was not



**Fig. 4.** Evoked  $\Delta rCBV$ . Top panel shows ROIs on sample structural image, and representative subacute control and FPI subject structural images with overlaid colored percent  $\Delta rCBV$  maps. Mean  $\pm$  SEM  $\Delta rCBV$  from ROIs in the hippocampus (Hipp), cortex (Ctx) and colliculus (Coll) in the subacute (PID 3–5) and chronic (PID 13–18) time points. Stimulation DCS dose induced a hippocampal-specific increase in rCBV in the controls that is abolished by FPI subacutely that recovery over time. \* $p < .05$  conveys significant difference between planned comparisons.

protective after injury to the immature brain (Bittigau et al., 1999; Pohl et al., 1999). NMDAR inhibition also failed to translate clinically, in some cases even leading to worsened outcome (Albers et al., 2001; Ikonomidou and Turski, 2002; Maas et al., 2010; Morris et al., 1999; Muir, 2006; Narayan et al., 2002). Due to the dynamic response of NMDAR activity post-injury, NMDAR antagonists may have been delivered during periods of already down-regulated NMDAR function that missed the critical window of receptor hyperactivity. Furthermore, during the recovery phase, excitatory synaptic activity may be necessary to facilitate optimal outcome (Ikonomidou and Turski, 2002).

In contrast, NMDAR agonists, such as D-cycloserine (DCS), promote neuroprotection and recovery after TBI (Adeleye et al., 2009; Biegon et al., 2004; Sta Maria et al., 2017; Temple and Hamm, 1996; Yaka et al., 2007). Administration of low doses of NMDA result in a pro-survival molecular response, indicating synaptic NMDARs are preferentially activated (Soriano et al., 2006). DCS is a partial NMDAR agonist binding to the glycine site in the GluN1 subunit. DCS is already approved for use in humans and has been shown to freely cross the blood-brain barrier (Baxter and Lanthorn, 1995). In mice, however, Wlaz et al. showed that after subcutaneous injection of DCS at a low dose (5 mg/kg), there were only trace amounts of DCS detected in the brain. After administration of a much higher dose (320 mg/kg) of DCS, only about 14% of the DCS plasma levels were detected in mice cortex during peak plasma levels of DCS (15 min post-injection). The half-life of DCS in mice is 30 min, and 1 h in rats, and most of the administered

DCS dose is excreted in rodents unchanged (Baxter and Lanthorn, 1995). DCS potentiates NMDAR-mediated processes and demonstrates NMDAR agonist properties (under 50 mg/kg in rats and mice; under 100 mg/kg in humans). At even higher doses (over 100 mg/kg in rodents and over 250 mg/kg in humans), DCS behaves more like an NMDAR antagonist (Baxter and Lanthorn, 1995).

DCS treatment at 3 mg/kg was shown to optimally enhance learning in a passive-shock avoidance task in adult rats (Monahan et al., 1989). After a closed head injury in adult mice, a 10 mg/kg dose of DCS was demonstrated to significantly improve neurological outcome (Adeleye et al., 2009; Yaka et al., 2007). DCS restored hippocampal CA1 LTP after closed head injury, and fully restored reduced BDNF levels. Following FPI in adult rats, a DCS dose of 30 mg/kg administered intraperitoneally, rescued deficits in the Morris water maze task. When administered twice daily for four days beginning at 24 h post-FPI in P19 rats, 30 mg/kg DCS restored deficits in GluN2A subunit protein levels, increased GluR2, and recovered novel object recognition performance (Sta Maria et al., 2017), as well as rescued secondary messenger CAMKII (Buen et al., 2010), in the injured young rat. P19 rats treated with DCS and exposed to EE-rearing as soon as 24 h post TBI showed improved MWM performance in adulthood (Sta Maria et al., 2017).

Other co-agonists of synaptic NMDARs include glycine and D-serine. D-serine level is high in mammalian brain and is increased by DCS administration. D-serine is metabolized by D-amino acid oxidase (DAAO), and its metabolism may increase H<sub>2</sub>O<sub>2</sub>, causing oxidative

stress. A recent study utilized a novel approach to protect against TBI by using 6-chlorobenzo[*d*]isoxazol-3-ol (CBIO), an inhibitor of DAAO, to increase D-serine levels and reduce oxidative stress. Treatment with CBIO after mouse closed head injury improved neurocognitive outcomes, reduced inflammation and enhanced molecular pathways of plasticity related to glutamatergic neurotransmission (Liraz-Zaltsman et al., 2018).

#### 4.4. MRI of NMDAR-mediated glutamatergic transmission

The vascular and metabolic demands of neuronal activity underlying fMRI signals have been suggested to be mostly driven by glutamate and the effects on its receptor-mediated action on neurons and astrocytes (Bonvento et al., 2002). For instance, NMDAR activation has been shown to be the primary factor in BOLD responses. Using a somatosensory evoked potential stimulation, BOLD signal can be significantly reduced using NMDAR antagonist MK801 (Gsell et al., 2006). NMDAR mediated activation can also be elicited in the hippocampus by measuring the change in percent relative cerebral blood volume ( $\Delta$ rCBV) using a pharmacological stimulus targeting the NMDAR co-agonist glycine-binding site, which is not saturated in hippocampal neurons in culture or in slice preparations (Wilcox et al., 1996). Systemic administration of D-serine (Panizzutti et al., 2005) and DCS (Santa Maria et al., 2009) can increase  $\Delta$ rCBV in a region-specific manner in the hippocampus using pharmacological MRI (phMRI) in normal rats. In our hands, FPI in P19 rats abolished the DCS-induced increase in hippocampal  $\Delta$ rCBV (Fig. 4) in the subacute (PID 3–5) time point, which corresponds to diminished NMDAR function, impaired behavior and experience-dependent plasticity, in the absence of gross histological damage. DCS-induced  $\Delta$ rCBV is restored at the chronic (PID 13–18) time point, when spontaneous recovery of NMDAR neurotransmission, novel object preference, and experience-dependent plasticity are observed following P19 rat FPI (Giza et al., 2005, 2006; Li et al., 2005; Regeer et al., 2005; Sta Maria et al., 2017). These results suggest that DCS-induced  $\Delta$ rCBV can be used as a non-invasive surrogate biomarker of NMDAR-mediated transmission. Its clinical implications include the use of the phMRI  $\Delta$ rCBV signal as a diagnostic tool to assess the magnitude and duration of impaired NMDAR-mediated glutamatergic transmission after TBI.

## 5. Conclusions

Normal postnatal brain development is heralded by a complex sequence of biological events – timing is everything. Axonal outgrowth, synapse formation and adaptation, changes in energy metabolism, myelination and dendritic pruning are all normally orchestrated to result in a unique and individualized final product. Biomechanical injury during development alters this construction, with remarkable capacity for plasticity and recovery but also potential long-term alterations in developmental trajectory. In addition to the complexities of brain-building, the immature brain also has distinct responses to the injury itself – metabolic patterns are perturbed, synapses retract and change in sensitivity, excitatory-inhibitory balance can be disrupted, neurovascular responses may uncouple and inflammatory cascades are initiated. Clinical observations such as delay or arrest of brain maturation after TBI (Dennis et al., 2017b; Ewing-Cobbs et al., 2008), and the double-hazard of young age and injury severity on pediatric TBI outcomes (Anderson et al., 2005; Keenan et al., 2019), are rooted in this seemingly tangled web of neurobiological mechanisms. Given the high incidence of pediatric TBI and the potential for long-term developmental consequences, understanding the fundamental mechanisms of brain development, injury and recovery will be critical to devising future targeted therapeutic interventions. Promising directions for developmental-specific treatments include metabolic therapy, behavioral and pharmacological neural activation and modulation of inflammatory responses.

## Acknowledgements

CCG acknowledges support from NIH NINDS (R01 NS27544), NCAA, U.S. Dept of Defense, UCLA Brain Injury Research Center, UCLA Steve Tisch BrainSPORT program, Easton Clinic for Brain Health, Avansir (research grant 2017-2018), NINDS Neural Analytics SBIR grant (NS092209 2016-2018). NGH acknowledges support by NIH NINDS NS065877, UG3NS106945, The UCLA Brain Injury Research Center. The project described was also supported in part by the MRI Core of the Semel Institute of Neuroscience at UCLA which is supported by Intellectual Development and Disabilities Research Center grant number U54HD087101-01 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development.

## References

- Adeleye, A., Shohami, E., Nachman, D., Alexandrovich, A.G., Trembovler, V., Yaka, R., Shoshan, Y., Dhawan, J., Biegan, A., 2009. D-cycloserine improves functional outcome after traumatic brain injury with wide therapeutic window. *Eur. J. Pharmacol.* 629 (1-3), 25–30.
- Ajao, D.O., Pop, V., Kamper, J.E., Adami, A., Rudbeck, E., Huang, L., Vilkovinsky, R., Hartman, R.E., Ashwal, S., Obenaus, A., Badaut, J., 2012. Traumatic brain injury in young rats leads to progressive behavioral deficits coincident with altered tissue properties in adulthood. *J. Neurotrauma* 29, 2060–2074.
- Albers, G.W., Goldstein, L.B., Hall, D., Lesko, L.M., 2001. Aptiganel hydrochloride in acute ischemic stroke. *J. Am. Med. Assoc.* 286, 2673–2682.
- Anderson, V., Catroppa, C., Morse, S., Haritou, F., Rosenfeld, J., 2005. Functional plasticity or vulnerability after early brain injury? *Pediatrics* 116, 1374–1382.
- Anderson, V., Godfrey, C., Rosenfeld, J.V., Catroppa, C., 2012. Predictors of cognitive function and recovery 10 years after traumatic brain injury in young children. *Pediatrics* 129, e254–e261.
- Anderson, V.A., Spencer-Smith, M.M., Coleman, L., Anderson, P.J., Greenham, M., Jacobs, R., Lee, K.J., Leventer, R.J., 2014. Predicting neurocognitive and behavioural outcome after early brain insult. *Dev. Med. Child Neurol.* 56, 329–336.
- Araki, Y., Zeng, M., Zhang, M., Haganir, R.L., 2015. Rapid dispersion of SynGAP from synaptic spines triggers AMPA receptor insertion and spine enlargement during LTP. *Neuron* 85, 173–189.
- Arbogast, K.B., Curry, A.E., Pfeiffer, M.R., Zonfrillo, M.R., Haarbauer-Krupa, J., Breiding, M.J., Coronado, V.G., Master, C.L., 2016. Point of health care entry for youth with concussion within a large pediatric care network. *JAMA Pediatr.* 170, e160294.
- Atkins, C.M., 2011. Decoding hippocampal signaling deficits after traumatic brain injury. *Transl. Stroke Res.* 2, 546–555.
- Atkins, C.M., Selcher, J.C., Petraitis, J.J., Trzaskos, J.M., Sweatt, J.D., 1998. The MAPK Cascade is required for mammalian associative learning. *Nat. Neurosci.* 1, 602–609.
- Atkins, C.M., Chen, S., Alonso, O.F., Dietrich, W.D., Hu, B.-R., 2006. Activation of calcium/calmodulin-dependent protein kinases after traumatic brain injury. *J. Cereb. Blood Flow Metab.* 26, 1507–1518.
- Atkins, C.M., Faló, M.C., Alonso, O.F., Bramlett, H.M., Dietrich, W.D., 2009. Deficits in ERK and CREB activation in the Hippocampus after traumatic brain injury. *Neurosci. Lett.* 459, 52–56.
- Babikyan, T., Prins, M.L., Cai, Y., Barkhoudarian, G., Hartonian, I., Hovda, D.A., Giza, C.C., 2010. Molecular and physiological responses to juvenile traumatic brain injury: focus on growth and metabolism. *Dev. Neurosci.* 32, 431–441.
- Babikyan, T., Alger, J.R., Ellis-Blied, M.U., Giza, C.C., Dennis, E., Olsen, A., Mink, R., Babbitt, C., Johnson, J., Thompson, P.M., Asarnow, R.F., 2018. Whole brain magnetic resonance spectroscopic determinants of functional outcomes in pediatric moderate/severe traumatic brain injury. *J. Neurotrauma* 35, 1637–1645.
- Baker, E.W., Kinder, H.A., Hutcheson, J.M., Duberstein, K.J.J., Platt, S.R., Howerth, E.W., West, F.D., 2019. Controlled cortical impact severity results in graded cellular, tissue, and functional responses in a piglet traumatic brain injury model. *J. Neurotrauma* 36, 61–73.
- Basser, P.J., 1997. New histological and physiological stains derived from diffusion-tensor MR images. *Ann. N. Y. Acad. Sci.* 820, 123–138.
- Baxter, M.G., Lanthorn, T.H., 1995. Modulation of the NMDA receptor complex by D-Cycloserine. *CNS Drug Rev.* 1, 74–90.
- Bell, J.D., Ai, J., Chen, Y., Baker, A.J., 2007. Mild in vitro trauma induces rapid GluR2 endocytosis, robustly augments calcium permeability and enhances susceptibility to secondary excitotoxic insult in cultured Purkinje cells. *Brain* 130, 2528–2542.
- Bell, J.D., Park, E., Ai, J., Baker, A.J., 2009. PICK1-mediated GluR2 endocytosis contributes to cellular injury after neuronal trauma. *Cell Death Differ.* 16, 1665–1680.
- Benarroch, E.E., 2011. NMDA receptors. Recent insights and clinical correlations. *Neurology* 76, 1750–1757.
- Bentzer, Peter, Davidsson, Helen, Grande, Per-Olof, 2000. *J. Neurotrauma* 17 (5), 441–447.
- Bergsneider, Marvin, Hovda, David A., Shalmon, Ehud, Kelly, Dnaiel F., Vespa, Paul M., Martin, Neil A., Phelps, Michael E., McArthur, David L., Caron, Michael J., Kraus, Jess F., Becker, Donald P., 1997. *J. Neurosurg.* 86, 241–251.
- Biegan, A., Fry, P.A., Paden, C.M., Alexandrovich, A., Tsenter, J., Shohami, E., 2004. Dynamic changes in N-methyl-D-aspartate receptors after closed head injury in mice: implications for treatment of neurological and cognitive deficits. *Proc. Natl. Acad. Sci. U. S. A.* 101, 5117–5122.

- Bittigau, P., Siffringer, M., Pohl, D., Stadthaus, D., Ishimaru, M., Shimizu, H., Ikeda, M., Lang, D., Speer, A., Olney, J.W., Ikonomidou, C., 1999. Apoptotic Neurodegeneration following trauma is markedly enhanced in the immature brain. *Ann. Neurol.* 45, 724–735.
- Blumbergs, P.C., Scott, G., Manavis, J., Wainwright, H., Simpson, D.A., McLean, A.J., 1994. Staining of amyloid precursor protein to study axonal damage in mild head injury. *Lancet* 344, 1055–1056.
- Bondi, C.O., Klitsch, K.C., Leary, J.B., Kline, A.E., 2014. Environmental enrichment as a viable neurorehabilitation strategy for experimental traumatic brain injury. *J. Neurotrauma* 31, 873–888.
- Bonnelle, V., Ham, T.E., Leech, R., Kinnunen, K.M., Mehta, M.A., Greenwood, R.J., Sharp, D.J., 2012. Salience network integrity predicts default mode network function after traumatic brain injury. *Proc. Natl. Acad. Sci.* 109, 4690–4695.
- Bonvento, G., Sibson, N., Pellerin, L., 2002. Does glutamate image your thoughts? *Trends Neurosci.* 25, 359–364.
- Bartnik, Brenda L., Sutton, Richard L., Fukushima, Masamichi, Harris, Neil G., Hovda, David A., Lee, Stefan M., 2005. *J. Neurotrauma* 22 (10), 1052–1065.
- Buen, F., Cai, Y., Santa Maria, N.S., Hovda, D.A., Giza, C.C., 2010. Lateral fluid percussion injury in developing rats disrupts activation of CaMKII and ERK 1/2 [abstract]. *J. Neurotrauma* 26.
- Buki, A., Povlishock, J.T., 2006. All roads lead to disconnection?—traumatic axonal injury revisited. *Acta Neurochir.* 148, 181–193 (discussion 193–184).
- Burda, J.E., Bernstein, A.M., Sofroniew, M.V., 2016. Astrocyte roles in traumatic brain injury. *Exp. Neurol.* 275 (Pt 3), 305–315.
- Cai, D., Deng, K., Mellado, W., Lee, J., Ratan, R.R., Filbin, M.T., 2002. Arginase I and polyamines act downstream from cyclic AMP in overcoming inhibition of axonal growth MAG and myelin in vitro. *Neuron* 35, 711–719.
- Casella, E.M., Thomas, T.C., Vanino, D.L., Fellows-Mayle, W., Lifshitz, J., Card, J.P., Adelson, P.D., 2014. Traumatic brain injury alters long-term hippocampal neuron morphology in juvenile, but not immature, rats. *Childs Nerv. Syst.* 30, 1333–1342.
- Chen, S., Atkins, C.M., Liu, C.L., Alonso, O.F., Dietrich, W.D., Hu, B.R., 2007. Alterations in mammalian target of rapamycin signaling pathways after traumatic brain injury. *J. Cereb. Blood Flow Metab.* 27, 939–949.
- Chugani, H.T., 1998. A critical period of brain development: studies of cerebral glucose utilization with PET. *Prev. Med.* 27, 184–188.
- Comeau, W.L., McDonald, R.J., Kolb, B.E., 2010. Learning-induced alterations in prefrontal cortical dendritic morphology. *Behav. Brain Res.* 214, 91–101.
- Coronado, V.G., Hailley, T., Cheng, T.A., Bell, J.M., Haarbauer-Krupa, J., Lionberger, M.R., Flores-Herrera, J., McGuire, L.C., Gilchrist, J., 2015. Trends in sports- and recreation-related traumatic brain injuries treated in US emergency departments: the National Electronic Injury Surveillance System-all Injury Program (NEISS-AIP) 2001–2012. *J. Head Trauma Rehabil.* 30, 185–197.
- Cosi, Cristina, Marien, Marc, 1998. *Brain Res.* 809, 58–67.
- Cull-Candy, S., Brickley, S., Farrant, M., 2001. NMDA receptor subunits: diversity, development and disease. *Curr. Opin. Neurobiol.* 11, 327–335.
- Cull-Candy, S., Kelly, L., Farrant, M., 2006. Regulation of Ca<sup>2+</sup>-permeable AMPA receptors: synaptic plasticity and beyond. *Curr. Opin. Neurobiol.* 16, 288–297.
- Dash, P.K., Mach, S.A., Moore, A.N., 2002. The role of extracellular signal-regulated kinase in cognitive and motor deficits following experimental traumatic brain injury. *Neuroscience* 114, 755–767.
- Dennis, E.L., Ellis, M.U., Marion, S.D., Jin, Y., Moran, L., Olsen, A., Kernan, C., Babikian, T., Mink, R., Babbitt, C., Johnson, J., Giza, C.C., Thompson, P.M., Asarnow, R.F., 2015a. Callosal function in pediatric traumatic brain injury linked to disrupted white matter integrity. *J. Neurosci.* 35, 10202–10211.
- Dennis, E.L., Jin, Y., Villalon-Reina, J.E., Zhan, L., Kernan, C.L., Babikian, T., Mink, R.B., Babbitt, C.J., Johnson, J.L., Giza, C.C., Thompson, P.M., Asarnow, R.F., 2015b. White matter disruption in moderate/severe pediatric traumatic brain injury: advanced tract-based analyses. *Neuroimage Clin.* 7, 493–505.
- Dennis, E.L., Faskowitz, J., Rashid, F., Babikian, T., Mink, R., Babbitt, C., Johnson, J., Giza, C.C., Jahanshad, N., Thompson, P.M., Asarnow, R.F., 2017a. Diverging volumetric trajectories following pediatric traumatic brain injury. *Neuroimage Clin.* 15, 125–135.
- Dennis, E.L., Rashid, F., Ellis, M.U., Babikian, T., Vlasova, R.M., Villalon-Reina, J.E., Jin, Y., Olsen, A., Mink, R., Babbitt, C., Johnson, J., Giza, C.C., Thompson, P.M., Asarnow, R.F., 2017b. Diverging white matter trajectories in children after traumatic brain injury: the RAPBI study. *Neurology* 88, 1392–1399.
- Dennis, E.L., Babikian, T., Alger, J., Rashid, F., Villalon-Reina, J.E., Jin, Y., Olsen, A., Mink, R., Babbitt, C., Johnson, J., Giza, C.C., Thompson, P.M., Asarnow, R.F., 2018a. Magnetic resonance spectroscopy of fiber tracts in children with traumatic brain injury: a combined MRS - diffusion MRI study. *Hum. Brain Mapp.* 39 (9), 3759–3768.
- Dennis, E.L., Babikian, T., Giza, C.C., Thompson, P.M., Asarnow, R.F., 2018b. Neuroimaging of the injured pediatric brain: methods and new lessons. *Neuroscientist* 24, 652–670.
- Deng-Bryant, Ying, Prins, Mayumi L., Hovda, David A., Harris, Neil G., 2011. *J. Neurotrauma* 28, 1812–1825.
- Dewan, M.C., Mumma, N., Wellons 3rd, J.C., Bonfield, C.M., 2016. Epidemiology of global pediatric traumatic brain injury: qualitative review. *World Neurosurg.* 91, 497–509 (e491).
- Dietrich, W.D., Alonso, O.F., Busto, R., Ginsberg, M.D., 1994. Widespread metabolic depression and reduced somatosensory circuit activation following traumatic brain injury in rats. *J. Neurotrauma* 11, 629–640.
- Diez, I., Drijckoningen, D., Stramaglia, S., Bonifazi, P., Marinazzo, D., Gooijers, J., Swinnen, S.P., Cortes, J.M., 2017. Enhanced prefrontal functional-structural networks to support postural control deficits after traumatic brain injury in a pediatric population. *Netw. Neurosci.* 1, 116–142.
- Dolle, J.P., Jaye, A., Anderson, S.A., Ahmadzadeh, H., Shenoy, V.B., Smith, D.H., 2018. Newfound sex differences in axonal structure underlie differential outcomes from in vitro traumatic axonal injury. *Exp. Neurol.* 300, 121–134.
- Ellis, M.U., DeBoard Marion, S., McArthur, D.L., Babikian, T., Giza, C., Kernan, C.L., Newman, N., Moran, L., Akarakiyan, R., Houshijarnejad, A., Mink, R., Johnson, J., Babbitt, C.J., Olsen, A., Asarnow, R.F., 2016. The UCLA study of children with moderate-to-severe traumatic brain injury: event-related potential measure of inter-hemispheric transfer time. *J. Neurotrauma* 33, 990–996.
- English, J.D., Sweatt, J.D., 1997. A requirement for the mitogen-activated protein kinase cascade in hippocampal long term potentiation. *J. Biol. Chem.* 272, 19103–19106.
- Ewing-Cobbs, L., Hasan, K.M., Prasad, M.R., Kramer, L., Bachevalier, J., 2006. Corpus callosum diffusion anisotropy correlates with neuropsychological outcomes in twins discordant for traumatic brain injury. *Am. J. Neuroradiol.* 27, 879–881.
- Ewing-Cobbs, L., Prasad, M.R., Swank, P., Kramer, L., Cox Jr., C.S., Fletcher, J.M., Barnes, M., Zhang, X., Hasan, K.M., 2008. Arrested development and disrupted callosal microstructure following pediatric traumatic brain injury: relation to neurobehavioral outcomes. *Neuroimage* 42, 1305–1315.
- Farrant, M., Nusser, Z., 2005. Variations on an inhibitory theme: phasic and tonic activation of GABA(a) receptors. *Nat. Rev. Neurosci.* 6, 215–229.
- Fineman, I., Giza, C.C., Nahed, B.V., Lee, S.M., Hovda, D.A., 2000. Inhibition of neocortical diffusion during development by a moderate concussive brain injury. *J. Neurotrauma* 17, 739–749.
- Finkbeiner, S., 2000. Calcium regulation of the brain-derived neurotrophic factor gene. *Cell. Mol. Life Sci.* 57, 394–401.
- Folkerts, M.M., Parks, E.A., Dedman, J.R., Kaetzel, M.A., Lyeth, B.G., Berman, R.F., 2007. Phosphorylation of calcium calmodulin-dependent protein kinase II following lateral fluid percussion brain injury in rats. *J. Neurotrauma* 24, 638–650.
- Gao, X., Deng, P., Xu, Z.C., Chen, J., 2011. Moderate traumatic brain injury causes acute dendritic and synaptic degeneration in the hippocampal dentate gyrus. *PLoS One* 6, e24566.
- Gilmore, J.H., Knickmeyer, R.C., Gao, W., 2018. Imaging structural and functional brain development in early childhood. *Nat. Rev. Neurosci.* 19, 123–137.
- Giza, C.C., Prins, M.L., 2006. Is being plastic fantastic? Mechanisms of altered plasticity after developmental traumatic brain injury. *Dev. Neurosci.* 28, 364–379.
- Giza, C.C., Griesbach, G.S., Hovda, D.A., 2005. Experience-dependent behavioral plasticity is disturbed following traumatic injury to the immature brain. *Behav. Brain Res.* 157, 11–22.
- Giza, C.C., Santa Maria, N.S., Hovda, D.A., 2006. N-methyl-D-aspartate receptor subunit changes after traumatic injury to the developing brain. *J. Neurotrauma* 23, 950–961.
- Giza, C.C., Kolb, B., Harris, N.G., Asarnow, R.F., Prins, M.L., 2009. Hitting a moving target: basic mechanisms of recovery from acquired developmental brain injury. *Dev. Neurorehabil.* 12, 255–268.
- Glazewski, S., Giese, K.P., Silva, A., Fox, K., 2000. The role of a-CaMKII Autophosphorylation in neocortical experience-dependent plasticity. *Nat. Neurosci.* 3, 911–918.
- Goldberg, J.L., 2003. How does an axon grow? *Genes Dev.* 17, 941–958.
- Griesbach, G., 2004. Voluntary exercise following traumatic brain injury: brain-derived neurotrophic factor upregulation and recovery of function. *Neuroscience* 125, 129–139.
- Griesbach, G.S., Hovda, D.A., 2015. Cellular and molecular neuronal plasticity. *Handb. Clin. Neurol.* 128, 681–690.
- Grinan-Ferre, C., Puigoriol-Illamola, D., Palomera-Avalos, V., Perez-Caceres, D., Companys-Aleman, J., Camins, A., Ortuno-Sahagun, D., Rodrigo, M.T., Pallas, M., 2016. Environmental enrichment modified epigenetic mechanisms in SAMP8 mouse Hippocampus by reducing oxidative stress and Inflammation and achieving neuroprotection. *Front. Aging Neurosci.* 8, 241.
- Gsell, W., Burke, M., Wiedermann, D., Bonvento, G., Silva, A.C., Dauphin, F., Buhrle, C., Hoehn, M., Schwindt, W., 2006. Differential effects of NMDA and AMPA glutamate receptors on functional magnetic resonance imaging signals and evoked neuronal activity during forepaw stimulation of the rat. *J. Neurosci.* 26, 8409–8416.
- Guerriero, R.M., Giza, C.C., Rotenberg, A., 2015. Glutamate and GABA imbalance following traumatic brain injury. *Curr. Neurol Neurosci. Rep.* 15, 27.
- Hagmann, P., Sporns, O., Madan, N., Cammoun, L., Pieni, V.J., Meuli, R., Thiran, J.P., Grant, P.E., 2010. White matter maturation reshapes structural connectivity in the late developing human brain. *Proc. Natl. Acad. Sci. U. S. A.* 107, 19067–19072.
- Han, R.Z., Hu, J.J., Weng, Y.C., Li, D.F., Huang, Y., 2009. NMDA receptor antagonist MK-801 reduces neuronal damage and preserves learning and memory in a rat model of traumatic brain injury. *Neurosci. Bull.* 25, 367–375.
- Hardingham, G.E., Fukunaga, Y., Bading, H., 2002. Extrasynaptic NMDARs oppose synaptic NMDARs by triggering CREB shut-off and cell death pathways. *Nat. Neurosci.* 5, 405–414.
- Hill, C.S., Coleman, M.P., Menon, D.K., 2016. Traumatic axonal injury: mechanisms and translational opportunities. *Trends Neurosci.* 39, 311–324.
- Hirata, T., 2009. Axon outgrowth. In: Binder, M.D., Hirokawa, N., Windhorst, U. (Eds.), *Encyclopedia of Neuroscience*. Springer, Berlin, Heidelberg, pp. 311–313.
- Hoffman, A.N., Malena, R.R., Westergom, B.P., Luthra, P., Cheng, J.P., Aslam, H.A., Zafonte, R.D., Kline, A.E., 2008a. Environmental enrichment-mediated functional improvement after experimental traumatic brain injury is contingent on task-specific neurobehavioral experience. *Neurosci. Lett.* 431, 226–230.
- Hoffman, A.N., Malena, R.R., Westergom, B.P., Luthra, P., Cheng, J.P., Aslam, H.A., Zafonte, R.D., Kline, A.E., 2008b. Environmental enrichment-mediated functional improvement after experimental traumatic brain injury is contingent on task-specific neurobehavioral experience. *Neurosci. Lett.* 431, 226–230.
- Huang, H., Shu, N., Mishra, V., Jeon, T., Chalak, L., Wang, Z.J., Rollins, N., Gong, G., Cheng, H., Peng, Y., Dong, Q., He, Y., 2015. Development of human brain structural networks through infancy and childhood. *Cereb. Cortex* 25, 1389–1404.

- Huh, J.W., Widing, A.G., Raghupathi, R., 2008. Midline brain injury in the immature rat induces sustained cognitive deficits, bihemispheric axonal injury and neurodegeneration. *Exp. Neurol.* 213, 84–92.
- Hyder, A.A., Wunderlich, C.A., Puvanachandra, P., Gururaj, G., Kobusingye, O.C., 2007. The impact of traumatic brain injuries: a global perspective. *NeuroRehabilitation* 22, 341–353.
- Ikonomidou, C., Turski, L., 2002. Why did NMDA receptor antagonists fail clinical trials for stroke and traumatic brain injury? *Lancet Neurol.* 1, 383–386.
- Ip, E.Y., Giza, C.C., Griesbach, G.S., Hovda, D.A., 2002. Effects of enriched environment and fluid percussion injury on dendritic Arborization within the cerebral cortex of the developing rat. *J. Neurotrauma* 19, 573–585.
- Isaac, J.T.R., Ashby, M.C., McBain, C.J., 2007. The role of the GluR2 subunit in AMPA receptor function and synaptic plasticity. *Neuron* 54, 859–871.
- Ishii, T., 1993. Molecular characterization of the family of N-methyl-D-aspartate receptor subunits. *J. Biol. Chem.* 268, 2836–2843.
- Johnson, V.E., Stewart, W., Smith, D.H., 2013. Axonal pathology in traumatic brain injury. *Exp. Neurol.* 246, 35–43.
- Karver, C.L., Wade, S.L., Cassidy, A., Taylor, H.G., Stancin, T., Yeates, K.O., Walz, N.C., 2012. Age at injury and long-term behavior problems after traumatic brain injury in young children. *Rehabil. Psychol.* 57, 256–265.
- Karver, C.L., Kurowski, B., Semple, E.A., Stancin, T., Taylor, H.G., Yeates, K.O., Walz, N.C., Wade, S.L., 2014. Utilization of behavioral therapy services long-term after traumatic brain injury in young children. *Arch. Phys. Med. Rehabil.* 95, 1556–1563.
- Katayama, Y., Becker, D.P., Tamura, T., Hovda, D.A., 1990. Massive increases in extracellular potassium and the indiscriminate release of glutamate following concussive brain injury. *J. Neurosurg.* 73, 889–900.
- Keenan, H.T., Presson, A.P., Clark, A.E., Cox, C.S., Ewing-Cobbs, L., 2019. Longitudinal developmental outcomes after traumatic brain injury in young children: are infants more vulnerable than toddlers? *J. Neurotrauma* 36, 282–292.
- Kenzie, E.S., Parks, E.L., Bigler, E.D., Lim, M.M., Chesnut, J.C., Wakeland, W., 2017. Concussion as a multi-scale complex system: an interdisciplinary synthesis of current knowledge. *Front. Neurol.* 8, 513.
- Kharlamov, E.A., Lepsveridze, E., Meparishvili, M., Solomon, R.O., Lu, B., Miller, E.R., Kelly, K.M., Mchedlishvili, Z., 2011. Alterations of GABA(a) and glutamate receptor subunits and heat shock protein in rat hippocampus following traumatic brain injury and in posttraumatic epilepsy. *Epilepsy Res.* 95, 20–34.
- Kilbaugh, Todd J., Bhandare, Sunita, Lorom, David H., Saraswati, Manda, Robertson, Couthney L., Margulies, Susan S., 2011. *J. Neurotrauma* 28, 763–774.
- Kochunov, P., Williamson, D.E., Lancaster, J., Fox, P., Cornell, J., Blangero, J., Glahn, D.C., 2012. Fractional anisotropy of water diffusion in cerebral white matter across the lifespan. *Neurobiol. Aging* 33, 9–20.
- Kolb, B., Cioe, J., 2003. Recovery from early cortical damage in rats. IX. Differential behavioral and anatomical effects of temporal cortex lesions at different ages of neural maturation. *Behav. Brain Res.* 144, 67–76.
- Kolb, B., Gibb, R., 2015. Plasticity in the prefrontal cortex of adult rats. *Front. Cell. Neurosci.* 9, 15.
- Kolb, B., Tomie, J.-A., 1988. Recovery from early cortical damage in rats. IV. Effects of hemidecortication at 1, 5 or 10 days of age on cerebral anatomy and behavior. *Behav. Brain Res.* 28.
- Kolb, B., Gibb, R., Gorny, G., 2000. Cortical plasticity and the development of behavior after early frontal cortical injury. *Dev. Neuropsychol.* 18, 423–444.
- Kolb, B., Gibb, R., Gorny, G., 2003. Experience-dependent changes in dendritic arbor and spine density in neocortex vary qualitatively with age and sex. *Neurobiol. Learn. Mem.* 79, 1–10.
- Kolb, B., Muhammad, A., Gibb, R., 2011. Searching for factors underlying cerebral plasticity in the Normal and injured brain. *J. Commun. Disord.* 44, 503–514.
- Kumar, A., Zou, L., Yuan, X., Long, Y., Yang, K., 2002. N-methyl-D-aspartate receptors: transient loss of NR1/NR2A/NR2B subunits after traumatic brain injury in a rodent model. *J. Neurosci. Res.* 67, 781–786.
- Lau, C.G., Zukin, R.S., 2007. NMDA receptor trafficking in synaptic plasticity and neuropsychiatric disorders. *Nat. Rev. Neurosci.* 8, 413–426.
- Laurie, D.J., Bartke, I., Schoepfer, R., Naujoks, K., Seeburg, P.H., 1997. Regional, developmental and interspecies expression of the four NMDAR2 subunits, examined using monoclonal antibodies. *Mol. Brain Res.* 51, 23–32.
- Lee, Stefan M., Wong, Monica D., Samii, Amir, Hovda, David A., 1999. *Annals NY Acad. Sci.* 893 (1), 337–340.
- Li, Q., Spigelman, I., Hovda, D.A., Giza, C.C., 2005. Decreased NMDA receptor mediated synaptic currents in CA1 neurons following fluid percussion injury in developing rats. abstract. *J. Neurotrauma* 20, 1249.
- Liraz-Zaltsman, S., Slusher, B., Atrakchi-Baranes, D., Rosenblatt, K., Friedman Levi, Y., Kesner, E., Silva, A.J., Biegon, A., Shohami, E., 2018. Enhancement of brain d-serine mediates recovery of cognitive function after traumatic brain injury. *J. Neurotrauma* 35, 1667–1680.
- Lisman, J., Shulman, H., Cline, H., 2002. The molecular basis of CaMKII function in Synaptic and Behavioural memory. *Nat. Rev. Neurosci.* 3, 175–190.
- Liu, D., Day, J.C., Francis, D.D., Meaney, M.J., 2000. Maternal care, hippocampal synaptogenesis and cognitive development in rats. *Nat. Neurosci.* 3, 799–806.
- Liu, X., Qiu, J., Alcon, S., Hashim, J., Meehan, W.P.I., Mannix, R., 2017. Environmental enrichment mitigates deficits after repetitive mild traumatic brain injury. *J. Neurotrauma* 34, 2445–2455.
- Loane, D.J., Byrnes, K.R., 2010. Role of microglia in neurotrauma. *Neurotherapeutics* 7, 366–377.
- Low, C.M., Wee, K.S.L., 2010. New insights into the not-so-new NR3 subunits of N-methyl-D-aspartate receptor: localization, structure, and function. *Mol. Pharmacol.* 78, 1–11.
- Lyons, D.N., Vekaria, H., Macheda, T., Bakshi, V., Powell, D.K., Gold, B.T., Lin, A.L., Sullivan, P.G., Bachstetter, A.D., 2018. A mild traumatic brain injury in mice produces lasting deficits in brain metabolism. *J. Neurotrauma* 35, 2435–2447.
- Maas, A., Roozenbeek, B., Manley, G.T., 2010. Clinical trials in traumatic brain injury: past Experience and current developments. *Neurotherapeutics* 7, 115–126.
- Marklund, N., Hillered, L., 2011. Animal modelling of traumatic brain injury in pre-clinical drug development: where do we go from here? *Br. J. Pharmacol.* 164, 1207–1229.
- Matter, A.M., Fowlie, K.A., Curatolo, L.M., Kline, A.E., 2011. Temporal effects of environmental enrichment-mediated functional improvement after experimental traumatic brain injury in rats. *Neurorehabil. Neural Repair* 25, 558–564.
- Menon, V., 2013. Developmental pathways to functional brain networks: emerging principles. *Trends Cogn. Sci.* 17, 627–640.
- Miller, L.P., Lyeth, B.G., Jenkins, L.W., Oleniak, L., Panchision, D., Hamm, R.J., Phillips, L.L., Dixon, C.E., Clifton, G.L., Hayes, R.L., 1990. Excitatory amino acid receptor subtype binding following traumatic brain injury. *Brain Res.* 526, 103–107.
- Monahan, J.B., Handelman, G.E., Hood, W.F., Cordi, A.A., 1989. D-Cycloserine, a positive modulator of the N-methyl-D-aspartate receptor, enhances performance of learning tasks in rats. *Pharmacol. Biochem. Behav.* 34, 649–653.
- Morris, G.F., Bullock, M.R., Marshall, S.B., Marmarou, A., Maas, A., The Selfotel Investigators, Marshall, L.F., 1999. Failure of the competitive N-methyl-D-aspartate antagonist Selfotel (CGS 19755) in the treatment of severe head injury: results of two phase III clinical trials. *J. Neurosurg.* 91, 737–743.
- Muir, K.W., 2006. Glutamate-based therapeutic approaches: clinical trials with NMDA antagonists. *Curr. Opin. Pharmacol.* 6, 53–60.
- Mychasiuk, R., Hehar, H., Ma, I., Kolb, B., Esser, M.J., 2015. The development of lasting impairments: a mild pediatric brain injury alters gene expression, dendritic morphology, and synaptic connectivity in the prefrontal cortex of rats. *Neuroscience* 288, 145–155.
- Narayan, R.K., Michel, M.E., GROUP, T.C.T.I.I.S., 2002. Clinical trials in head injury. *J. Neurotrauma* 19, 503–557.
- Nehlig, A., Porrino, L.J., Crane, A.M., Sokoloff, L., 1985. Local cerebral glucose utilization in Normal female rats: variations during the estrous cycle and comparison with males. *J. Cereb. Blood Flow Metab.* 5, 393–400.
- Ogawa, H., Hovda, D.A., Becker, D.P., 1992. Cerebral protein synthesis in the rat: further studies following fluid percussion injury [abstract]. *J. Neurotrauma* 9.
- Oh, W.C., Smith, K.R., 2019. Activity-dependent development of GABAergic synapses. *Brain Res.* 1707, 18–26.
- Oni, M.B., Wilde, E.A., Bigler, E.D., McCauley, S.R., Wu, T.C., Yallampalli, R., Chu, Z., Li, X., Hunter, J.V., Vasquez, A.C., Levin, H.S., 2010. Diffusion tensor imaging analysis of frontal lobes in pediatric traumatic brain injury. *J. Child Neurol.* 25, 976–984.
- Osteen, C.L., Moore, A.H., Prins, M.L., Hovda, D.A., 2001. Age-dependency of 45-calcium accumulation following lateral fluid percussion: acute and delayed patterns. *J. Neurotrauma* 18, 141–162.
- Osteen, C.L., Giza, C.C., Hovda, D.A., 2004. Injury-induced alterations in N-methyl-D-aspartate receptor subunit composition contribute to prolonged <sup>45</sup>Calcium accumulation following lateral fluid percussion. *Neuroscience* 128, 305–322.
- Owens, D.F., Kriegstein, A.R., 2002. Is there more to GABA than synaptic inhibition? *Nat. Rev. Neurosci.* 3, 715–727.
- Panizzutti, R., Rausch, M., Zurbrugg, S., Baumann, D., Beckmann, N., Rudin, M., 2005. The pharmacological stimulation of NMDA receptors via co-agonist site: an fMRI study in the rat brain. *Neurosci. Lett.* 380, 111–115.
- Paoletti, P., Neyton, J., 2007. NMDA receptor subunits: function and pharmacology. *Curr. Opin. Pharmacol.* 7, 39–47.
- Papadia, S., Stevenson, P., Hardingham, N.R., Bading, H., Hardingham, G.E., 2005. Nuclear Ca<sup>2+</sup> and the cAMP response element-binding protein family mediate a late phase of activity-dependent neuroprotection. *J. Neurosci.* 25, 4279–4287.
- Pettus, E.H., Povlishock, J.T., 1996. Characterization of a distinct set of intra-axonal ultrastructural changes associated with traumatically induced alteration in axolemmal permeability. *Brain Res.* 722, 1–11.
- Vespa, Paul, 2005. *J. Clin. Neurophysiol.* 22 (2), 99–106.
- Pickard, L., Noel, J., Henley, J.M., Coolingridge, G.L., Molnar, E., 2000. Developmental changes in synaptic AMPA and NMDA receptor distribution and AMPA receptor subunit composition in living hippocampal neurons. *J. Neurosci.* 20, 7922–7931.
- Pohl, D., Bittigau, P., Ishimaru, M.J., Stadthaus, D., Hubner, C., Olney, J.W., Turski, L., Ikonomidou, C., 1999. N-methyl-D-aspartate antagonists and apoptotic cell death triggered by head trauma in developing rat brain. *Proc. Natl. Acad. Sci.* 96, 2508–2513.
- Pressler, R., Auvin, S., 2013. Comparison of brain maturation among species: an example in translational research suggesting the possible use of bumetanide in newborn. *Front. Neurol.* 4, 36.
- Prins, M.L., Hovda, D.A., 2009. The effects of age and ketogenic diet on local cerebral METabolic rates of glucose after controlled cortical impact injury in rats. *J. Neurotrauma* 26, 1063–1093.
- Prins, M.L., Alexander, D., Giza, C.C., Hovda, D.A., 2013. Repeated mild traumatic brain injury: mechanisms of cerebral vulnerability. *J. Neurotrauma* 30, 30–38.
- Prins, Mayumi L., 2012. *Epilepsy Res.* 100, 218–223.
- Przekwas, A., Somayaji, M.R., Gupta, R.K., 2016. Synaptic mechanisms of blast-induced brain injury. *Front. Neurol.* 7, 2.
- Purves, D., Augustine, G.J., Fitzpatrick, D., Hall, W.C., LaMantia, A.-S., McNamara, J.O., Williams, S.M., 2004. *Neuroscience*, Third ed. Sinauer Associates, Inc.
- Quinlan, E.M., Olstein, D.H., Bear, M.F., 1999a. Bidirectional, experience-dependent regulation of N-methyl-d-aspartate receptor subunit composition in the rat visual cortex during postnatal development. *Proc. Natl. Acad. Sci.* 96, 12876–12880.
- Quinlan, E.M., Philpot, B.D., Hagan, R.L., Bear, M.F., 1999b. Rapid, experience-dependent expression of synaptic NMDA receptors in visual cortex in vivo. *Nat. Neurosci.* 2, 352–357.

- Raghupathi, R., 2003. Acute activation of mitogen-activated protein kinases following traumatic brain injury in the rat: implications for posttraumatic cell death. *Exp. Neurol.* 183, 438–448.
- Raghupathi, R., 2004. Cell death mechanisms following traumatic brain injury. *Brain Pathol.* 14, 215–222.
- Rao, V.L., Dogan, A., Todd, K.G., Bowen, K.K., Dempsey, R.J., 2001. Neuroprotection by Mementine, a non-competitive NMDA receptor antagonist after traumatic brain injury in rats. *Brain Res.* 911, 96–100.
- Reeves, T.M., Phillips, L.L., Povlishock, J.T., 2005. Myelinated and unmyelinated axons of the corpus callosum differ in vulnerability and functional recovery following traumatic brain injury. *Exp. Neurol.* 196, 126–137.
- Reger, M.L., Gurkoff, G.G., Hovda, D.A., Giza, C.C., 2005. The novel object recognition task detects a transient cognitive deficit after developmental TBI. abstract. *J. Neurotrauma* 20, 1206.
- Risen, S.R., Barber, A.D., Mostofsky, S.H., Suskauer, S.J., 2015. Altered functional connectivity in children with mild to moderate TBI relates to motor control. *J. Pediatr. Rehabil. Med.* 8, 309–319.
- Roberts, M.A., Manshadi, F.F., Bushnell, D.L., Hines, M.E., 1995. Neurobehavioural dysfunction following mild traumatic brain injury in childhood: a case report with positive findings on positron emission tomography (PET). *Brain Inj.* 9, 427–436.
- Robertson, C.L., Saraswati, M., Scaffidi, S., Fiskum, G., Casey, P., McKenna, M.C., 2013. Cerebral glucose metabolism in an immature rat model of pediatric traumatic brain injury. *J. Neurotrauma* 30, 2066–2072.
- Rosenzweig, M.R., Bennett, E.L., 1996. Psychobiology of plasticity: effects of training and experience on brain and behavior. *Behav. Brain Res.* 78, 57–65.
- Rosenzweig, M.R., Krech, D., Bennett, E.L., Diamond, M.C., 1962. Effects of environmental complexity and training on brain chemistry and anatomy: a replication and extension. *J. Comp. Physiol. Psychol.* 55, 429–437.
- Roskoski Jr., R., 2012. ERK1/2 MAP kinases: structure, function, and regulation. *Pharmacol. Res.* 66, 105–143.
- Saab, A.S., Tzvetavona, I.D., Trevisiol, A., Baltan, S., Dibaj, P., Kusch, K., Mobius, W., Goetze, B., Jahn, H.M., Huang, W., Steffens, H., Schomburg, E.D., Perez-Samartin, A., Perez-Cerda, F., Bakhtiari, D., Matute, C., Lowel, S., Griesinger, C., Hirrlinger, J., Kirchhoff, F., Nave, K.A., 2016. Oligodendroglial NMDA receptors regulate glucose import and axonal energy metabolism. *Neuron* 91, 119–132.
- Samorajski, T., Friede, R.L., 1968. A quantitative electron microscopic study of myelination in the pyramidal tract of rat. *J. Comp. Neurol.* 134.
- Sampedro-Piquero, P., Begega, A., 2017. Environmental enrichment as a positive behavioral intervention across the lifespan. *Curr. Neuropharmacol.* 15, 459–470.
- Santa Maria, N.S., Harris, N.G., Hovda, D.A., Giza, C.C., 2009. Reduced glutamatergic activation after developmental fluid percussion injury mapped by pharmacological MRI [abstract]. *J. Neurotrauma* 26 (A-44).
- Scheller, A., Bai, X., Kirchhoff, F., 2017. The role of the oligodendrocyte lineage in acute brain trauma. *Neurochem. Res.* 42, 2479–2489.
- Schonbach, J., Hu, K.H., Friede, R.L., 1968. Cellular and chemical changes during myelination: histologic, autoradiographic, histochemical and biochemical data on myelination in the pyramidal tract and corpus callosum of rat. *J. Comp. Neurol.* 134.
- Schreiber, S., Lin, R., Haim, L., Baratz-Goldstien, R., Rubovitch, V., Vaisman, N., Pick, C.G., 2014. Enriched environment improves the cognitive effects from traumatic brain injury in mice. *Behav. Brain Res.* 271, 59–64.
- Schuchat, A., Houry, D., Baldwin, G., 2016. Report to Congress - The Management of Traumatic Brain Injury in Children: Opportunities for Action.
- Schumann, J., Alexandrovich, A.G., Biegan, A., Yaka, R., 2008. Inhibition of NR2B phosphorylation restores alterations in NMDA receptor expression and improves functional recovery following traumatic brain injury in mice. *J. Neurotrauma* 25, 945–957.
- Schwarzbach, E., Bonislowski, D.P., Xiong, G., Cohen, A.S., 2006. Mechanisms underlying the inability to induce area CA1 LTP in the mouse after traumatic brain injury. *Hippocampus* 16, 541–550.
- Semple, B.D., Blomgren, K., Gimlin, K., Ferriero, D.M., Noble-Haesslein, L.J., 2013. Brain development in rodents and humans: identifying benchmarks of maturation and vulnerability to injury across species. *Prog. Neurobiol.* 106–107, 1–16.
- Semple, B.D., Noble-Haesslein, L.J., Jun Kwon, Y., Sam, P.N., Gibson, A.M., Grissom, S., Brown, S., Adahman, Z., Hollingsworth, C.A., Kwakye, A., Gimlin, K., Wilde, E.A., Hanten, G., Levin, H.S., Schenk, A.K., 2014. Sociosexual and communication deficits after traumatic injury to the developing murine brain. *PLoS One* 9, e103386.
- Sheline, Christian T., Behrens, M., Margarita, Choi, Dennis W., 2000. *J. Neurosci.* 20 (9), 3139–3146.
- Ying, Weihai, Garnier, Philippe, Swanson, Raymond A., 2003. *Biochem. Biophys. Res. Commun.* 308, 809–813.
- Simeone, T.A., Donevan, S.D., Rho, J.M., 2003. Molecular biology and ontogeny of gamma-aminobutyric acid (GABA) receptors in the mammalian central nervous system. *J. Child Neurol.* 18, 39–48 (discussion 49).
- Soriano, F.X., Papadia, S., Hofmann, F., Hardingham, N.R., Bading, H., Hardingham, G.E., 2006. Preconditioning doses of NMDA promote neuroprotection by enhancing neuronal excitability. *J. Neurosci.* 26, 4509–4518.
- Sozda, C.N., Hoffman, A.N., Olsen, A.S., Cheng, J.P., Zafonte, R.D., Kline, A.E., 2010. Empirical comparison of typical and atypical environmental enrichment paradigms on functional and histological outcome after experimental traumatic brain injury. *J. Neurotrauma* 27, 1047–1057.
- Sta Maria, N.S., Reger, M.L., Cai, Y., Baquing, M.A.T., Buen, F., Ponnaluri, A., Hovda, D.A., Harris, N.G., Giza, C.C., 2017. D-Cycloserine restores experience-dependent neuroplasticity after traumatic brain injury in the developing rat brain. *J. Neurotrauma* 34, 1692–1702.
- Stephens, J.A., Salorio, C.E., Gomes, J.P., Nebel, M.B., Mostofsky, S.H., Suskauer, S.J., 2017. Response inhibition deficits and altered motor network connectivity in the chronic phase of pediatric traumatic brain injury. *J. Neurotrauma* 34, 3117–3123.
- Stephens, J.A., Salorio, C.F., Barber, A.D., Risen, S.R., Mostofsky, S.H., Suskauer, S.J., 2018. Preliminary findings of altered functional connectivity of the default mode network linked to functional outcomes one year after pediatric traumatic brain injury. *Dev. Neurorehabil.* 21, 423–430.
- Stiles, J., Jernigan, T.L., 2010. The basics of brain development. *Neuropsychol. Rev.* 20, 327–348.
- Stranahan, A.M., Mattson, M.P., 2008. Impact of energy intake and expenditure on neuronal plasticity. *NeuroMolecular Med.* 10, 209–218.
- Tang-Schomer, M.D., Johnson, V.E., Baas, P.W., Stewart, W., Smith, D.H., 2012. Partial interruption of axonal transport due to microtubule breakage accounts for the formation of periodic varicosities after traumatic axonal injury. *Exp. Neurol.* 233, 364–372.
- Taylor, C.A., Bell, J.M., Breiding, M.J., Xu, L.X., 2017. In: Prevention, C.f.D.C.a (Ed.), *Traumatic Brain Injury-Related Emergency Department Visits, Hospitalizations, and Deaths - United States, 2007 and 2013*.
- Temple, M.D., Hamm, R.J., 1996. Chronic, post-injury administration of D-cycloserine, an NMDA partial agonist, enhances cognitive performance following experimental brain injury. *Brain Res.* 741, 246–251.
- Thomas, G.M., Hagan, R.L., 2004. MAPK Cascade Signalling and synaptic plasticity. *Nat. Rev. Neurosci.* 5, 173–183.
- Thomas, S., Prins, M.L., Samii, M., Hovda, D.A., 2000. Cerebral metabolic response to traumatic brain injury sustained early in development: a 2-deoxy-D-glucose autoradiographic study. *J. Neurotrauma* 17, 649–665.
- van Tilborg, E., de Theije, C.G.M., van Hal, M., Wagenaar, N., de Vries, L.S., Benders, M.J., Rowitch, D.H., Nijboer, C.H., 2018. Origin and dynamics of oligodendrocytes in the developing brain: implications for perinatal white matter injury. *Glia* 66, 221–238.
- de la Tremblaye, P.B., Wellcome, J.L., de Witt, B.W., Cheng, J.P., Skidmore, E.R., Bondi, C.O., Kline, A.E., 2017. Rehabilitative success after brain trauma by augmenting a subtherapeutic dose of environmental enrichment with Galantamine. *Neurorehabil. Neural Repair* 31, 977–985.
- Weihmuller, F.B., Bruno, J.P., 1989. Age-dependent plasticity in the dopaminergic control of sensorimotor development. *Behav. Brain Res.* 35, 95–109.
- Weil, Z.M., Karelina, K., Gaier, K.R., Corrigan, T.E., Corrigan, J.D., 2016. Juvenile traumatic brain injury increases alcohol consumption and reward in female mice. *J. Neurotrauma* 33, 895–903.
- Wheble, J.L., Menon, D.K., 2016. TBI-the most complex disease in the most complex organ: the CENTER-TBI trial-a commentary. *J. R. Army Med. Corps* 162, 87–89.
- Wilcox, K.S., Fitzsimonds, R.M., Johnson, B., Dichter, M.A., 1996. Glycine regulation of Synaptic NMDA receptors in hippocampal neurons. *J. Neurophysiol.* 76, 3415–3424.
- Wilde, E.A., Newsome, M.R., Bigler, E.D., Pertab, J., Merkle, T.L., Hanten, G., Scheibel, R.S., Li, X., Chu, Z., Yallampalli, R., Hunter, J.V., Levin, H.S., 2011. Brain imaging correlates of verbal working memory in children following traumatic brain injury. *Int. J. Psychophysiol.* 82, 86–96.
- de Witt, B.W., Ehrenberg, K.M., McAloon, R.L., Panos, A.H., Shaw, K.E., Raghavan, P.V., Skidmore, E.R., Kline, A.E., 2011. Abbreviated environmental enrichment enhances neurobehavioral recovery comparably to continuous exposure after traumatic brain injury. *Neurorehabil. Neural Repair* 25, 343–350.
- Worley, G., Hoffman, J.M., Paine, S.S., Kalman, S.L., Claerhout, S.J., Boyko, O.B., Kandt, R.S., Santos, C.C., Hanson, M.W., Oakes, W.J., et al., 1995. 18-Fluorodeoxyglucose positron emission tomography in children and adolescents with traumatic brain injury. *Dev. Med. Child Neurol.* 37, 213–220.
- Wu, A., Ying, Z., Gomez-Pinilla, F., 2009. Vitamin E protects against oxidative damage and learning disability after mild traumatic brain injury in rats. *Neurorehabil. Neural Repair* 24, 290–298.
- Wu, A., Ying, Z., Gomez-Pinilla, F., 2011. The salutary effects of DHA dietary supplementation on cognition, neuroplasticity, and membrane homeostasis after brain trauma. *J. Neurotrauma* 28, 2113–2122.
- Xiong, Y., Mahmood, A., Chopp, M., 2013. Animal models of traumatic brain injury. *Nat. Rev. Neurosci.* 14, 128–142.
- Yaka, R., Biegan, A., Grigoriadis, N., Simeonidou, C., Grigoriadis, S., Alexandrovich, A.G., Matzner, H., Schumann, J., Trembovler, V., Tsenter, J., Shohami, E., 2007. Cycloserine improves functional recovery and reinstates Long-term potentiation (LTP) in a mouse model of closed head injury. *FASEB J.* 21, 2033–2041.
- Yoshino, A., Hovda, D.A., Kawamata, T., Katayama, Y., Becker, D.P., 1991. Dynamic changes in local cerebral glucose utilization following cerebral concussion in rats: evidence of a hyper- and subsequent hypometabolic state. *Brain Res.* 561, 106–119.
- Yuan, W., Treble-Barna, A., Sohlberg, M.M., Harn, B., Wade, S.L., 2017a. Changes in structural connectivity following a cognitive intervention in children with traumatic brain injury. *Neurorehabil. Neural Repair* 31, 190–201.
- Yuan, W., Wade, S.L., Quatman-Yates, C., Hugentobler, J.A., Gubanich, P.J., Kurowski, B.G., 2017b. Structural connectivity related to persistent symptoms after mild TBI in adolescents and response to aerobic training: preliminary investigation. *J. Head Trauma Rehabil.* 32, 378–384.
- Zhang, Z., Saraswati, M., Koehler, R.C., Robertson, C., Kannan, S., 2015. A new rabbit model of pediatric traumatic brain injury. *J. Neurotrauma* 32, 1369–1379.
- Zhang, T.Y., Keown, C.L., Wen, X., Li, J., Voudsen, D.A., Anacker, C., Bhattacharyya, U., Ryan, R., Diorio, J., O'Toole, N., Lerch, J.P., Mukamel, E.A., Meaney, M.J., 2018. Environmental enrichment increases transcriptional and epigenetic differentiation between mouse dorsal and ventral dentate gyrus. *Nat. Commun.* 9, 298.