



# A longitudinal examination of associations between age and chemotherapy-induced peripheral neuropathy in patients with gynecologic cancer

Hailey W. Bulls <sup>a</sup>, Aasha I. Hoogland <sup>a,b</sup>, Brittany Kennedy <sup>a</sup>, Brian W. James <sup>b</sup>, Bianca L. Arboleda <sup>b</sup>, Sachin Apte <sup>a</sup>, Hye Sook Chon <sup>a</sup>, Brent J. Small <sup>b</sup>, Brian D. Gonzalez <sup>a</sup>, Heather S.L. Jim <sup>a,\*</sup>

<sup>a</sup> Moffitt Cancer Center, Tampa, FL, United States of America

<sup>b</sup> University of South Florida, Tampa, FL, United States of America

## HIGHLIGHTS

- Chemotherapy-induced peripheral neuropathy is commonly reported.
- On average, neuropathy significantly decreases for younger patients after treatment.
- Older adults report chronic neuropathy, extending well past the end of treatment.
- No differences in treatment modifications by age group were observed.

## ARTICLE INFO

### Article history:

Received 15 September 2018

Received in revised form 29 November 2018

Accepted 1 December 2018

Available online 14 December 2018

### Keywords:

Chemotherapy-induced peripheral neuropathy  
Age-related differences

## ABSTRACT

**Objective.** Increasing age has been associated with higher risk of chemotherapy-related toxicities, often resulting in treatment disruptions or discontinuations. Age has also been evaluated as a potential risk factor for chemotherapy-induced peripheral neuropathy (CIPN), but current understanding of recovery from CIPN in older adults after treatment is limited. The goal of the current study was to: 1) evaluate longitudinal change in patient-reported CIPN symptoms from the start of chemotherapy to one year post-chemotherapy; and 2) examine treatment modifications in older ( $\geq 65$  years) and younger patients ( $< 65$  years).

**Methods.** As part of a larger ongoing study, gynecologic cancer patients ( $n = 90$ ) treated with cytotoxic chemotherapy reported their CIPN symptoms via the EORTC-CIPN20 three times during active treatment and at 6 and 12 months post-treatment. Medical record reviews were conducted to abstract clinical information during active treatment.

**Results.** Piecewise mixed models revealed that older and younger patients reported similar increases in CIPN during the active treatment phase. However, older patients did not recover from CIPN after treatment completion, whereas younger patients exhibited significant declines in CIPN symptoms post-treatment. No age differences were observed in the presence of provider-recorded sensory neuropathy and pain; neuropathy-related treatment delays, changes in chemotherapy dose, regimen, or discontinuations; or falls (all  $p$ -values  $> 0.05$ ).

**Conclusions.** Results from the current study indicate that older adults are at higher risk for chronic CIPN. Older survivors may require additional education and treatment for chronic CIPN symptoms. Additional studies are needed to explore novel interventions to manage chronic CIPN in older cancer survivors.

© 2018 Elsevier Inc. All rights reserved.

## 1. Introduction

More than half of new cancer cases are diagnosed in adults over the age of 65 [1]. Moreover, cancer incidence among older adults is

projected to increase by 67% by 2030 [2]. Older cancer patients are at high risk for chemotherapy-related toxicities, with over 50% of patients over the age of 65 developing severe, disabling, or fatal treatment-related toxicities [3–6]. For example, older cancer patients treated with chemotherapy are significantly more likely to experience life-threatening toxicity (i.e., grade 4), to report treatment disruptions (e.g., treatment discontinuations due to such toxicity), and to die from treatment-related complications when compared to their younger

\* Corresponding author at: 12902 Magnolia Drive, MRC-PSY, Tampa, FL 33612, United States of America.

E-mail address: [heather.jim@moffitt.org](mailto:heather.jim@moffitt.org) (H.S.L. Jim).

counterparts [7,8]. As a result, even low-grade chemotherapy toxicities (i.e.,  $\leq$ grade 2) are considered clinically significant in older adults and can be sufficient cause for modifying or discontinuing chemotherapy [9].

One common chemotherapy-related toxicity in adult patients regardless of age is peripheral neuropathy. Chemotherapy-induced peripheral neuropathy (CIPN) is characterized by a variety of sensory and motor symptoms such as numbness, tingling, reduced sense of touch, reduced proprioception, pain, weakness, balance disturbances, and deficits in motor skills [10]. From a patient perspective, CIPN is one of the least expected and most distressing side effects of chemotherapy [11]. Symptoms typically increase in severity during treatment, but at least partially remit after treatment is completed [12]. However, a subset of patients experience chronic CIPN symptoms that continue past treatment and into survivorship [13,14]. Chronic CIPN is particularly common following treatment with cytotoxic chemotherapy agents [15]. At 6 months post-treatment, up to 30% of patients treated with chemotherapy report chronic CIPN symptoms [16]. Chronic CIPN has a negative impact on patients' quality of life long after treatment completion [17]. CIPN may also result in gait instability, an area of particular concern for older adults who are at higher risk of falls and subsequent injury [10,18].

Identification of patients at high risk for chronic CIPN is important in order to educate patients, manage expectations, and enact interventions. Age has been evaluated as a potential risk factor for CIPN, but methodological limitations constrain current understanding of recovery from CIPN in older adults. No known studies have longitudinally examined the progression of patient-reported CIPN symptomatology in older adults from before chemotherapy to the survivorship period. Among studies examining CIPN during active treatment or shortly thereafter (i.e., within 3 months), older patients typically report similar CIPN severity compared to younger patients [19–22]. Literature evaluating CIPN symptomatology after treatment in older adults is sparse. In one study with ovarian cancer survivors, older age was associated with higher incidence of patient-reported CIPN [23], and a second study with testicular cancer survivors found that older age was associated with greater physician-assessed CIPN severity [24]. However, these studies were cross-sectional and it is unclear when these associations emerged. Thus, further research is needed to better understand the longitudinal course of CIPN in older versus younger patients.

The goal of the current study was to evaluate longitudinal change in patient-reported CIPN in older ( $\geq 65$  years) and younger patients ( $< 65$  years) from the start of chemotherapy to one year post-chemotherapy. In this study, gynecologic cancer patients reported their CIPN symptoms via questionnaire three times during active treatment and at 6 and 12 months post-treatment. We anticipated that older and younger adults would report similar CIPN symptom severity during active treatment, but older patients would demonstrate poor recovery from CIPN symptoms after treatment compared to their younger counterparts. We also expected that older participants would 1) report more treatment disruptions (e.g., delays, regimen changes, and discontinuations) due to CIPN, and 2) sustain more falls during the active treatment phase than their younger counterparts.

## 2. Methods

### 2.1. Participant eligibility and recruitment

Participants were recruited as part of an ongoing, larger IRB-approved study examining side effects of cytotoxic chemotherapy in patients with gynecologic cancer. To be eligible, participants had to: (1) be 18–89 years of age; (2) be diagnosed with gynecologic cancer; (3) be scheduled to undergo intravenous (IV) or intraperitoneal (IP) cytotoxic chemotherapy at Moffitt Cancer Center (MCC); (4) have not received chemotherapy or radiation in the month prior to enrollment; (5) have no documented or observable psychiatric or neurological

disorders that would interfere with study participation (e.g., dementia, psychosis); (6) have no reported or documented diagnosis of immune-related disease (7) not be pregnant; (8) be able to speak and read English; and (9) be able to provide informed consent. Participants were recruited between August 2013 and January 2018.

Potential participants were identified via medical chart review in consultation with their treating physician at MCC and were contacted via phone by a trained research assistant in order to determine initial eligibility and interest in the study. Eligible women who wished to participate provided informed consent during an outpatient clinic visit prior to the start of chemotherapy. A total of 127 patients consented to participate in the study. Because the EORTC-CIPN20 was added later, 90 patients provided complete demographic data and at least one measure of CIPN, and were included in analyses. Participants completed CIPN and other questionnaires at five timepoints: (1) following chemotherapy cycle 1 (post-cycle 1), (2) following chemotherapy cycle 3 (post-cycle 3), (3) following chemotherapy cycle 6 (post-cycle 6), (4) 6 months post-chemotherapy (6 mo post-chemo), and (5) 12 months post-chemotherapy (12 mo post-chemo). A study flow diagram is presented in Fig. 1.




Fig. 1. Study flow diagram (CONSORT).

## 2.2. Measures

### 2.2.1. Demographic and clinical data

Sociodemographic characteristics were assessed via patient self-report prior to beginning chemotherapy. These characteristics included age, race/ethnicity, marital status, education level, and income. Medical comorbidities, including presence of diabetes, were ascertained via patient report on the Charlson Comorbidity Index [25]. Clinical characteristics were obtained by a medical record review, which included cancer type, stage, previous chemotherapy, and provider-recorded CIPN symptoms. ASCO-recommended medications for CIPN [26], including duloxetine, gabapentin, nortriptyline/SSRIs, and/or a topical gel comprised of baclofen, amitriptyline, and ketamine were also recorded if present at any time, for any reason, during treatment. Deviations from the planned regimen (e.g., treatment delays, dose and regimen changes, and discontinuations), patient-reported falls, and reasons for each were abstracted from clinical notes. Treatment modifications were considered to be possibly related to CIPN if they were 1) recorded as occurring due to CIPN symptoms, or 2) occurred in a patient with CIPN with no defined reason. Definitions of treatment modifications are as follows: Treatment delays were defined as delays in scheduled chemotherapy infusions due to CIPN symptoms. Dose changes were recorded when the dosage of chemotherapy drugs were lowered due to the presence of neuropathy. In both treatment delays and dose changes, the chemotherapy agent remained stable. In contrast, regimen changes were noted when the chemotherapy agent was changed to another drug in response to increasing neuropathy. Instances in which one agent in a combination was removed (e.g., removing taxanes from a platinum-taxane combination regimen) were also recorded as a regimen change. Finally, treatment discontinuations were recorded when chemotherapy was discontinued entirely due to CIPN symptoms, with no plans for subsequent treatment.

### 2.2.2. Chemotherapy-induced peripheral neuropathy

The European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire, Chemotherapy-Induced Peripheral Neuropathy-20 (EORTC QLQ-CIPN20) is a 20-item quality of life questionnaire developed to assess symptoms and functional limitations related to CIPN [27]. Items assess sensory, motor, and autonomic symptoms. Participants rate their symptoms over the past week on a 1 (not at all) to 4 (very much) scale. Summed scores are converted to a 0–100 scale for analysis, with higher scores indicating worse CIPN. Studies assessing the QLQ-CIPN20 indicate that the measure distinguishes between contrasting groups (e.g., those who have received neurotoxic chemotherapy vs. those who have not), has acceptable validity and reliability, and demonstrates sensitivity to change over time [28].

### 2.3. Statistical analysis

Patients were categorized into one of two age groups (<65 and ≥65 years old), consistent with previous epidemiological and clinical literature defining 65 as the cut-off age for older adults [19]. Patients' demographic and clinical characteristics were summarized using descriptive statistics. *t*-Tests and chi-square tests were used to examine age group differences in potential sociodemographic and clinical covariates. Sociodemographic and clinical characteristics that differed at a significance of  $p < 0.10$  were included as covariates in multivariate models. Mean CIPN scores by age group were plotted to visually inspect symptom trajectories during active treatment and post-treatment. SAS PROC MIXED was used to examine age-related longitudinal changes in CIPN from treatment initiation to one year post-chemotherapy [29]. Mixed models use all available data at every timepoint, making it an appropriate choice for longitudinal designs. Given that there are two distinct phases being assessed – on chemotherapy and post-chemotherapy – we used piecewise models [30] with post-cycle 6 as the cut-point. This analytic approach yielded information about

associations between age and self-reported CIPN change during chemotherapy (at post-cycle 1, post-cycle 3, and post-cycle 6), and assessed separate relationships between age and CIPN change during the post-treatment phase (from post-cycle 6 to 6 m and 12 m post-chemo). Interactions were then decomposed using follow-up mixed models separately within age group. Chi-square analyses or Fisher's exact tests were used to assess group differences in provider-reported sensory neuropathy and pain, treatment disruptions, and falls. All statistical analyses were performed using SAS 9.4 (Cary, NC).

## 3. Results

### 3.1. Sample characteristics

Sociodemographic and clinical characteristics of the 90 participants are shown in Table 1. The majority of the sample was non-Hispanic, white, and married. College degrees were held by 42% of the sample, and almost three-quarters of patients had a household income of at least \$40,000 per year. Patients were most often diagnosed with ovarian cancer (56%), followed by endometrial cancer (22%). Approximately 70% of patients had advanced (stages III and IV) cancer. The majority of patients (96%) received intravenous (IV) chemotherapy. Of the 4 patients who received intraperitoneal (IP) treatment, one patient received three IV/IP infusions (of six total) at the conclusion of her treatment, one patient received four IV/IP infusions (of six total) prior to switching to IV-only infusions due to increasing neuropathy, one patient received two IV/IP infusions and was still on study at the time of analysis, and one patient received a single IV/IP infusion before being discharged to hospice. The majority of patients were chemotherapy-naïve (49/90, 54%) and were receiving a taxane-platinum combination regimen (59/90, 66%). Information about current chemotherapy regimen and cycle length is shown in Table 1.

**Table 1**  
Participant characteristics.

	Overall N = 90	<65 y old n = 59	≥65 y old n = 31	Comparison p-value
Age: M (SD)	60.1 (10.9)	54.2 (8.0)	71.5 (4.8)	
Race: N (% white) <sup>a</sup>	84 (93)	53 (90)	31 (100)	0.07
Ethnicity: N (% non-Hispanic)	87 (97)	58 (98)	29 (94)	0.33
Marital status: N (% married)	60 (67)	41 (69)	19 (63)	0.83
Education: N (% college grad)	38 (42)	27 (45)	11 (35)	0.35
Household income: N (% > \$40k)	52 (73)	40 (78)	12 (60)	0.12
Cancer stage				0.13
I	17 (20)	14 (26)	3 (10)	
II	8 (9)	7 (13)	1 (3)	
III	47 (55)	27 (50)	20 (67)	
IV	13 (15)	7 (13)	6 (20)	
Chemotherapy regimen <sup>a</sup>				<0.0001
Platinum-taxane	59 (66)	44 (75)	15 (48)	
Platinum-containing	19 (21)	13 (22)	6 (19)	
Taxane-containing	4 (4)	0 (0)	4 (13)	
Other (non-platinum or taxane)	8 (9)	2 (6)	6 (19)	
Chemotherapy cycle				0.38
21 day	66	45 (76)	21 (68)	
28 day	24	14 (24)	10 (32)	
Diabetes: N (%)				0.18
Yes	9 (10)	4 (7)	5 (16)	
No	81 (90)	55 (93)	26 (84)	
Previous chemotherapy <sup>a</sup>				0.08
Yes	41 (46)	23 (39)	18 (58)	
No	49 (54)	36 (61)	13 (42)	

<sup>a</sup> Denotes group differences at  $p < 0.10$ , which were considered for inclusion in subsequent models as covariates.

**Table 2**

Piecewise mixed model unstandardized parameter estimates demonstrating an age group \* time interaction in the post-chemo phase, adjusting for race and prior chemotherapy.

Effect	Estimate	p-Value
Intercept	21.29	<0.01
Race	−9.91	0.13
Prior chemotherapy	2.73	0.39
Age group	0.47	0.89
OnChemo	2.22	0.13
OnChemo * race	0.54	0.73
OnChemo * prior chemotherapy	−2.07	0.01
OnChemo * age group	0.58	0.49
PostChemo	−0.46	0.30
PostChemo * race	0.86	0.06
PostChemo * prior chemotherapy	−0.35	0.10
Post-Chemo * age group	0.57	0.01

Chi-square analyses indicated that older participants were marginally more likely to be white ( $X^2(1) = 3.38, p = 0.07$ ), to have had prior chemotherapy ( $X^2(1) = 2.98, p = 0.08$ ), and were less likely to have received a platinum/taxane combination regimen ( $p < 0.0001$ ). Thus, race and receipt of prior chemotherapy were included in subsequent models. Because regimen type was confounded with receipt of prior chemotherapy (e.g., platinum/taxane given almost exclusively first-line), inclusion of prior chemotherapy as a covariate also largely controlled for regimen. No other demographic or clinical characteristics were significantly different between the age groups, including ethnicity, marital status, education, household income, and presence of diabetes (all  $p$  values  $> 0.10$ ).

**3.2. Piecewise mixed models**

Adjusted analyses indicated that during chemotherapy, there were no age by time interactions during the active phase of treatment above and beyond the influence of prior chemotherapy, suggesting that younger and older patients have similar increases in CIPN during this time ( $p > 0.05$ ). CIPN trajectories varied as a result of prior chemotherapy ( $p = 0.01$ ) but not race ( $p > 0.05$ ).

In the post-treatment assessment period, CIPN trajectories over time varied as a result of age group ( $p = 0.01$ ), while neither race nor prior chemotherapy significantly impacted the trajectory of CIPN symptoms after the conclusion of chemotherapy (both  $p$  values  $> 0.05$ ). Post-hoc analyses indicate that older patients did not exhibit changes in their CIPN symptoms post-treatment ( $p > 0.05$ ), whereas younger patients exhibited significant declines in CIPN symptoms ( $p < 0.01$ ). Results of


the piecewise mixed models analyses are shown in Table 2 and displayed in Fig. 2.

**3.3. CIPN-related treatment modifications**

Presence of provider-recorded sensory neuropathy and pain, treatment disruptions possibly due to CIPN (delays, dose changes, regimen changes, and discontinuations without subsequent chemotherapy), and falls are shown by age group in Table 3. Sensory neuropathy and pain were commonly documented in the medical records, with the majority of patients in both age groups reporting these symptoms at least once during treatment. Four patients were prescribed ASCO-recommended medication at any point in their treatment, including three patients using gabapentin (all in the younger group) and one patient using duloxetine (in the older group). Presence of medication management did not differ by age group. Treatment modifications (e.g., dose and regimen changes, delays, and discontinuations) due to CIPN were infrequent, with the most common modification being chemotherapy regimen changes. Falls were rare, with only one older patient reporting a fall during treatment. Chi-square analyses and Fisher’s exact tests revealed no statistically significant differences in the frequency of provider-reported sensory neuropathy and pain, treatment modification, or falls between the age groups (all  $p$ -values  $> 0.05$ ).

**4. Discussion**

The goals of this study were to evaluate longitudinal change in patient-reported CIPN in older ( $\geq 65$  years) and younger patients ( $< 65$  years) from the start of chemotherapy to one year post-chemotherapy. Results indicated that during active treatment, both older and younger patients reported similar CIPN trajectories. However, age differences were apparent in the post-treatment phase above and beyond the influence of prior chemotherapy and race, such that older adults continued to experience CIPN symptoms in the year following the completion of chemotherapy while younger patients exhibited a significant decline in CIPN symptoms. Though we anticipated that older patients may have a more difficult treatment course, resulting in more changes to planned treatment, there were no differences in the occurrence of treatment delays, dose or regimen changes, or treatment discontinuations during the active treatment phase between the age groups. Sensory neuropathy and pain were commonly reported symptoms across both age groups, while treatment changes and falls were relatively infrequent. Taken together, these data suggest that older patients are at higher risk of chronic CIPN than their younger counterparts,



**Fig. 2.** Piecewise CIPN scores, by age group.

**Table 3**  
Provider-reported symptomatology, treatment changes, and falls.

	Younger (<65 y old) n = 59	Older (≥65 y old) n = 31	p-Value
Sensory neuropathy & pain: N (%)	44 (74.6)	25 (80.1)	0.52
CIPN medication management: N (%)	3 (5)	1 (3)	1.00
Treatment modifications possibly related to CIPN:			
Delay: N (%)	2 (3.4)	2 (6.5)	0.61
Dose change: N (%)	0 (0)	2 (6.5)	0.12
Regimen change: N (%)	9 (15.3)	3 (9.7)	0.53
Treatment discontinuation: N (%)	2 (3.4)	2 (6.5)	0.61
Falls: N (%)	0 (0)	1 (3.2)	0.34

though their symptoms may be similar during active treatment. To our knowledge, this is the first longitudinal study to assess acute and chronic CIPN symptoms in the same cohort, allowing for a comparison of the CIPN symptom trajectory across age groups. That these age differences emerge shortly after the completion of treatment (e.g., 6 months post-chemotherapy) and continue until a year after treatment is a new finding.

The timing of age-related differences is notable, as older patients may not be aware that CIPN can continue or even worsen after the completion of treatment. Prior studies in this area were conducted in only the during- or post-treatment phase and were cross-sectional, which limited the ability to identify when age-related associations emerge. [19–23] This study adds to the literature by examining CIPN during and after treatment, identifying the post-treatment phase as an important period for age-related differences in CIPN. Thus, focused education outlining CIPN trajectories for older adults should be considered. Unfortunately, interventions to manage CIPN are limited. In this study, treatment modification decisions were made by each physician, based on the individualized needs of the patient and dependent on clinical variables such as the severity of CIPN, disease status, and preferences and goals of each patient. In terms of national guidelines for CIPN management, duloxetine is the best supported medication for use in patients with CIPN [26]. In light of the results of the current study, CIPN symptoms in patients over 65 years of age may warrant increased consideration of duloxetine in order to manage their CIPN. ASCO guidelines also tentatively support a few other medications (such as tricyclic antidepressants, gabapentin, and a compounded topical gel) based on their utility in other neuropathic pain conditions, but warn that it is unclear whether these agents are efficacious in treating CIPN specifically. Notably, medications were rarely used in this sample to manage CIPN; per medical record reviews, only four patients (4%) were using ASCO guideline-consistent medications at any time during treatment to manage their CIPN. Rather, the more common approach to CIPN management was to change treatment regimens (12/90, 13%). Older patients may also consider engaging in strength and balance training to help manage symptoms of CIPN that impact gait stability [10]. More research is needed to establish the efficacy of these interventions in older cancer survivors with CIPN.

Strengths of this study include a longitudinal design, allowing for CIPN symptom assessment through the entire active treatment phase and into a year post-treatment, and the availability of treatment records, allowing for the examination of treatment-related events such as delays, dose changes, and regimen changes. Additionally, the innovative use of piecewise analyses allowed us to examine CIPN changes in separate assessment periods (e.g., during active treatment and post-treatment), while the mixed models approach allow for the use of all available data. Limitations of the study include a limited sample size and a relatively homogenous patient group comprised of primarily white, married, well-educated patients. Additionally, there were no differences in comorbidities between the age groups, suggesting that this sample of older adult patients may have been more healthy than average. CIPN trajectories and subsequent impact on treatment changes may have been more pronounced in a more representative sample. These factors may limit the generalizability of findings.

Despite limitations, this study advances our understanding of CIPN trajectories in older cancer patients. Given the rapidly rising age of the population, combined with the increase in effective treatments and higher survival rates, CIPN is likely to become an ever-increasing burden to older cancer survivors. Therefore, additional studies are needed to explore novel interventions to manage chronic CIPN in older cancer survivors.

### Author contributions

**HWB:** Conceptualization, data curation and analysis, visualization, writing (original draft), writing (review and editing). **AIH:** Data curation and analysis, writing (review and editing). **BK:** Participant recruitment, data collection and curation, writing (review and editing); **BWJ:** Data collection and curation, writing (review and editing); **BA:** Data collection and curation, writing (review and editing); **SA:** Supervision, participant recruitment, writing (review and editing); **HSC:** Supervision, participant recruitment, writing (review and editing); **BJS:** Supervision, data curation and analysis, writing (review and editing); **BDG:** Supervision, writing (review and editing). **HSLJ:** Conceptualization, funding acquisition, methodology, project administration, supervision, writing (review and editing). The authors also wish to acknowledge the contributions of the Moffitt Cancer Center Survey Methods Shared Resource.

### Conflicts of interest

**HWB:** No conflicts of interest to disclose; **AIH:** No conflicts of interest to disclose; **BK:** No conflicts of interest to disclose; **BWJ:** No conflicts of interest to disclose; **BA:** No conflicts of interest to disclose; **SA:** No conflicts of interest to disclose; **HSC:** No conflicts of interest to disclose; **BJS:** No conflicts of interest to disclose; **BDG:** No conflicts of interest to disclose; **HSLJ:** No conflicts of interest to disclose.

### Funding sources

This study was supported by National Cancer Institute grants R01 CA164109 (PI: Jim), R25 CA090314 (PI: Brandon), and P30 CA076292 (PI: Sellers).

### References

- [1] National Cancer Institute, Age and Cancer Risk, Available from: <https://www.cancer.gov/about-cancer/causes-prevention/risk/age> 2015.
- [2] B.D. Smith, G.L. Smith, A. Hurria, G.N. Hortobagyi, T.A. Buchholz, Future of cancer incidence in the United States: burdens upon an aging, changing nation, *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*. 27 (17) (2009 Jun 10) 2758–2765, PubMed PMID 19403886 (Epub 2009/05/01).
- [3] T. Aparicio, J.L. Jouve, L. Teillet, D. Gargot, F. Subtil, V. Le Brun-Ly, et al., Geriatric factors predict chemotherapy feasibility: ancillary results of FFCO 2001–02 phase III study in first-line chemotherapy for metastatic colorectal cancer in elderly patients, *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*. 31 (11) (2013 Apr 10) 1464–1470, PubMed PMID 23460711 (Epub 2013/03/06).
- [4] A. Hurria, S. Mohile, A. Gajra, H. Klepin, H. Muss, A. Chapman, et al., Validation of a prediction tool for chemotherapy toxicity in older adults with cancer, *Journal of Clinical Oncology*. 34 (20) (2016) 2366–2371 2016/07/10.

- [5] A. Hurria, K. Togawa, S.G. Mohile, C. Owusu, H.D. Klepin, C.P. Gross, et al., Predicting chemotherapy toxicity in older adults with cancer: a prospective multicenter study, *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 29 (25) (2011 Sep 1) 3457–3465 , PubMed PMID 21810685 (PubMed Central PMCID: PMC3624700. Epub 2011/08/04).
- [6] T. Kalsi, G. Babic-Illman, P.J. Ross, N.R. Maisey, S. Hughes, P. Fields, et al., The impact of comprehensive geriatric assessment interventions on tolerance to chemotherapy in older people, *British Journal of Cancer*, 112 (2015) 1435 04/14/online.
- [7] H.B. Muss, D.A. Berry, C. Cirincione, D.R. Budman, I.C. Henderson, M.L. Citron, et al., Toxicity of older and younger patients treated with adjuvant chemotherapy for node-positive breast cancer: the cancer and leukemia group b experience, *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 25 (24) (2007 Aug 20) 3699–3704 , PubMed PMID 17704418 (Epub 2007/08/21).
- [8] S.M. Lichtman, A. Hurria, C.T. Cirincione, A.D. Seidman, E. Winer, C. Hudis, et al., Paclitaxel efficacy and toxicity in older women with metastatic breast cancer: combined analysis of CALGB 9342 and 9840, *Ann. Oncol.* 23 (3) (2012 Mar) 632–638 , PubMed PMID 21693770 (PubMed Central PMCID: PMC3331731. Epub 2011/06/23).
- [9] T. Kalsi, G. Babic-Illman, P. Fields, S. Hughes, N. Maisey, P. Ross, et al., The impact of low-grade toxicity in older people with cancer undergoing chemotherapy, *British Journal of Cancer*, 111 (2014) 2224 09/30/online.
- [10] T.F. Marshall, G.P. Zipp, F. Battaglia, R. Moss, S. Bryan, Chemotherapy-induced-peripheral neuropathy, gait and fall risk in older adults following cancer treatment, *Journal of Cancer Research and Practice*, 4 (4) (2017) 134–138 2017/12/01.
- [11] M. Sasane, T. Tencer, A. French, T. Maro, K. Beusterien, Patient-reported outcomes in chemotherapy-induced peripheral neuropathy: A review, *J Support Oncol.* 8 (6) (2010) e15–e21.
- [12] N. Kerckhove, A. Collin, S. Condé, C. Chaletix, D. Pezet, D. Balayssac, Long-term effects, pathophysiological mechanisms, and risk factors of chemotherapy-induced peripheral neuropathies: a comprehensive literature review, *Frontiers in Pharmacology*, 8 (2017) 86 (PubMed PMID: PMC5323411).
- [13] G. Cavaletti, P. Alberti, B. Frigeni, M. Piatti, E. Susani, Chemotherapy-induced neuropathy, *Curr. Treat. Options Neurol.* 13 (2) (2011 Apr) 180–190 , PubMed PMID 21191824 (Epub 2010/12/31).
- [14] A.A. Argyriou, J. Bruna, P. Marmiroli, G. Cavaletti, Chemotherapy-induced peripheral neurotoxicity (CIPN): an update, *Crit. Rev. Oncol. Hematol.* 82 (1) (2012 Apr) 51–77 , PubMed PMID 21908200 (Epub 2011/09/13).
- [15] W. Grisold, G. Cavaletti, A.J. Windebank, Peripheral neuropathies from chemotherapeutics and targeted agents: Diagnosis, treatment, and prevention, *Neuro-Oncology*, 14 (Suppl. 4) (2012) iv45–iv54 (PubMed PMID: PMC3480245).
- [16] T. Bao, C. Basal, C. Seluzicki, S.Q. Li, A.D. Seidman, J.J. Mao, Long-term chemotherapy-induced peripheral neuropathy among breast cancer survivors: prevalence, risk factors, and fall risk, *Breast Cancer Research and Treatment*, 159 (2) (2016) 327–333 08/10. (PubMed PMID: PMC5509538).
- [17] A.A. Beijers, F. Mols, W. Dercksen, C.M.L. Driessen, G. Vreugdenhil, Chemotherapy-induced peripheral neuropathy and impact on quality of life 6 months after treatment with chemotherapy, *J. Community Support Oncol.* 12 (11) (2014) 401–406, <https://doi.org/10.12788/jcso.0086>.
- [18] K.M. Winters-Stone, F. Horak, P.G. Jacobs, P. Trubowitz, N.F. Dieckmann, S. Stoyles, et al., Falls, functioning, and disability among women with persistent symptoms of chemotherapy-induced peripheral neuropathy, *J. Clin. Oncol.* 35 (23) (2017) 2604–2612 , PubMed PMID 28586243.
- [19] A.A. Argyriou, Panagiotis Polychronopoulos, Angelos Koutras, Gregoris Ionomou, Philippos Gourzis, Konstantinos Assimakopoulos, Kalofonos, P. Haralabos, Elisabeth Chroni, Is advanced age associated with increased incidence and severity of chemotherapy-induced peripheral neuropathy? *Support. Care Cancer* 14 (3) (2006 March 01) 223–229.
- [20] D.W. Kiser, T.B. Greer, M.C. Wilmoth, J. Dmochowski, R.W. Naumann, Peripheral neuropathy in patients with gynecologic cancer receiving chemotherapy: patient reports and provider assessments, *Oncol. Nurs. Forum* 37 (6) (2010 Nov) 758–764 , PubMed PMID 21059587 (Epub 2010/11/10).
- [21] K.A. Griffith, D.J. Couture, S. Zhu, N. Pandya, M.E. Johantgen, G. Cavaletti, et al., Evaluation of chemotherapy-induced peripheral neuropathy using current perception threshold and clinical evaluations, *Supportive Care in Cancer: Official Journal of the Multinational Association of Supportive Care in Cancer*, 22 (5) (2014 12/21) 1161–1169 (PubMed PMID: PMC3969764).
- [22] B.P. Schneider, F. Zhao, M. Wang, V. Stearns, S. Martino, V. Jones, et al., Neuropathy is not associated with clinical outcomes in patients receiving adjuvant taxane-containing therapy for operable breast cancer, *Journal of Clinical Oncology*, 30 (25) (2012) 3051–3057 (PubMed PMID: PMC3732004).
- [23] N.P. Ezendam, B. Pijlman, C. Bhugwandass, J.F. Pruijt, F. Mols, M.C. Vos, et al., Chemotherapy-induced peripheral neuropathy and its impact on health-related quality of life among ovarian cancer survivors: results from the population-based profiles registry, *Gynecol. Oncol.* 135 (3) (2014 Dec) 510–517 , PubMed PMID 25281491.
- [24] J.L. Glendenning, Y. Barbachano, A.R. Norman, D.P. Dearnaley, A. Horwich, R.A. Huddart, Long-term neurologic and peripheral vascular toxicity after chemotherapy treatment of testicular cancer, *Cancer* 116 (10) (2010 May 15) 2322–2331 , PubMed PMID 20225230 (Epub 2010/03/13).
- [25] M. Charlson, T.P. Szatrowski, J. Peterson, J. Gold, Validation of a combined comorbidity index, *J. Clin. Epidemiol.* 47 (11) (1994) 1245–1251.
- [26] D.L. Hershman, C. Lacchetti, R.H. Dworkin, E.M. Lavoie Smith, J. Bleeker, G. Cavaletti, et al., Prevention and management of chemotherapy-induced peripheral neuropathy in survivors of adult cancers: American Society of Clinical Oncology Clinical Practice Guideline, *J. Clin. Oncol.* 32 (18) (2014 Jun 20) 1941–1967 , PubMed PMID 24733808.
- [27] T.J. Postma, N.K. Aaronson, J.J. Heimans, M.J. Muller, J.G. Hildebrand, J.Y. Delattre, et al., The development of an eortc quality of life questionnaire to assess chemotherapy-induced peripheral neuropathy: the QLQ-CIPN20, *European Journal of Cancer (Oxford, England: 1990)* 41 (8) (2005 May) 1135–1139 , PubMed PMID 15911236 (Epub 2005/05/25).
- [28] E.M. Lavoie Smith, D.L. Barton, R. Qin, P.D. Steen, N.K. Aaronson, C.L. Loprinzi, Assessing patient-reported peripheral neuropathy: the reliability and validity of the european organization for research and treatment of cancer QLQ-CIPN20 questionnaire, *Qual. Life Res.* 22 (10) (2013 December 01) 2787–2799.
- [29] R.C. Littell, G.A. Milliken, W.W. Stroup, R.D. Wolfinger, O. Schabenberger, *SAS for Mixed Models*, 2nd ed. SAS Press, Cary, NC, 2006.
- [30] J.J. McArdle, Latent variable modeling of differences and changes with longitudinal data, *Annu. Rev. Psychol.* 60 (2009) 577–605 , PubMed PMID 18817479 (Epub 2008/09/27).