

Post-Mortem Imaging Adjudicated Sudden Death: Causes and Controversies



Giuseppe Femia, BSc, MBBS^{a,c*},
Christopher Semsarian, MBBS, PhD, MPH^{a,c,d}, Neil Langlois, MBBS, MD^e,
Mark McGuire, MMed, PhD^{a,c}, James Raleigh, MBBS^b,
Andrew Taylor, BA, MD^{f,g}, Rajesh Puranik, MBBS, PhD^{a,c}

^aDepartment of Cardiology, Royal Prince Alfred Hospital, Sydney, NSW, Australia

^bDepartment of Radiology, Royal Prince Alfred Hospital, Sydney, NSW, Australia

^cFaculty of Medicine, University of Sydney, Sydney, NSW, Australia

^dAgnes Ginges Centre for Molecular Cardiology, Centenary Institute, Sydney, NSW, Australia

^eSchool of Anatomy and Physiology, University of Adelaide, Frome Road, Adelaide, SA, Australia

^fUCL Institute of Cardiovascular Science, London, UK

^gGreat Ormond Street Hospital for Children, NHS Foundation Trust, London, UK

Sudden cardiac death (SCD) is a catastrophic complication of many cardiac conditions often occurring without warning. In these cases, a post-mortem examination is required to elucidate the cause of death and is regarded as the 'gold standard'. However, in circumstances of certain religious/cultural beliefs and advanced body decomposition an alternative non-invasive approach would be preferred. Although a developing field, post-mortem imaging using computed tomography (pmCT) or magnetic resonance imaging (pmMR) provides a non-invasive and accurate alternative to traditional post-mortem in specific circumstances. In particular, pmMR has an important role in younger decedents while pmCT is more suited to examination of adults with SCD. Despite encouraging results from several preliminary studies, more research is needed to determine the most appropriate role for post-mortem imaging in the clinical algorithm for investigation of SCD.

Keywords

Sudden cardiac death (SCD) • Conventional autopsy • Post-mortem imaging • Post-mortem CT • Post-mortem MR • Post-mortem CT (pmCT) • Post-mortem MR (pmMR)

Introduction

Sudden cardiac death (SCD) is an unexpected and devastating tragedy associated with major implications for the surviving family and community [1]. SCD can be defined as a sudden pulseless condition in the absence of a non-cardiac condition [2]. The estimated annual incidence for those aged 1–35 years is 0.5 to 3.2 per 100,000 person years with those aged 31 to 35 years having the highest incidence (3.2 per 100,000 person years) [3,4]. The lifetime risk of SCD in the adult population greater than 45 years of age ranges from 2.4 to 3.4% for women and 6.7 to 11.2% for men [5]. For the adult population, the most frequent underlying cause of SCD is

coronary artery disease (CAD) and cardiomyopathy. For the younger population, the causes of SCD are diverse and may be categorised into structural (e.g. inherited cardiomyopathies and Marfan syndrome) and non-structural, regarded as arrhythmogenic conditions (Brugada, long and short QT syndromes and catecholaminergic polymorphic ventricular tachycardia) [6,7].

The post-mortem or conventional autopsy is widely considered the "gold standard" in delineating the cause of SCD. In the majority of SCD cases attributable to a cardiac cause, structural heart disease is detected by conventional autopsy [8]. In contrast, the cause remains unclear in approximately 40% of young SCD cases, despite detailed histopathological

*Corresponding author at: SMRI level 1 100 Carillon Avenue Newtown Australia., Email: femia82@gmail.com

and toxicological analyses [3]. Despite its proven capabilities in post mortem diagnostics, there is a growing reluctance amongst families and some health care professionals to refer for conventional autopsies and, as a result, the rate of autopsy referrals has declined globally [9]. For reportable deaths, coroners must reconcile the public interest and potential value to the family of the deceased being fully investigated balanced against consideration of specific cultural and religious customs. Hence, there is an emerging role for an alternative, non-invasive post-mortem examination.

Recently, the emergence of post-mortem computed tomography (pmCT) and post-mortem magnetic resonance imaging (pmMR) addresses the need for a non-invasive and expeditious post-mortem investigation that maintains diagnostic accuracy. For both pmCT and pmMR, there have been substantial improvements in imaging technology, image acquisition and processing domains. When compared to conventional autopsy, a number of studies have shown promising evidence in support of post-mortem imaging. In some situations, the information obtained from post-mortem imaging or autopsy could be complemented by genetic testing. In both autopsy positive and negative cases, the addition of genetic testing may add information to identify the cause of death or provide an opportunity to screen family members for inheritable conditions [3,10]. In this review, we will discuss the potential and emerging roles for post-mortem imaging in the diagnosis of SCD in children and adults.

Causes of Sudden Cardiac Death

The causes of SCD can be divided into two categories: inherited and acquired. Inherited conditions may be further characterised as structural or arrhythmogenic. Structural conditions may be genotype and/or phenotype positive and include the inherited cardiomyopathies, such as hypertrophic cardiomyopathy (HCM), dilated cardiomyopathy (DCM), arrhythmogenic right ventricular cardiomyopathy (ARVC) and Marfan syndrome. The inherited arrhythmogenic syndromes are characteristically genotype positive but phenotypically have structurally normal hearts. These conditions include Brugada syndrome (BrS), long QT (LQTS) and short QT syndromes (SQTS) and catecholaminergic polymorphic ventricular tachycardia (CPVT). Acquired causes of SCD include myocarditis and CAD. Non-inherited structural conditions also include coronary artery anomalies as a cause for SCD. The most common cause of SCD in those less than 35 years of age is 'un-ascertained, presumed primary arrhythmogenic disorder' where the heart is structurally normal [7]. For those older than 35, the most common cause of SCD is ischaemic heart disease [11]. That said, structural conditions account for about 50% of deaths in those less than 5 years of age and 75% for those aged between 5 and 35 years of age.

Conventional Autopsy

Conventional autopsy involves an extensive and thorough external examination as well as an in-depth assessment of

internal organs with blood and tissue sampling for histological, microbiological and toxicological examination. The coronial process, including a detailed autopsy can be long and arduous, potentially adding substantial emotional stress to families as well as postponing funeral and religious ceremonies. Religious and cultural groups have different views on autopsy with some being opposed to the practice. As such, the referral rate for conventional autopsy has declined, particularly for the fetal and paediatric populations [12]. In addition to the reduction in referrals, several issues deplete the accuracy of conventional autopsy including the variability in tissue sampling practices and the lack of standardised examination protocols across forensic facilities.

Given the current sentiment regarding conventional autopsies, there is a growing need to develop alternative strategies for the investigation of sudden and unexpected death. One strategy uses post-mortem imaging as a 'triage' tool to determine the need for further, more conventional, histopathological examination. In some cases, imaging may provide a cause of death without the need for an invasive autopsy. One group performed whole-body computed tomography (CT) scans and external body examinations on 318 consecutive cases and found CT scanning has advantages in cases of accidental or suicidal death [13]. In these cases, a lower proportion of cases proceeded to formal autopsy (31–50%) than usual. Another strategy being tested is a minimally invasive autopsy facilitated by post-mortem imaging. In a recent study, full body pmMR at 1.5Tesla (T) in conjunction with a minimally invasive autopsy (blood sample without dissection or tissue histology) was compared to a conventional autopsy in fetuses and children aged less than 16 years of age [14]. In total there were 400 cases, where congenital ($n = 89$; 22%) and infective/inflammatory ($n = 82$; 21%) pathologies accounted for the most common diagnoses. A minimally invasive approach had excellent concordance for detection of major pathological abnormalities (89.3%) related to death compared to conventional autopsy but there was some variation amongst age groups. In addition, with a minimally invasive approach, a full autopsy may not have been necessary in 165 cases (41%); in these cases, the concordance rate between the two approaches was 99.4%. Minimally invasive autopsy had the highest concordance rate in fetuses and newborns; 94.6% for fetuses at 24 weeks' gestation, 81% for fetuses more than 24 weeks' gestation, 81% for newborns aged 1 month or younger and 84.9% for infants from 1 month to 1 year. However, the minimally invasive approach was less accurate (53.6%) for children from 1 year to 16 years of age. Using the pmMR and minimally invasive autopsy approach, sepsis was the most commonly missed cause of death (24 of 83 cases; 29%). In terms of cardiac pathologies (with or without relation to death), the combined approach failed to detect four cardiac conditions in the fetal group (Tetralogy of Fallot (ToF), ventricular septal defect (VSD) and myocardial infarction) and six in the older group (myocarditis). In addition, 10 cardiac abnormalities were incorrectly

diagnosed in the fetal group (aortic stenosis, anomalous venous drainage, co-arcuation of aorta and VSD) and two in the children group (ASD). Using pmMR alone, the concordance rate with conventional autopsy was lower than the combined approach (55.5%) with sepsis and placental pathology most commonly missed by pmMR. This study suggests that pmMR combined with a minimally invasive autopsy may provide a reliable alternative to conventional autopsy, especially in those cases aged under 1 year.

Post-Mortem Imaging—Current Evidence for Determining Cause of Death

Post-mortem imaging is a non-invasive tool that complements the conventional armamentarium and aids in the diagnosis of SCD. Both pmCT and pmMR are capable of high resolution cross sectional imaging with established utility in the post-mortem setting. Modern CT scanners have up to 320 slices and provide excellent spatial resolution, however, many post-mortem facilities only have access to 16- or 32-slice scanners limiting the technical quality and image resolution. The lower image resolution particularly becomes an issue when attempting to visualise small calibre vascular anatomy such as the coronary arteries [15]. In comparison to pmCT, pmMR does not use radiation but, instead, relies on strong magnetic fields and radiofrequency energy to alter the spin of protons within tissue. The image quality of MR depends on the signal to noise ratio and differs with various field strengths where current facilities use either 3T or 1.5T. Imaging at 3T generates greater signal than at 1.5T and can potentially provide higher resolution images with shorter acquisition times, making it ideal for visualising small bones, breast tissue, brain, spine and cardiac structures [16]. Imaging at 3T is, however, more susceptible to image artifacts and potentially more expensive than 1.5T scanners [17]. In terms of acquisition speed, CT is faster than MR and allows for whole body imaging within a few minutes. Magnetic resonance is also potentially dangerous in the presence of ferromagnetic foreign bodies such as “unsecured” cardiac devices or medical pumps. In these cases, caution is recommended, especially where such a device may be free rather than fixed and thus be attracted directly to the magnet. Overall, pmCT has excellent visualisation of the skeletal system, gas, fluids and foreign bodies while pmMR has excellent soft tissue contrast for most solid organs [16]. These technical differences provide each modality with distinct strengths and weaknesses in post-mortem imaging, summarised in Table 1.

Post-Mortem Imaging in Fetuses and Children

There is an increasing reluctance amongst health care professionals and parents to refer fetuses and children for

conventional autopsy. To address this issue, the development of alternative imaging strategies has become a priority, especially with sudden and unexpected deaths. As a diagnostic tool in this population, pmMR has been shown to be highly specific with a high negative predictive value. One study compared three-dimensional pmMR images at 1.5T to conventional autopsy from 185 fetuses younger than 24 weeks gestation, 92 fetuses older than 24 weeks gestation and 123 newborns and children younger than 16 years of age with SCD [18]. Compared to autopsy, pmMR correctly diagnosed 19 of 22 [86%] structural cardiac abnormalities from the pre-natal group and 13 of 13 (100%) from the post-natal group but missed three cases of structural heart disease (two ToFs and one VSD) in the fetal group and eight cases of myocarditis in the post-natal group. In addition, pmMR overcalled 13 abnormalities. Overall pmMR had a sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of 72.7%, 96.2%, 72.7% and 96.2%, respectively for any cardiac pathology; the accuracy was even higher for major structural abnormalities. This study highlights the difficulty differentiating myocarditis from post-mortem oedema using pmMR. Differentiating the two conditions poses a diagnostic dilemma and may rely on image guided biopsy to make a histological diagnosis.

In a prospective study comparing the diagnostic performance of perinatal and paediatric post-mortem imaging, 53 fetuses and 29 children underwent whole body pmCT using a 64-slice multi-detector system and pmMR at 1.5T prior to autopsy [19]. Comparing pmCT and pmMR to conventional autopsy, pmCT was found to have limited value in young children and fetuses. In this group of patients, pmMR was superior to pmCT with a higher diagnostic accuracy (43.6% vs. 32.7%). In addition, significantly more pmCT than pmMR examinations were non-diagnostic (18/82 (21.9%) vs. 4/82 (4.9%)). This difference was enhanced in the fetal group less than 24 weeks gestation (40% vs. 11%). Overall, in this cohort, pmMR was more accurate at detecting cardiac and central nervous system abnormalities.

A third study compared pmCT with a 64-slice scanner and pmMR at 1.5T to conventional autopsy in 17 patients aged 1–35 years of age with sudden death [20]. In this cohort, pmMR correctly identified all 12 (100%) cases with positive findings at autopsy. In addition, pmMR had no false negative cases compared to three (25%) cases with pmCT. However, pmMR incorrectly diagnosed one HCM case, one ARVC case and one pneumonia case; the misdiagnosis of HCM is a common issue described by others. In this population, pmMR was more accurate than pmCT and was found to be highly sensitive (100%) with a high negative (100%) and positive (80%) predictive value.

Lastly, in a systematic review examining the diagnostic accuracy of pmMR, data was extracted from nine studies that included 146 fetuses, 11 children and 24 adults [21]. In accurately identifying the final cause of death, pmMR had a sensitivity and specificity of 69% and 95% in fetuses and 28% and 64% in children and adults. These studies suggest that pmMR is highly accurate for imaging structural heart

Table 1 Comparing Post-Mortem Computed Tomography (pmCT) and Post-Mortem Magnetic Resonance Imaging (pmMR).

	pmCT	pmMR
Availability	Good	Limited
Maintenance Costs	High	Very High
Acquisition Time (whole body)	Short ~5 minutes	Long ~60 minutes
Spatial Resolution	High	High
Indications	Skeletal system injuries/trauma, detection of foreign bodies, detection of air/gas/fluid	Natural deaths, excellent soft tissue contrast for solid organs
Adult Cases	Preferred	Limited indications
Coronary Artery Calcification	Yes	No
Coronary Stenosis/Thrombosis	Possible	No
Acute Myocardial Infarction	No	Potential role
Fetal and Paediatric cases	Limited indications	Preferred
Anomalous Coronary Artery	Yes	No
Hypertrophic Cardiomyopathy	No	Technically difficult
Arrhythmogenic Right Ventricular Cardiomyopathy	No	Yes
Dilated Cardiomyopathy	Yes	Yes
Marfan Syndrome	Yes	Yes
Arrhythmogenic Conditions	No	Use as a "rule out" test for structural heart disease

disease in younger patients, but may be less suitable for investigating SCD in adults. As demonstrated by two studies in this review, pmMR may be less suited to diagnosing coronary artery disease [22,23]. Figure 1 demonstrates three conditions suitable for diagnosis with pmMR and confirmed on histopathology: acute myocardial infarction, ARVC and aortic dissection.

Post-Mortem Imaging in Adults

Compared to the younger population, there is less published data for post-mortem imaging in the adult SCD population. The largest study to date (n = 182), compared whole body pmCT using a 16-slice scanner and pmMR at 1.5-T to conventional autopsy [24]. Within this population, pmCT was the more accurate imaging modality with a higher concordance to conventional autopsy (68% vs 57%; p = 0.54). The most common conditions resulting in diagnostic discrepancies between autopsy and imaging were CAD, pulmonary emboli, pneumonia and intra-abdominal lesions. Using autopsy, 86 deaths were attributed to CAD and 10 to pulmonary emboli however post-mortem imaging failed to diagnose 12 cases (14%) of CAD and 10 cases (100%) of pulmonary emboli. It is not surprising that these two conditions were the most frequently missed given the difficulty in diagnosing arterial occlusions and infarctions using the current post-mortem techniques. In their systematic review, Thayyil et al. found that pmMR was less accurate in

identifying the final cause of death for children and adults (n = 35) compared to fetuses (n = 146) with lower sensitivity and specificity; 28% and 64% vs. 69% and 95%, respectively [14]. The most commonly missed diagnoses in the adult population were coronary artery disease and acute myocardial infarction. Although the published data is limited to small, heterogeneous studies, they highlight the potential need for using different imaging modalities in different age groups (Table 1).

Role of Post-Mortem Imaging in the Investigation of Inherited Cardiac Causes of Sudden Death

The effectiveness of post-mortem imaging in the diagnosis of inherited cardiac diseases is variable depending on the underlying conditions. In these circumstances, there may be significant value from combining post-mortem imaging, genetic testing and more invasive investigations to assist in diagnosing the cause of death. We propose the following schema categorising conditions based on the ability to make a diagnosis with post-mortem imaging: (i) definite phenotype, (ii) borderline phenotype or (iii) absent phenotype.

Definite Phenotype

For certain inherited structural conditions such as syndromic aortopathies, ARVC and premature CAD with characteristic




Figure 1 Post-mortem magnetic resonance imaging with correlating histopathology.

Acute myocardial infarction; (A–B) Axial T2 STIR and short axis T2 STIR MR images demonstrating regional hyperintense signal of the distal lateral wall, consistent with acute myocardial infarction/oedema. (C) Movat pentachrome stain, 2 x objective showing culprit left anterior descending coronary artery atherosclerosis with acute occlusive thrombus.

Arrhythmogenic Right Ventricular Cardiomyopathy; (A–B) Four-chamber and short axis b-fast field echo (FFE) MR image demonstrating severe RV dilation. (C) H&E stain, 10x objective showing right ventricular outflow tract focal fibrosis, scanty lymphocytic infiltration and extensive fatty infiltration.

Aortic Dissection; (A–B) 3-D “whole-heart” MR imaging reconstructed in axial and sagittal planes demonstrating severe ascending aortic dilatation and aortic dissection posteriorly with an associated large haemopericardium. (C) Movat pentachrome stain, 10x objective of aorta showing marked cystic medial degeneration with large pools of acid mucin.

Abbreviations: STIR, short tau inversion recovery; MR, magnetic resonance; RV, right ventricular, 3D, three dimensional

signs of familial hypercholesterolaemia a definitive phenotype can often be identified with post-mortem imaging. In these cases, post-mortem imaging may prevent the need for more invasive investigations such as conventional autopsy. Securing an accurate clinical diagnosis in these cases provides an important opportunity to offer clinical screening and/or cardiac imaging to family members.

Borderline Phenotype

For some inherited conditions such as HCM, post-mortem imaging may have limited capabilities to confirm a diagnosis. The difficulty with diagnosing HCM with post-mortem imaging is differentiating pathological asymmetric myocardial hypertrophy from concentric left ventricular hypertrophy, physiological remodelling in athletes and post-mortem myocardial oedema leading to increased wall thickness. Establishing a diagnosis in this setting is critical as HCM is a genetically based condition with an autosomal dominant pattern of inheritance and so has implications for other family members who may be at risk. Further, HCM is the most common inheritable heart disease with a prevalence of 1 in 200 people and an important cause of SCD in the young [25,26]. Clearly, in the scenario where there is clinical or imaging suspicion of HCM as the cause of SCD i.e. unexplained hypertrophy on post-mortem imaging, more

invasive autopsy investigations and/or genetic testing may be necessary to secure a diagnosis.

Absent Phenotype

Arrhythmogenic syndromes such as the channelopathies are most often associated with structurally normal hearts. This presents an important opportunity for post-mortem cardiac imaging to exclude structural heart disease i.e. relying on a good negative predictive value. In the situation where the heart can confidently be deemed structurally normal on post-mortem imaging, genetic testing may be required to identify the cause of SCD. One common approach to genetic screening in this situation involves testing for four genes; *KCNQ1* (LQTS1), *KCNH2* (LQTS2), *SCN5A* (LQTS3 and BrS) and *RYR2* (CPVT1) [27,28]. This testing regimen has shown variable success with a detection rate in selected tertiary referral populations up to 35% [29]. More recent advances in sequencing technologies have allowed the expansion of up to 200 detectable genes. Using genetic testing of 59 cardiac genes in the first ever prospective, population-based study of SCD in the young in Australia and New Zealand, a clinically relevant cardiac gene mutation was identified in 27% of autopsy negative cases [3]. In SCD cases where genetic testing is necessary, it is recommended that such testing be undertaken in a specialised multidisciplinary clinic setting [30].

Challenges for Post-Mortem Imaging

Despite encouraging results from the previously mentioned studies, post-mortem imaging has several limitations relating to image acquisition and access. In clinical practice, current protocols do not routinely use radio-opaque contrast for the visualisation of vascular anatomy, mostly owing to the need for a spontaneous circulation. Without a contrast agent, the effectiveness of both pmCT and pmMR as diagnostic modalities are somewhat compromised. Calcification of coronary arteries can be detected by pmCT but without a radio-opaque contrast the modality is limited in its capacity to diagnose pathological coronary lesions namely soft plaque and/or luminal thrombus. In ante-mortem cardiac MR imaging, scar is detected with the administration of the contrast agent gadolinium however, without the diagnostic capabilities of gadolinium in pmMR certain cardiac conditions such as myocarditis and myocardial infiltration are difficult to accurately diagnose.

Modern techniques are being established for post-mortem CT angiography (pmCTA) using various contrast agents and perfusion devices. One study selected 23 cases based on clinical history as well as the results of a baseline pmCT and evaluated the diagnostic value of pmCTA for investigating ischaemic heart disease as the cause of SCD [31]. The authors describe better visualisation of the coronary arteries and were able to better evaluate coronary lesions. Post-mortem CTA identified 11 of 14 coronary thrombi detected at autopsy with one identified as a chronic occlusion with collaterals and one as post-mortem clot. Despite studies demonstrating the utility of pmCTA in the diagnosis of ischaemic heart disease, it remains a research tool and will require further investigation in order to become part of routine practice. One group has proposed a novel strategy that uses air as a negative contrast agent in addition to a conventional positive contrast agent for targeted pmCTA of the coronary arteries [32]. In a small study (n = 25) using a multi-detector CT, this group described a simple and cost-effective targeted post-mortem cardiac angiography method.

Putrefaction and post-mortem cooling can alter the composition and structure of organs as well as the quality of post-mortem imaging. One particular issue is the variability that exists in the post-mortem interval to imaging as the precise time of death is often unknown.

Changes in body temperature, particularly due to cooling, can substantially affect pmMR imaging by prolonging the T1 value and shortening the T2 value producing reduced signal to noise on both T1 and T2 sequences [33]; pmCT is less susceptible to temperature changes above 0 degrees Celsius. These factors all contribute to making post-mortem imaging different from ante-mortem imaging and as a result, they require specialised image optimisation and sequence modification. In addition, any diagnostic criteria of cardiac pathologies that relies on the difference between systolic and diastolic dimensions or functional abnormalities cannot be

used. This does, however, present an important opportunity for the development of new diagnostic criteria in the post-mortem setting.

Although post-mortem imaging has been shown to be highly accurate in certain circumstances, access to scanners and reporting expertise remains challenging, particularly if the scanning facility is not co-located with a mortuary. In order to achieve specialised reporting, collaboration of local imaging facilities with trained professionals is recommended [34]. Moving forward, standardised training programs may become necessary. Currently, there are no established protocols for post-mortem imaging but some image parameter recommendations are available but will often be modality and vendor specific [15] [35]. Of note, the literature to date has often used different imaging modalities, acquisition protocols and reporting expertise that ultimately confounds conclusions from the available published data. The need for larger studies with uniform imaging protocols and reporting is necessary to further understand the diagnostic potential of post-mortem imaging in SCD.

Conclusions and Future Directions

In the evaluation of SCD, post-mortem imaging with either MR or CT may be combined with conventional autopsy and histological examination to identify the cause of death. Based on current knowledge, post-mortem imaging provides an effective alternative to conventional autopsy in specific circumstances. Post-mortem MR has important utility within the younger population especially with structural cardiac abnormalities. In contrast, post-mortem CT has an emerging role in the adult population, potentially assisting with the diagnosis of coronary artery disease. The addition of genetic testing in certain circumstances of suspected inherited heart disease may also add to overall diagnostic and ultimately screening capabilities.

In order for post-mortem imaging to become a routine component for the investigation of SCD, there remain important challenges of cost, resource availability and the development of specialised reporting and imaging protocols. However, in situations where conventional autopsy cannot be performed due to cultural or religious reasons or significant body decomposition, post-mortem imaging has definite advantages. More research is necessary to determine appropriate clinical pathways utilising post-mortem imaging in the investigation of SCD.

References

- [1] Semsarian C, Hamilton RM. Key role of the molecular autopsy in sudden unexpected death. *Heart Rhythm* 2012;9:145–50.
- [2] Siscovick DS, Weiss NS, Fletcher RH, Lasky T. The incidence of primary cardiac arrest during vigorous exercise. *N Engl J Med* 1984;311:874–7.
- [3] Bagnall RD, Weintraub RG, Ingles J, Dufflou J, Yeates L, Lam L, et al. A prospective study of sudden cardiac death among children and young adults. *Engl J Med* 2016;374:2441–52.

- [4] Risgaard B, Winkel BC, Jabbari R, Behr ER, Ingemann-Hansen O, Thomsen JL, et al. Burden of sudden cardiac death in persons aged 1 to 49 years: nationwide study in Denmark. *Circ Arrhythm Electrophysiol* 2014;7:205–11.
- [5] Bogle BM, Ning H, Mehrotra S, Goldberger JJ, Lloyd-Jones DM. Lifetime risk for sudden cardiac death in the community. *J Am Heart Assoc* 2016;5(7).
- [6] Drory Y, Turetz Y, Hiss Y, Lev B, Fisman EZ, Pines A, et al. Sudden unexpected death in persons less than 40 years of age. *Am J Cardiol* 1991;68:1388–92.
- [7] Doolan A, Langlois S, Semsarian C. Causes of sudden cardiac death in young Australians. *Med J Aust* 2004;180(3):110–2.
- [8] van der Werf C, van Langen IM, Wilde AA. Sudden death in the young: what do we know about it and how to prevent. *Circ Arrhythm Electrophysiol* 2010;3:96–104.
- [9] O'Grady G. Death of the teaching autopsy. *BMJ* 2003;327:802–3.
- [10] Bagnall RD, Das KJ, Duflou J, Semsarian C. Exome analysis-based molecular autopsy in cases of sudden unexplained death in the young. *Heart Rhythm* 2014;11:655–62.
- [11] Eckart RE, Shry EA, Burke AP, McNear JA, Appel DA, Castillo-Rojas LM, et al. Sudden death in young adults: an autopsy-based series of a population undergoing active surveillance. *J Am Coll Cardiol* 2011;58:1254–61.
- [12] Kumar P, Angst DB, Taxy J, Mangurten HH. Neonatal autopsies: a 10-year experience. *Arch Pediatr Adolesc Med* 2000;154:38–42.
- [13] Bedford PJ. Routine CT scan combined with preliminary examination as a new method in determining the need for autopsy. *Forensic Sci Med Pathol* 2012;8:390–4.
- [14] Thayyil S, Sebire NJ, Chitty LS, Wade A, Chong W, Olsen O, et al. Post-mortem MRI versus conventional autopsy in fetuses and children: a prospective validation study. *Lancet* 2013;382:223–33.
- [15] Gascho D, Thali MJ, Niemann T. Post-mortem computed tomography: technical principles and recommended parameter settings for high-resolution imaging. *Med Sci Law* 2018;58:70–82.
- [16] Grabherr S, Egger C, Vilarino R, Campana L, et al. Jotterand M. Modern post-mortem imaging: an update on recent developments. *Forensic Sciences Research* 2017;2(2):52–64.
- [17] R. Wood , K. Bassett , V. Foester , C. Spry , L. Tong . 1.5 Tesla Magnetic Resonance Imaging Scanners Compared with 3.0 Tesla Magnetic Resonance Imaging Scanners: Systematic Review of Clinical Effectiveness: Pilot Project, Canadian Agency for Drugs and Technologies in Health
- [18] Taylor AM, Sebire NJ, Ashworth MT, Schievano S, Scott RJ, Wade A, et al. Post-mortem cardiovascular magnetic resonance imaging in fetuses and children a masked comparison study with conventional autopsy. *Circulation* 2014;129:1937–44.
- [19] Arthurs OJ, Guy A, Thayyil S, Wade A, Jones R, Norman W, et al. Comparison of diagnostic performance for perinatal and paediatric post-mortem imaging: CT versus MRI. *Eur Radiol* 2016;26:2327–36.
- [20] Puranik R, Gray B, Lackey H, Yeates L, Parker G, Duflou J, et al. Comparison of conventional autopsy and magnetic resonance imaging in determining the cause of sudden death in the young. *J Cardiovasc Magn Reson* 2014;16:44.
- [21] Thayyil S, Chandrasekaran M, Chitty LS, Wade A, Skordis-Worrall J, Bennett-Britton I, et al. Diagnostic accuracy of post-mortem magnetic resonance imaging in fetuses, children and adults: a systematic review. *Eur J Radiol* 2010;75:e142–148.
- [22] Bisset RA, Thomas NB, Turnbull IW, Lee S. Post-mortem examinations using magnetic resonance imaging: four year review of a working service. *BMJ* 2002;324:1423–4.
- [23] Patriquin L, Kassarjian A, Barish M, Casserley L, O'Brien M, Andry C, Eustace S. Post-mortem whole-body magnetic resonance imaging as an adjunct to autopsy: preliminary clinical experience. *J Magn Reson Imaging* 2001;13:277–87.
- [24] Roberts IS, Benamore RE, Benbow EW, Lee SH, Harris JN, Jackson A, et al. Post-mortem imaging as an alternative to autopsy in the diagnosis of adult deaths: a validation study. *Lancet* 2012;379:136–42.
- [25] Semsarian C, Ingles J, Maron MS, Maron BJ. New perspectives on the prevalence of hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2015;65:1249–54.
- [26] Maron BJ, Maron MS, Semsarian C. Genetics of hypertrophic cardiomyopathy after 20 years: clinical perspectives. *J Am Coll Cardiol* 2012;60:705–15.
- [27] Ackerman MJ, Priori SG, Willems S, Berul C, Brugada R, Calkins H, et al. HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies. This document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Heart Rhythm* 2011;8:1308–39.
- [28] Wilde AA, Behr ER. Genetic testing for inherited cardiac disease. *Nat Rev Cardiol* 2013;10:571–83.
- [29] Tester DJ, Medeiros-Domingo A, Will ML, Haglund CM, Ackerman MJ. Cardiac channel molecular autopsy: insights from 173 consecutive cases of autopsy-negative sudden unexplained death referred for post-mortem genetic testing. *Mayo Clin Proc* 2012;87:524–39.
- [30] Semsarian C, Ingles J, Wilde AA. Sudden cardiac death in the young: the molecular autopsy and a practical approach to surviving relatives. *Eur Heart J* 2015;36:1290–6.
- [31] Michaud K, Grabherr S, Doenz F, Mangin P. Evaluation of post-mortem MDCT and MDCT-angiography for the investigation of sudden cardiac death related to atherosclerotic coronary artery disease. *Int J Cardiovasc Imaging* 2012;28:1807–22.
- [32] Saunders SL, Morgan B, Raj V, Robinson CE, Ruddy GN. Targeted post-mortem computed tomography cardiac angiography: proof of concept. *Int J Legal Med* 2011;125:609–16.
- [33] Kobayashi T, Isobe T, Shiotani S, Saito H, Saotome K, Kaga K, et al. Post-mortem magnetic resonance imaging dealing with low temperature objects. *Magn Reson Med Sci* 2010;9:101–8.
- [34] O'Donnell C, Woodford N. Post mortem radiology: a new sub-speciality. *Clin Radiol* 2008;63:1189–94.
- [35] Savage DD, Siedes SF, Maron BJ, Myers DJ, Epstein SE. Prevalence of arrhythmias during 24-hour electrocardiographic monitoring and exercise testing in patients with obstructive and nonobstructive hypertrophic cardiomyopathy. *Circulation* 1979;59:866–75.