



## Corrigendum

## Corrigendum to “Single-center experience with percutaneous lead extraction of cardiac implantable electric devices” [J. Cardiol. 71 (2018) 192–196]



Ayako Okada (MD, PhD, FJCC) \*, Morio Shoda (MD, PhD), Hiroaki Tabata (MD), Wataru Shoin (MD), Hideki Kobayashi (MD), Takahiro Okano (MD), Koji Yoshie (MD), Yasutaka Oguchi (MD), Takahiro Takeuchi (MD), Ken Kato (MD), Koichiro Kuwahara (MD, PhD, FJCC)

Department of Cardiovascular Medicine, Shinshu University School of Medicine, Nagano, Japan

There is an error in “Table 2. Extraction outcomes.”  
Correctly, the rate of blood transfusion is 2/29 (7%).  
The table should be revised as:

**Table 2**  
Extraction outcomes.

Result	(n = 29, leads = 61) Complete success: 28 (60 leads), 97%; Partial success: 1 (1 lead)
Anesthesia time (hours)	5.1 (4.6–5.5)
Procedure time (hours)	3.7 (3.2–4.1)
Length of ICU stay (days)	1 day (No. 24: 30 days)
Bleeding (mL)	64.8 (48.6–81.0)
Blood transfusion	2 (7%)
Excimer laser sheath	26 (58 leads)
Mechanical sheath	7 (12 leads)
Only mechanical sheath	1 (2 leads)
Manual extraction	1 (2 leads)
Complications: major/minor	0/0
Device re-implantation	27 (93%)
Infection recurrence	0 (0%)

DOI of original article: <https://doi.org/10.1016/j.jjcc.2017.07.011>

\* Corresponding author at: Department of Cardiovascular Medicine, Shinshu University School of Medicine, 3-1-1 Asahi, Matsumoto, Nagano 390-8621, Japan.

<https://doi.org/10.1016/j.jjcc.2018.12.001>

0914-5087/© 2018 Published by Elsevier Ltd on behalf of Japanese College of Cardiology.