



Association of pro-inflammatory cytokines and monocyte subtypes in older and younger patients on clinical outcomes after mechanical circulatory support device implantation

Joanna M. Schaenman^{a,*}, Maura Rossetti^b, Tiffany Sidwell^b, Victoria Groysberg^b, Gemalene Sunga^b, Emily Liang^b, Sitaram Vangala^c, Eleanor Chang^d, Maral Bakir^d, Galya Bondar^d, Martin Cadeiras^d, Murray Kwon^e, Elaine F. Reed^b, Mario Deng^d

^a Division of Infectious Diseases, Department of Medicine, David Geffen School of Medicine, Los Angeles, CA 90095, United States

^b Department of Pathology and Laboratory Medicine, David Geffen School of Medicine, Los Angeles, CA 90095, United States

^c UCLA Department of Medicine Statistics Core, David Geffen School of Medicine, Los Angeles, CA 90095, United States

^d Division of Cardiology, Department of Medicine, David Geffen School of Medicine, Los Angeles, CA 90095, United States

^e Department of Cardiothoracic Surgery, David Geffen School of Medicine, Los Angeles, CA 90095, United States

ARTICLE INFO

Keywords:

Immunosenescence
Inflammation
Monocytes
Aging
Heart failure
Ventricular assist device

ABSTRACT

Noninvasive immunologic analysis of peripheral blood holds promise for explaining the mechanism of development of adverse clinical outcomes, and may also become a method for patient risk stratification before or after mechanical circulatory support device (MCS) implantation. Dysregulation of the innate immune system is associated with increased patient age but has yet to be evaluated in the older patient with advanced heart failure undergoing MCS surgery.

Patients pre- and post-MCS implantation had peripheral blood mononuclear cells (PBMC) and serum isolated. Multiparameter flow cytometry was used to analyze markers of innate cell function, including monocyte subtypes. Multiplex cytokine analysis was performed. MELD-XI and SOFA scores were utilized as surrogate markers of outcomes.

Increased levels of pro-inflammatory cytokines including IL-15, TNF- α , and IL-10 were associated with increased MELD-XI and SOFA scores. IL-8, TNF- α , and IL-10 were associated with risk of death after MCS implantation, even with correction for patient age. Increased frequency of ‘classical’ monocytes (CD14 + CD16 –) were associated with increased MELD-XI and SOFA scores.

This suggests that inflammation and innate immune system activation contribute to progression to multiorgan system failure and death after MCS surgery. Development of noninvasive monitoring of peripheral blood holds promise for biomarker development for candidate selection and patient risk stratification.

1. Introduction

For the growing numbers of older patients with advanced heart failure, mechanical circulatory support device implantation can be an effective intervention for patients not manageable by medical therapy alone, either as a bridge to heart transplantation or as destination therapy [1–4]. However, older patients experience increased rates of death and inability to bridge to heart transplantation compared with younger patients with similar severity of heart failure [4]. Inflammation and pro-inflammatory changes in the innate immune system are known to be associated with normal aging, and may be part of the mechanism of progression of heart failure and atherosclerosis [5,6]. It has been

described that MCS implantation leads to inflammation and innate immune changes [7–11]. Given the many deleterious effects of inflammation, sometimes termed ‘inflammaging’ in the geriatrics literature [12,13], we propose that measurement of inflammation and innate immune changes before and after implantation will provide insight into the mechanism of development of adverse outcomes after MCS, and may provide help with patient risk stratification in combination with currently validated clinical tools such as INTERMACS score.

Our previous work has demonstrated increased frequency of immunosenescent and terminally differentiated CD8 + T cells in older patients and in patients with adverse outcomes after MCS implantation [14]. We have additionally observed an association between

* Corresponding author.

E-mail address: jschaenman@mednet.ucla.edu (J.M. Schaenman).

<https://doi.org/10.1016/j.humimm.2018.11.004>

Received 4 May 2018; Received in revised form 29 October 2018; Accepted 7 November 2018

Available online 14 November 2018

0198-8859/© 2018 Published by Elsevier Inc. on behalf of American Society for Histocompatibility and Immunogenetics.

clinical outcomes and changes in gene regulation in PBMC with important immune functions including T cell differentiation, KIR expression, and the TFG-beta receptor [15].

Development of additional tools for patient candidacy evaluation is especially important to consider given the observation that older and frailer patients are at increased risk for death after MCS D implantation [4,16–19]. The potential link between markers of inflammation including cytokines and monocyte subtypes and adverse outcomes after MCS D in younger and older patients has not been previously examined. Monocyte subtypes as defined by CD14 and CD16 expression include the classical (CD14+/CD16-), intermediate CD14+/CD16+), and non-classical CD14+/CD16++), of which the classical is most strongly associated with inflammation and response to innate immune system stimuli including pro-inflammatory cytokines [20,21]. We hypothesized that increased levels of pro-inflammatory cytokines and innate immune system dysregulation are associated with adverse clinical outcomes after MCS D implantation.

2. Methods

2.1. Patients and samples

We enrolled patients undergoing evaluation for MCS D from the Ronald Reagan Medical Center with advanced heart failure, as described previously [14]. This observational study was approved by the UCLA Institutional Review Board. All patients signed informed consent. Older patients were defined as those \geq age 60. Blood was collected for serum and peripheral blood mononuclear cell (PBMC) isolation within 24 h prior to MCS D implantation and on Days 1, 3, 5, 8, 10, 14 and 17, \pm 1 day after surgery. 27 patients were enrolled who had PBMC available after MCS D implantation for analysis, and completed at least 6 months of clinical follow-up between September 2012 and March 2015, with the last point of clinical review as of July 1, 2016. 24 patients had PBMC available prior to implantation. Our previous studies have confirmed the reproducibility of frozen and thawed samples for immunologic testing [14,15]. The following FDA-approved durable MCS D were included in our analysis: HeartMate II, HeartWare, Thoratec paracorporeal ventricular assist device, CentriMag, or Total Artificial Heart, but not percutaneously inserted devices such as extracorporeal membranous oxygenation, TandemHeart, or Impella pumps. PBMC were isolated using previously published techniques [22], and frozen for storage until batched analysis could be performed. The majority (75%) of samples were collected between Days 0 and 8, with 57% with 5 samples and 71% with \geq 5 samples available for analysis.

2.2. Flow cytometry

Viable cells were identified using a fluorescent live/dead marker (Life Technologies). Innate cell subsets were evaluated using a cocktail of fluorochrome-conjugated antibodies. CD14 and CD16 were used to define monocyte subsets as previously described [23]. CD56 and KLRG1 were utilized to define NK cells, and CD284 (Toll like receptor) to define possession of antibacterial properties and ability to recognize bacterial LPS [24]. The complete set of subtypes analyzed were as follows: CD14, CD56, CD14+/CD16-, CD14+/CD16+ +, CD14-/CD16+, CD14-/CD16-, CD14+/CD284+, CD14+/CD16-CD284+, CD14+/CD16+/CD284+, CD14-/CD16+/CD284+, CD14-/CD16-CD284+, CD14+/CD284+, CD14-/CD284+, CD14+/CD16+ + of monocytes, and CD14-/CD16+ of monocytes. Antibodies were obtained from either BD Biosciences or Biolegend. Fluorescence from viable cells was measured by the BD LSRFortessa (BD Biosciences) with analysis via FCS Express software (DeNovo Software).

2.3. Multiplex cytokine analysis

Human 38-plex magnetic cytokine/chemokine kits were purchased from EMD Millipore and used per manufacturer's instructions. The following analytes were detected: G-CSF, GM-CSF, IFN-gamma, IFN-a2, IL-1beta, IL-1Ra, IL-4, IL-5, IL-6, IL-8, IL-10, IL-12 (p40), IL-15, IL-17A, MCP-1, MIP-1alpha, MIP-1beta, CD40L, MDC, TNF-alpha, EGF, FGF-2, GRO, Eotaxin, Fractalkine, FLT-3L, and VEGF. Fluorescence was quantified using a Luminex 200™ instrument. Cytokine values measured in pg/ml.

2.4. Clinical data collection

Prospectively collected data was used to calculate MELD-XI (Model for End-Stage Liver Disease eXcluding INR) and SOFA (Sequential Organ Failure Assessment) scores, used as a surrogate for multiorgan dysfunction [25,26]. For MELD-XI calculation, an appropriate measure for patients on anticoagulation, serum creatinine was set to 1.0 for patients with creatinine levels of $<$ 1.0 to prevent calculation of negative numbers, generating a minimum score of 9.44, following previously published guidelines, and as described previously [14,26]. Records were reviewed for 3 month prior to and 6 months after MCS D implantation for evidence of infection, including sepsis syndrome, bacteremia, driveline infection, pneumonia, and urinary tract infection following standard definitions [27,28]. Severe infection was defined as requiring intravenous antibiotic treatment and/or leading to extension of hospital stay or death. Information on bypass time and peri-operative blood transfusion was not available for 3 patients (2 older and 1 younger). Median values were used to divide patients into "High" or "Low" MELD-XI and SOFA groups either prior to or after MCS D implantation. The median value and High/Low cutoff prior to MCS D implantation was 18; after MCS D implantation the median value and High/Low cutoff was 16. The median follow-up time was 474 days (IQR 91 to 779 days).

2.5. Statistical analysis

Individual comparisons between immune subtypes were performed as a screening analysis using JMP Pro 11 (SAS Software). Differences between continuous values (SOFA or MELD-XI, chronologic age, frequencies of immunologic subtypes) were compared by nonparametric 2-sample test (Mann-Whitney U-Test), while differences between categorical variables were compared by Fisher exact test. Standard least squares regression was used to compare numeric variables.

To correct for the issue of repeated measures and patient-to-patient variability, linear mixed effects models were used to evaluate the association between immune phenotype and clinical outcomes utilizing MELD-XI and SOFA scores, including random patient effects. Unadjusted associations between immune phenotypes and clinical outcomes were evaluated to identify candidate variables for inclusion into the multivariable regression models. Among those which were statistically significant, subsets were selected to limit the number of variables in each model based on a priori plausible clinical pathways [6,12,13]. Time to infection and death were analyzed using Cox proportional hazards models, with immune phenotypes included as time-varying covariates. Two models were constructed for data analysis: a static regression model, using actual variable values, and a dynamic predictor model, modeling each variable relative to its value at time 0 (MCS D implant). These analyses were performed using R v 3.3.2 (<http://www.r-project.org/>).

3. Results

3.1. Patient characteristics at time of MCS D implantation

Twenty-seven patients with advanced heart failure underwent

Table 1A
Demographic characteristics of all study participants (n = 27) % (n) for each variable.

Age (yrs)(median)(range)	61 (25–81)
Older (≥ age 60)	55.6% (15)
Sex (% male)	81.5% (22)
Nonischemic CMY	70.4% (19)
Intended bridge to transplantation	85.2% (23)
HeartMate II device	70.4% (20)
RVAD*	33.3% (9)
INTERMACS 1/2	59.3% (16)
MELD-XI at Day 0, median (range)	17.1 (9.4–28.8)
SOFA at Day 0, median (range)	7 (3–16)
Creatinine, mg/dl, median (range)	1.7 (1–3.7)
Total bilirubin, mg/dl, median (range)	1.6 (1–6.6)
Mean arterial pressure, median (range)	81 (54–107)

* RVAD includes right-sided Centrimag or ventricular support from PVAD or TAH.

Table 1B
Demographic characteristics of by Age ≥ 60 (Older) (n = 15) compared with patients < 60 (Younger) (n = 12) at time of implantation. *comparison not performed as attribute used to define groups.

Characteristic	Older	Younger	p-value
Age (yrs) (median) (range)	67.0 (61–81)	42.0 (26–59)	N/A*
Sex (% male)	93.3%	66.7%	0.139
Nonischemic CMY	53.3%	92.7%	0.043
Intended bridge to transplantation	58.3%	83.3%	0.371
HeartMate II device	80.0%	66.7%	0.327
RVAD*	20.0%	50.0%	0.127
INTERMACS 1/2	73.3%	26.7%	0.130
MELD-XI at Day 0, median (range)	20.1 (13.0–28.8)	13.7 (9.4–21.5)	0.006
SOFA at Day 0, median (range)	8 (4–16)	5.5 (3–13)	0.047
Creatinine, mg/dl, median (range)	2.0	1.4	0.008
Total bilirubin, mg/dl, median (range)	2.0	1.5	0.111
Mean arterial pressure, median (range)	81 (54–107)	081 (69–95)	0.864

testing enrolled in as described above (Table 1A). Patient age ranged from 25 to 81 years old. Primary indication for implantation was non-ischemic heart failure and the primary device utilized was HeartMate II. Median bypass time was 111 min in younger patients and 88.5 in older patients (p = 0.273). The number of PBMC transfused was 4 units in younger patients and 6 units in older patients (p = 0.371). Patients were divided into high (≥ 18) and low (< 18) MELD-XI group at the time of implant. Older patients (median age 67) were less likely to have nonischemic heart disease as the cause of heart failure, and as previously noted, older patients were more likely to have higher MELD-XI and SOFA scores at the time of MCS implantation compared with younger patients (median age 42) [14] (Table 1B). INTERMACS scores tended to be more severe in older compared with younger patients, although this did not reach statistical significance.

3.2. Patient outcomes after MCS implantation

Infectious complications were observed both prior to and after MCS implantation (Table 2A). These included bacterial infections such as pneumonia and sepsis, candida infection, and viral pneumonia. 22.2% of all patients (n = 6) died by 3 months after implant. Of the 9 patients who died by one year post implant, the majority (n = 9) died due to multiorgan failure, often with sepsis. We observed a statistically significant association between older patient age and death 3 months after MCS implantation (p = 0.020). Older patients demonstrated increased incidence of infection and severe infection in this cohort (Table 2B). There was no significant difference in bypass time in patients with or without severe infection (107 versus 101 min, p = 0.977), or in PBMC transfused (5 versus 6 units, p = 0.975).

Table 2A
Clinical outcomes of study participants (n = 27) % (n) for each variable.

Infection pre- or post-implantation	77.8% (21)
Severe infection post-implantation	63.0% (17)
Successful bridge to transplant*	73.9% (17)*
Death 30 days	11.1% (3)
Death 3 months	22.2% (6)
Death 1 year	33.3% (9)
Days to death (median) (range)	51 (18–736)
Infection post-MCSD (≤ 6 months)	61.5% (16)

* Excluding 4 patients with MCS implant as destination therapy.

Table 2B
Clinical outcomes of study participants (n = 27) by Age ≥ 60 (Older) (n = 15) compared with patients < 60 (Younger) (n = 12) after implantation % (n) for each variable.

Characteristic	Older	Younger	p-value
Infection pre- or post-implantation	86.9%	60.5%	0.003
Severe infection post-implantation	75.4%	44.2%	0.002
Successful bridge to transplant*	58.3%	90.9%	0.156
Death 30 days	18.0%	0.0%	0.002
Death 3 months	39.3%	0.0%	< 0.001
Death 1 year	39.3%	25.6%	0.206
Days to death (median) (range)	32 (18–77)	137 (134–736)	0.037
Infection post-MCSD (≤ 6 months)	60.0% (9)	58.3% (7)	NS

* Excluding 4 patients with MCS implant as destination therapy.

3.3. Monocyte characteristics and patient age

Older patients demonstrated a trend towards increased frequency of classical CD14+ + CD16- monocytes prior to MCS implant as compared with younger patients, with a median frequency of 31.5% as compared with 14.0% (p = 0.224). There were no statistically significant differences prior to MCS implant for intermediate CD14 + CD16 + monocytes (2.2% compared with 2.0%, p = 0.385) or non-classical CD14-CD16++ monocytes (2.0% compared with 2.2%, p = 0.505) between older and younger patients, respectively.

After MCS implantation (excluding pre-implant values), differences were observed in monocyte subtypes between older and younger patients, with increased frequency of classical CD14+ + CD16- monocytes as compared with younger patients, with a median frequency of 28.7% as compared with 15.5% (p = 0.001) (Fig. 1). There were also statistically significant differences observed for intermediate CD14 + CD16 + monocytes (2.5% compared with 0.91%, p = 0.010) and nonclassical CD14-CD16++ monocytes (1.9% compared with 1.2%, p = 0.009) between older and younger patients, respectively. Median values for older and younger patients are shown in Supplementary Tables 1A and 1B.

Differences in monocyte subtypes did not vary by INTERMACS score, measured either before or after MCS implantation (data not shown).

3.4. TLR4+ expression and patient age

We also noted differences in TLR4+ (CD284) expression, a surface molecule important in innate immune function, which binds bacterial LPS. Older patients demonstrated decreased frequency of monocytes expressing this surface protein, with median frequency of TLR4 + expression of 5.3% for older patients compared with 48.6% in younger patients after MCS implantation (p < 0.001) (Fig. 2A). This observation was interesting given that older patients displayed an increased percentage of total monocytes (48.6% compared with 29.1% in younger patients, p < 0.001) (Fig. 2A).

Repeating this analysis by monocyte subtypes revealed similar results, with older patients demonstrating decreased frequency of TLR4 + expression in classical (2.4% compared with 48.6% in younger




Fig. 1. Frequency of monocyte subtypes by patient age. PBMC from time points after MCS D implantation were analyzed for classical (CD14 + +/CD16 –), intermediate (CD14 + /CD16 +), and nonclassical (CD14 –/CD16 +) monocytes, expressed as a percentage of the total number of PBMC. Each dot or triangle corresponds to a sample; bars indicate median. p-values as indicated by nonparametric testing.

patients, $p < 0.001$), intermediate (33.0% compared with 86.8% in younger patients, $p < 0.001$), and nonclassical monocytes (14.8% compared with 54.0% in younger patients, $p < 0.001$) (Fig. 2B). Median values for older and younger patients are shown in Supplementary Tables 1A and 1B.

3.5. Plasma cytokines and patient age

Analysis of cytokine expression after MCS D implantation revealed multiple pro-inflammatory cytokines associated with the older patient. These include TNF- α ($p < 0.001$), IL-10 ($p = 0.002$), IL-15 ($p < 0.001$), TGF- α ($p = 0.013$), IL-8 ($p < 0.001$), MDC ($p = 0.006$), IP-10 ($p = 0.010$), and MCP-1 ($p < 0.001$) (Table 3). Increased levels of many of the pro-inflammatory cytokines were seen across all older patients regardless of progression to severe infection. However, we found that MCP-1 was uniquely elevated only in older patients with severe infection (median 638), while for older patients without severe

infection, it was significantly lower, similar to levels seen in younger patients (median 465) ($p = 0.024$). Similarly, there were several cytokines that were significantly elevated in younger patients with severe infection but lower in younger patients without severe infection including TNF- α ($p < 0.001$), IL-8 ($p = 0.009$), and IP-10 ($p = 0.018$). These cytokines are also analyzed under Multivariate Analysis below demonstrating the independent impact of age as well as levels of proinflammatory cytokines in predicting outcomes after MCS D implantation (Tables 4 and 5).

3.6. Innate immunity and MELD-XI/SOFA score

After MCS D implantation, an association between MELD-XI score and monocyte phenotype was observed: Categorizing each MELD-XI score as ‘high’ (≥ 16) or ‘low’ (< 16) (divided by level post-implant) revealed higher frequencies of classical CD14 + +CD16 – monocytes with high MELD-XI (31.4%) as compared with low MELD-XI (19.4%)




Fig. 2A. Frequency of CD14 + and CD14 + CD284 + (TLR4 +) monocytes by patient age. PBMC from time points after MCS D implantation were analyzed, expressed as a percentage of the total number of PBMC or CD14 + cells, as indicated. Each dot or triangle corresponds to a sample; bars indicate median. p-values as indicated by nonparametric testing.

B




Fig. 2B. Frequency of CD14 + CD284 + (TLR4 +) monocyte subtypes by patient age. PBMC from time points after MCS D implantation were analyzed for percentage of CD284 + expression within the classical (CD14 + /CD16 –), intermediate (CD14 + /CD16 +), and nonclassical (CD14 – /CD16 +) monocyte subtypes. Each dot or triangle corresponds to a sample; bars indicate median. p-values as indicated by nonparametric testing.

Table 3

Median cytokine values in older (n = 15) as compared with younger patients (n = 12) after MCS D implantation. P values < 0.05 in bold.

Cytokine	Median (young)	Median (old)	p-value
IL-12(p40)	13.3	9.4	0.201
IL-12(p70)	11.9	13.5	0.855
IFN-g	67.5	51.2	0.857
TNF-a	20.8	31.1	< 0.001
TNF-b	5.8	12.5	0.042
IL-4	3.5	6.7	0.041
IL-5	2.7	3.4	0.103
IL-9	2.9	3.0	0.018
IL-10	16.6	28.2	0.002
IL-13	2.4	4.1	0.041
IL-17A	15.3	12.7	0.564
IL-1a	8.2	23.4	0.062
IL-1b	2.7	2.7	0.445
IL-2	2.9	2.4	0.853
IL-3	3.0	3.0	0.831
IL-6	36.9	58.2	0.054
IL-15	4.8	10.3	< 0.001
TGF-a	4.5	6.1	0.013
IFN-a2	41.8	46.4	0.686
IL-8	39.9	61.8	< 0.001
GRO	339.6	334.0	0.596
Eotaxin	120.8	170.5	< 0.001
MDC	579.0	436.5	0.006
IP-10	605.0	809.5	0.010
MCP-1	380.8	587.0	< 0.001
MCP-3	21.0	25.6	0.147
Fractalkine	78.0	90.9	0.593
MIP-1a	6.3	8.2	0.458
MIP-1b	32.1	44.9	< 0.001
GM-CSF	16.4	20.5	0.083
IL-7	5.3	5.9	0.493
G-CSF	61.9	76.8	0.135
VEGF	266.0	242.0	0.365
EGF	32.2	26.5	0.493
FGF-2	99.9	122.0	0.098
Flt-3L	2.5	2.5	0.399
IL-1RA	94.0	124.5	0.014
sCD40L	1113.0	1279.5	0.339

(p < 0.001) (Fig. 3). There was also an increased frequency of intermediate CD14 + CD16 + monocytes with high MELD-XI (3.2%) compared with low MELD-XI (1.1%) (p < 0.001) and nonclassical CD14-CD16 + monocytes (2.0% compared with 1.6%, p = 0.036) (Fig. 3).

Similarly, higher frequencies of classical CD14 + /CD16 – monocytes were observed with high SOFA (29.2%) as compared with low SOFA scores (< 6) (19.7%) (p = 0.006). There was also an increased frequency of intermediate CD14 + /CD16 + monocytes with high SOFA (2.6%) compared with low SOFA (1.1%) (p = 0.012). However, there was no significant difference between nonclassical CD14-CD16 + monocytes and SOFA score (1.8% compared with 2.0%, p = 0.686).

Analysis of TLR4 + expression on monocytes did not demonstrate an association between MELD-XI score, although there was an increased frequency of monocytes observed in the high MELD-XI group (48.6% compared with 34.5%, p = 0.010). (Fig. 4). A similar pattern was observed for SOFA score, with no association by TLR + expression but an association with high SOFA score and increased frequency of monocytes (47.7% versus 35.4%, p = 0.007). There was however, no significant association between MELD-XI or SOFA score in non-monocyte TLR4 + cells (data not shown).

3.7. Innate immunity and clinical outcomes

Frequency of monocyte subtypes was not significantly associated with infection after MCS D implantation (data not shown). Using this nonparametric analysis approach, there was similarly no observed association between monocyte subtypes and death or successful bridge to transplant after MCS D implantation (data not shown).

For TLR4 + expression, a significant association was observed between infection and decreased expression of TLR4 on monocytes, with 6.5% in those with infection compared with 34.4% in those without (p = 0.012). This association was also seen for non-monocyte cells, with 4.5% frequency of CD14 – TLR4 + cells in those with infection compared with 29.4% in those without (p = 0.005). In contrast, the expression of total monocytes was not different between groups (45.5% compared with 42.3%, p = 0.818). An association was also observed between TLR4 + monocytes subtypes, with patients with infection demonstrating decreased frequency of TLR4 + expression in classical (5.0% compared with 37.8%, p = 0.009), intermediate (37.2% compared with 88.8%, p = 0.008), and nonclassical monocytes (18.1% compared with 53.8%, p = 0.024) compared with patients without infection.

However, no association was observed between frequency of total monocytes or monocytes expressing TLR4 + and death or successful bridge to transplant by nonparametric analysis (data not shown).

Table 4
Static regression models. Only variables with p < 0.05 in preliminary analyses included.

Effect	MELD-XI		SOFA		Time to infection		Time to death	
	Diff. (95% CI)	p	Diff. (95% CI)	p	Diff. (95% CI)	p	Diff. (95% CI)	p
Age	0.08 (-0.02–0.19)	0.120	0.01 (-0.04–0.05)	0.809	0.99 (0.96–1.02)	0.564	1.02 (0.97–1.08)	0.423
EGF	ND		ND		0.99 (0.96–1.01)	0.200	ND	
TNF-α	0.03 (-0.01–0.08)	0.120	ND		ND		1.02 (0.98–1.07)	0.383
IL-8	ND		0.03 (0.02–0.04)	< 0.001	ND		1.01 (1.00–1.03)	0.128
IL-10	0.00 (0.00–0.01)	0.085	0.00 (0.00–0.01)	0.176	ND		ND	
IL-15	0.26 (0.08–0.44)	0.005	ND		ND		ND	
MCP-1	ND		0.00 (0.00–0.00)	0.015	ND		ND	
MDC	ND		ND		1.00 (1.00–1.00)	0.223	ND	
GCSF	0.00 (-0.01–0.01)	0.530	ND		ND		ND	
CD14 + CD16- (classical monocytes)	0.08 (0.03–0.13)	0.001	0.06 (0.02–0.09)	0.001	ND		ND	
CD14-CD284+	ND		ND		ND		1.03 (1.00–1.06)	0.023

Diff.: Difference; ND, not done.

3.8. Monocyte subtypes and T cell immunosenescence

Our previous studies demonstrated an association between T cell immunosenescence and adverse clinical outcomes after MCS D implantation [14]. We found that the intermediate monocyte subtype CD14 + CD16+ was significantly associated with CD8+ T cell immunosenescent or exhaustion subtypes including KLRG1+ (p = 0.002), KLRG1+/CD38+ (p < 0.001), and KLRG1 + /PD1+ (p = 0.006). There was no association, however, between CD8+ T cell immunosenescence or exhaustion and expression of TLR4 (data not shown.)

3.9. Multivariable analysis combining demographic, innate immunity

To address the issue of repeated measures and individual patient effects on the association between immune phenotype and clinical outcomes, a linear mixed effects model was used. This allows for correction for random effects at the individual patient level and provides flexibility to model varying slopes and intercepts, providing an ideal analysis approach for sequential observations across a group of patients. In addition, we used this approach to assess the predictive impact of multiplex cytokine analysis, which are less amenable to individual analysis due to issues with repeated measures. Multivariable mixed effect models predicting MELD-XI score revealed a statistically significant association between one pro-inflammatory cytokine, IL-15, (p = 0.005) and the pro-inflammatory classical monocyte subtype

(p = 0.001) (Table 3). Interestingly, for SOFA score, the classical monocyte subtype was again significantly associated (p = 0.001), but two different pro-inflammatory cytokines were identified, IL-8 (p < 0.001) and MCP-1 (p = 0.015), although the effect size for MCP-1 was relatively small (Table 3).

Multivariable analysis of time to infection after MCS D implantation did not identify any significant associations, although in unadjusted analysis a trend was observed for age and the EGF and MDC cytokines. For prediction of time to death, the TLR4+ expressing non-monocytes population demonstrated a statistically significant association (p = 0.023), while a trend towards association was observed for age and the TNF-α and IL-8 cytokines (Table 3).

In order to analyze how the change in levels of the immunologic predictors predicts the change in the level of the outcomes measured, we additionally performed a dynamic analysis using a similar mixed model approach to correct for repeated measurements. Analyses were repeated with values expressed as change from baseline to attempt to better capture the dynamic nature of changes in the innate immune system and cytokine expression after MCS D implantation. Dynamic multivariable mixed effect models predicting MELD-XI score revealed a statistically significant association between change in IL-8, over time (p < 0.001) and trend towards significance for change in GCSF (p = 0.067) and IL-5 (p = 0.069) (Table 4). Interestingly, for SOFA score, change of the classical monocyte subtype over time was significantly associated with change in SOFA score (p = 0.021) as seen in the static models (Table 4). There were also several pro-inflammatory

Table 5
Fully dynamic models based on variable change from baseline. Only variables with p < 0.05 in preliminary analyses included.

Effect	MELD-XI		SOFA	
	Diff. (95% CI)	p	Diff. (95% CI)	p
Age	0.03 (-0.12–0.17)	0.712	0.01 (-0.10–0.12)	0.865
EGF	0.04 (-0.01–0.10)	0.123	ND	
TGF	ND		-0.14 (-0.32–0.03)	0.111
TNF-α	-0.07 (-0.18–0.04)	0.209	-0.02 (-0.11–0.07)	0.628
IL-1 α	-0.01 (-0.03–0.00)	0.154	ND	
IL-5	0.16 (-0.01–0.34)	0.069	ND	
IL-6	-0.01 (-0.02–0.01)	0.393	0.02 (0.00–0.03)	0.019
IL-8	0.07 (0.03–0.10)	< 0.001	0.02 (0.00–0.05)	0.098
IL-10	0.00 (0.00–0.01)	0.973	0.00 (0.00–0.01)	0.440
IL-15	0.17 (-0.11–0.46)	0.239	0.26 (0.01–0.48)	0.021
MCP-1	ND		0.00 (0.00–0.00)	0.482
MIP-1 α	0.02 (-0.05–0.08)	0.615	0.03 (-0.02–0.08)	0.224
MIP-1β	-0.02 (-0.08–0.03)	0.395	ND	
MDC	ND		-0.01 (-0.01–0.00)	0.027
GCSF	-0.01 (-0.03–0.00)	0.067	0.00 (-0.19–0.16)	0.590
CD14 + CD16- (classical monocytes)	0.03 (-0.03–0.10)	0.296	0.06 (0.01–0.12)	0.021
CD14-CD16+ of monocytes (nonclassical monocytes)	-0.13 (-0.34–0.09)	0.247	-0.01 (-0.19–0.16)	0.877

Diff.: Difference; ND, not done.




Fig. 3. Frequency of monocyte subtypes by patient high or low MELD-XI score. PBMC from time points after MCS D implantation were analyzed for classical (CD14++/CD16–), intermediate (CD14+/CD16+), and nonclassical (CD14–/CD16+) monocytes, expressed as a percentage of the total number of PBMC. Each dot or triangle corresponds to a sample; bars indicate median. p-values as indicated by nonparametric testing.

cytokines that were identified in the dynamic model: IL-6 (p = 0.019), IL-15 (p = 0.021), and MDC (p = 0.027) (Table 4). A trend towards statistical significance was also seen for IL-8 (p = 0.098).

4. Discussion

Inflammation is known to play a role in adverse outcomes in older patients, however, less is known about the role of pro-inflammatory monocytes and cytokines in the setting of MCS D implantation. Previously assessed markers such as C-reactive protein and procalcitonin have not added significantly to prediction of adverse outcomes after MCS D, suggesting the need for more immunologically oriented assessment [29,30]. In this analysis of MCS D recipients, we found that despite similar etiologies of heart failure and types of interventions, older patients demonstrated increased frequency of the pro-inflammatory CD14++/CD16– classical monocyte. In contrast, older patients had decreased frequency of TLR4+ monocytes compared with younger patients, which may explain the mechanism of increased

frequency of sepsis and death in older patients after MCS D implantation. Increased frequency of the pro-inflammatory CD14++/CD16– classical monocyte was also correlated with clinical outcomes as measured by MELD-XI and SOFA scores. Mixed effect analysis correcting for repeated measures further supported the association between inflammation and adverse clinical outcomes, with statistically significant association between monocyte subtype and expression of pro-inflammatory cytokines (Table 3). Repeating this analysis as dynamic change in immune phenotype and cytokines revealed a similar pattern of association between inflammation and adverse clinical outcomes as represented by increased MELD-XI and SOFA scores (Table 4). This dynamic analysis suggests that it is not just pre-implant inflammation related to advanced heart failure that leads to adverse outcomes, or the immediate post-surgical stress, but rather that ongoing or worsening inflammation occurring after MCS D implantation drives adverse outcomes as measured by increased MELD-XI and SOFA scores, and ultimately leading to multiorgan system failure and patient death.

These findings demonstrate that patient age is associated with




Fig. 4. Frequency of CD14+ and CD14+CD284+ (TLR4+) monocytes by high or low MELD-XI score. PBMC from time points after MCS D implantation were analyzed, expressed as a percentage of the total number of PBMC, as indicated. Each dot or triangle corresponds to a sample; bars indicate median. p-values as indicated by nonparametric testing.

differences in innate immune phenotype and cytokine expression, revealing a mechanism for the increased rates of adverse outcomes including sepsis and multiorgan dysfunction observed in older patients. This immune profile is an important predictor of adverse outcomes independent of patient age in both the static and dynamic models (Tables 3 and 4), suggesting the potential benefit of immune profiling in conjunction with currently utilized methods of patient risk profiling such as INTERMACS score, as suggested in our previous analysis of T cell immune senescence and change in gene expression and clinical outcomes after MCS implantation [14,15]. Although there is necessarily some overlap between innate immune system changes seen in all older patients and those seen in patients who experience adverse clinical outcomes, our analyses demonstrate that several cytokines such as IL-8 and MCP-1 may possess the ability to discriminate between patients with typical age-associated immunologic changes and those at increased risk for infection or death. Validation of this approach could lead to a model of noninvasive testing of patients with advanced heart failure to predict outcomes after MCS placement and allow for non-invasive post-operative monitoring of patients after device implantation in both biologically and chronologically older patients.

Our previous work demonstrated a significant association between immunosenescence and increased patient age as well as adverse clinical outcomes including increased MELD-XI and SOFA scores, infection, and death [31]. Interestingly we found an association between immunosenescence and the intermediate monocyte subtype (CD14+/CD16+), suggesting a connection between the adaptive and innate immune system that may have implications both for both understanding mechanism of immune dysfunction after MCS and to increase the number of markers available for patient risk stratification as described above.

This pilot study demonstrates the promise of immunologic assessment in patients undergoing MCS implantation. Study limitations include the relatively small sample size and diversity of devices utilized. In addition, as there were relatively few patients experiencing death during the period of study observation, we were not able to establish a strong association between markers of inflammation or lack of TLR4 expression and death. The large overlap between older patients and those experiencing infectious complications makes it difficult to separate out the potentially differing impacts of these two clinical variables in a relatively small cohort, especially given that both increased age and infection favor a pro-inflammatory immunologic milieu. However, we did observe that younger patients who experienced infection demonstrated patterns of cytokine expression similar to that of older patients, suggesting the utility of this approach in identifying at risk patients. In addition, the number of patients included is similar to other studies of immune dysfunction after MCS implantation [7,8,10]. As a single center study, we are able to include patients receiving similar medical care with similar waiting time for transplantation. In addition, our work confirms previous studies that have also identified IL-6, IL-8, TNF- α , and MDC as associated with heart failure and MCS implantation [6,29,32,33]. Future multicenter studies will be important to validate our pilot findings and evaluate whether pre-implant as well as post-implant assessment is associated with clinical outcomes.

For the growing numbers of older patients with advanced heart disease, MCS remains an important intervention for either destination therapy or as a bridge to heart transplantation. Our finding that older patients demonstrate increased proinflammatory monocyte subtypes and decreased expression of TLR4, an important surface marker of ability to control bacterial infections suggests a possible mechanism for the increased frequency of adverse outcomes in older MCS recipients. In addition, the association between proinflammatory monocyte subtypes and proinflammatory cytokine expression in patients with increases in the clinical markers MELD-XI and SOFA, suggests that increased inflammation is associated with adverse clinical outcomes including multiorgan immune dysfunction and death after MCS implantation. Noninvasive assessment of innate immune phenotype and

cytokine expression may prove to be an important tool for patient risk stratification and identification of risk factors for progression to adverse clinical complications in older patients with advanced heart disease.

Funding information

This research was supported through R21 HL 120040-01A1 NHLB (M.D.) and the NIH National Center for Advancing Translational Science (NCATS) UCLA CTSI Grant Number UL1TR001881 (M.K.).

Acknowledgement

We appreciate the contributions of laboratory scientist Fadi Kandarian to this work.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.humimm.2018.11.004>.

References

- [1] J.K. Kirklin, D.C. Naftel, F.D. Pagani, R.L. Kormos, L.W. Stevenson, E.D. Blume, et al., Seventh INTERMACS annual report: 15,000 patients and counting, *J. Heart Lung Transplant.* 34 (2015) 1495–1504, <https://doi.org/10.1016/j.healun.2015.10.003>.
- [2] L.W. Miller, M. Guglin, Patient selection for ventricular assist devices, *Jac* 61 (2013) 1209–1221, <https://doi.org/10.1016/j.jacc.2012.08.1029>.
- [3] D. Mozaffarian, E.J. Benjamin, A.S. Go, D.K. Arnett, M.J. Blaha, M. Cushman, et al., Heart disease and stroke statistics-2016 update: a report from the American Heart Association, *Circulation* 133 (2016) e38–60, <https://doi.org/10.1161/CIR.0000000000000350>.
- [4] J.K. Kirklin, F.D. Pagani, R.L. Kormos, L.W. Stevenson, E.D. Blume, S.L. Myers, et al., Eighth annual INTERMACS report: special focus on framing the impact of adverse events, *J. Heart Lung Transplant.* 36 (2017) 1080–1086, <https://doi.org/10.1016/j.healun.2017.07.005>.
- [5] D. Weiskopf, B. Weinberger, B. Grubeck-Loebenstien, The aging of the immune system, *Transplant Int.* 22 (2009) 1041–1050, <https://doi.org/10.1111/j.1432-2277.2009.00927.x>.
- [6] M.A. Moro-García, A. Echeverría, M.C. Galán-Artínez, F.M. Suárez-García, J.J. Solano-Jaurieta, P. Avanzas-Fernández, et al., Immunosenescence and inflammation characterize chronic heart failure patients with more advanced disease, *Int. J. Cardiol.* 174 (2014) 590–599, <https://doi.org/10.1016/j.ijcard.2014.04.128>.
- [7] S. Itescu, J.-H. Ankersmit, A.A. Kocher, M.D. Schuster, Immunobiology of left ventricular assist devices, *Prog. Cardiovasc. Dis.* 43 (2000) 67–80, <https://doi.org/10.1053/pcad.2000.7191>.
- [8] M. Rothenburger, M. Wilhelm, D. Hammel, C. Schmid, G. Plenz, T.D. Tjan, et al., Immune response in the early postoperative period after implantation of a left-ventricular assist device system, *Tps* 33 (2001) 1955–1957.
- [9] H.J. Ankersmit, S. Tugulea, T. Spanier, A.D. Weinberg, J.H. Artrip, E.M. Burke, et al., Activation-induced T-cell death and immune dysfunction after implantation of left-ventricular assist device, *Lancet* 354 (1999) 550–555.
- [10] P.M. Kimball, M. Flattery, F. McDougan, V. Kasirajan, Cellular immunity impaired among patients on left ventricular assist device for 6 months, *Ann. Thoracic Surg.* 85 (2008) 1656–1661, <https://doi.org/10.1016/j.athoracsur.2008.01.050>.
- [11] L. Grosman-Rimon, F. Billia, A. Fuks, I. Jacobs, M.A. McDonald, D.Z. Cherney, et al., New therapy, new challenges: the effects of long-term continuous flow left ventricular assist device on inflammation, *Int. J. Cardiol.* 215 (2016) 424–430, <https://doi.org/10.1016/j.ijcard.2016.04.133>.
- [12] C. Franceschi, J. Campisi, Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases, *J. Gerontol. A: Biol. Sci. Med. Sci.* 69 (2014) S4–S9, <https://doi.org/10.1093/geron/glu057>.
- [13] D. Frasca, B.B. Blomberg, Inflammaging decreases adaptive and innate immune responses in mice and humans, *Biogerontology* (2015) 1–13, <https://doi.org/10.1007/s10522-015-9578-8>.
- [14] J.M. Schaanman, M. Rossetti, Y. Korin, T. Sidwell, V. Groysberg, E. Liang, et al., T cell dysfunction and patient age are associated with poor outcomes after mechanical circulatory support device implantation, *Hum. Immunol.* (2018), <https://doi.org/10.1016/j.humimm.2018.01.011>.
- [15] G. Bondar, R. Togashi, M. Cadeiras, J. Schaanman, R.K. Cheng, L. Masukawa, et al., Association between preoperative peripheral blood mononuclear cell gene expression profiles, early postoperative organ function recovery potential and long-term survival in advanced heart failure patients undergoing mechanical circulatory support, *PLoS One* 12 (2017) e0189420, <https://doi.org/10.1371/journal.pone.0189420>.
- [16] W.L. Holman, R.L. Kormos, D.C. Naftel, M.A. Miller, F.D. Pagani, E. Blume, et al., Predictors of death and transplant in patients with a mechanical circulatory support device: a multi-institutional study, *J. Heart Lung Transplant.* 28 (2009) 44–50, <https://doi.org/10.1016/j.healun.2008.10.011>.

- [17] S.M. Dunlay, S.J. Park, L.D. Joyce, R.C. Daly, J.M. Stulak, S.M. McNallan, et al., Frailty and outcomes after implantation of left ventricular assist device as destination therapy, *J. Heart Lung Transplant.* 33 (2014) 359–365, <https://doi.org/10.1016/j.healun.2013.12.014>.
- [18] S.M. Joseph, J.L. Manghelli, J.M. Vader, T. Keeney, E.L. Novak, J. Felius, et al., Prospective assessment of frailty using the fried criteria in patients undergoing left ventricular assist device therapy, *Am. J. Cardiol.* 120 (2017) 1349–1354, <https://doi.org/10.1016/j.amjcard.2017.07.074>.
- [19] S.R. Jha, M.K. Hannu, P.J. Newton, K. Wilhelm, C.S. Hayward, A. Jabbour, et al., Reversibility of frailty after bridge-to-transplant ventricular assist device implantation or heart transplantation, *Transplant. Direct.* 3 (2017) e167, <https://doi.org/10.1097/TXD.0000000000000690>.
- [20] L. Ziegler-Heitbrock, P. Ancuta, S. Crowe, M. Dalod, V. Grau, D.N. Hart, et al., Nomenclature of monocytes and dendritic cells in blood, *Blood* 116 (2010) e74–e80, <https://doi.org/10.1182/blood-2010-02-258558>.
- [21] Y. Oishi, I. Manabe, Macrophages in age-related chronic inflammatory diseases, *NPJ Aging Mech Dis.* 2 (2016) 16018, <https://doi.org/10.1038/npjamd.2016.18>.
- [22] G. Freer, L. Rindi, Intracellular cytokine detection by fluorescence-activated flow cytometry: basic principles and recent advances, *Methods* 61 (2013) 30–38, <https://doi.org/10.1016/j.ymeth.2013.03.035>.
- [23] E.C. Liang, M. Rossetti, T. Sidwell, V. Groysberg, G. Sunga, Y. Korin, et al., Differences in proinflammatory cytokines and monocyte subtypes in older as compared with younger kidney transplant recipients, *Transplant. Direct.* 1 (2018), <https://doi.org/10.1097/TXD.0000000000000762>.
- [24] B.S. Park, J.-O. Lee, Recognition of lipopolysaccharide pattern by TLR4 complexes, *Exp. Mol. Med.* 45 (2013) e66, <https://doi.org/10.1038/emm.2013.97>.
- [25] L. Minne, A. Abu-Hanna, E. de Jonge, Evaluation of SOFA-based models for predicting mortality in the ICU: a systematic review, *Crit Care.* 12 (2009) R161, <https://doi.org/10.1186/cc7160>.
- [26] J.C. Grimm, A.S. Shah, J.T. Magruder, A. Kilic, V. Valero, S.P. Dungan, et al., MELD-XI score predicts early mortality in patients after heart transplantation, *Ann. Thoracic Surg.* 100 (2015) 1737–1743, <https://doi.org/10.1016/j.athoracsur.2015.07.026>.
- [27] S. Husain, M.L. Mooney, L. Danziger-Isakov, F. Mattner, N. Singh, R. Avery, et al., A 2010 working formulation for the standardization of definitions of infections in cardiothoracic transplant recipients, *J. Heart Lung Transplant.* 30 (2011) 361–374, <https://doi.org/10.1016/j.healun.2011.01.701>.
- [28] M.M. Hannan, S. Husain, F. Mattner, L. Danziger-Isakov, R.J. Drew, G.R. Corey, et al., Working formulation for the standardization of definitions of infections in patients using ventricular assist devices, *J. Heart Lung Transplant.* 30 (2011) 375–384, <https://doi.org/10.1016/j.healun.2011.01.717>.
- [29] R. Caruso, J. Campolo, A. Verde, L. Botta, L. Cozzi, M. Parolini, et al., Research article relationship between early inflammatory response and clinical evolution of the severe multiorgan failure in mechanical circulatory support-treated patients, *Mediat. Inflamm.* (2014) 1–9, <https://doi.org/10.1155/2014/281790>.
- [30] T. Ahmad, T. Wang, E.C. O'Brien, M.D. Samsky, J.A. Pura, Y. Likhnygina, et al., Effects of left ventricular assist device support on biomarkers of cardiovascular stress, fibrosis, fluid homeostasis, inflammation, and renal injury, *JACC: Heart Fail.* 3 (2015) 30–39, <https://doi.org/10.1016/j.jchf.2014.06.013>.
- [31] J.M. Schaeffer, M. Rossetti, T. Sidwell, V. Groysberg, G. Sunga, Y. Korin, et al., Increased T cell immunosenescence and accelerated maturation phenotypes in older kidney transplant recipients, *Hum. Immunol.* (2018), <https://doi.org/10.1016/j.humimm.2018.06.006>.
- [32] R. Caruso, L. Botta, A. Verde, F. Milazzo, I. Vecchi, M.G. Trivella, et al., Relationship between pre-implant interleukin-6 levels, inflammatory response, and early outcome in patients supported by left ventricular assist device: a prospective study, *PLoS One* 9 (2014) e90802, <https://doi.org/10.1371/journal.pone.0090802.t003>.
- [33] L. Grosman-Rimon, I. Jacobs, L.C. Tumiati, M.A. McDonald, S.P. Bar-Ziv, A. Fuks, et al., Longitudinal assessment of inflammation in recipients of continuous-flow left ventricular assist devices, *Can. J. Cardiol.* 31 (2015) 348–356, <https://doi.org/10.1016/j.cjca.2014.12.006>.