



Original contribution

Validation of the International Tumor Budding Consensus Conference 2016 recommendations on tumor budding in stage I-IV colorectal cancer[☆]



Heather Dawson MD^{a,*},¹, Francesca Galuppini MD^{b,1}, Peter Träger^c,
Martin D. Berger MD^d, Peter Studer MD, PhD^e, Lukas Brügger MD^e, Inti Zlobec PhD^a,
Daniel Inderbitzin MD^f, Alessandro Lugli MD^a

^aInstitute of Pathology, University of Bern, Bern, 3008 Switzerland

^bPathology Unit, Department of Medicine (DIMED), University of Padova, Padova, 35121 Italy

^cCareanaesth AG, Zurich, 3006 Switzerland

^dDepartment of Medical Oncology, Inselspital, Bern University Hospital, University of Bern, Bern, 3010 Switzerland

^eClinic of Visceral Surgery, Inselspital, Bern University Hospital, University of Bern, Bern, 3010 Switzerland

^fDepartment of General Surgery and Visceral Surgery, Bürgerspital Solothurn, Solothurn, 4500 Switzerland

Received 23 August 2018; revised 15 October 2018; accepted 18 October 2018

Keywords:

Tumor budding;
Colorectal cancer;
Consensus conference;
Validation study;
Prognostic factor

Summary Tumor budding is a robust prognostic parameter in colorectal cancer and can be used as an additional factor to guide patient management. Although backed by large bodies of data, a standardized scoring method is essential for integrating tumor budding in reporting protocols. The International Tumor Budding Consensus Conference (ITBCC) 2016 has proposed such a scoring system. The aim of this study is to validate the ITBCC method of tumor budding assessment on a well-characterized colorectal cancer cohort. Three hundred seventy-nine patients with resected stage I-IV colorectal cancer were entered into the study. Tumor budding was scored by 2 pathologists according to the ITBCC recommendations on hematoxylin and eosin-stained slides and scored as BD1 (low grade), BD2 (intermediate grade), and BD3 (high grade). Analysis was performed using a 3-tier approach, a 2-tier approach (BD1 + 2 versus BD3) and budding as a continuous variable. High-grade tumor budding was associated with adverse clinicopathological features including higher pT, higher pN stage, and higher TNM stage (all $P < .001$) and poorer overall survival on univariate analysis ($P = .0251$ for BD1/2/3, $P = .0106$ for BD1 + 2 versus BD3, and $P = .0195$ for continuous scores; hazard ratio, 1.023 [95% confidence interval, 1.004-1.043 per bud]). In stage II cancers, BD3 was associated with poorer disease-free survival ($P < .01$). Tumor budding assessed by the method proposed by the ITBCC is applicable to colorectal cancer resection specimens and can be used for widespread reporting in routine.

© 2018 Elsevier Inc. All rights reserved.

[☆] Disclosures: The authors report no competing interest. This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

* Corresponding author at: Institute of Pathology, University of Bern, Murtenstrasse 31, PO Box 62, CH-3008 Bern, Switzerland.

E-mail address: heather.dawson@pathology.unibe.ch (H. Dawson).

¹ Equally contributing first authors.

1. Introduction

In colorectal cancer (CRC), TNM staging is the most important factor in determining prognosis and patient management [1]. However, a wide biological heterogeneity of individual tumors may account for considerable differences in tumor behavior seen in patients within the same stage. Therefore, additional biomarkers to better stratify outcome are sought after within certain patient groups. In the avid search for such potential biomarkers in CRC, tumor budding has emerged as an especially robust and promising candidate.

Tumor budding, defined as single tumor cells or small clusters of ≤ 4 tumor cells at the invasive front of CRC [2], is a morphologically visible sign of tumor dissemination and has been linked to epithelial-to-mesenchymal-like processes [3]. Most recently, transcriptome profiling studies have linked tumor buds with the mesenchymal type (Consensus Molecular Classification type 4) of CRC [4]. It is therefore not surprising that large bodies of data have consistently demonstrated tumor budding to be an independent adverse prognostic marker in CRC and associated with unfavorable clinicopathological features, nodal and distant metastases [5-7].

Because of its potential clinical implications, tumor budding has been especially well studied in 2 scenarios. In endoscopically resected pT1 CRC, tumor budding has been demonstrated to be a strong and independent predictor of nodal metastases and may help select patients for segmental resection with lymphadenectomy [2,8-11]. In stage II CRC patients, high-grade budding tumors have been shown to behave aggressively [2,12] and similarly to stage III CRC [13-15], with shorter survival times and higher rates of recurrence;

therefore, these patients may be offered adjuvant chemotherapy.

Although the data in the literature would certainly support the integration of tumor budding in reporting protocols, the lack of a standardized scoring system has been one of the major barriers to the routine reporting of tumor budding in CRC. The objective of the International Tumor Budding Consensus Conference (ITBCC) was to provide an evidence-based set of recommendations for such a standardized method [2]. Since published, the ITBCC guidelines have been included as an additional reporting parameter in the protocol of the College of American Pathologists with the recommendation to report tumor budding in pT1 and stage II CRC [16].

Although largely based on the method proposed by Ueno et al [10], further validation studies should provide the basis for more solid implementation of the ITBCC method in CRC reporting. Therefore, the aim of this study was to validate the ITBCC method of tumor budding assessment in a large, well-characterized CRC cohort.

2. Materials and methods

2.1. Patient cohort

Three hundred seventy-nine patients with primary CRC resected between 2002 and 2014 were entered into this study. Hematoxylin and eosin (H&E)-stained slides were re-reviewed by 2 expert gastrointestinal pathologists (A. L., H. D.) according to the Union for International Cancer Control TNM seventh edition [1] for pathological features. In addition




Fig. 1 Tumor budding as assessed by the ITBCC 2016 guidelines. No tumor budding (A) and low-grade tumor budding (B; both BD1, 0-4 tumor buds/0.785 mm²). C, Intermediate-grade tumor budding (BD2, 5-9 tumor buds/0.785 mm²). D, High-grade tumor budding (BD3, ≥ 10 tumor buds/0.785 mm²). Arrows point to selected tumor buds in all images.

Table 1 Patient characteristics (n = 379) and association of tumor budding (BD category) with clinicopathological features

Features	Frequency, n (%)	BD category			P		P
		BD1	BD2	BD3	BD1, 2, 3	BD1 + 2 vs 3	Continuous score
Age (y; n = 379), mean	69.6	70.0	69.3	69.4	.9714	.9008	.4777
Sex (n = 379)							
Female	150 (39.6)	59 (39.6)	32 (31.7)	59 (45.7)	.0964	.0782	.4839
Male	229 (60.4)	90 (60.4)	69 (68.3)	70 (54.3)			
Histologic subtype (n = 378)							
Adenocarcinoma	319 (84.4)	121 (81.8)	86 (85.2)	112 (86.8)	.5738	.6446	.3979
Mucinous	52 (13.8)	25 (16.9)	12 (11.9)	15 (11.6)			
Other	7 (1.9)	2 (1.4)	3 (3.0)	2 (1.6)			
Tumor location (n = 379)							
Left	229 (60.4)	91 (61.1)	60 (59.4)	78 (60.5)	.9655	.9902	.6809
Right	150 (39.6)	58 (38.9)	41 (40.6)	51 (39.5)			
Postoperative therapy (n = 232)							
None	172 (74.1)	81 (84.4)	36 (75.0)	55 (62.5)	.0032 ^a	.0016 ^a	.0004 ^a
Treated	60 (25.9)	15 (15.6)	12 (25.0)	33 (37.5)			
pT (n = 378)							
pT1	30 (7.9)	20 (13.4)	6 (5.9)	4 (3.1)	<.0001 ^a	<.0001 ^a	<.0001 ^a
pT2	47 (12.4)	23 (15.4)	14 (13.9)	10 (7.8)			
pT3	190 (50.1)	84 (56.4)	49 (48.5)	57 (44.2)			
pT4	111 (29.3)	21 (14.2)	32 (31.7)	58 (45.0)			
pN (n = 370)							
pN0	185 (50.0)	98 (67.1)	52 (52.5)	35 (28.0)	<.0001 ^a	<.0001 ^a	<.0001 ^a
pN1	115 (31.1)	38 (26.0)	35 (35.4)	42 (33.6)			
pN2	70 (18.9)	10 (6.9)	12 (12.1)	48 (38.4)			
TNM stage (n = 347)							
I	57 (16.4)	34 (24.8)	16 (17.2)	7 (6.0)	<.0001 ^a	<.0001 ^a	<.0001 ^a
II	109 (31.4)	54 (39.4)	31 (33.3)	24 (20.5)			
III	111 (32.0)	37 (27.0)	32 (34.4)	42 (35.9)			
IV	70 (20.2)	12 (8.8)	14 (15.1)	44 (37.6)			
Tumor grade (n = 364)							
G1	21 (5.8)	16 (11.4)	5 (5.0)	0 (0.0)	.002 ^a	.002 ^a	<.0001 ^a
G2	264 (72.5)	97 (68.8)	76 (76.0)	91 (74.0)			
G3	79 (21.7)	28 (19.9)	19 (19.0)	32 (26.0)			
Lymphatic invasion (n = 362)							
L0	146 (40.3)	94 (65.7)	36 (38.3)	16 (12.8)	<.0001 ^a	<.0001 ^a	<.0001 ^a
L1	216 (59.7)	49 (34.3)	58 (61.7)	109 (87.2)			
Venous invasion (n = 365)							
V0	213 (58.4)	107 (74.3)	56 (58.3)	50 (40.0)	<.0001 ^a	<.0001 ^a	<.0001 ^a
V1	152 (41.6)	37 (25.7)	40 (41.7)	75 (60.0)			
Perineural invasion (n = 346)							
Pn0	276 (79.8)	127 (93.4)	78 (85.7)	71 (59.7)	<.0001 ^a	<.0001 ^a	<.0001 ^a
Pn1	70 (20.2)	9 (6.6)	13 (14.3)	48 (40.3)			
MMR status (n = 258)							
MMR deficient	36 (14.0)	10 (10.9)	11 (14.9)	15 (16.3)	.5479	.4172	.3876
MMR proficient	222 (86.1)	82 (89.1)	63 (85.1)	77 (83.7)			
Klintrup-Mäkinen (n = 304)							
0	15 (4.9)	5 (3.9)	1 (1.3)	9 (9.0)	.0206 ^a	.0048 ^a	.0258 ^a
1	137 (45.1)	61 (48.0)	37 (48.1)	39 (39.0)			
2	116 (38.2)	45 (35.4)	25 (32.5)	46 (46.0)			
3	36 (11.8)	16 (12.6)	14 (18.2)	6 (6.0)			
Expanding tumor border (%) (n = 298), mean	46.5	61.5	42.4	30.9	<.0001 ^a	<.0001 ^a	<.0001 ^a
OS time (mo; n = 379), median (95% CI)	127.6 (99-147)	103.2 + -5.7	93.1 + -6.6	67 + -4.2	.0251 ^a	.0106 ^a	.0195 ^{a,b}

^a Statistically significant values ($P < .05$).^b Increased HR per bud, 1.023 (95% CI, 1.004-1.043).

to standard reporting elements, this included assessment of peritumoral inflammation (Klintrup-Mäkinen score) and tumor border configuration. The Klintrup-Mäkinen score was obtained by evaluating the overall inflammatory reaction at the invasive margin on a 4-degree scale ranging from 0 (no increase of inflammatory cells) to 3 (very prominent inflammatory reaction with invariable and frequent destruction of cancer cell islets) [17]. Tumor border configuration was assessed according to Karamitopoulou et al [18] by scoring the percentage of infiltrating tumor margin in 5% increments. Tumors were categorized as right (cecum, ascending colon, hepatic flexure, and transverse colon), left (splenic flexure, descending colon, sigmoid), or rectal based on surgical and pathological reports. Exclusion criteria included preoperative chemoradiotherapy, previous endoscopic resection, tumors with pure signet ring cell and mucinous histology in accordance with the ITBCC guidelines, and patients with time of death within 1 month after surgery. Patient follow-up was scheduled according to the recommendations of the Swiss Society of Gastroenterology for surgically resected colorectal tumors [19]. Follow-up data including overall survival (OS) time

Table 2 Multivariate analysis of tumor budding category along with TNM stage and postoperative therapy

Feature	HR (95% CI)	<i>P</i>
BD category		
BD1 + 2	1.0	.0911
BD3	1.51 (0.94-2.42)	
TNM stage		
IV	1.0	.0041
I	0.23 (0.08-0.62)	
II	0.62 (0.33-1.15)	.127
III	0.75 (0.43-1.3)	.3214
Postoperative therapy		
None	1.0	.1022
Treated	0.65 (0.38-1.09)	

(available for all patients; median, 127.6 months; 95% confidence interval [CI], 99-147) and disease-free survival (DFS) time (available for 78 patients in subset analysis of stage II patients; mean, 37.3 months) were obtained. Ethics approval was




Fig. 2 Survival curves of CRC patients stratified by budding status. A, OS in all patients (n = 379) with low- (BD1), intermediate- (BD2), and high-grade budding (BD3). B, OS in all patients (n = 379) stratified by BD1 + 2 versus BD3. C, DFS in stage II patients (n = 78) stratified by BD1 + 2 versus BD3.

granted by the Cantonal Ethics Commission of Bern (KEK 2017-01803; October 24, 2017).

2.2. Assessment of tumor budding

Tumor budding was scored by 2 observers (H. D. and F. G.) according to the ITBCC 2016 recommendations [2]: H&E-stained sections were scanned at medium power ($\times 10$) to identify the densest area of budding at the tumor front (“hot-spot”). Tumor buds were counted in this area at $\times 20$ magnification (Nikon Eclipse E600, 750 objective, field diameter of 0.25 mm, area of 1.227 mm²; Nikon AG Instruments, Egg, Switzerland). The bud count was divided by the normalization

factor (1.563) relative to the specific microscope eyepiece field number diameter to determine the tumor bud count per 0.785 mm². The final bud count and the budding category (Bd 1, 0-4 buds; Bd2, 5-9 buds; Bd3, ≥ 10 buds; Fig. 1) were recorded. Cases with pure mucinous or signet ring cell morphology were excluded from analysis. Selected cases were discussed at a multiheader microscope to obtain a consensus score.

2.3. Mismatch repair protein status

Immunohistochemistry for mismatch repair (MMR) proteins MLH1, MSH2, MSH6, and PMS2 was available for

Table 3 Association of tumor budding (BD category) with clinicopathological features in stage II patients (n = 109)

Features	Frequency, n (%)	BD category			P		P
		BD1	BD2	BD3	BD1, 2, 3	BD1 + 2 vs 3	Continuous score
Age (y) (n = 109), mean	69.5	69.5	69.0	69.8	.9471	.7755	.5387
Sex (n = 109)							
Female	48 (44.0)	23 (43.6)	11 (35.5)	14 (58.3)	.2281	.1101	.422
Male	61 (56.0)	31 (57.4)	20 (64.5)	10 (41.7)			
Histologic subtype (n = 109)							
Adenocarcinoma	89 (81.7)	44 (81.5)	25 (80.7)	20 (83.3)	.6262	.8593	.5759
Mucinous	19 (17.4)	10 (18.5)	5 (16.1)	4 (16.7)			
Other	1 (0.9)	0 (0.0)	1 (3.2)	0 (0.0)			
Tumor location (n = 109)							
Left	56 (51.4)	33 (61.1)	12 (38.7)	11 (45.8)	.1145	.5384	.1633
Right	53 (48.6)	21 (38.9)	19 (61.3)	13 (54.2)			
Postoperative therapy (n = 68)							
None	57 (83.8)	32 (88.9)	13 (81.3)	12 (75.0)	.4321	.2731	.1723
Treated	11 (16.2)	4 (11.1)	3 (18.8)	4 (25.0)			
pT (n = 109)							
pT3	88 (80.7)	48 (88.9)	24 (77.4)	16 (66.7)	.0614	.0478	.0846
pT4	21 (19.3)	6 (11.1)	7 (22.6)	8 (33.3)			
Tumor grade (n = 105)							
G1	6 (5.7)	4 (8.0)	2 (6.5)	0 (0.0)	.1851	.0497	.5626
G2	83 (79.1)	41 (82.0)	25 (80.7)	17 (70.8)			
G3	16 (15.2)	5 (10.0)	4 (12.9)	7 (29.2)			
Lymphatic invasion (n = 103)							
L0	69 (67.0)	41 (82.0)	19 (63.3)	9 (39.1)	.0013	.0013	.0003
L1	34 (33.0)	9 (18.0)	11 (36.7)	14 (60.9)			
Venous invasion (n = 104)							
V0	79 (76.0)	44 (86.3)	19 (63.3)	16 (69.6)	.0472	.416	.0572
V1	25 (24.0)	7 (13.7)	11 (36.7)	7 (30.4)			
Perineural invasion (n = 99)							
Pn0	90 (90.9)	47 (97.9)	27 (93.1)	16 (72.7)	.0027	.0008	.0023
Pn1	9 (9.1)	1 (2.1)	2 (6.9)	6 (27.3)			
MMR status (n = 66)							
MMR deficient	12 (18.2)	3 (9.7)	5 (27.8)	4 (23.5)	.229	.507	.1581
MMR proficient	54 (81.8)	28 (90.3)	13 (72.2)	13 (76.5)			
Klintrup-Mäkinen (n = 89)							
0	2 (2.3)	0 (0.0)	0 (0.0)	2 (10.5)	.1818	.0429	.0306
1	38 (42.7)	22 (47.8)	9 (37.5)	7 (36.8)			
2	40 (45.0)	19 (41.3)	12 (50.0)	9 (47.4)			
3	9 (10.1)	5 (10.9)	3 (12.5)	1 (5.3)			
Expanding tumor border (%; n = 109), mean	56.7	67.4	42.8	47.3	.0023	.1526	<.0001

342 patients and performed as previously described [20]. Markers were considered positive if any nuclear expression in tumor cells was seen. Tumors negative for at least 1 marker were considered MMR deficient, and MMR proficient if all markers were expressed in the tumor. Because information on family history was unavailable, no attempt was made to further subdivide patients into Lynch syndrome or sporadic microsatellite instable tumors (MSI).

2.4. Statistical analysis

Descriptive statistics for categorical and continuous variables were analyzed. The differences between categorical histopathologic features and BD categories were carried out using the χ^2 test. The association with continuous budding counts was analyzed using the Wilcoxon rank sum test. For age and percentage of expanding tumor border configuration, the Pearson correlation analysis was performed. Log-rank test and Kaplan-Meier method were used for univariate survival time analysis. OS and DFS were the end points of interest. Multivariate survival analysis was performed using Cox regression analysis. Hazard ratios (HRs) and 95% CI were used to determine effect size. *P* values were all 2-sided and considered significant when *P* < .05.

3. Results

3.1. Associations of tumor budding with clinicopathological features and survival

Across all stages (*n* = 379), tumor budding was low grade (BD1) in 149 patients, intermediate grade (BD2) in 101 patients, and high-grade (BD3) in 129 patients (illustrated in Fig. 1A-D). Frequencies of each variable and associations with BD category (BD1, BD2, and BD3 separately), a 2-tier classification (BD1 + 2 versus BD3), and continuous scores are included in Table 1. As expected, higher tumor budding grades were seen in patients having received postoperative therapy (BD1/2/3: *P* = .032, BD1/2 versus BD3: *P* = .0016 and *P* = .0044 for continuous scores, respectively), patients with higher pT-stage (*P* < .001, all), nodal metastases (*P* < .001, all), more advanced TNM stage (*P* < .001, all), higher tumor grade (BD1/2/3: *P* = .002, BD1/2 versus BD3: *P* = .002 and *P* < .001 for continuous scores, respectively), lymphatic invasion (*P* < .001, all), venous invasion (*P* < .001, all), and perineural invasion (*P* < .001, all). Higher tumor budding was inversely associated with expanding tumor border configuration (*P* < .001, all) and higher peritumoral inflammation (Klintrup-Mäkinen score; *P* = .026, *P* = .0048, and *P* = .0258, respectively). No significant differences in BD scores were seen in different tumor sites and in MMR-proficient versus MMR-deficient tumors.

On univariate analysis, tumor budding was significantly associated with shorter OS times in a 3-tier (BD1 versus BD2

versus BD3, *P* = .0251; Fig. 2A) and 2-tier approach (BD1-2 versus BD3, *P* = .0106; Fig. 2B) and when taken as a continuous variable (HR, 1.023 [95% CI, 1.004-1.043] per increased bud; *P* = .0195). However, significant associations between tumor budding and survival were lost in multivariate analysis (Table 2) including BD category (*P* = .0911), TNM stage (stage IV: *P* = .0041), and postoperative therapy (*P* = .1022).

3.2. Analysis of tumor budding, clinicopathological features, and survival in stage II patients

In stage II patients (*n* = 109,) tumor budding was low grade (BD1) in 54 (49.5%) patients, intermediate grade (BD2) in 31 (28.4%) patients, and high grade (BD3) in 24 (22.0%) patients. Frequencies of each variable and associations with BD category (BD1, BD2, and BD3 separately), a 2-tier classification (BD1 + 2 versus BD3), and continuous scores can be found in Table 3. In univariate analysis, tumor budding was not associated with OS time but with 5-year DFS (*P* = .0084; Fig. 2C). Because only 11 patients received postoperative therapy, no conclusions regarding possible associations between tumor budding grade, postoperative therapy, and survival could be made.

4. Discussion

In this study, we provide a validation of the ITBCC scoring method on a large, well-characterized mixed-stage CRC cohort. Assessed by this method and in line with previous results in the literature using similar and other scoring systems, tumor budding was associated with adverse clinicopathological features and survival both in the entire cohort and in a subset of stage II patients.

Although several different tumor budding scoring methods have been proposed in the literature, the ITBCC consensus scoring method is largely based on the system proposed by Ueno et al [10]. Key aspects of this proposal (hotspot, $\times 20$ magnification, and 0.785 mm²) were adopted by the Japanese Society for Cancer of the Colon and the Rectum before the ITBCC, and routine reporting of tumor budding in pT1 tumors according to this method has been performed since 2009 [21]. Indeed, this system has been used in studies including large Japanese patient cohorts with pT1 CRC [11,22] and by others with slight variations, such as an adapted high-power field size of 0.95 mm² [23-25]. However, a standardized consensus approach such as provided by the ITBCC guidelines is essential for tumor budding to be validated to compare study results and ultimately be used as a biomarker in routine diagnostics.

One important aspect of the ITBCC recommendations is the separate reporting of tumor budding grade (BD 1, 2, or 3) and the raw tumor bud count. As a numerical variable on a biological spectrum, continuous tumor bud counts are expected to provide more precise risk stratification than cutoffs alone (as seen in Table 1). It must also be emphasized that

relevant cutoffs will vary according to the clinical end point of a certain scenario (nodal metastases for pT1 tumors where BD2 and BD3 are considered high risk, and recurrence/survival in advanced tumors where only BD3 is considered a risk factor) [2]. Hence, a 2-tier classification (BD1 + 2 versus BD3) was preferred to better stratify patients in this cohort, where pT1 tumors were underrepresented (n = 30) and therefore not analyzed separately.

In this study using a mixed-stage cohort including patients with distant metastases, tumor budding was not an independent prognostic factor on multivariate analysis including TNM stage. This expected result emphasizes the importance and relevance of accurate clinical and pathological staging in routine as a basis for complementary biomarker studies. In a separate analysis of stage II patients, tumor budding was significantly associated with shorter 5-year DFS but not with shorter OS. This finding must be interpreted with caution because only few events were observed in this relatively small patient group, but underlines DFS as a more meaningful survival parameter in stage II tumors.

Tumor budding in CRC is a robust biomarker that is simple to use and can be assessed using routine light microscopy on H&E-stained slides. The standardized method as proposed by the ITBCC consensus recommendations will hopefully pave the way of integrating tumor budding in reporting protocols, as has recently been the case for the latest update of the College of American Pathologists CRC checklist. Further validation studies such as we present here aim to promote more widespread integration of the ITBCC consensus method in standardized reporting of tumor budding in CRC.

References

- [1] Brierly JD, GM, Wittekind C. TNM Classification of Malignant Tumors. 7th edition Oxford, UK: Wiley Blackwell; 2010.
- [2] Lugli A, Kirsch R, Ajioka Y, et al. Recommendations for reporting tumor budding in colorectal cancer based on the International Tumor Budding Consensus Conference (ITBCC) 2016. *Mod Pathol* 2017;30:1299-311.
- [3] Dawson H, Lugli A. Molecular and pathogenetic aspects of tumor budding in colorectal cancer. *Front Med (Lausanne)* 2015;2:11.
- [4] De Smedt L, Palmans S, Andel D, et al. Expression profiling of budding cells in colorectal cancer reveals an EMT-like phenotype and molecular subtype switching. *Br J Cancer* 2017;116:58-65.
- [5] Rogers AC, Winter DC, Heeney A, et al. Systematic review and meta-analysis of the impact of tumour budding in colorectal cancer. *Br J Cancer* 2016;115:831-40.
- [6] Mitrovic B, Schaeffer DF, Riddell RH, Kirsch R. Tumor budding in colorectal carcinoma: time to take notice. *Mod Pathol* 2012;25:1315-25.
- [7] van Wyk HC, Park J, Roxburgh C, Horgan P, Foulis A, McMillan DC. The role of tumour budding in predicting survival in patients with primary operable colorectal cancer: a systematic review. *Cancer Treat Rev* 2015;41:151-9.
- [8] Beaton C, Twine CP, Williams GL, Radcliffe AG. Systematic review and meta-analysis of histopathological factors influencing the risk of lymph node metastasis in early colorectal cancer. *Colorectal Dis* 2013;15:788-97.
- [9] Bosch SL, Teerenstra S, de Wilt JH, Cunningham C, Nagtegaal ID. Predicting lymph node metastasis in pT1 colorectal cancer: a systematic review of risk factors providing rationale for therapy decisions. *Endoscopy* 2013;45:827-34.
- [10] Ueno H, Mochizuki H, Hashiguchi Y, et al. Risk factors for an adverse outcome in early invasive colorectal carcinoma. *Gastroenterology* 2004;127:385-94.
- [11] Kawachi H, Eishi Y, Ueno H, et al. A three-tier classification system based on the depth of submucosal invasion and budding/sprouting can improve the treatment strategy for T1 colorectal cancer: a retrospective multicenter study. *Mod Pathol* 2015;28:872-9.
- [12] Petrelli F, Pezzica E, Cabiddu M, et al. Tumour budding and survival in stage II colorectal cancer: a systematic review and pooled analysis. *J Gastrointest Cancer* 2015;46:212-8.
- [13] Nakamura T, Mitomi H, Kanazawa H, Ohkura Y, Watanabe M. Tumor budding as an index to identify high-risk patients with stage II colon cancer. *Dis Colon Rectum* 2008;51:568-72.
- [14] Okuyama T, Nakamura T, Yamaguchi M. Budding is useful to select high-risk patients in stage II well-differentiated or moderately differentiated colon adenocarcinoma. *Dis Colon Rectum* 2003;46:1400-6.
- [15] Wang LM, Kevans D, Mulcahy H, et al. Tumor budding is a strong and reproducible prognostic marker in T3N0 colorectal cancer. *Am J Surg Pathol* 2009;33:134-41.
- [16] Kakar S, Shi C, Berho ME, et al. Protocol for the examination of specimens from patients with primary carcinoma of the colon and rectum. . Edition 4.0.0.1 College of American Pathologists; 2017 <http://www.cap.org/ShowProperty?nodePath=/UCMCon/Contribution%20Folders/WebContent/pdf/cp-colon-17protocol-4001.pdf>, Accessed date: 22 May 2017.
- [17] Klinttrup K, Makinen JM, Kauppila S, et al. Inflammation and prognosis in colorectal cancer. *Eur J Cancer* 2005;41:2645-54.
- [18] Karamitopoulou E, Zlobec I, Koelzer VH, Langer R, Dawson H, Lugli A. Tumour border configuration in colorectal cancer: proposal for an alternative scoring system based on the percentage of infiltrating margin. *Histopathology* 2015;67:464-73.
- [19] Dorta G, MC. Nachsorge nach koloskopischer Polypektomie und entferntem Kolorektalkarzinom (article in German and French). *Swiss Med Forum* 2016;16:164-7.
- [20] Dawson H, Koelzer VH, Lukesch AC, et al. Loss of Cdx2 expression in primary tumors and lymph node metastases is specific for mismatch repair-deficiency in colorectal cancer. *Front Oncol* 2013;3:265.
- [21] Watanabe T, Itabashi M, Shimada Y, et al. Japanese Society for Cancer of the C, Rectum. Japanese Society for Cancer of the Colon and Rectum (JSCCR) guidelines 2010 for the treatment of colorectal cancer. *Int J Clin Oncol* 2012;17:1-29.
- [22] Ueno H, Hase K, Hashiguchi Y, et al. Novel risk factors for lymph node metastasis in early invasive colorectal cancer: a multi-institution pathology review. *J Gastroenterol* 2014;49:1314-23.
- [23] Graham RP, Vierkant RA, Tillmans LS, et al. Tumor budding in colorectal carcinoma: confirmation of prognostic significance and histologic cutoff in a population-based cohort. *Am J Surg Pathol* 2015;39:1340-6.
- [24] Betge J, Kornprat P, Pollheimer MJ, et al. Tumor budding is an independent predictor of outcome in AJCC/UICC stage II colorectal cancer. *Ann Surg Oncol* 2012;19:3706-12.
- [25] Max N, Harbaum L, Pollheimer MJ, Lindtner RA, Kornprat P, Langner C. Tumour budding with and without admixed inflammation: two different sides of the same coin? *Br J Cancer* 2016;114:368-71.