



## Original contribution

# Plasmacytoid/diffuse urothelial carcinoma: a single-institution immunohistochemical and molecular study of 69 patients <sup>☆, ☆ ☆, ★</sup>



Carmen M. Perrino MD<sup>a,\*</sup>, John Eble MD<sup>a</sup>, Chia-Sui Kao MD<sup>b</sup>, Rumeal D. Whaley MD<sup>a</sup>,  
Liang Cheng MD<sup>a</sup>, Mohammad Idrees MD<sup>a</sup>, Neda Hashemi-Sadraei MD<sup>c</sup>,  
M. Francesa Monn MD<sup>d</sup>, Hristos Z. Kaimakliotis MD<sup>d</sup>, Elhaam Bandali MS<sup>d</sup>,  
David Grignon MD<sup>a</sup>

<sup>a</sup>Indiana University, Department of Pathology and Laboratory Medicine, Indianapolis, IN 46202

<sup>b</sup>Stanford University, Department of Pathology, Stanford, CA 94305

<sup>c</sup>Indiana University, Department of Medicine, Division of Hematology and Oncology, Indianapolis, IN 46202

<sup>d</sup>Indiana University, Department of Urology, Indianapolis, IN 46202

Received 28 January 2019; revised 13 April 2019; accepted 18 April 2019

**Keywords:**

Urothelial carcinoma;  
Plasmacytoid;  
Signet ring;  
Diffuse;  
Bladder

**Summary** Accurate diagnosis of plasmacytoid urothelial carcinoma (PUC) is important given its poor prognosis and frequent presentation at high stage. We aim to assess the clinicopathological features, molecular aberrations, and follow-up data in a series of PUC cases from a single tertiary cancer center. Seventy-two urinary bladder, ureteral, and renal pelvic specimens with urothelial carcinoma with plasmacytoid differentiation were identified. Immunohistochemical stains were performed on 48 cases. Among urinary bladder origin markers, GATA3 was most sensitive (96%). Breast carcinoma markers (estrogen receptor, mammaglobin) were usually negative, but progesterone receptor stained 1 case (4%). Neuroendocrine markers CD56 and TTF-1 were each positive in 1 case (4% and 4%, respectively). Gastrointestinal adenocarcinoma marker CDX2 was positive in 4 cases (15%), but nuclear  $\beta$ -catenin was negative in all cases. CD138 was positive in 83% and E-cadherin expression was lost in 57% of cases. Fluorescence in situ hybridization using the UroVysion Bladder Cancer Kit and *FGFR3* mutational analysis using polymerase chain reaction were performed on 15 cases; deletion of chromosome 9p21 was common (60%), and *FGFR3* mutations were detected in 60% of cases (5 cases had both deletion 9p21 and *FGFR3* mutations). Cases were divided into 3 morphologic groups: classic (29%), desmoplastic (35%), and pleomorphic (36%). The 3 morphologic subtypes had distinct survival outcomes ( $P = .083$ ), with median survival for all patients 18 being months versus 10 months for the desmoplastic group.

© 2019 Elsevier Inc. All rights reserved.

<sup>☆</sup> Competing interests: The authors have no conflicts of interest to disclose.

<sup>☆☆</sup> Funding/Support: All funding was provided by the Indiana University Department of Pathology and Laboratory Medicine (Indianapolis, IN).

<sup>\*</sup> Previous presentation: Portions of this data were presented at the United States and Canadian Academy of Pathology 2017 Annual Meeting as an abstract and a poster presentation.

<sup>\*</sup> Corresponding author at: Rhode Island Hospital Department of Pathology and Laboratory Medicine, 593 Eddy St APC 12, Providence, RI 02903.

*E-mail address:* cperrino@lifespan.org (C. M. Perrino).

## 1. Introduction

Urothelial carcinoma accounts for approximately 90% of primary urinary bladder carcinomas and has a tremendous propensity for morphologic variations [1]. Accurate identification of variants is important for a variety of reasons, as some variants have prognostic significance and others may be mistaken for a benign entity or another tumor [1]. In the most recent edition of the *World Health Organization (WHO) Classification of Tumours of the Urinary System and Male Genital Organs*, the plasmacytoid variant is combined with signet ring (lacking extracellular mucin) and diffuse variants [1]. The plasmacytoid variant in particular has been associated with presentation at a higher pathologic T stage, a higher rate of positive surgical margins, and lower overall survival [2]. Awareness of this variant is crucial when interpreting ureteral and urethral margins on frozen section given its propensity to infiltrate the soft tissue surrounding the urothelium rather than involving the overlying urothelium itself [3]. In addition, being alert to the unique pattern of discohesive, single cells that may infiltrate the sinusoids of lymph nodes is important to identify metastases [4].

One of the earliest references to primary signet ring cell carcinoma of the bladder in the literature is by Saphir in 1955, which describes cells with intracytoplasmic mucin that compressed the nuclei and gave them a crescent shape, cells with basophilic cytoplasm and eccentric nuclei, and cells resembling monocytes. He references another report of tumors composed almost exclusively of similar signet ring cells which were diffusely infiltrative and were associated with distant metastases [5]. In a subsequent study of signet ring cell carcinoma of the urinary bladder, Grignon et al included linitis plastica-like growth in the definition and described similar cells to those reported by Saphir [6]. Since that time, several case series have described the morphologic features of plasmacytoid urothelial carcinoma, the largest and most recent of which included 49 cases from The University of Texas MD Anderson Cancer Center [7-18]. Although these studies describe plasmacytoid differentiation as having a central or eccentric nucleus and infiltrative architecture consisting of discohesive, single cells and linear arrangements of cells, they rarely offer much additional detail [1]. Some authors mention a few other morphologic observations such as the presence of a stromal response in a minority of cases; however, these additional descriptions are minimal and not a focal point of any particular study [12,13,19-21]. Some cases of urothelial carcinoma have been described as having rhabdoid differentiation; however, the morphologic features and images in these studies fall within the spectrum of plasmacytoid differentiation outlined in most other papers [22-25]. Lastly, reports of conventional urothelial carcinoma with an infiltrative growth pattern as single cells or cords of cells often with associated desmoplasia have shown a particularly poor prognosis in those cases [26].

We aim to describe the detailed morphologic features, immunohistochemical expression pattern, molecular aberrations, and clinical follow-up of a large series of urothelial carcinoma with plasmacytoid/diffuse differentiation from a single

institution. Previously reported immunohistochemical stain results have been reported by variety of authors. However, we provide a description of the staining pattern for pertinent stains when applied to a large series of tumors which were all processed by 1 laboratory, thus eliminating interlaboratory technical issues as a possible explanation for aberrant staining. Likewise, we hope to expand upon previously reported information regarding the link between molecular aberrations detected in conventional urothelial carcinoma and the plasmacytoid variant.

## 2. Materials and methods

A search of the pathology database at Indiana University (January 1996-September 2016) for in-house specimens from the urinary bladder, renal pelvis, or ureter with a final diagnosis of urothelial carcinoma with plasmacytoid or signet ring cell differentiation was performed. Pathology reports were reviewed, and data including patient demographics (age, sex), types of specimens, estimated quantity of plasmacytoid differentiation specified, pathologic American Joint Committee on Cancer (AJCC) TNM stage, and clinical follow-up data were collected. In addition, all available slides from each case were re-reviewed, and the cases were divided into 3 different morphologic groups (please refer to the "Results" section to see criteria used for distinction).

### 2.1. Immunohistochemistry

A representative block from 48 cases of urothelial carcinoma with plasmacytoid differentiation was selected. Unstained slides were cut at 5  $\mu$ m in thickness. The antibodies used are listed in Table 1. Immunohistochemical stains were performed as previously described [14]. Twenty-six cases were initially stained with all 16 antibodies. An additional 22 cases were selected because of unique morphologic features (classic morphology, pleomorphic cells, desmoplastic stromal reaction, etc) and were stained with E-cadherin and CD138. Positivity was interpreted as nuclear for  $\beta$ -catenin, CDX2, estrogen receptor (ER), GATA3, p53, p63, progesterone receptor (PR), and TTF-1. Positivity was cytoplasmic for 34 $\beta$ E12, keratin 7, keratin 20, and mammaglobin. Positivity was membranous for CD56, CD138, E-cadherin, thrombomodulin, and uroplakin III. Staining was scored for intensity (0: no staining; 1: weak; 2: moderate; 3: strong) and proportion of cells stained (0: 0%; 1: >0% to  $\leq$ 25%; 2: >25% to  $\leq$ 50%; 3: >50% to  $\leq$ 75%; 4: >75 to 100%).

### 2.2. UroVysion fluorescence in situ hybridization

Unstained sections were prepared from available tissue blocks from 15 cases and were deparaffinized with xylene for fluorescence in situ hybridization (FISH) using the UroVysion probe set. Slides were treated with absolute ethanol, then

**Table 1** Antibodies used in this study

Antibody	Clone	Dilution	Company
$\beta$ -Catenin	14	Prediluted	Cell Marque, Rocklin, CA
CD56	123C3	Prediluted	Dako, Santa Clara, CA
CD138	M115	Prediluted	Dako, Santa Clara, CA
CDX2	DAK-CDX2	Prediluted	Dako, Santa Clara, CA
Keratin 7	OV-TL	Prediluted	Dako, Santa Clara, CA
Keratin 20	K <sub>s</sub> 20.8	Prediluted	Dako, Santa Clara, CA
E-cadherin	NCH-38	Prediluted	Dako, Santa Clara, CA
Estrogen receptor	EP1	1:50	Dako, Santa Clara, CA
GATA3	L50-823	Prediluted	Biocare Medical, Concord, CA
Mammaglobin	31A5	Prediluted	Cell Marque, Rocklin, CA
P53	DO-7	Prediluted	Dako, Santa Clara, CA
P63	BC4A4	1:100	Dako, Santa Clara, CA
Progesterone receptor	PgR 636	Prediluted	Dako, Santa Clara, CA
Thrombomodulin	1009	1:100	Dako, Santa Clara, CA
TTF-1	86763/1	Prediluted	Dako, Santa Clara, CA
Uroplakin	AU-1	1:30	Cell Marque, Rocklin, CA

air-dried and boiled in a glass staining-jar with 1× citrate buffer (pH 6.0) (Zymed, South San Francisco, CA) within a beaker filled with distilled water on a hot block for 10 minutes. Slides were washed with distilled water and transferred to 2× sodium citrate buffer for 5 minutes. The slides were air dried and digested with 0.75-mL pepsin (5 mg/mL in 0.01 N HCl with 0.9% NaCl; Sigma, St Louis, MO) at 37°C for 40 minutes. The slides were then washed with distilled water and 2× sodium citrate buffer, followed by air-drying. Chromosome enumeration probes (CEPs) for chromosomes 3, 7, and 17, and the locus specific indicator probe for 9p21 were labeled with fluorophores. CEP3, CEP7, and CEP17 probes were labeled with Spectrum Red, Spectrum Green, and Spectrum Aqua, respectively. Locus specific indicator p16 (9p21) was labeled with Spectrum Gold (Vysis, Downers Grove, IL). The probes were diluted 1:10 with tDenHyb2 (Insitus, Albuquerque, NM). Five microliters of diluted probes was added to each slide in reduced light conditions. The slides were then covered with a 22 × 22 cover slip, sealed with rubber cement, and put into an opaque plastic box wrapped with aluminum foil. The slides were denatured at 83°C for 12 minutes and hybridized at 37°C overnight. After hybridization, the slides were washed and counterstained with 10  $\mu$ L DAPI (Insitus, Albuquerque, NM) and sealed with a 50 × 22 cover slip.

The stained slides were observed and documented using MetaSystem software (Belmont, MA) under 100× oil objective using filters SP-100 for DAPI, FITC MF-101 for Spectrum Green, Gold 31003 for Spectrum Gold, Aqua 31036 V2 for Spectrum Aqua, and Texas Red Sp103 for Spectrum Red signals. Five sequential focus stacks with 0.4- $\mu$ m intervals were acquired and integrated into a single image to reduce thickness-related artifacts.

For each case, 200 nuclei were counted. Each cell was simultaneously analyzed for the signals of chromosomes 3, 7, and 17, and 9p21. Chromosomal gain or loss was defined based on the Gaussian model and relative to normal controls.

The cutoff values were set at the mean  $\pm$ 3 SD of the number of disomic cells in control individuals. The mean  $\pm$ 3 SD represents a specificity of 99.9%. Any tumor cases with a score beyond the cutoff value were considered to have either a gain or a loss of the designated chromosome(s).


### 2.3. *FGFR3* mutation analysis

*FGFR3* mutational analysis was performed on 15 cases. For the *FGFR3* gene, exons 7, 10, and 15 were amplified by polymerase chain reaction (PCR) using previously established primers. PCR was performed with 3 mL of isolated genomic DNA in a final volume of 50 mL containing 2.3 mmol/L MgCl<sub>2</sub>, 10 mmol/L Tris-HCl (pH 8.3), 50 mmol/L KCl, 2 mmol/L deoxynucleotide triphosphates, 2 mmol/L each primer, and 2 U Taq DNA polymerase (Bio-Rad, Hercules, CA, USA). Each PCR protocol had an initial denaturing step of 95°C for 5 minutes followed by 40 cycles of 95°C for 30 seconds, 55°C (for exons 7 and 15) or 58°C (for exon 10) for 30 seconds, and 72°C for 30 seconds. There was a single final extension step at 72°C for 7 minutes. The PCR products were purified using the QIAquick PCR Purification kit (Qiagen Sciences, Germantown, MD). DNA concentration of PCR products was measured and adjusted to 20 ng per microliter. Sequencing of the purified PCR product was then performed using an ABI Prism 3100 Genetic Analyzer (Applied Biosystems, Foster City, CA).

## 3. Results

### 3.1. Epidemiology

A total of 72 specimens from 69 patients were identified. Among the 69 patients, 56 were male and 13 female. The



**Figure 1** Plasmacytoid urothelial carcinoma: 3 morphologic groups. A, Three cells types are present, including plasmacytoid (circles), monocytoid (arrows), and signet ring (arrowheads) (hematoxylin and eosin [H&E], original magnification  $\times 400$ ). B, A solid expansile nest of tumor cells is present beneath the luminal surface of the urinary bladder (H&E  $\times 100$ ). C, Classic category: small cells with discohesive cells and cells arranged in a linear pattern are present (H&E  $\times 200$ ). D, Pleomorphic category: pleomorphic cells with bizarre atypia are present (H&E  $\times 400$ ). E, Desmoplastic category: an exuberant desmoplastic response is present surrounding tumor cells (H&E,  $\times 100$ ). F, Pleomorphic cells with marked nuclear atypia with eccentric, eosinophilic cytoplasm, bordering on rhabdoid differentiation (H&E  $\times 400$ ).

median patient age was 68 years (range 36-91 years). Among the 72 specimens, 5 were biopsies, 13 were transurethral resections (TURs), 50 were radical cystectomies (RCs), 2 were nephroureterectomies, and 2 were ureterectomies. Among the 50 radical cystectomy cases, there were 41 radical cystoprostatectomies in men, 8 radical cystectomies with pelvic exenteration in women, and 1 radical cystectomy with vaginal margins in a woman who was already status post

hysterectomy. Among the biopsy/TUR specimens, 7 patients went on to have an RC. Of those 7 RC cases, 3 had plasmacytoid morphology and were included in the series. Four cases were excluded from analysis because 2 had no residual disease, 1 had urothelial carcinoma in situ only (no plasmacytoid morphology), and slides from 1 case (from 1996) could not be located for re-review and plasmacytoid/signet ring morphology was not mentioned in the pathology report.

**Table 2** Clinicopathological features of 72 cases of urothelial carcinoma with plasmacytoid differentiation

Total patients	69
Male	56 (81)
Female	13 (19)
Age median (range)	68 y (36-91 y)
Total cases (n)	72
Biopsy	5 (7)
TUR	13 (18)
Radical cystectomy	50 (69)
Nephroureterectomy	2 (3)
Ureterectomy	2 (3)
pT category <sup>a</sup>	
pT1	1 (2)
pT2 <sup>b</sup>	1 (2)
pT2a	2 (4)
pT2b	4 (7)
pT3a	7 (13)
pT3b	13 (24)
pT4 <sup>c</sup>	1 (2)
pT4a	23 (42)
pT4b	2 (4)
pN category <sup>a</sup>	
pN0	14 (26)
pN1	5 (9)
pN2	31 (58)
pN; not applicable	4 (7)
pM category <sup>a</sup>	
pM1	2 (4)
pM; not applicable	52 (96)
Positive ureteral margin (n = 54 resections)	14 (26)
Invasive carcinoma	9 (64)
Dysplasia	2 (14)
Soft tissue invasion	3 (22)
Positive urethral margin (n = 50 cystectomies)	2 (4)
Urothelial carcinoma in situ	1 (50)
Soft tissue invasion	1 (50)
Involvement of adjacent organs (n = 50 cystectomies)	26 (52)
Prostate	15 (58)
Seminal vesicle	15 (58)
Both prostate and seminal vesicle	10 (38)
Prostate only	5 (19)
Seminal vesicle only	5 (19)
Vagina	1 (4)
Cervix	2 (8)
Uterus	2 (8)
Ovary	1 (4)
Fallopian tube	2 (8)
Rectum	3 (12)
Skeletal muscle	1 (4)
Spermatic Cord	1
Median % plasmacytoid differentiation (range)	30% (5%-100%)
Other variant morphology present	23 (32)
Clear cell	1 (4)
Giant cell	1 (4)

**Table 2 (continued)**

Total patients	69
Glandular	3 (13)
Micropapillary	3 (13)
Nested	3 (13)
Sarcomatoid	7 (31)
Small cell	1 (4)
Squamous	4 (18)
Tumor size median (range)	4.7 cm (0.5-14.4 cm)
Concomitant urothelial carcinoma in situ present	31 (43)
Lymphovascular space invasion present	40

<sup>a</sup> Pathologic pTNM categories assigned to resection specimens (n = 54).

<sup>b</sup> Tumor involved muscularis propria only on initial transurethral resection (not on subsequent radical cystectomy), therefore not possible to reliably designate as T4a versus T4b.

<sup>c</sup> Nephroureterectomy, therefore no subcategories of T4 staging in the AJCC seventh edition.

<sup>d</sup> Spermatic cord was biopsied during a ureterectomy and was involved by tumor.

### 3.2. Gross findings

Among resection cases, 83% (45/54) had a mass and 17% (9/54) had no mass present. Three cases (5%, 3/54) had a scar, all of which did not have a grossly identifiable mass. The presence or absence of a scar was not stated in the remaining 94% (51/54) of cases. Among cystectomies, 34% (17/50) had a thickened bladder wall. Of note, the 2 ureterectomy specimens were described as having no gross abnormality. The 2 nephroureterectomies each had a mass described, but no scar or thickening of the renal pelvis wall/proximal ureter was noted.

### 3.3. Histopathology and morphologic groups

After reviewing all of the cases, 3 main morphologic groups were recognized. The first group was dubbed the “classic” group and was composed of cells somewhat resembling invasive lobular carcinoma of the breast. The cells had small to medium nuclei (generally 2 to 3 times larger than a lymphocyte), some of which were eccentrically located (plasmacytoid) and others were located in the center of the cell (monocytoid) (Fig. 1). Cells with signet ring–like morphology were also a feature of this group. Overall, the cells were singly scattered and discohesive or in loose aggregates forming linear cords. The second group was the “pleomorphic” group, in which cases were composed of cells with pleomorphic nuclei which were larger and had more atypia than the cells in the classic category (nuclei were generally at least 4 times the size of a lymphocyte). A subset of these cases had markedly bizarre nuclear atypia which at times was rhabdoid in appearance (prominent nucleolus, eccentric eosinophilic cytoplasmic inclusion). This group was usually seen in association with more typical plasmacytoid cells. The third group of cases was the “desmoplastic” group. Cells in this category were surrounded

by an exuberant, desmoplastic stromal response. Whereas a minority of the cells in the desmoplastic cases closely resembled the cells in the classic group, most of the cells were pleomorphic with more abundant cytoplasm. At times, the cells in this third group more closely resembled conventional high-grade urothelial carcinoma.

Overall, 29% (21/72) of cases were in the classic category, 35% (25/72) were in the exuberant desmoplasia category, and 36% (26/72) of cases were in the pleomorphic category. Among cases with an exuberant, desmoplastic stromal response, 84% (21/25) were composed of pleomorphic cells and 16% (4/25) were composed of cells with classic morphology. A total of 6 cases were noted to have marked/bizarre nuclear atypia, 3 of which were included in the category with exuberant desmoplasia and 3 of which were included in the pleomorphic category. Interestingly, a few of the cases in the classic category consisted of closely packed cells resembling a discohesive sheet, often located beneath the luminal surface of the urinary bladder.

All cases were initially diagnosed as urothelial carcinoma with plasmacytoid (93%, 67/72) or signet ring (7%, 5/72) differentiation. The amount of plasmacytoid differentiation ranged from 5% to 100% (median 25%). Additional variant differentiation was identified in 32% (23/72) of cases; additional variants included clear cell (4%, 1/23), giant cell (4%, 1/23), glandular (13%, 3/23), micropapillary (13%, 3/23), nested (13%, 3/23), sarcomatoid (31%, 7/23), small cell (4%, 1/23), and squamous differentiation (18%, 4/23). Concomitant urothelial carcinoma in situ (UCIS) was present in 43% (31/72) of specimens.

Overall, among resections, the pathologic T category was most commonly pT4 (48%, 26/54), followed by pT3 (37%, 20/54), pT2 (13%, 7/54), and pT1 (2%, 1/54) (Table 2). In the classic category, 14% (3/21) of cases were pT1, 33%

(7/21) pT2, 29% (6/31) pT3, and 24% (5/21) pT4. In the desmoplastic category, 0% (0/25) were pT1, 12% (3/25) pT2, 36% (9/25) pT3, and 52% (13/25) pT4. In the pleomorphic category, 11% (3/26) were pT1, 35% (9/26) pT2, 23% (6/26) pT3, and 31% (8/26) pT4.

A positive ureteral margin was present in of all resection cases (9 invasive carcinoma arising in the urothelium, 3 invasive in soft tissue, 2 urothelial dysplasia). When analyzed by morphologic group, positive ureteral margins were present in 25% (4/16) of classic resection cases, 32% (7/22) of desmoplastic resection cases, and 19% (3/16) of pleomorphic resection cases; of these cases, 2 classic cases, 1 desmoplastic case, and no pleomorphic cases were positive based on soft tissue invasion surrounding the ureteral lumen. The ureter was involved in 39% (21/54) of all resection cases (12 invasive in soft tissue only, 1 invasive arising from the urothelium only, 5 UCIS only, 1 urothelial dysplasia only, 1 invasive arising from the urothelium and in the soft tissue, and 1 invasive in soft tissue and UCIS). A positive urethral margin was present in 4% (2/50) of radical cystectomies (1 invasive carcinoma, 1 UCIS). The urethra was involved in 10% (5/50) of radical cystectomies (2 invasive carcinoma, 1 invasive in soft tissue, 2 UCIS).

Among resection cases, 93% (50/54) included lymph nodes, 72% (36/50) of which had lymph node metastases. Most cases with positive lymph nodes were N2 (83%, 30/36), and a smaller number were N1 (14%, 5/36) and N3 (3%, 1/36). The median number of positive lymph nodes was 4 (range 1-32), the median size of metastatic deposits was 1.3 cm (range 0.1-3.9 cm), and extranodal extension was present in 64% (23/36) of cases.

Among radical cystoprostatectomies/cystectomies, 52% (26/50) had involvement of adjacent organs (Table 3). Among radical cystoprostatectomies, 37% (15/41) had involvement of


**Table 3** Immunohistochemical stains performed in this study

Stain	Intensity	Proportion	Negative n (%)	Positive n (%)
	0 1 2 3	0 1 2 3 4		
β-Catenin (nuclear)	26 0 0 0	26 0 0 0 0	26 (100)	0 (0)
CD56	25 1 0 0	25 1 0 0 0	25 (96)	1 (4)
CD138	8 15 14 11	8 14 7 9 10	8 (17)	40 (83)
CDX2 (nuclear)	22 3 1 0	22 3 1 0 0	22 (85)	4 (15)
Keratin 7	4 3 8 11	4 7 1 2 12	4 (15)	22 (85)
Keratin 20	6 4 11 5	6 8 4 4 4	6 (23)	20 (77)
E-cadherin	27 9 6 6	27 10 3 3 5	27 (56)	21 (44)
Estrogen receptor	26 0 0 0	26 0 0 0 0	26 (100)	0 (0)
GATA3	1 4 4 17	1 4 3 1 17	1 (4)	25 (96)
Mammaglobin	26 0 0 0	26 0 0 0 0	26 (100)	0 (0)
P53	1 5 6 14	1 4 5 4 12	1 (4)	25 (96)
P63	10 1 10 5	10 8 2 2 4	10 (38)	16 (62)
Progesterone receptor	25 1 0 0	25 1 0 0 0	25 (96)	1 (4)
Thrombomodulin	21 2 3 0	21 5 0 0 0	21 (81)	5 (19)
TTF-1	25 0 1 0	25 0 1 0 0	25 (96)	1 (4)
Uroplakin	14 4 7 1	14 7 1 1 3	14 (54)	12 (46)


the prostatic stroma and 37% (15/41) of the seminal vesicles (10 cases with both prostate and seminal vesicles; 5 cases prostate only; 5 cases seminal vesicles only). In women who underwent radical cystectomy and pelvic exenteration, 25% (2/8) had involvement of the cervix, 25% (2/8) of the uterus, 25% (2/8) of the fallopian tube, 13% (1/8) of the ovary, and 13% (1/8) of the vagina. Among radical cystectomies, 6% (3/50) had involvement of the rectum and 2% (1/50) of the skeletal muscle of the abdominal wall. One patient who underwent ureterectomy had involvement of the spermatic cord (spermatic cord was biopsied during surgery).

### 3.4. Immunohistochemical stains

For this study, 14 immunohistochemical markers were performed on 26 cases (Table 3). Two additional stains, CD138 and E-cadherin, were performed on 48 cases. Among urothelial origin markers, GATA3 was the most sensitive (96%, 25/26), followed by p63 (62%, 16/26), uroplakin III (46%), and thrombomodulin (19%, 5/19). Keratin 7 (85%, 22/26) and keratin 20 (77%, 20/26) were positive in most cases. Breast carcinoma markers were usually negative; PR stained 1 case (intensity 1, proportion 1), whereas mammaglobin



**Figure 2** Immunohistochemical findings. A-C, Classic category (A, H&E  $\times 400$ ), with positive expression of CD138 (B, CD138  $\times 400$ ) and loss of E-cadherin (C, E-cadherin  $\times 400$ ). D, Desmoplastic category (D, H&E  $\times 200$ ), with retention of E-cadherin (D inset, E-cadherin  $\times 200$ ). E-F, Desmoplastic category (H&E  $\times 200$ ), with an absence of staining for CD138 (E inset, CD138  $\times 200$ ) and retention of staining for E-cadherin (F, E-cadherin,  $\times 400$ ).



**Figure 3** Overall survival for patients with plasmacytoid urothelial carcinoma. A, Overall survival (OS) for all patients with plasmacytoid urothelial carcinoma (median [IQR] OS: 20.99 [8.87-82.79], 69 total patients). B, OS of patients with plasmacytoid urothelial carcinoma by morphologic category ( $P = .0336$ ; median [IQR] OS classic: 20.63 [10.41-NR], 21 patients; median [IQR] OS desmoplastic: 11.56 [7.69-22.93], 25 patients; median [IQR] OS pleomorphic: 82.79 [17.74-85.55], 23 patients).

and ER were negative in all cases. Neuroendocrine markers CD56 and TTF-1 were each positive in 1 case (intensity 1, proportions 1 and 2, respectively). Gastrointestinal adenocarcinoma marker CDX2 showed nuclear positivity in 4 cases, whereas nuclear  $\beta$ -catenin staining was absent in all cases. Nuclear p53 staining was present in 96% (25/26) of cases. CD138 was positive in 83% (40/48), and E-cadherin was positive in 44% (21/48) of cases.

Notably, cases with associated exuberant desmoplasia were more likely to have retained expression of E-cadherin (68%, 13/19) (Fig. 2). In contrast, E-cadherin expression was retained only in a minority of classic (36%, 5/14) and pleomorphic (20%, 3/15) cases. Otherwise, differential staining between the morphologic categories was not noted. CD138 was positive in most cases (classic 10/14, desmoplastic

18/19, pleomorphic 12/15). Similarly, p63 was positive in most cases (classic 3/9, desmoplastic 7/8, pleomorphic 6/9).

### 3.5. Molecular

UroVysion FISH detected polysomy of chromosomes 3 (13%, 2/15), 7 (20%, 3/15), and 17 (20%, 3/15) and deletion of chromosome 9p21 (60%, 9/15). *FGFR3* mutations were detected by PCR in 60% (9/15) of cases (exon 7 in 6 cases, exon 10 in 6 cases, and exon 15 in 8 cases).

### 3.6. Clinicopathological results

A total of 69 patients were diagnosed with plasmacytoid urothelial cancer. Five patients presented with metastatic

disease, whereas 64 patients were diagnosed with presumed clinically localized plasmacytoid urothelial bladder cancer. Among patients with locoregional disease, 58 underwent radical cystectomy, and the remainder elected for bladder preservation approaches. At the time of cystectomy, 6 patients were found to have pathologically organ-confined disease ( $\leq$ pT2 N0), and 51 were diagnosed with non-organ-confined disease (pT3, T4, or N+), 2 of which were noted to have diffuse peritoneal carcinomatosis. Twenty-eight (40.5%) patients had positive surgical margins, most of which had extensive infiltrative disease not amenable to complete excision.

Neoadjuvant chemotherapy appeared to have little effect on this cohort. Of the 14 patients who received neoadjuvant cisplatin-based chemotherapy, 10 patients (71.4%) had lymph node involvement and 12 (85.7%) had pT3/4 disease at cystectomy. Only 1 patient was found to have a pathologic complete response with pT0N0. Thirteen patients received adjuvant chemotherapy, 2 of which received neoadjuvant chemotherapy as well.

Median survival for all patients with plasmacytoid urothelial cancer from the time of initial diagnosis was 18 months (interquartile range [IQR] 8–78 months) (Fig. 3A) and was 14 (IQR 5–74) months from the time of cystectomy for patients undergoing radical cystectomy. Of note, there appeared to be 3 morphologic subtypes of plasmacytoid disease: classic, desmoplastic, and pleomorphic, with distinct survival outcomes ( $P = .083$ ). Patients with desmoplastic disease tended to have the worst clinical outcomes (median survival 10 months), whereas pleomorphic and classic subtypes had somewhat better survival outcomes (Fig. 3B).

#### 4. Discussion

The plasmacytoid/diffuse variant is included with a number of variants with established prognostic or potential clinical significance outlined in the latest edition of the *WHO Classification of Tumours of the Urinary System and Male Genital Organs* [1]. The WHO, International Consultation on Urologic Disease, European Association of Urology, AJCC, and the College of American Pathologists have all recommended documenting the presence of divergent differentiation/variant morphology in pathology reports [1,27–29]. However, the difficulty of detecting the plasmacytoid variant both in community practices and in large academic medical centers is well described in the literature [30,31]. With this in mind, we report the clinicopathological features of a large, single-institution case series of urothelial carcinoma with plasmacytoid/diffuse differentiation.

A spectrum of morphologic features exists in these cases, a finding which has been suggested but not extensively explored in the literature. After reviewing all of the cases for this study, 3 main morphologic groups emerged: (1) classic (29% of cases); (2) desmoplastic (35% of cases); and (3) pleomorphic (36% of cases). Some authors have previously applied the term *rhabdoid* to cases with marked pleomorphism [22–25,32].

Notably, a total of 6 cases in our series were noted to have marked/bizarre nuclear atypia, but we considered these changes to be within the spectrum of plasmacytoid differentiation (3 placed in the desmoplastic category, 3 in the pleomorphic category).

We applied an extensive immunohistochemical panel to a large number of cases processed in the same laboratory. Caution should be exercised when using immunohistochemistry to determine whether or not the tumor is urothelial in origin because some stains such as PR for breast carcinoma (1 case; intensity 1 and proportion 1) and CDX2 for gastrointestinal carcinoma (4 cases; intensity 1–2 and proportion 1–2) can occasionally be positive in the plasmacytoid variant. Staining for ER, mammaglobin, and nuclear  $\beta$ -catenin was notably absent in all cases. Finally, all tested urothelial origin markers were positive, with GATA3 (96%) being the most frequent. A recent study showed similar findings to ours (occasional positive CDX2 and PR), and in addition, they report staining for GCDFP-15 (24%, 11/45 cases positive) and p-CEA (49%, 22/45 cases) [33].

Loss of E-cadherin and positive staining for CD138 have been touted as useful findings suggestive of plasmacytoid urothelial carcinoma. However, a recent study calls into question the utility of CD138 staining given that positivity in benign urothelium and a variety of benign and malignant urothelial tumors was found [34]. We observed a loss of E-cadherin in the majority of cases (57%) and CD138 positivity in 83% of cases. Notably, cases with associated exuberant desmoplasia were more likely to have retained expression of E-cadherin (68%, 13/19). In contrast, E-cadherin expression was retained only in a minority of classic (36%, 5/14) and pleomorphic (20%, 3/15) cases. Otherwise, differential staining between the morphologic categories was not noted.

The median overall survival for the entire cohort was 21 months. Subsequent analysis of overall survival for each morphologic category showed that there was a statistically significant difference between the groups. Cases within the desmoplastic category had the worst overall survival (median 12 months). Notably, 88% (22/25) of cases within the desmoplastic category were high stage (pT3 or pT4) in comparison to 48% (10/21) of cases in the classic category and 54% (14/26) of cases in the pleomorphic category. Although conclusions are difficult to draw given that the number of cases in each category is small, this raises the question of whether cases with desmoplasia inherently have a worse outcome or if they have a similar prognosis when matched stage for stage with the other groups. The association of worse outcome with certain morphologic features has been alluded to in a prior study by Jimenez et al [26]. The authors identified 3 architectural patterns (nodular, trabecular, infiltrative) within the invasive component of urothelial carcinoma and found that cases with any amount of an infiltrative pattern (narrow cords or single cells, desmoplasia, necrosis) had a worse outcome (median survival 29 months versus 85 months for cases without the infiltrative pattern) [26]. Although their findings did not reach statistical significance, they do offer further support that the presence

of desmoplasia in urothelial carcinoma could suggest a worse prognosis.

The molecular aberrations that give rise to urothelial carcinoma with plasmacytoid differentiation are still largely unknown. One study that used UroVysion FISH found a homozygous deletion of chromosome 9p21 in 15% of cases and a relative deletion in 70% of cases; therefore, the authors suggested that chromosome 9 may play an important role in the pathogenesis of this variant [8]. We performed UroVysion FISH on 15 cases and found chromosome 9 to be most commonly aberrant, with a deletion of chromosome 9p21 in 60% of cases. A prior study using mutation analysis via capillary electrophoresis did not show any mutations of *FGFR3*; however, we detected *FGFR3* mutations in 60% of cases [8]. The significance of these findings needs to be further elucidated in future studies.

In conclusion, it is important for the practicing pathologist to be aware of the range of morphologic features of the plasmacytoid variant and to be aware of the pitfalls related to immunohistochemical staining. Finally, further studies regarding the association between desmoplasia and outcome as well as the molecular aberrations leading to the plasmacytoid variant are warranted.

## References

- Moch H, Humphrey PA, Ulbright TM, Reuter VE. World Health Organization classification of tumours of the urinary system and male genital organs. Lyon: International Agency for Research on Cancer; 2016.
- Monn MF, Kaimakliotis HZ, Pedrosa JA, et al. Contemporary bladder cancer: variant histology may be a significant driver of disease. *Urol Oncol* 2015;33(18.e):15-20.
- Kaimakliotis HZ, Monn MF, Cheng L, et al. Plasmacytoid bladder cancer: variant histology with aggressive behavior and a new mode of invasion along fascial planes. *Urology* 2014;83:1112-6.
- Rice KR, Koch MO, Kao CS, et al. Lymph node metastases in patients with urothelial carcinoma variants: influence of the specific variant on nodal histology. *Urol Oncol* 2015;33(20):e23-9.
- Saphir O. Signet-ring cell carcinoma of the urinary bladder. *Am J Pathol* 1955;31:223-31.
- Grignon DJ, Ro JY, Ayala AG, et al. Primary signet-ring cell carcinoma of the urinary bladder. *Am J Clin Pathol* 1991;95:13-20.
- Fritsche HM, Burger M, Denzinger S, et al. Plasmacytoid urothelial carcinoma of the bladder: histological and clinical features of 5 cases. *J Urol* 2008;180:1923-7.
- Keck B, Stoehr R, Wach S, et al. The plasmacytoid carcinoma of the bladder—rare variant of aggressive urothelial carcinoma. *Int J Cancer* 2011;129:346-54.
- Kim B, Kim G, Song B, et al. Her2 protein overexpression and gene amplification in plasmacytoid urothelial carcinoma of the urinary bladder. *Dis Markers* 2016;2016:8463731.
- Liang Y, Heitzman J, Kamat AM, et al. Differential expression of gata-3 in urothelial carcinoma variants. *HUM PATHOL* 2014;45:1466-72.
- Lim MG, Adsay NV, Grignon DJ, et al. E-cadherin expression in plasmacytoid, signet ring cell and micropapillary variants of urothelial carcinoma: comparison with usual-type high-grade urothelial carcinoma. *Mod Pathol* 2011;24:241-7.
- Lopez-Beltran A, Requena MJ, Montironi R, et al. Plasmacytoid urothelial carcinoma of the bladder. *HUM PATHOL* 2009;40:1023-8.
- Nigwekar P, Tamboli P, Amin MB, et al. Plasmacytoid urothelial carcinoma: detailed analysis of morphology with clinicopathologic correlation in 17 cases. *Am J Surg Pathol* 2009;33:417-24.
- Paner GP, Annaiah C, Gulmann C, et al. Immunohistochemical evaluation of novel and traditional markers associated with urothelial differentiation in a spectrum of variants of urothelial carcinoma of the urinary bladder. *HUM PATHOL* 2014;45:1473-82.
- Raspolini MR, Sardi I, Giunti L, et al. Plasmacytoid urothelial carcinoma of the urinary bladder: Clinicopathologic, immunohistochemical, ultrastructural, and molecular analysis of a case series. *HUM PATHOL* 2011;42:1149-58.
- Ro JY, Shen SS, Lee HI, et al. Plasmacytoid transitional cell carcinoma of urinary bladder: a clinicopathologic study of 9 cases. *Am J Surg Pathol* 2008;32:752-7.
- Fox MD, Xiao L, Zhang M, et al. Plasmacytoid urothelial carcinoma of the urinary bladder: a clinicopathologic and immunohistochemical analysis of 49 cases. *Am J Clin Pathol* 2017;147:500-6.
- Keck B, Wach S, Kunath F, et al. Nuclear e-cadherin expression is associated with the loss of membranous e-cadherin, plasmacytoid differentiation and reduced overall survival in urothelial carcinoma of the bladder. *Ann Surg Oncol* 2013;20:2440-5.
- Shimada K, Nakamura M, Ishida E, et al. Urothelial carcinoma with plasmacytoid variants producing both human chorionic gonadotropin and carbohydrate antigen 19-9. *Urology* 2006;68(891):e7-10.
- Mai KT, Park PC, Yazdi HM, et al. Plasmacytoid urothelial carcinoma of the urinary bladder report of seven new cases. *Eur Urol* 2006;50:1111-4.
- Cockerill PA, Cheville JC, Boorjian SA, et al. Outcomes following radical cystectomy for plasmacytoid urothelial carcinoma: defining the need for improved local cancer control. *Urology* 2017;102:143-7.
- Parwani AV, Herawi M, Volmar K, et al. Urothelial carcinoma with rhabdoid features: report of 6 cases. *HUM PATHOL* 2006;37:168-72.
- Inagaki T, Nagata M, Kaneko M, et al. Carcinosarcoma with rhabdoid features of the urinary bladder in a 2-year-old girl: possible histogenesis of stem cell origin. *Pathol Int* 2000;50:973-8.
- Duvdevani M, Nass D, Neumann Y, et al. Pure rhabdoid tumor of the bladder. *J Urol* 2001;166:2337.
- Harris M, Eyden BP, Joglekar VM. Rhabdoid tumour of the bladder: a histological, ultrastructural and immunohistochemical study. *Histopathology* 1987;11:1083-92.
- Jimenez RE, Gheiler E, Oskanian P, et al. Grading the invasive component of urothelial carcinoma of the bladder and its relationship with progression-free survival. *Am J Surg Pathol* 2000;24:980-7.
- Amin MB, Smith SC, Reuter VE, et al. Update for the practicing pathologist: the International Consultation on Urologic Disease—European Association of Urology consultation on bladder cancer. *Mod Pathol* 2015;28:612-30.
- Bernard H, Bochner DEH, Efstathiou Jason A, et al. Urinary bladder. In: Amin MB, editor. *AJCC cancer staging manual*. Springer; 2016. p. 757-65.
- Amin MB. Protocol for the examination of specimens from patients with carcinoma of the urinary bladder. College of American Pathologists; 2013.
- Shah RB, Montgomery JS, Montie JE, et al. Variant (divergent) histologic differentiation in urothelial carcinoma is under-recognized in community practice: impact of mandatory central pathology review at a large referral hospital. *Urol Oncol* 2013;31:1650-5.
- Linder BJ, Boorjian SA, Cheville JC, et al. The impact of histological reclassification during pathology re-review—evidence of a Will Rogers effect in bladder cancer? *J Urol* 2013;190:1692-6.
- Perez-Montiel D, Wakely PE, Hes O, et al. High-grade urothelial carcinoma of the renal pelvis: clinicopathologic study of 108 cases with emphasis on unusual morphologic variants. *Mod Pathol* 2006;19:494-503.
- Borhan WM, Cimino-Mathews AM, Montgomery EA, et al. Immunohistochemical differentiation of plasmacytoid urothelial carcinoma from secondary carcinoma involvement of the bladder. *Am J Surg Pathol* 2017;41:1570-5.
- Goto K. Cd138 expression is observed in the urothelial epithelium and in various urothelial carcinomas, and cannot be evidence for plasmacytoid urothelial carcinoma. *Int J Surg Pathol* 2016;24:614-9.