

**Results:** The cost of managing open lower limb fractures was £13,959/patient in 2016 and £12,005/patient in 2017. Remuneration in 2016 was £5,196/patient; whilst remuneration in 2017 was £10,707/patient. The clinically dominant procedure determined by HRG4+ was in orthopaedic surgery 70.3% in 2016 and 44.1% in 2017; yet, they received 40.5% of the income in 2016 and 44.2% in 2017. Simulation revealed that income under new HRG4+ tariffs increased by 25.1%, where improving coding accuracy yielded a further increase of 29.1% in income.

**Conclusions:** Providing complex lower limb reconstruction for open fractures in a multidisciplinary unit is resource intensive and associated with high costs. Current hospital remuneration for providing orthoplastic services is insufficient and inadequate. To ensure that orthoplastic units across NHS England can operate efficiently and to improve patient care, more accurate coding and an increase in national tariffs are essential. Remuneration goes to a single clinical specialty despite a multidisciplinary service provision and local assessment of parity is required.

<https://doi.org/10.1016/j.injury.2018.11.051>

### Best Poster Presentation

#### Increasing capture of patient-reported outcomes in trauma research



Grace Turner<sup>1</sup>, Ameeta Retzer<sup>1</sup>, Anita Slade<sup>1</sup>, Christel McMullan<sup>1,\*</sup>, Derek Kyte<sup>1</sup>, Karen Piper<sup>2</sup>, Tony Belli<sup>2</sup>, Melanie Calvert<sup>1</sup>

<sup>1</sup> University of Birmingham, Birmingham, United Kingdom

<sup>2</sup> National Institute for Health Research Surgical Reconstruction and Microbiology Research Centre, Birmingham, United Kingdom

**Background:** In order to understand how peoples' quality of life is affected following major trauma and the effects of that injury on their health and wellbeing, it is important to capture patients' perspectives of their own health. Patient Reported Outcome Measures (PROMs) can capture patients' own experience of their health such as symptoms, mobility, mental health and social function.

The aim of this research is to establish the impact of trauma on quality of life/symptoms and to explore views on using PROMs to support clinical care and research.

**Methods:** One-to-one, semi-structured interviews will be conducted with: (i) people who have experienced a major trauma, (ii) their family members/carers; (iii) healthcare professionals working in trauma related clinical areas; (iv) trauma researchers; and (v) staff members/volunteers from third sector organisations who support trauma patients and their families/carers.

**Results:** This is an ongoing study based at the Centre for Patient Reported Outcomes Research (CPROR) funded by the National Institute for Health Research Surgical Reconstruction and Microbiology Research Centre (NIHR SRMRC).

Findings will be used to inform the development of a pathway for the electronic capture of PROMs for inclusion within routine clinical care of trauma patients and trauma research. Future research will test the feasibility and acceptability of this ePROM system.

The research programme is being delivered in close collaboration with key stakeholders, including patient partners, trauma clinicians, trauma researchers and the ministry of defence.

**Conclusion:** The rising number of major trauma survivors has driven the need for improvements in rehabilitation to enable patients to return to functional activities, work and education after complex re-enablement and reconstructive surgery. PROMs

are essential to deliver patient-centred healthcare and research which is informed by patient-focused priorities/outcomes. The programme will also increase capacity for trauma-specific knowledge and expertise in relation to PROMs.

<https://doi.org/10.1016/j.injury.2018.11.052>

### Best Student Presentation

#### Analysis of the relationship between Vitamin D levels and Infection in Orthopaedic patients



Alexander Zargaran<sup>1</sup>, Alex Trompeter<sup>2,\*</sup>

<sup>1</sup> St. George's, University of London, London, United Kingdom

<sup>2</sup> St George's Hospital, London, United Kingdom

**Background:** One in four of the United Kingdom population suffers from low Vitamin D levels. Vitamin D has immunomodulatory properties, but its precise role in Orthopaedic infection is unclear. This study aimed to quantify average Vitamin D levels among Orthopaedic patients and elucidate the relationship between Vitamin D levels and incidence of infection.

**Methods:** A convenience sample of 187 Orthopaedic patients was taken in our institution, which is a tertiary referral centre for Orthopaedic infection. 25OHD concentration and infection status were recorded, and a Mann-Whitney *U* test for non-parametric data was performed on the means. The relationship was then validated through a bivariate correlation analysis (Spearman's rho).

**Results:** 104 patients had infection. Mean ages were 64.8 years in patients with infection and 63.0 years in patients without infection. Gender split was approximately equal in both groups. There was no significant difference in age or gender between both groups. Mean 25OHD concentration was 39.8 nmol/L for patients with infection and 59.6 nmol/L for patients without infection ( $p < 0.00$ ). Overall mean 25OHD concentration for Orthopaedic patients was 48.6 nmol/L. The correlation coefficient between 25OHD levels and infection incidence was  $-0.3$  ( $p < 0.00$ ).

**Conclusion:** There was a negative correlation between 25OHD concentration and infection, suggesting that Vitamin D could have a protective effect against infection. Furthermore, patients without infection had a mean of 19.8 nmol/L higher concentration of 25OHD than patients with infection. Patients with infection had 25OHD insufficiency, whilst patients without infection had normal 25OHD levels. Future RCTs are needed to determine whether Vitamin D supplementation reduces incidence of infection and leads to improved outcomes in Orthopaedic patients.

**Implications:** These findings suggest a potential future role for prophylactic Vitamin D supplementation to help combat the Vitamin D insufficiency prevalent in Orthopaedic patients, as well as in the prevention of infection during the hospital stay.

<https://doi.org/10.1016/j.injury.2018.11.053>

### Best Scientific Presentation

#### Intra-medullary Nail Insertion Accuracy: A comparison of the Infra-patellar and Supra-patellar approach



Thomas Anderson\*, Philip Beak, Alex Trompeter

Trauma & Orthopaedics, St George's University Hospitals NHS Foundation Trust, London, United Kingdom

**Aims:** The anatomical safe zone for intra-medullary nail insertion through the tibial plateau is small, insertion outside of this area