



Corrigendum

Corrigendum to “Mast cells improve functional recovery of transected peripheral nerve: A novel preliminary study” [Injury 48/7 (2017) 1480-1485]

Behrooz Ilkhanizadeh^a, Leila Zarei^{b,*}, Negin Farhad^c, Mehran Bahrami-Bukani^c, Rahim Mohammadi^d

^a Department of Pathology, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran

^b Department of Anatomical Sciences, Faculty of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

^c Solid Tumor Research Center, Urmia University of Medical Sciences, Urmia, Iran

^d Department of Surgery and Diagnostic Imaging, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

The authors regret that Fig. 2 was incorrect. Please see below for the correct figure and figure legend. The authors would like to apologise for any inconvenience caused.




Fig. 2. Characterization of bone marrow derived mast cells (BMMC) after 3 weeks. Flow cytometric analyses of cell surface markers showed that the cells were positive for BMMC-related antigens and for BMMC-related double positive cells.

DOI of original article: <http://dx.doi.org/10.1016/j.injury.2017.05.015>

* Corresponding author at: Solid Tumor Research Center, Urmia University of Medical Sciences, Urmia, Iran.

E-mail address: Zarei.l@lums.ac.ir (L. Zarei).

<http://dx.doi.org/10.1016/j.injury.2018.11.012>

0020-1383/© 2018 Elsevier Ltd. All rights reserved.