

## Review

## New pathologic mechanisms in nucleotide repeat expansion disorders

Rodriguez C.M.<sup>a,b</sup>, Todd P.K.<sup>a,c,\*</sup><sup>a</sup> Department of Neurology, University of Michigan, Ann Arbor, MI, USA<sup>b</sup> Department of Genetics, Stanford University, Stanford, CA, USA<sup>c</sup> VA Ann Arbor Healthcare System, Ann Arbor, MI, USA

## A B S T R A C T

Tandem microsatellite repeats are common throughout the human genome and intrinsically unstable, exhibiting expansions and contractions both somatically and across generations. Instability in a small subset of these repeats are currently linked to human disease, although recent findings suggest more disease-causing repeats await discovery. These nucleotide repeat expansion disorders (NREs) primarily affect the nervous system and commonly lead to neurodegeneration through toxic protein gain-of-function, protein loss-of-function, and toxic RNA gain-of-function mechanisms. However, the lines between these categories have blurred with recent findings of unconventional Repeat Associated Non-AUG (RAN) translation from putatively non-coding regions of the genome. Here we review two emerging topics in NREs: 1) The mechanisms by which RAN translation occurs and its role in disease pathogenesis and 2) How nucleotide repeats as RNA and translated proteins influence liquid-liquid phase separation, membraneless organelle dynamics, and nucleocytoplasmic transport. We examine these topics with a particular eye on two repeats: the CGG repeat expansion responsible for Fragile X syndrome and Fragile X-associated Tremor Ataxia Syndrome (FXTAS) and the intronic GGGGCC repeat expansion in *C9orf72*, the most common inherited cause of amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD). Our thesis is that these emerging disease mechanisms can inform a broader understanding of the native roles of microsatellites in cellular function and that aberrations in these native processes provide clues to novel therapeutic strategies for these currently untreatable disorders.

## 1. Repetitive elements in the genome

Repetitive regions of DNA comprise over half of the human genome, and can span millions of bases (Treangen and Salzberg, 2011; Lander et al., 2001). The exact composition and variation of repeat DNA within the genome is unknown, as current sequencing technology is sub-optimal for resolving these sequences (Treangen and Salzberg, 2011). Within this repetitive genome (the “repeatome”) exist two broad categories of repetitive elements: Tandem repeats and interspersed repeats (Hannan, 2018b). Tandem repeats are regions of DNA with sequential blocks of patterns of nucleotides confined to a specific locus (Dumbovic et al., 2017). An estimated 17% of human genes contain tandem repeats within open reading frames (Gemayel et al., 2010) and they are common components of intragenic regions such as introns and 5' and 3' untranslated regions (UTRs). As such, tandem repeats are of particular interest as a potential source of genetic variation and function. The other, larger category of repetitive DNA, interspersed repeats, are noncontiguous structures dispersed throughout the genome, and include transposons and retropseudo genes (Smit, 1999). Interspersed repeats will not be further discussed in this review.

Satellite DNA is a subcategory of tandem repeats. The term “satellite” was given based on the experimental finding that a distinct band of DNA separated from the principle DNA band after density gradient

centrifugation of mouse genomic DNA (Kit, 1962). It was hypothesized that the distinct “satellite” band arose due to differences in base composition, which is directly related to the buoyant density of native DNA (Kit, 1962; Jones, 1973). Through biochemical analysis, these satellite sequences were discovered to be largely repetitive (Britten and Kohne, 1968; Waring and Britten, 1966). It was soon concluded that repetitive regions were common but species-specific in prokaryotes and eukaryotes, including humans, and had discrete subcellular localization patterns at sites of heterochromatin (Britten and Kohne, 1968; Yunis and Yasmin, 1971; Holmquist and Dancis, 1979). Satellite DNA is further categorized into microsatellites, minisatellites, and macrosatellites by size of the repeat unit (Dumbovic et al., 2017). Macrosatellites primarily occupy telomeres and centromeres (Jones, 1970; Pardue and Gall, 1970), while minisatellites and microsatellites are present throughout the genome (Lander et al., 2001).

Human genome sequencing allowed for further characterization and quantification of these repetitive regions (Lander et al., 2001). Microsatellites (which we define here as 1-8 or more bases per repeat unit) reside in both coding and noncoding regions, and comprise around 3% of the genome (Lander et al., 2001; Richard et al., 2008). These repetitive units are also known as short tandem repeats (STR) or simple sequence repeats (SSR). Microsatellites likely arose through polymerase template slippage during transcription, and they are considered

\* Corresponding author.

E-mail address: [pertertod@umich.edu](mailto:pertertod@umich.edu) (P.K. Todd).<https://doi.org/10.1016/j.nbd.2019.104515>

Received 19 February 2019; Received in revised form 7 June 2019; Accepted 19 June 2019

Available online 21 June 2019

0969-9961/ Published by Elsevier Inc.

unstable due to their tendency to change size both somatically and between generations (Lander et al., 2001; Kruglyak et al., 1998). Thus, distinct microsatellites are often polymorphic within populations and can be expanded or contracted within families. The degree of variance and instability across all microsatellites in humans is currently unknown. Microsatellite polymorphisms in coding regions of eukaryotic genes have been linked to a number of roles from modulating cell-cell interactions in *S. cerevisiae*, circadian rhythms in *D. melanogaster*, body morphology in canines, and more recently an association between the length of a CAG trinucleotide repeat in the huntingtin gene (HTT) and cognitive performance in humans (Verstrepen et al., 2005; Gemayel et al., 2010; Lee et al., 2018; Hannan, 2018a; Fondon 3rd and Garner, 2004). While microsatellites in promoter regions of genes can greatly influence gene expression (Gemayel et al., 2010; Vines et al., 2009; Sawaya et al., 2013), the full range of mechanisms by which variations in repeat size influence gene and genome biology is an area of active study (Bagshaw, 2017; Usdin, 2008).

2. Microsatellites in neurodegenerative disease: shared and divergent molecular mechanisms that depend on repeat size, content and context

Instability and expansion of specific microsatellites is the genetic basis of over 40 nucleotide repeat expansion diseases (NREDs) (Fig. 1). The vast majority of NREDs affect the nervous system, often leading to neurodegeneration (Paulson, 2018a). Disease causing microsatellites are traditionally categorized into three groups: dominantly inherited repeat expansions within protein coding sequences, dominantly inherited repeat expansions outside of known protein coding sequences, and recessive or X-linked repeat expansions that impact the expression of the genes in which they reside. Most of the disease-causing repeats present within protein coding exons are CAG trinucleotides encoding for glutamine. These polyglutamine diseases include Huntington's Disease (HD), spinal and bulbar muscular atrophy (SBMA), and several spinocerebellar ataxias (SCA). The resulting polyglutamine-containing proteins form insoluble aggregates, which place a proteotoxic burden on cells (La Spada et al., 1991; Trotter et al., 1995a, 1995b; Trotter et al., 1995a, 1995b; Paulson, 2018a; Bates, 1996). Microsatellite expansion mutations can also result in RNA mediated toxicity, with

myotonic dystrophy type 1 (DM1) as the standout example. Symptoms and severity in DM1 patients range from mild myotonia and cataracts to congenital forms which are often fatal due to respiratory deficits (Kamsteeg et al., 2012). In DM1, a CTG repeat resides in the 3' UTR of the DMPK gene. This mutation does not meaningfully alter the abundance of DMPK, rather it causes RNA foci formation and sequestration of key proteins (Pettersson et al., 2015; Miller et al., 2000; Mankodi et al., 2001; Jiang et al., 2004). Microsatellite expansions can also elicit a loss of function in the genes in which they reside. Friedreich's ataxia (FRDA) is the most common autosomal recessive ataxia, characterized by early adulthood onset of symptoms such as progressive ataxia with sensory loss (Paulson, 2018a). In Friedreich's ataxia, a GAA repeat expansion in an intron of the *FXN* gene leads to a change in the epigenetic state of the *FXN* locus that strongly suppresses the mRNA transcription (Campuzano et al., 1996). This leads to a partial, but not complete, loss of *FXN* mRNA and Frataxin protein, and impaired mitochondrial function and iron homeostasis (Akbar and Ashizawa, 2015). Unlike the dominant disorders described above, Friedreich's ataxia requires expansion or mutations in both *FXN* alleles, and the size of the repeat correlates inversely with the degree of mRNA expression and the severity of the clinical phenotype (Castaldo et al., 2008).

While certain microsatellite expansions likely share underlying pathologic mechanisms, the convergence and divergence of phenotypes and genotypes across a range of microsatellites presents intriguing questions about these diverse disorders. The 28 dominantly inherited spinocerebellar ataxias are all progressive disorders characterized by cerebellar degeneration and clinically by loss of motor coordination (Soong and Paulson, 2007). Six of the SCAs are caused by a CAG repeat expansion in coding regions of the genes in which they reside (SCA1, SCA2, SCA3, SCA7, DRPLA and SCA17) leading to polyglutamine expansions within six different protein coding genes, and an additional two SCAs are caused by CAG/CTG repeat expansions in putatively noncoding regions (SCA8 and SCA12) (Paulson et al., 2017). Early in the course of disease, specific clinical features can be used to separate the six polyglutamine SCAs phenotypically with varying degrees of certainty (Shakkottai and Fogel, 2013). In addition, classic late-onset HD exhibits a very different clinical and pathological phenotype compared to the SCAs. As all of these genes are expressed ubiquitously within the nervous system, their divergent phenotypes are thought to




Fig. 1. Nucleotide repeat expansion disorders. Disorders are listed by their genomic location on an illustrated simplified gene. Each disorder is accompanied by the unstable repeat sequence that elicits disease. The prevailing disease mechanism is listed below each group to illustrate that genomic location of the expansion influences but does not completely determine the potential effects of any given repeat. Abbreviations: Fragile X-associated tremor/ataxia syndrome (FXTAS), Fragile X-primary ovarian insufficiency (FXPOI), progressive myoclonic epilepsy type 1/Unverricht-Lundborg disease (EPM1), spinocerebellar ataxia (SCA), neuronal intranuclear inclusion disease (NIID), dentatorubral-pallidolysian atrophy (DRPLA), spinal-bulbar muscular atrophy (SBMA), oculopharyngeal muscular dystrophy (OPMD), Huntington disease-like 2 (HDL2), amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), myotonic dystrophy (DM), Fuchs endothelial corneal dystrophy (FECD), cerebellar ataxia, neuropathy and vestibular areflexia syndrome (CANVAS), benign adult familial myoclonic epilepsy (BAFME), and FRA2A, FRA7A and FRA7A refer to disease associated fragile sites in the genome that result from CGG repeat expansions.

emerge through specific aberrations in the native functions of the proteins in which they reside. Molecular evidence for this arises from differences in the repeat length thresholds for disease between different CAG repeat loci as well as the ability of specific mutations within these same proteins (most notably Ataxin 1) to recapitulate key features of disease absent expansion of the repeat element (Duvick et al., 2010). Perhaps the clearest example of native gene context influence on CAG repeat toxicity is seen in SBMA, where expansions within the androgen receptor gene only elicit toxicity in the setting of exposure to the receptor ligand, testosterone (Lieberman et al., 2019; Katsuno et al., 2003; Katsuno et al., 2002). However, all polyglutamine NREDS share similar pathological features, including intranuclear ubiquitinated neuronal inclusions of the polyglutamine protein, and inclusions are often observed in brain regions which do not exhibit marked neurodegeneration or significant clinical phenotypes (Lieberman et al., 2019). Moreover, at larger repeat sizes, a series of shared clinical features (dystonia, dementia, and spasticity) emerge in both HD and SCAs (Paulson, 2018b). This convergence of clinical phenotypes may reflect a larger contribution of gene context-independent toxicity elicited by very large polyglutamine stretches. Consistent with this, expression of large polyglutamine expansions outside of a native gene context is sufficient to elicit inclusions and neurodegeneration across a series of model systems (Morley et al., 2002; Warrick et al., 1998; Ordway et al., 1997). Thus, for each polyglutamine NRED, it is likely the combinatorial effects of gene context dependent toxicity and polyglutamine expansion context independent toxicity that drive neurodegeneration, with the relative influence of each component dependent on repeat size and the strength of the effects of the native gene context.

Like DM1, DM2 (myotonic dystrophy type 2) is caused by a microsatellite repeat expansion in a noncoding region, however in this case it is a CCTG repeat in the first intron of the CNBP gene (Liquori et al., 2001). In their adult forms, there is significant overlap in the clinical features of DM2 and DM1, with some specific differences (such as the distribution of muscle weakness and dystrophy) usually allowing for clinical distinction of the two conditions. However, molecularly, the length of the DM1 CUG repeat expansion directly correlates with disease severity, which shows marked genetic anticipation and qualitatively different phenotypes in congenital cases with large expansions (Tsiflidis et al., 1992). This is not the case for DM2 which has a mean expansion length of 5,000 repeats but no clear congenital or juvenile form of the disease and a much less clear relationship between repeat size and age of onset (Kamsteeg et al., 2012; Paulson, 2018a). Despite this clear clinical discrepancy, the mechanism of toxicity in DM2 is thought to be the same as that for DM1, with the repeat binding to and sequestering the RNA binding protein Muscleblind (Margolis et al., 2006). Similar splicing changes are observed in both conditions and in disease models expressing the two repeats (Nakamori et al., 2013). The underlying cause of the differences in these two disorders despite apparently shared disease mechanisms remains unclear, with potential roles of the neighboring genes in each repeat locus as well as non-muscleblind interacting partners for each repeat postulated as possible contributors (Meola and Cardani, 2015).

A striking example of genetic convergence with phenotypic divergence is the G<sub>4</sub>C<sub>2</sub> hexanucleotide expansion in *C9orf72* which is the most prevalent genetic cause of the two clinically separate diseases amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) (Renton et al., 2011; DeJesus-Hernandez et al., 2011). ALS is a progressive neurodegenerative disease affecting neurons in the brain and spinal cord, whereas clinical phenotypes of FTD are most often forms of cognitive impairment such as disinhibition, apathy, changes in personality, and language dysfunction (Orr, 2011; Paulson, 2018a). These two diseases were known to be linked prior to this genetic finding through the study of families with ALS-FTD, and are also genetically linked through single nucleotide polymorphisms not associated with a microsatellite in the *UBQLN2* gene (Deng et al., 2011). We will discuss major findings associated with the G<sub>4</sub>C<sub>2</sub> hexanucleotide expansion in

this review, but it is worth noting that the molecular underpinnings of this clinical divergence between ALS and FTD remains mysterious.

While much focus has been placed on the pathogenicity of microsatellites, the fact that their portion of the genome is about three-times greater than that of protein-coding regions suggests that microsatellites are likely to also influence the normal functions of the genes in which they reside (Consortium, ENCODE project, 2012). Our knowledge and understanding of microsatellites has grown recently with new discoveries related to the ways in which these repetitive regions can affect cellular function. This includes the topics of liquid-liquid phase separation to dictate membraneless organelle dynamics, nucleocytoplasmic transport and Repeat Associated Non-AUG (RAN) translation mediated production of cryptic proteins from repeat containing transcripts. In this review, we will outline the recent research on disease causing microsatellites, and discuss how this research has expanded our understanding of how microsatellites can function in the cell.

### 3. RAN translation of microsatellite repeats in non-coding and coding regions of mRNA

Microsatellites are capable of forming stable structures in RNA. While this can confer a toxic gain-of-function through direct binding with proteins (Renoux and Todd, 2012; Todd and Paulson, 2010), it also supports a newly recognized unconventional mode of translation initiation. This process is termed Repeat Associated Non-AUG (RAN) translation, whereby proteins are synthesized from microsatellite containing mRNA in the absence of an AUG start codon (Zu et al., 2011). This discovery was paradigm shifting as diseases caused by microsatellite expansions in noncoding regions were now looked at as having the potential to encode repetitive stretches of amino acids like those in polyglutamine disorders. RAN translation was first described by Zu et al. (2011) through work on spinocerebellar ataxia 8 (SCA8) where expansion of a CTG repeat in the 3' UTR of *ATXN8* is transcribed into both sense and antisense transcripts (Zu et al., 2011; Daughters et al., 2009; Moseley et al., 2006). The antisense transcript from *ATXN8* (*ATXN8OS*) contains a short AUG-driven ORF encompassing the CAG repeat and encoding polyglutamine. Removal of the AUG or placement of stop codons just upstream of the repeat did not preclude synthesis of the polyglutamine product. Instead, translation persisted in all three repeat reading frames to produce polyalanine and polyserine products, in addition to a non-AUG initiated polyglutamine protein (Zu et al., 2011). The polyglutamine, polyalanine, and polyserine products were found to accumulate in cerebellar Purkinje cells of SCA8 patients (Zu et al., 2011; Ayhan et al., 2018). RAN translation of all three products increased with repeat size, and replacement of the CAG repeat with CAA prevented RAN translation in all three frames (Zu et al., 2011). Similarly, polyglutamine, polyalanine, and polyserine products were detected in the context of the antisense DM1 transcript (in the CAG orientation), where deletion of the only in-frame AUG (for polyserine) did not preclude translation. Polyglutamine aggregates were detected in myoblasts, skeletal muscle, and blood of DM1 patients (Zu et al., 2011). Since this initial finding, unconventional translation of repeats has been described in a variety of other nucleotide repeat expansion disorders, including *C9orf72* in ALS/FTD, *HTT* in HD, *TCF4* in Fuchs' corneal dystrophy, along with several (but not all) of the spinocerebellar ataxia genes (Green et al., 2016; Ash et al., 2013; Mori et al., 2013; Banez-Coronel et al., 2015; Soragni et al., 2018; Ishiguro et al., 2017; Scoles et al., 2015). The exact mechanism by which this unconventional translation occurs is an area of active research and studies to date suggest that multiple mechanisms for initiation may occur at different repeats, or even in different reading frames of the same repeat (Gao et al., 2017). For the purposes of this review, we refer to all of these unconventional translation initiation events as RAN translation, with the understanding that future studies will be needed to truly delineate the mechanism(s) by which this process occurs.



### 3.1. RAN translation of CGG repeats in FXTAS


The gene *FMR1* contains a polymorphic CGG repeat in its 5' leader. This repeat is conserved among mammals, and increases in size in higher order primates (Eichler et al., 1995). In humans, this CGG element can expand to intermediate or "premutation" lengths of between 55 and 200 repeats. When this occurs, there is an enhancement in *FMR1* transcription, but inefficient Fragile X protein (FMRP) production (Tassone et al., 2000a, 2000b). Clinically, premutation expansions result in the neurodegenerative disorder Fragile X-associated Tremor/Ataxia Syndrome (FXTAS) (Hagerman, 2013). FXTAS occurs in ~1:5000 men over the age of 50 and in a lower percentage of female premutation carriers. In addition, premutation sized repeat expansions also cause premature ovarian failure, or Fragile X-associated Primary Ovarian Insufficiency (FXPOI), which is the most common genetic cause of early menopause. In both cases, the repeat is thought to elicit disease primarily through gain-of-function mechanisms, as expression of the repeat element in isolation is sufficient to elicit toxicity and enhancing or suppressing FMRP expression fails to modulate repeat elicited phenotypes in model systems (Lu et al., 2012; Arocena et al., 2005; Jin et al., 2007; Hashem et al., 2009). In contrast to FXTAS and FXPOI, Fragile X Syndrome (FXS) results from expansion of CGG repeats in *FMR1* to greater than 200. This leads to hypermethylation of the promoter region and CGG repeats and heterochromatinization of the *FMR1* locus. This epigenetic change leads to a partial or complete transcriptional shut down of the *FMR1* locus (Wang et al., 2010). FXS is the most common known monogenic cause of autism and intellectual disability. FXS results from the loss of FMRP, as point mutations and deletions in *FMR1* also elicit the clinical syndrome independent of repeat expansions. There is no evidence for neurodegeneration as a major feature in FXS cases, which suggests that the loss of FMRP is not a driver of the clinical or pathological features observed in FXTAS. Thus, while these diseases are allelic and result from expansion of the same repeat, they appear to have largely separate pathogenic mechanisms.

The *FMR1* CGG repeat was found to support RAN translation in all three potential reading frames on the sense strand in the absence of an AUG (Krans et al., 2016; Todd et al., 2013; Kearsse et al., 2016), albeit with differing efficiencies. RAN translation also occurs at CCG repeats generated by antisense transcription through the beginning of the *FMR1* gene (Krans et al., 2016). Distinct from what was described in *ATXN8*, RAN translation in two of the three potential reading frames of the 5' leader of *FMR1* relies on initiation at near-cognate start codons just upstream of the repeat (Kearsse et al., 2016; Todd et al., 2013). On the

sense strand, the 0-frame RAN initiation event uses a ACG initiation site, and the +1-frame uses both an ACG and a GUG for initiation (Fig. 2) (Kearsse et al., 2016; Todd et al., 2013). Peaks upstream of the repeat were detected in RP datasets from mouse and man (Todd et al., 2013). Interestingly, an earlier study found that the *Drosophila* homologue to *FMR1*, *dFMR1*, encodes a functional N-terminal expansion that initiates at a CUG start site (Beerman and Jongens, 2011). In CGG RAN translation, the +2-frame does not appear to rely on a start codon upstream of the repeat, making it more akin to what is observed at CAG repeat expansions with initiation occurring within the repeat (Kearsse et al., 2016). Interestingly, both the +1-frame and +2-frame products increase as a function of increased repeat size, and 0-frame RAN translation is attenuated at normal and expanded repeat lengths (Todd et al., 2013; Kearsse et al., 2016).

The scanning mechanism of translation initiation broadly refers to the process whereby the small ribosomal subunit with its associated factors interacts with the modified guanosine cap (m<sup>7</sup>G) on the 5' end of mRNA transcripts, then progresses in the 3' direction to a start codon where it joins with the large subunit to initiate translation (Hinnebusch, 2011; Hinnebusch et al., 2016). This process begins when a free 40S ribosomal subunit binds the ternary complex which is composed of the methionine initiator tRNA bound to eIF2, forming the 43S preinitiation complex (PIC). The PIC is recruited to the 5' end of the transcript through interaction with the eIF4F complex, which binds the m<sup>7</sup>G cap. The PIC complex then scans along the mRNA, proceeding in a 5' to 3' direction, while sampling the sequence for pairing with the initiator methionine tRNA at an AUG initiation codon. Prior to experimentation, it was unknown whether RAN translation used a cap-dependent mechanism or an internal ribosome initiation site (IRES)-like mechanism for initiation (Kearsse and Todd, 2014). IRES initiation occurs on highly structured transcripts, and directly recruits the 43S PIC to the transcript independent of the cap (Pelletier and Sonenberg, 1988; Jang et al., 1988; Pestova et al., 1996a, 1996b; Pestova et al., 1996a, 1996b). While originally described for viral elements, this same process also occurs from a subset of native mRNA transcripts in human cells (Ray et al., 2006; Vagner et al., 1995; Coldwell et al., 2000; Gan and Rhoads, 1996). Kearsse et al. (2016) showed that translation of CGG repeats in RAN reporter transcripts required a functional 5' cap and interaction with the cap-binding protein eIF4E, a component of the eIF4F complex that binds to the cap, and that blocking eIF4A, the helicase critical for PIC scanning, significantly impaired RAN translation in all three reading frames (Kearsse et al., 2016). These data point to a canonical scanning mechanism of translation initiation at the CGG repeat, with a

#### Model mRNA transcript for RAN translation



**Fig. 2.** RAN translation in repeat expansion disorders. Repeat associated non-AUG translation at three repeat expansions are laid out schematically. Experimentally determined initiation sites are listed for individual reading frames where determined, although in all cases initiation may also occur within the repeat itself. A) A G<sub>4</sub>C<sub>2</sub> hexanucleotide repeat in the first intron of *C9orf72* is bidirectionally transcribed. Whether the repeat is translated as a spliced intron or as part of a retained intron or other aberrant transcript is not known. A CUG located just 5' to the repeat is utilized for initiation of poly(GA). It may also be utilized for poly(GR) translation with subsequent frameshifting. AUG codons reside upstream of poly(PR) and poly(PG) in the most common antisense transcripts. B) A CGG repeat in the 5' leader/5' UTR of *FMR1* is also bidirectionally transcribed. An AUG in the poly(P) reading frame of *ASFMR1* is utilized for initiation in this reading frame but it can also undergo RAN translation in this reading frame. FMRpolyR expression is nearly undetectable from reporter constructs. C) The CAG repeat in Huntington's disease is located in the coding region of the *HTT* transcript and is also bidirectionally


transcribed. ? : initiation site unknown, # : initiation in this frame exhibits cap-dependency.

primary failure in start codon fidelity leading to RAN initiation (Green et al., 2016).

There is significant evidence supporting a role for RAN translation in the pathogenesis of FXTAS. RAN translation of the +1-frame CGG RAN protein, FMRpolyG, as a GFP fusion construct elicits formation of large intranuclear ubiquitinated inclusions in multiple model systems, including *Drosophila*, rodent neurons, mice and human cells (Sellier et al., 2017; Todd et al., 2013). These inclusions are highly reminiscent of the pathological hallmark of the disease observed in patients (Greco et al., 2006). Three different groups have now generated antibodies against different epitopes on FMRpolyG, which contains an expanded polyglycine stretch near its N-terminus. These antibodies recognize recombinant FMRpolyG protein and stain inclusions in multiple mouse and cell based models of the disease and, more importantly, in pathological samples from FXTAS cases (Todd et al., 2013; Sellier et al., 2017; Buijsen et al., 2016). FMRpolyG aggregates are also identified in ovarian stromal cells from FXPOI patients (Buijsen et al., 2016). FMRpolyG staining co-localizes with p62 and ubiquitin positive inclusions in patient brains, whereas pre-immune sera or peptide blocked sera do not recognize such inclusions (Todd et al., 2013; Buijsen et al., 2016; Sellier et al., 2017) (Amy Krans, personal communication).

In mice, ubiquitous transgenic expression or neuronal specific expression of the 5'UTR of *FMR1* leads to neuronal inclusion formation, motoric dysfunction, Purkinje cell loss, and a decrease in survival (Sellier et al., 2017). In contrast, expression of this same CGG repeat sequence lacking the near-AUG initiation sites that support FMRpolyG production has no phenotype, no ubiquitinated inclusions and normal survival (Sellier et al., 2017). Similar results are observed in *Drosophila* models of FXTAS and in cultured neurons (Todd et al., 2013). Driving expression of FMRpolyG causes impaired proteasome function in *Drosophila* and human cells (Oh et al., 2015) and nuclear lamina integrity is affected by FMRpolyG expression, which directly interacts with the inner nuclear membrane protein Lap2 $\beta$  (Sellier et al., 2017). In CGG repeat expansion “knock-in” mouse models, only those with 5'UTR sequences that support FMRpolyG production form neuronal ubiquitinated inclusions (Todd et al., 2013). Taken together, these data suggest that FMRpolyG production is necessary for CGG repeats to elicit neuronal toxicity and pathology across a variety of model systems. However, as CGG knock-in mice lack significant neurodegeneration (Krans and Todd, unpublished results), a direct link between RAN translation and inclusion formation with toxicity in endogenous systems lacking overexpression has not yet been established.

*FMR1* RAN translation is also interesting because it appears to occur at normal repeat lengths in the unaffected human population. This is supported by both ribosome profiling data in cell lines with a normal repeat length and reporter studies (Todd et al., 2013; Kears et al., 2016). This is important for two reasons: 1) it suggests that RAN proteins from this transcript can be synthesized under normal conditions and thus could have normal functions, and 2) translation in the 5' leader of an mRNA transcript could have a significant negative impact on the translation of the main protein-coding ORF, suggesting that RAN translation on *FMR1* could act as an upstream open reading frame (uORF) (Fig. 3). uORFs can implement a regulatory role on its resident transcript through several overarching mechanisms: through sequence-dependent interactions with the translation machinery, through cis- and trans-interactions with uORF-derived peptides, and through influencing RNA stability (Hinnebusch et al., 2016). Considering the translational efficiency of *FMR1* and the levels of FMRP are both known to be decreased in FXTAS where RAN translation occurs most, RAN translation may act as a uORF to repress downstream FMRP synthesis (Tassone et al., 2000a, 2000b; Feng et al., 1995). uORF translation is intricately tied to environmental stressors that act upon translation machinery. Moreover, uORFs have been implicated in neuronal response to receptor activity as the transcripts that show a higher level of translation after pharmacological activation of metabotropic glutamate receptors are enriched in uORFs (Di Prisco et al., 2014). *FMR1* is an important



**Fig. 3.** RAN translation as a regulatory element. A) Translation from the *FMR1* transcript produces both RAN translation products and FMRP. Reporter constructs and ribosome profiling data suggests that this process occurs at the normal repeat size in humans. As both cistrons utilize a scanning- and cap-dependent initiation mechanism, preinitiation complexes must bypass RAN initiation sites and the CGG repeat to make FMRP. As such, RAN translation of FMRpolyG might serve a regulatory upstream open reading frame (uORF) to control FMRP synthesis. B) At expanded repeats, RAN translation is more efficient, which correlates with less FMRP translation per transcript.

neuronal transcript whose regulation is likely central to normal neuronal function and plasticity. Thus elucidating how its 5' leader might regulate its own translation and produce novel synaptic proteins could reveal a lot about gene expression in the nervous system.

### 3.2. RAN translation in HD and ALS/FTD

Two other nucleotide repeat expansion disorders where RAN translation may contribute to overall pathology are HD and ALS. HD is a polyglutamine disease caused by CAG repeat expansions in the coding region of *HTT*, the Huntingtin gene. Patients generally exhibit involuntary movements and cognitive decline (Group, The Huntington's Disease Collaborative Research, 1993; Dickey and La Spada, 2018). Individuals with larger expanded repeats are prone to earlier onset HD (Paulson, 2018a; Andrew et al., 1993). While the polyglutamine containing protein defines the pathology of this disease, Banez-Coronel et al. (2015) showed that it is not the only homopolymeric expansion protein derived from this locus. RAN translation was shown to occur in both the sense and antisense direction through the CAG/CTG repeat, despite its residence within a protein coding open reading frame. In addition to the polyglutamine product, both poly-serine and poly-alanine products (sense) as well as poly-leucine and poly-cysteine products (antisense) were detected (Banez-Coronel et al., 2015). The RAN peptides aggregate in patient brains, and are increased in abundance in early onset HD (Banez-Coronel et al., 2015). Intriguingly, HD RAN peptides are linked to nucleocytoplasmic export disruption in neurons (Grima et al., 2017).

In 2011, expansion of the hexanucleotide repeat G<sub>4</sub>C<sub>2</sub> in the gene C9orf72 was discovered to be the most common known mutation underlying both ALS and FTD (Renton et al., 2011; DeJesus-Hernandez et al., 2011). Unaffected individuals normally have up to around 23 repeats, whereas individuals with ALS can have hundreds or thousands of this hexanucleotide repeat. G<sub>4</sub>C<sub>2</sub> bearing transcripts are transcribed in the sense and antisense direction, and the repeat is present in an intron or promoter region of sense strand RNA transcripts. RAN translation through this repeat produces five potential dipeptide repeat (DPR) tract proteins—poly-glycine/alanine (GA), -glycine/arginine (GR), -proline/alanine (PA), -proline/arginine (PR), and -glycine/

proline (GP) (Green et al., 2016; Ash et al., 2013; Gendron et al., 2013; Zu et al. 2013; Mori et al., 2013). C9orf72-associated RAN proteins accumulate in the brains of patients in aggregates distinct from the TDP-43 positive aggregates classically seen in ALS/FTD. Several studies have concluded that GA-DPRs are the most abundant of the G<sub>4</sub>C<sub>2</sub> RAN proteins pathologically (Gendron et al., 2015; Mackenzie et al., 2015; Mann et al., 2013). In vitro and cell based assays similarly demonstrate that translation in the GA reading frame of the sense strand repeat is the most efficient, with lesser expression in the GR and GP reading frames (Green et al., 2017; Sonobe et al., 2018; Westergard et al., 2019; Tabet et al., 2018). In addition, each frame exhibits different length-dependency and differential impedances based on initiation and elongation (Green et al., 2016; Tabet et al., 2018; Cheng et al., 2018). G<sub>4</sub>C<sub>2</sub> RAN translation from linear mRNA transcripts appears to mainly depend on cap-binding of eIF4E and a canonical scanning mechanism involving eIF4A, similar to what has been reported for CGG repeats (Green et al., 2017; Tabet et al., 2018). However, G<sub>4</sub>C<sub>2</sub> RAN can occur to at least some degree from bicistronic constructs and transcripts, suggesting a mechanism more aligned with internal ribosome entry (IRES) (Cheng et al., 2018; Sonobe et al., 2018). Translation in the GA reading frame primarily utilizes a near-cognate CUG codon 5' to the repeat for initiation, as well as an AGG closer to the repeat (Green et al., 2017; Tabet et al., 2018). The dependency on a single start site for multiple reading frames suggests that frame-shifting may occur in certain contexts (Tabet et al., 2018). Frameshifting was previously reported at CAG repeats (Toulouse et al., 2005; Wills and Atkins, 2006; Stochmanski et al., 2012; Girstmair et al., 2013; Wojciechowska et al., 2014; Saffert et al., 2016) but did not appear to be a major contributor to RAN translation product generation from CAG repeats on ATXN8OS (Zu et al., 2011). Thus, while many nucleotide repeat expansions appear to support RAN, the underlying mechanisms appear to have significant differences—not only between repeats but even in different frames of the same repeat.

The Integrated Stress Response (ISR) is activated by specific stimuli, namely ER stress, viral infection, oxidative stress and amino acid deprivation (Pakos-Zebrucka et al., 2016; Young and Wek, 2016). These stimuli each act on specific kinases that phosphorylate eIF2 $\alpha$ . eIF2 is comprised of three subunits. The  $\gamma$  subunit binds GTP. When GTP is hydrolyzed to GDP, eIF2 cannot bind methionine initiator tRNA. Phosphorylation of the  $\alpha$  subunit of eIF2 inhibits the guanine exchange factor (GEF) eIF2B, which is responsible for adding GTP to eIF2 (Hinnebusch, 2011). Thus, phosphorylation of the  $\alpha$  subunit leads to a limiting number of ternary complexes and a global decrease in protein synthesis (Harding et al., 2003; Pakos-Zebrucka et al., 2016; Harding et al., 2000). Despite a global translational arrest, a subset of transcripts show a paradoxical increase in translation, and many of the gene products are associated with cellular response to mitigate stress and maintain homeostasis (Harding et al., 2000; Vattem and Wek, 2004). Intriguingly, under ER stress and phosphorylation of eIF2 $\alpha$ , CGG RAN translation of the +1- and +2-frames increases despite the global decrease in translation (Green et al., 2017). This increase was dependent on the presence of a near-cognate start site, and was also observed at the ALS/FTD G<sub>4</sub>C<sub>2</sub> repeat (Green et al., 2017; Cheng et al., 2018; Sonobe et al., 2018; Westergard et al., 2019). Importantly, expression of either expanded CGG or G<sub>4</sub>C<sub>2</sub> repeats induces stress granule formation in cells, which is dependent on eIF2 $\alpha$  phosphorylation (Green et al., 2017; Cheng et al., 2018; Rossi et al., 2015). In addition, expression of GA DPRs triggers ISR activation independent of RNA repeats through a PERK-dependent mechanism (Zhang et al., 2014). This ISR activation, in concert likely with direct effects of DPRs on nucleolar function and ribosomes, leads to a suppression of global protein translation in cells expressing repeats or arginine-rich DPRs (Kwon et al., 2014; Zhang et al., 2014; Rossi et al., 2015; Tao et al., 2015; Yamakawa et al., 2015; Kanekura et al., 2016; Lee et al., 2016; Green et al., 2017; Mizielinska et al., 2017; Hartmann et al., 2018; Zhang et al., 2018a, 2018b; Moens et al., 2019; Haeusler et al., 2014). Translation of the GA reading frame

of the G<sub>4</sub>C<sub>2</sub> repeat was also found to be partly dependent on eIF2A, an initiation factor with roles in unconventional translation initiation (Sonobe et al., 2018; Starck et al., 2012). In iPS spinal motor neurons derived from patients with the G<sub>4</sub>C<sub>2</sub> expansion both cellular stress and neuron-specific excitotoxic stress, through application of homocysteine or excess glutamate, increased endogenous poly(GA), (GP), and (GR) at variable levels (Westergard et al., 2019). This was also observed in rat cortical neurons expressing reporters for each protein (Westergard et al., 2019). Taken together, these data support the model of toxicity where RAN translation is upregulated in the presence of cellular stress. In turn, both repeats as RNA and translated RAN proteins independently induce stress, creating a feed-forward loop that leads to further RAN translation and a global suppression of protein translation that contributes to neurodegeneration (Green et al., 2017).

#### 4. Microsatellites impact on liquid-liquid phase separation of membraneless organelles

Membraneless organelles are organizational centers formed within the cell to perform a variety of functions in close proximity and shielded from the rest of the cellular milieu but in the absence of a structural membrane. For example, Cajal bodies and the nucleolus are nuclear organelles that perform specialized tasks which require a very specific set of molecular components within a restricted area (Boisvert et al., 2007; Gall, 2003). Membraneless organelles are forged through liquid-liquid phase separation (LLPS), the physical process by which two liquids, when mixed together, spontaneously separate. The simplest way to explain LLPS is the example of oil and vinegar (Hyman et al., 2014). When oil and vinegar are added together, discrete droplets of each liquid form through de-mixing to accomplish a lower free energy state (Gomes and Shorter, 2018). In the cell, local molecular interactions can drive phase separation between the cytosol and its various components, which are classified as liquids (Hyman and Brangwynne, 2011). These liquids can take many forms such as gels and colloidal solutions (Hyman et al., 2014). LLPS ultimately both drives the formation of and maintains specialized compartments that can perform specific cellular tasks.

Proteins susceptible to phase separation commonly have intrinsically disordered regions (IDR). IDRs typically have low sequence complexity, and are enriched in amino acids that promote low hydrophobicity and a high net charge including Ala, Arg, Gly, Gln, Ser, Glu, Lys, and Pro (Uversky and Dunker, 2010). An important outcome of such attributes is a failure to form rigid 3-D structures, and as a result they maintain a high level of flexibility (Uversky and Dunker, 2010). IDRs permit a wide range of molecular interactions, and thus allow for both homomeric and heteromeric protein complex formations (Dyson and Wright, 2005). In the crowded cytoplasm of a cell, binding of an IDR-containing protein to its molecular interactors drives up the local concentration of these proteins, which in turn favors phase separation a mechanism of reaching higher thermodynamic stability through multivalent interactions (Boeynaems et al., 2018). Importantly, some proteins with IDRs can self-assemble (Malinowska et al., 2013). However, when the local concentration of IDR-containing proteins is too high, or their relative affinities and interactions are stronger, they form more rigid conformations that are less dynamic and harder to disassemble (Dogan et al., 2009; Crick et al., 2006; Vitalis et al., 2007; Walters and Murphy, 2009). As discussed above, a hallmark of diseases in which microsatellite repeats are encoded into proteins is the presence of aggregates. It is thought that pathologically expanded repeat proteins prevent proper liquid-liquid phase transitions by forming solid, insoluble aggregates (Malinowska et al., 2013; Dobra et al., 2018; Boeynaems et al., 2018). Intramolecular interactions may also play a role. For example, as the size of a polyglutamine stretch within a protein expands, the monomer is more likely to adopt a structured conformation, increasing the propensity to form aggregates both with itself and with other proteins (Walters and Murphy, 2009; Haran, 2012;



Vitalis and Pappu, 2011; Williams and Paulson, 2008).

Ribonucleoprotein (RNP) granules make up one category of membraneless organelles. These are dense bodies of RNA and RNA binding proteins (RBP) (Spector, 2006). Messenger RNP (mRNP) granules form in the cytoplasm and are composed of translationally-silenced mRNA (Anderson and Kedersha, 2009). In neurons, mRNPs are important for regulation of translation and trafficking of mRNAs out to synapses. The fragile X protein FMRP is an RNA binding protein and a key component of neuronal mRNP granules within dendrites. FMRP binds to ~4% of the brain transcriptome, with enrichment for transcripts related to neuronal function and whose dysfunction are associated with autism and intellectual disability (Siomi et al., 1994; Siomi et al., 1993; Brown et al., 2001; Darnell et al., 2011; Ascano Jr. et al., 2012). Loss of FMRP in models of FXS leads to altered global protein translation and mRNA transport within neurons, and as a consequence, dysregulated activity-dependent translation at synapses (Thomson et al., 2017; Darnell et al., 2001; Laggenbauer et al., 2001; Li et al., 2001; Todd et al., 2003; Brown et al., 2001; Zalfa et al., 2003; Huber et al., 2002; Liu et al., 2018; Udagawa et al., 2013; Banerjee et al., 2018; Ifrim et al., 2015; Gross and Bassell, 2012; Liu-Yesucevitz et al., 2011; Gross et al., 2011). This uncoupling of synaptic activity and local protein synthesis is believed to underlie learning and memory deficits in FXS. Ribosome profiling, a next generation sequencing assay to survey the location of all translating ribosomes, was implemented in adult neural stem cells of *FMR1* knockout mice and compared to wild-type cells (Liu et al., 2018). This work corroborated aspects of FMRP's proposed role as a translational repressor of mRNA transcripts to which it binds by describing increased translational efficiency of FMRP CLIP (cross-linking immunoprecipitation) targets in *FMR1* knock-out cells. However, global translational efficiency changes were largely buffered at the mRNA level suggesting that compensatory changes may bypass some of FMRP's roles as a translational regulator (Liu et al., 2018). What remains to be explored is how the absence of FMRP and this compensated state influences activity-dependent translation shifts in specific cellular compartments such as dendrites and axons.

Stress granules are a specialized type of mRNP with roles in a diverse set of neurodegenerative diseases (Wolozin, 2012; Kedersha et al., 1999). They contain mRNAs stalled at translation initiation, along with small ribosomal subunits, select initiation factors, and RBPs such as G3BP, DDX3X, FMRP and TIA1 (Protter and Parker, 2016; Buchan and Parker, 2009; Teixeira et al., 2005; Hilliker et al., 2011). Stress granules form as a consequence of cellular stress that induces eIF2 $\alpha$  phosphorylation and translational inhibition. This blockade of initiation without a concomitant blockade of elongation creates "naked" stretches of mRNA on protein-coding transcripts that would normally be coated with translating ribosomes. This unassociated RNA facilitates specific protein-protein interactions presumably by creating a nidus upon which abundant RBPs with low sequence binding specificity can assemble (Molliex et al., 2015; Lin et al., 2015). RNA-RNA interactions, RNA-protein interactions as well as protein-protein mediated phase separation of factors such as G3BP through IDR domains all contribute to this process (Molliex et al., 2015; Lin et al., 2015).


Stress granules are generally thought of as a way for cells to redirect their energy expenditure elsewhere in the setting of a transient insult to regain their healthy set point. While stress granules are generally adaptive, alterations in their assembly and disassembly dynamics may contribute meaningfully to multiple neurodegenerative disorders (Li et al., 2013; Wolozin, 2012; Becker and Gitler, 2018). Overexpression of C9 FTD/ALS DPRs modulates SG granule formation and dynamics (Zhang et al., 2014; Zhang et al., 2018a, 2018b; Lee et al., 2016; Kanekura et al., 2016; Tao et al., 2015; Rossi et al., 2015), with poly(GA) contributing to SG activation and poly(GR) and poly(PR) impacting SG stability and disassembly (Fig. 4).

#### 4.1. The polyglutamine tract in the ataxin-2 protein regulates membraneless organelles in the cytoplasm

Ataxin-2 is a common eukaryotic mRNP granule component that can bind directly to RNA and other RBPs (Kozlov et al., 2010; Swisher and Parker, 2010; Nonhoff et al., 2007; Lastres-Becker et al., 2016). It helps promote, but is not required for, stress granule assembly in yeast and human cells, and is necessary for proper formation of RNP granules positive for the stress granule marker Me3B1 in *Drosophila* projection neurons (Buchan et al., 2008; Becker et al., 2017; Bakthavachalu et al., 2018). Ataxin-2 has a polyglutamine tract in its IDR. Deletion of the IDR in the ataxin-2 protein, including the polyglutamine tract, disrupts Me3B1-positive RNP granule formation (Bakthavachalu et al., 2018). The ataxin-2 polyglutamine tract, encoded by a CAG microsatellite, is polymorphic in humans and can become expanded as a result of somatic and intergenerational instability (Pulst et al., 1993; Sanpei et al., 1996; Pulst et al., 1996; Imbert et al., 1996). The mean repeat size in the population is 25 CAGs. Expansions beyond 34 CAGs causes spinocerebellar ataxia type 2 (SCA2), a late onset neurodegenerative disorder characterized by cerebellar dysfunction, spasticity, and some motor neuron disease-like features (Pulst, 1993). In contrast, expansion of the repeat in the intermediate range (~27-33 CAGs) increases the risk for ALS without any association with SCA2 (Elden et al., 2010). Ataxin-2 interacts with FUS and TDP-43, RBPs that are commonly mutated and mislocalized in ALS. Intermediate expansions in ataxin-2 can enhance FUS mislocalization from the nucleus to the cytoplasm and ER-Golgi disruption in the setting of an ALS-causing FUS mutation (Farg et al., 2013). Deletion of the disordered region of ataxin-2, including the polyglutamine tract, partially rescues the eye degeneration phenotype in a *Drosophila* model expressing human FUS (Bakthavachalu et al., 2018). Additionally, intermediate length repeat expansions increase caspase 3 activation in cells expressing mutant TDP-43, and patient motor neurons with an intermediate length repeat show increased activated caspase 3 linking this moderate microsatellite repeat expansion directly to motor neuron death (Hart and Gitler, 2012). Decreasing wild-type *Ataxin-2* levels rescues mutant TDP-43 toxicity in yeast, fly, and mouse models of ALS (Elden et al., 2010; Becker et al., 2017) while decreasing expression of expanded *Ataxin-2* alleles suppresses SCA2-relevant phenotypes in mice (Scoles et al., 2017). While the exact mechanism by which intermediate length expansion increases the risk for motor neuron disease is yet to be deciphered, insight from research on ataxin-2 in various RNP granules makes disrupted RNP granule dynamics a likely contributor (Zhang et al., 2018a, 2018b).

#### 4.2. Transcribed and translated C9ORF72 hexanucleotide repeat impairs nuclear phase separation

In the setting of the C9ORF72 expanded G<sub>4</sub>C<sub>2</sub> repeat, the arginine-containing RAN proteins, poly(GR) and poly(PR), contribute to impaired membraneless organelle dynamics. Poly(GR) and poly(PR) peptides are extremely toxic to cells when put directly into culture medium or expressed from a plasmid (Kwon et al., 2014; Wen et al., 2014; Tao et al., 2015; Lee et al., 2016; Flores et al., 2016), and cause neurodegeneration in *Drosophila* models (Mizielinska et al., 2014; Lee et al., 2016; Wen et al., 2014). Poly(PR) and poly(GR) cellular interactors were identified by mass spectrometry; their interactomes were enriched in low-complexity domains, a common attribute of IDRs that can contribute to liquid-liquid phase separation, and many poly(PR) interactors fall into the category of membraneless cellular puncta (Lee et al., 2016; Lin et al., 2016; Zhang et al., 2018a, 2018b; Hartmann et al., 2018). One such protein was hnRNPA1, an RBP component of stress granules with a prion-like domain in which mutations can lead to impaired phase transition and aberrant fibrillization (Guo et al., 2018; Lin et al., 2016; Kim et al., 2013). Through mutational analysis, the low-complexity domain of hnRNPA1 was determined to be necessary and sufficient for this poly(PR) interaction providing evidence that poly



**Fig. 4.** Mechanisms by which transcribed repeats elicit neurodegeneration. Expanded repeats can be toxic as RNA or as protein. 1) poly(GR) and poly(PR) containing RAN DPR proteins undergo phase separation (1a) into liquid-like droplets (LLPS) *in vitro* and *in vivo*. Poly(GA), poly(GR), and poly(PR) RAN proteins can also alter stress granule assembly and dynamics (1b) and form insoluble aggregates (1c). In the nucleus, RAN proteins can impair the dynamics of Cajal bodies and membraneless organelles (1d), and disrupt nucleolar function and ribosomal biogenesis (1e). They can also disrupt the nuclear envelope and nuclear pore complex (1f). Repeat containing RNAs can sequester important RNA-binding proteins in the nucleus and can themselves undergo LLPS, perhaps in concert with RBPs (2a). Repeat RNAs that exit the nucleus can be exported to neuronal processes where they can impair neurite outgrowth and potentially alter mRNP dynamics (2b).

(PR) can directly interact with these domains (Lin et al., 2016). Poly (PR) and poly(GR) binding also promotes hnRNP A1 phase separation *in vitro* (Lee et al., 2016). Both RAN peptides promote the assembly of aberrant stress granules, and lower the critical concentration of hnRNP A1 and TIA-1, another stress granule protein, required for phase separation *in vitro* (Lee et al., 2016; Boeynaems et al., 2017). Poly(GA) peptides also independently trigger SG formation (Tao et al., 2015), and their co-expression with arginine rich repeat influences the toxicity and behavior of these charged DPRs (Yang et al., 2015; Darling et al., 2019; Lee et al., 2017).

Overexpression of poly(GR) and poly(PR) localize directly to the nucleolus, and poly(GR) also localizes to the cytoplasm (Lee et al., 2016; Wen et al., 2014; Tao et al., 2015; Kwon et al., 2014). Poly(GR) and poly(PR) impair the behavior of proteins that help organize the nucleolus through liquid-liquid phase transition. Ribosome biogenesis is also impaired in their presence, which takes place in the granular component of the nucleolus where both RAN peptides localize (Lee et al., 2016; Tao et al., 2015; Suzuki et al., 2018). Additionally, poly (GR) and poly(PR) alter the assembly of two other nuclear membraneless organelles: Cajal bodies and nuclear speckles (Lee et al., 2016). Taken together, these data demonstrate that arginine-containing RAN peptides can drive toxicity in cells and neurons by disrupting the phase separation properties of membraneless organelles. However, it is important to acknowledge that almost all studies to date have been done with constructs where poly(PR) and poly(GR) are generated from AUG-driven translation in the absence of the repeat element and usually fused to a fluorescent protein tag such as GFP. Given that both of these proteins are present at very low levels in C9 ALD/FTD neurons, these overexpression systems may not recapitulate what happens endogenously. In postmortem patient brain tissue, aggregates of poly(PR) and poly(GR) appear mostly perinuclear (Zu et al. 2013; Mori et al., 2013). Moreover, there is evidence that the other DPRs generated by RAN translation may counteract or influence the behavior of these arginine rich proteins in terms of their distribution and relative toxicity (Darling et al., 2019). Thus, future studies are needed which selectively decrease expression of these arginine-rich DPRs from endogenous loci to better delineate impact they have in the disease state.

G<sub>4</sub>C<sub>2</sub> repeat RNAs form G-quadruplex structures *in vitro* and there is some evidence to suggest that these same structures form *in vivo* (Conlon et al., 2016; Fay et al., 2017; Reddy et al., 2013; Haeusler et al., 2014). G<sub>4</sub>C<sub>2</sub> repeat-containing RNAs can also undergo phase separation

(Moens et al., 2018). Purified (G<sub>4</sub>C<sub>2</sub>)<sub>n</sub> RNA form gels *in vitro* in a repeat-dependent manner, and G<sub>4</sub>C<sub>2</sub> RNAs form foci when transfected into cells similar to what is observed for (CAG)<sub>n</sub> and (CUG)<sub>n</sub> RNA (Jain and Vale, 2017; Fay et al., 2017). Foci are also present in patient cells, including brain tissue and motor neurons (Lee et al., 2013; Lagier-Tourenne et al., 2013; Sareen et al., 2013; Donnelly et al., 2013). Of note, another study used the MS2 system to tag both (G<sub>4</sub>C<sub>2</sub>)<sub>n</sub> and (CAG)<sub>n</sub> RNA, both showed repeat-dependent neurite localization with the (CAG)<sub>n</sub> RNA increasing in neurite localization at larger repeats, with the opposite pattern of localization occurring with expansion of (G<sub>4</sub>C<sub>2</sub>)<sub>n</sub> RNA (Burguete et al., 2015). Neuritic expanded G<sub>4</sub>C<sub>2</sub> and CAG repeat-containing RNA was shown to be actively transported in mRNP granules, and was correlated with neurite branching defects. Intriguingly, neurite-localized G<sub>4</sub>C<sub>2</sub> RNA puncta were detected in iPSC-derived neurons from C9orf72 hexanucleotide expansion carriers (Burguete et al., 2015).

Sequestration of RBPs is a common pathologic mechanism underlying microsatellite expansions categorized as RNA gain-of-function mutations (Todd and Paulson, 2010). G<sub>4</sub>C<sub>2</sub> nuclear RNA foci sequester a number of RBPs including ADARB2, SRSF2, hnRNP H1/F, Aly/REF, hnRNP A1, Pur α, hnRNP H, among others (Donnelly et al., 2013; Cooper-Knock et al., 2014; Conlon et al., 2016; Xu et al., 2013; Sareen et al., 2013; Lee et al., 2013). Sequestration of hnRNP H, specifically, can lead to mis-splicing of its targets (Lee et al., 2013; Conlon et al., 2016). However, the exact mechanism by which such repetitive elements might bind to and sequester RNA binding proteins and thus drive cellular dysfunction is less clear (Renoux and Todd, 2012). The working model for RNA dominant toxicity has been that sequestered proteins would need to have a very high affinity for the repeat element sequence, otherwise their interactions would be dynamic and thus functional sequestration would be unlikely. One would anticipate that highly abundant proteins would in this context be very difficult to functionally sequester. Yet, in most contexts, the interactions between repeat RNAs and their target RBPs are reversible *in vitro* and with binding affinities that would make functional sequestration difficult to achieve through classically understood RNA-protein binding interaction relationships. Proteins such as Muscle blind and hnRNP H are abundant in cells. In this context, recent studies demonstrating that repeat RNAs themselves can phase separate are quite intriguing (Jain and Vale, 2017; Fay et al., 2017). Moreover, RNA-mediated phase separation can intersect with protein-mediated phase separation as the addition of



polyuridine RNA to purified poly(PR) protein *in vitro* promotes demixing in a dose-dependent manner, and this relationship influences stress granule formation and dynamics (Boeynaems et al., 2017; Van Treeck et al., 2018). If microsatellite expansion mutations drive phase separation of RNAs, then perhaps functional sequestration is achieved through a synergistic combination of classic repetitive RNA motif-protein interactions and phase separation that could be aided by both the intrinsically disordered domains of the protein and the phase separation propensities of the RNA. In such a context, even moderate affinity interactions with the repetitive motif sequence could potentially push RNA-protein complexes into a phase separated or even gelled state that would be very stable, making the protein component inaccessible for normal cellular functions.

#### 4.3. Aberrant phase transitions cause nucleocytoplasmic transport defects

The central transport system between the cytoplasm and the nucleus is the nuclear pore complex (NPC). It functions as a highly regulated channel to pass macromolecules between the two segregated compartments (Hoelz et al., 2011). Its function is central to proper RNA trafficking, and thus proper gene expression. The NPC is a large macromolecular complex with a mass of around 66MDa in eukaryotes. It is composed of multiple molecules of approximately 30 different nucleoporins or nups that are classified by their position within the larger NPC (Reichelt et al., 1990). The greater structure of the NPC is formed by both symmetrical nucleoplasmic and cytoplasmic rings, and an inner ring complex. In the center is a channel which can dilate and contract to permit cargo through (Hoelz et al., 2011). The barrier to this channel is formed by phenylalanine-glycine repeat containing nups (FG-nups) with low sequence complexity that must remain flexible and responsive to cellular conditions (Kim and Taylor, 2017). The FG motifs interact with each other to create a meshwork barrier (Schmidt and Gorlich, 2016). The “selective phase model” describes the FG-nup barrier as a hydrogel that forms reversible crosslinks between FG domains to act as a sieve (Frey and Gorlich, 2007; Mohr et al., 2009; Kim and Taylor, 2017; Ribbeck and Gorlich, 2001; Schmidt and Gorlich, 2015, 2016). The FG-nups undergo phase transitions through multivalent interactions of their IDRs in order to maintain compartmental fidelity.

The FG motifs interact with nuclear transport receptors that shuttle through the pore to facilitate the translocation of larger molecules. Importins and exportins, both different types of nuclear transport receptors, regulate the transport of molecules in and out of the nucleus with the help of the RanGTP-RanGDP system. Ran protein, which stands for Ras-related nuclear protein, regulates the assembly and disassembly of transport complexes (Hoelz et al., 2011). Ran guanine exchange factor (RanGEF) protein is located in the nucleus, while Ran GTPase activating protein (RanGAP) is located exclusively in the cytoplasm (Kim and Taylor, 2017). Segregation of RanGEF and RanGAP sets up a gradient of RanGTP that dictates the directionality of transport through the NPC and the integrity of the FG barrier is crucial for maintaining this barrier (Debler et al., 2009).

A role for nucleocytoplasmic transport machinery in *C9orf72* ALS emerged from several coincident reports across model systems. First, a genetic screen was performed in a *Drosophila* model of *C9orf72* ALS that expressed a (G<sub>4</sub>C<sub>2</sub>)<sub>58</sub> repeat within the 5'UTR of GFP, leading to an eye degeneration phenotype (Freibaum et al., 2015). Two key NPC proteins modified the eye phenotype. Deletion of Nup50 enhanced degeneration, while deletion of Ref1 mitigated G<sub>4</sub>C<sub>2</sub> toxicity (Freibaum et al., 2015). Human NUP50 promotes nuclear import, whereas Aly/REF (the human orthologue of Ref1) assists in mRNP export (Makise et al., 2012; Kim and Taylor, 2017; Viphakone et al., 2012). Fly salivary gland cells expressing the (G<sub>4</sub>C<sub>2</sub>)<sub>58</sub> repeat exhibited a “wrinkled” nuclear envelope and an accumulation of nuclear RNA, which was also observed in patient iPSC-derived neurons (Freibaum et al., 2015). Work done by a separate group using a similar fly model with 30 repeats showed that a dominant, gain-of-function mutation in *RanGAP*, that

works similar to *RanGAP* overexpression, suppressed repeat-induced toxicity (Zhang et al., 2015). *RanGAP1* binds directly to G<sub>4</sub>C<sub>2</sub>-containing RNA, and *RanGAP1* is mislocalized in repeat expressing cells and in *C9orf72* patient iPSCs and brain tissue (Zhang et al., 2015). This mislocalization of *RanGAP* disrupts the nucleocytoplasmic Ran gradient and impairs nucleocytoplasmic transport (Zhang et al., 2015).

Studies on the effect of peptides synthesized through RAN (Repeat Associated Non-AUG) translation of the G<sub>4</sub>C<sub>2</sub> repeat in *C9orf72* revealed an interaction between several dipeptide repeat proteins and the nucleocytoplasmic transport machinery. A large-scale screen in *Saccharomyces cerevisiae* for modifiers of poly(PR) toxicity revealed importins as suppressors of toxicity when overexpressed (Jovicic et al., 2015). Further studies in a *Drosophila* model of RAN peptide toxicity whereby the transgene was codon-optimized to remove the G<sub>4</sub>C<sub>2</sub> repeat in the RNA but still encode the same amino acids similarly identified multiple nucleocytoplasmic transport factors as modifiers of poly(PR) toxicity, corroborating several of the previously mentioned genetic interactors such as Nup50 and *RanGAP1* (Boeynaems et al., 2016). Poly (PR) binds to the nup FG domain and disrupts the nuclear pore complex in *Xenopus* (Shi et al., 2017). In a mouse model overexpressing poly(GA) by viral injection, *RanGAP1* as well as the nuclear pore protein POM121 showed aberrant accumulation (Zhang et al., 2016). Both often formed aggregates colocalizing with poly(GA) (Zhang et al., 2016). These reports point to a RAN peptide-mediated disruption of the nucleocytoplasmic machinery in the setting of the expanded G<sub>4</sub>C<sub>2</sub> hexanucleotide repeat.

## 5. Conclusion

Over almost thirty years, research on a relatively rare series of NREs has played a tremendous role in adding to our understanding of many aspects of neurobiology and neurodegeneration. The discovery of the G<sub>4</sub>C<sub>2</sub> repeat in *C9orf72* in particular has been a recent catalyst for such research, some of which is highlighted here. However, our understanding of the native functions of tandem microsatellites themselves has seen only limited progress, despite a great potential for contributing to normal function and human disease. Current deep-sequencing technology do not adequately resolve most microsatellites, and thus we do not have an accurate map of the variance in repeating elements in the human genome to even begin predicting which repeats might be important. Given recent discoveries—even in the past year—of new repeats that cause human disease, it seems likely that we are overlooking many disease-causing microsatellites (LaCroix et al., 2019; Ishiura et al., 2018; Cortese et al., 2019; Sone et al., 2019).

Improved technologies for long read and single cell deep-sequencing may provide new insights into both somatic mosaicism and population level variance of repeat sizes in brain tissue (McConnell et al., 2017). In this context, many of the repeat-dependent effects described here could affect different cells in different ways, including actions mediated by more moderate “risk-factor” effects akin to what occurs with *Ataxin-2*. A greater understanding of the basics of repeat biology in normal cellular functions will be required to understand their aberrancy in disease pathogenesis. This becomes more apparent as new disease associated NREs are uncovered. For example, a dominantly inherited intronic AAGGG repeat expansion in the poly(A) tail of the *RFC1* gene was recently identified a major cause of cerebellar ataxia, neuropathy, and vestibular areflexia syndrome (CANVAS) (Cortese et al., 2019). Histological examination of patient tissue reveals Purkinje cell loss, however neither *RFC1* transcript nor protein abundance appears affected and no inclusions or RNA foci were identified (Cortese et al., 2019). Thus, at first pass, this new repeat does not seem to fit within our current conceptualization of how dominantly inherited repeat expansions cause disease. This mystery may open us to new insights into repeat biology, opening up exploration of how repeats may trigger distant epigenetic effects or alter the functions of non-coding RNAs. As with our evolving understanding of how repeats elicit toxic gain-of-function as RNA or

through RAN translation, it will be important to keep an open mind on the possibilities for how such a repeat might cause disease.

Along these lines, the finding that microsatellites in noncoding regions of RNA can in fact be translated has the potential to dramatically alter how we view a series of normal cellular processes. Not only does this change the way expansion diseases are viewed, it changes our understanding of mRNA translation generally. Structured regions are quite common in RNA, and their ability to promote translation has long been overlooked (Kozak, 1984). The finding that RAN translation is supported at shorter repeat lengths in reporter constructs suggests potential biological functions of these processes in healthy cells (Todd et al., 2013). Not only has this been detected with the CGG repeat, but certain reading frames of the *C9orf72* repeat as well (Green et al., 2017). As many of these transcripts are highly expressed in neurons, their roles in neuronal function may be particularly interesting. Such non-canonical ORFs might greatly expand the known human proteome and perform distinct functions. Given how pathological expansions of repetitive elements influence the assembly and disassembly of membraneless organelles, it will be particularly important to evaluate how transcribed and translated microsatellite repeats impact mRNP trafficking and temporal and spatial regulation of translational dynamics in neurons, which are highly compartmentalized and rely on precise regulation of such events to allow for circuit formation and synaptic plasticity.

Similarly, the ability of repetitive elements as RNA and protein to trigger phase separation and influence the formation of membraneless organelles has revealed an important component of cellular organization that becomes dysfunctional in NREDS. Yet, at normal repeat sizes these same elements likely enhance the ability of the proteins and transcripts in which they reside to become components of these complexes and thus may aid in their function rather than acting as a hindrance. For example, the CGG Repeat in FMR1 as RNA may influence its entry into mRNPs that allow its transport out into distal dendrites where it is translated in response to activity (Muslimov et al., 2011). As proteins, this is especially apparent in the nuclear pore, where phenylalanine-glycine repeat elements in specific nup proteins play important roles in the pore function that is actively impaired by charged poly(PR) proteins generated by RAN translation (Kim and Taylor, 2017). Understanding these native functions and interactions elicited by repeats will allow a clearer delineation of the mechanisms by which expanded repeats impair the behavior of these macromolecular complexes and thus what cellular compensatory mechanisms might be co-opted as therapeutic targets in disease.

Rapid progress in our understanding of microsatellite expansions has led to a series of recent clinical trials in patients, some of which appear quite promising. Knockdown of repeat expanded mRNA transcripts using antisense oligonucleotides has been quite successful in animal models and is currently in trials in multiple repeat expansion disorders where gain-of-function mechanisms are thought to drive disease pathogenesis (Kordasiewicz et al., 2012). However, in diseases like FXTAS, where loss of the patent transcript itself would likely be problematic, alternative approaches will need to be developed. Such cases highlight the importance of further research on these topics in order to develop therapeutics that act downstream on affected pathways and not solely on the mutated gene. For example, reduction of wild-type *Ataxin-2* in a mouse model of ALS provides one example of how targeted therapies aimed a more general component of the neurodegenerative process could prove helpful (Becker et al., 2017). With continued research, more discoveries like this will bring a greater understanding of microsatellite expansion diseases and novel strategies by which we might cure these diseases.

## Acknowledgements

This work was funded by grants from the VA BLRD (1I21BX001841 and 1I01BX003231), the NIH (R01NS099280 and R01NS086810) and

the Fragile X Association of Michigan to PKT. CMR was supported by the Ruth L. Kirchstein National Research Service Award F31NS090883.

## References

- Akbar, U., Ashizawa, T., 2015. Ataxia. *Neurol. Clin.* 33, 225–248.
- Anderson, P., Kedersha, N., 2009. RNA granules: post-transcriptional and epigenetic modulators of gene expression. *Nat. Rev. Mol. Cell Biol.* 10, 430–436.
- Andrew, S.E., Goldberg, Y.P., Kremer, B., Telenius, H., Theilmann, J., Adam, S., Starr, E., Squitieri, F., Lin, B., Kalchman, M.A., et al., 1993. The relationship between trinucleotide (CAG) repeat length and clinical features of Huntington's disease. *Nat. Genet.* 4, 398–403.
- Arocena, D.G., Iwahashi, C.K., Won, N., Beilina, A., Ludwig, A.L., Tassone, F., Schwartz, P.H., Hagerman, P.J., 2005. Induction of inclusion formation and disruption of lamin A/C structure by premutation CGG-repeat RNA in human cultured neural cells. *Hum. Mol. Genet.* 14, 3661–3671.
- Ascano Jr., M., Mukherjee, N., Bandaru, P., Miller, J.B., Nusbaum, J.D., Corcoran, D.L., Langlois, C., Munschauer, M., Dewell, S., Hafner, M., Williams, Z., Ohler, U., Tuschl, T., 2012. FMRP targets distinct mRNA sequence elements to regulate protein expression. *Nature* 492, 382–386.
- Ash, P.E., Bieniek, K.F., Gendron, T.F., Caulfield, T., Lin, W.L., DeJesus-Hernandez, M., van Blitterswijk, M.M., Jansen-West, K., Paul 3rd, J.W., Rademakers, R., Boylan, K.B., Dickson, D.W., Petrucelli, L., 2013. Unconventional translation of C9ORF72 GGGGCC expansion generates insoluble polypeptides specific to c9FTD/ALS. *Neuron* 77, 639–646.
- Ayhan, F., Perez, B.A., Shorrock, H.K., Zu, T., Banez-Coronel, M., Reid, T., Furuya, H., Clark, H.B., Troncoso, J.C., Ross, C.A., Subramony, S.H., Ashizawa, T., Wang, E.T., Yachnis, A.T., Ranum, L.P., 2018. CAG RAN polySer protein preferentially accumulates in white matter regions and is regulated by eIF3F. *EMBO J.* 37.
- Bagshaw, Andrew T.M., 2017. Functional mechanisms of microsatellite DNA in eukaryotic genomes. *Genome Biol. Evol.* 9, 2428–2443.
- Bakthavachalu, B., Huelsmeier, J., Sudhakaran, I.P., Hillebrand, J., Singh, A., Petrauskas, A., Thiagarajan, D., Sankaranarayanan, M., Mizoue, L., Anderson, E.N., Pandey, U.B., Ross, E., VijayRaghavan, K., Parker, R., Ramaswami, M., 2018. RNP-granule assembly via ataxin-2 disordered domains is required for long-term memory and neurodegeneration. *Neuron* 98 (754–66), e4.
- Banerjee, A., Ifrim, M.F., Valdez, A.N., Raj, N., Bassell, G.J., 2018. Aberrant RNA translation in fragile X syndrome: From FMRP mechanisms to emerging therapeutic strategies. *Brain Res.* 1693, 24–36.
- Banez-Coronel, M., Ayhan, F., Tarabochia, A.D., Zu, T., Perez, B.A., Tusi, S.K., Pletnikova, O., Borchelt, D.R., Ross, C.A., Margolis, R.L., Yachnis, A.T., Troncoso, J.C., Ranum, L.P., 2015. RAN translation in huntington disease. *Neuron* 88, 667–677.
- Bates, G., 1996. Expanded glutamines and neurodegeneration—a gain of insight. *Bioessays* 18, 175–178.
- Becker, L.A., Gitler, A.D., 2018. A neurodegenerative-disease protein forms beneficial aggregates in healthy muscle. *Nature* 563, 477–478.
- Becker, L.A., Huang, B., Bieri, G., Ma, R., Knowles, D.A., Jafar-Nejad, P., Messing, J., Kim, H.J., Soriano, A., Auburger, G., Pulst, S.M., Taylor, J.P., Rigo, F., Gitler, A.D., 2017. Therapeutic reduction of ataxin-2 extends lifespan and reduces pathology in TDP-43 mice. *Nature* 544, 367–371.
- Beerman, R.W., Jongens, T.A., 2011. A non-canonical start codon in the *Drosophila* fragile X gene yields two functional isoforms. *Neuroscience* 181, 48–66.
- Boeynaems, S., Bogaert, E., Michiels, E., Gijssels, I., Sieben, A., Jovicic, A., De Baets, G., Scheveneels, W., Steyaert, J., Cuijt, I., Verstrepen, K.J., Callaerts, P., Rousseau, F., Schymkowitz, J., Cruts, M., Van Broeckhoven, C., Van Damme, P., Gitler, A.D., Robberecht, W., Van Den Bosch, L., 2016. *Drosophila* screen connects nuclear transport genes to DPR pathology in c9ALS/FTD. *Sci. Rep.* 6, 20877.
- Boeynaems, S., Bogaert, E., Kovacs, D., Konijnenberg, A., Timmerman, E., Volkov, A., Guharoy, M., De Decker, M., Jaspers, T., Ryan, V.H., Janke, A.M., Baatsen, P., Vercruysee, T., Kolaitis, R.M., Daelemans, D., Taylor, J.P., Kedersha, N., Anderson, P., Impens, F., Sobott, F., Schymkowitz, J., Rousseau, F., Fawzi, N.L., Robberecht, W., Van Damme, P., Tompa, P., Van Den Bosch, L., 2017. Phase separation of c9orf72 dipeptide repeats perturbs stress granule dynamics. *Mol. Cell* 65 (1044–55), e5.
- Boeynaems, S., Alberti, S., Fawzi, N.L., Mittag, T., Polymenidou, M., Rousseau, F., Schymkowitz, J., Shorter, J., Wolozin, B., Van Den Bosch, L., Tompa, P., Fuxreiter, M., 2018. Protein phase separation: a new phase in cell biology. *Trends Cell Biol.* 28, 420–435.
- Boisvert, F.M., van Koningsbruggen, S., Navascues, J., Lamond, A.I., 2007. The multifunctional nucleolus. *Nat. Rev. Mol. Cell Biol.* 8, 574–585.
- Britten, R.J., Kohne, D.E., 1968. Repeated sequences in DNA. Hundreds of thousands of copies of DNA sequences have been incorporated into the genomes of higher organisms. *Science* 161, 529–540.
- Brown, V., Jin, P., Ceman, S., Darnell, J.C., O'Donnell, W.T., Tenenbaum, S.A., Jin, X., Feng, Y., Wilkinson, K.D., Keene, J.D., Darnell, R.B., Warren, S.T., 2001. Microarray identification of FMRP-associated brain mRNAs and altered mRNA translational profiles in fragile X syndrome. *Cell* 107, 477–487.
- Buchan, J.R., Parker, R., 2009. Eukaryotic stress granules: the ins and outs of translation. *Mol. Cell* 36, 932–941.
- Buchan, J.R., Muhlrad, D., Parker, R., 2008. P bodies promote stress granule assembly in *Saccharomyces cerevisiae*. *J. Cell Biol.* 183, 441–455.
- Buijsen, R.A., Visser, J.A., Kramer, P., Severijnen, E.A., Gearing, M., Charlet-Berguerand, N., Sherman, S.L., Berman, R.F., Willemsen, R., Hukema, R.K., 2016. Presence of inclusions positive for polyglycine containing protein, FMRpolyG, indicates that repeat-associated non-AUG translation plays a role in fragile X-associated primary

- ovarian insufficiency. *Hum. Reprod.* 31, 158–168.
- Burguete, A.S., Almeida, S., Gao, F.B., Kalb, R., Akins, M.R., Bonini, N.M., 2015. GGGGCC microsatellite RNA is neuritically localized, induces branching defects, and perturbs transport granule function. *Elife* 4, e08881.
- Campuzano, V., Montermini, L., Molto, M.D., Pianese, L., Cossee, M., Cavalcanti, F., Monros, E., Rodius, F., Duclos, F., Monticelli, A., Zara, F., Canizares, J., Koutnikova, H., Bidichandani, S.I., Gellera, C., Brice, A., Trouillas, P., De Michele, G., Filla, A., De Frutos, R., Palau, F., Patel, P.I., Di Donato, S., Mandel, J.L., Coccozza, S., Koenig, M., Pandolfo, M., 1996. Friedreich's ataxia: autosomal recessive disease caused by an intronic GAA triplet repeat expansion. *Science* 271, 1423–1427.
- Castaldo, I., Pinelli, M., Monticelli, A., Acquaviva, F., Giacchetti, M., Filla, A., Sacchetti, S., Keller, S., Avvedimento, V.E., Chiariotti, L., Coccozza, S., 2008. DNA methylation in intron 1 of the frataxin gene is related to GAA repeat length and age of onset in Friedreich ataxia patients. *J. Med. Genet.* 45, 808–812.
- Cheng, W., Wang, S., Mestre, A.A., Fu, C., Makarem, A., Xian, F., Hayes, L.R., Lopez-Gonzalez, R., Drenner, K., Jiang, J., Cleveland, D.W., Sun, S., 2018. C9ORF72 GGGGCC repeat-associated non-AUG translation is upregulated by stress through eIF2alpha phosphorylation. *Nat. Commun.* 9, 51.
- Coldwell, M.J., Mitchell, S.A., Stoneley, M., MacFarlane, M., Willis, A.E., 2000. Initiation of Apaf-1 translation by internal ribosome entry. *Oncogene* 19, 899–905.
- Conlon, E.G., Lu, L., Sharma, A., Yamazaki, T., Tang, T., Shneider, N.A., Manley, J.L., 2016. The C9ORF72 GGGGCC expansion forms RNA G-quadruplex inclusions and sequesters hnRNP H to disrupt splicing in ALS brains. *Elife* 5.
- Consortium, ENCODE project, 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature* 489, 57–74.
- Cooper-Knock, J., Walsh, M.J., Higginbottom, A., Robin Highley, J., Dickman, M.J., Edbauer, D., Ince, P.G., Wharton, S.B., Wilson, S.A., Kirby, J., Hautbergue, G.M., Shaw, P.J., 2014. Sequestration of multiple RNA recognition motif-containing proteins by C9orf72 repeat expansions. *Brain* 137, 2040–2051.
- Cortese, A., Simone, R., Sullivan, R., Vandrovcova, J., Tariq, H., Yan, Y.W., Humphrey, J., Jaunmuktane, Z., Sivakumar, P., Polke, J., Ilyas, M., Tribollet, E., Tomaselli, P.J., Devigili, G., Callegari, I., Versino, M., Salpietro, V., Efthymiou, S., Kaski, D., Wood, N.W., Andrade, N.S., Buglo, E., Rebelo, A., Rossor, A.M., Bronstein, A., Fratta, P., Marques, W.J., Zuchner, S., Reilly, M.M., Houlden, H., 2019. Biallelic expansion of an intronic repeat in RFC1 is a common cause of late-onset ataxia. *Nat. Genet.* 51, 649–658.
- Crick, S.L., Jayaraman, M., Frieden, C., Wetzel, R., Pappu, R.V., 2006. Fluorescence correlation spectroscopy shows that monomeric polyglutamine molecules form collapsed structures in aqueous solutions. *Proc. Natl. Acad. Sci. U. S. A.* 103, 16764–16769.
- Darling, A.L., Breydo, L., Rivas, E.G., Gebru, N.T., Zheng, D., Baker, J.D., Blair, L.J., Dickey, C.A., Koren 3rd, J., Uversky, V.N., 2019. Repeated repeat problems: combinatorial effect of C9orf72-derived dipeptide repeat proteins. *Int. J. Biol. Macromol.* 127, 136–145.
- Darnell, J.C., Jensen, K.B., Jin, P., Brown, V., Warren, S.T., Darnell, R.B., 2001. Fragile X mental retardation protein targets G quartet mRNAs important for neuronal function. *Cell* 107, 489–499.
- Darnell, J.C., Van Driesche, S.J., Zhang, C., Hung, K.Y., Mele, A., Fraser, C.E., Stone, E.F., Chen, C., Fak, J.J., Chi, S.W., Licatalosi, D.D., Richter, J.D., Darnell, R.B., 2011. FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell* 146, 247–261.
- Daughters, R.S., Tuttle, D.L., Gao, W., Ikeda, Y., Moseley, M.L., Ebner, T.J., Swanson, M.S., Ranum, L.P., 2009. RNA gain-of-function in spinocerebellar ataxia type 8. *PLoS Genet.* 5, e1000600.
- Deblor, E.W., Blobel, G., Hoelz, A., 2009. Nuclear transport comes full circle. *Nat. Struct. Mol. Biol.* 16, 457–459.
- DeJesus-Hernandez, M., Mackenzie, I.R., Boeve, B.F., Boxer, A.L., Baker, M., Rutherford, N.J., Nicholson, A.M., Finch, N.A., Flynn, H., Adamson, J., Kouri, N., Wojtas, A., Segedy, P., Hsiung, G.Y., Karydas, A., Seeley, W.W., Josephs, K.A., Coppola, G., Geschwind, D.H., Wszolek, Z.K., Feldman, H., Knopman, D.S., Petersen, R.C., Miller, B.L., Dickson, D.W., Boylan, K.B., Graff-Radford, N.R., Rademakers, R., 2011. Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 72, 245–256.
- Deng, H.X., Chen, W., Hong, S.T., Boycott, K.M., Gorrie, G.H., Siddique, N., Yang, Y., Fecto, F., Shi, Y., Zhai, H., Jiang, H., Hirano, M., Rampersaud, E., Jansen, G.H., Donkervoort, S., Bigio, E.H., Brooks, B.R., Ajroud, K., Sufit, R.L., Haines, J.L., Mugnaini, E., Pericak-Vance, M.A., Siddique, T., 2011. Mutations in UBQLN2 cause dominant X-linked juvenile and adult-onset ALS and ALS/dementia. *Nature* 477, 211–215.
- Di Prisco, G.V., Huang, W., Buffington, S.A., Hsu, C.C., Bonnen, P.E., Placzek, A.N., Sidrauski, C., Krnjevic, K., Kaufman, R.J., Walter, P., Costa-Mattioli, M., 2014. Translational control of mGluR-dependent long-term depression and object-place learning by eIF2alpha. *Nat. Neurosci.* 17, 1073–1082.
- Dickey, A.S., La Spada, A.R., 2018. Therapy development in Huntington disease: from current strategies to emerging opportunities. *Am. J. Med. Genet. A* 176, 842–861.
- Dobra, I., Pankivskiy, S., Samsonova, A., Pastre, D., Hamon, L., 2018. Relation between stress granules and cytoplasmic protein aggregates linked to neurodegenerative diseases. *Curr. Neurol. Neurosci. Rep.* 18, 107.
- Donnelly, C.J., Zhang, P.W., Pham, J.T., Haessler, A.R., Mistry, N.A., Vidensky, S., Daley, E.L., Poth, E.M., Hoover, B., Fines, D.M., Maragakis, N., Tienari, P.J., Petrucelli, L., Traynor, B.J., Wang, J., Rigo, F., Bennett, C.F., Blackshaw, S., Sattler, R., Rothstein, J.D., 2013. RNA toxicity from the ALS/FTD C9ORF72 expansion is mitigated by antisense intervention. *Neuron* 80, 415–428.
- Dougan, L., Li, J., Badilla, C.L., Berne, B.J., Fernandez, J.M., 2009. Single homopolymeric peptide chains collapse into mechanically rigid conformations. *Proc. Natl. Acad. Sci. U. S. A.* 106, 12605–12610.
- Dumbovic, G., Forcales, S.V., Peruch, M., 2017. Emerging roles of macrosatellite repeats in genome organization and disease development. *Epigenetics* 12, 515–526.
- Duvick, L., Barnes, J., Ebner, B., Agrawal, S., Andresen, M., Lim, J., Giesler, G.J., Zoghbi, H.Y., Orr, H.T., 2010. SCA1-like disease in mice expressing wild-type ataxin-1 with a serine to aspartic acid replacement at residue 776. *Neuron* 67, 929–935.
- Dyson, H.J., Wright, P.E., 2005. Intrinsically unstructured proteins and their functions. *Nat. Rev. Mol. Cell Biol.* 6, 197–208.
- Eichler, E.E., Kunst, C.B., Lugenbeel, K.A., Ryder, O.A., Davison, D., Warren, S.T., Nelson, D.L., 1995. Evolution of the cryptic FMR1 CGG repeat. *Nat. Genet.* 11, 301–308.
- Elden, A.C., Kim, H.J., Hart, M.P., Chen-Plotkin, A.S., Johnson, B.S., Fang, X., Armarkola, M., Geser, F., Greene, R., Lu, M.M., Padmanabhan, A., Clay-Falcone, D., McCluskey, L., Elman, L., Juhr, D., Gruber, P.J., Rub, U., Auburger, G., Trojanowski, J.Q., Lee, V.M., Van Deerlin, V.M., Bonini, N.M., Gitler, A.D., 2010. Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for ALS. *Nature* 466, 1069–1075.
- Farg, M.A., Soo, K.Y., Warraich, S.T., Sundaramoorthy, V., Blair, I.P., Atkin, J.D., 2013. Ataxin-2 interacts with FUS and intermediate-length polyglutamine expansions enhance FUS-related pathology in amyotrophic lateral sclerosis. *Hum. Mol. Genet.* 22, 717–728.
- Fay, M.M., Anderson, P.J., Ivanov, P., 2017. ALS/FTD-associated C9ORF72 repeat RNA promotes phase transitions in vitro and in cells. *Cell Rep.* 21, 3573–3584.
- Feng, Y., Zhang, F., Lokey, L.K., Chastain, J.L., Lakkis, L., Eberhart, D., Warren, S.T., 1995. Translational suppression by trinucleotide repeat expansion at FMR1. *Science* 268, 731–734.
- Flores, B.N., Dulchavsky, M.E., Krans, A., Sawaya, M.R., Paulson, H.L., Todd, P.K., Barmada, S.J., Ivanova, M.I., 2016. Distinct C9orf72-associated dipeptide repeat structures correlate with neuronal toxicity. *PLoS ONE* 11, e0165084.
- Fondon 3rd, J.W., Garner, H.R., 2004. Molecular origins of rapid and continuous morphological evolution. *Proc. Natl. Acad. Sci. U. S. A.* 101, 18058–18063.
- Freibaum, B.D., Lu, Y., Lopez-Gonzalez, R., Kim, N.C., Almeida, S., Lee, K.H., Badders, N., Valentine, M., Miller, B.L., Wong, P.C., Petrucelli, L., Kim, H.J., Gao, F.B., Taylor, J.P., 2015. GGGGCC repeat expansion in C9orf72 compromises nucleocytoplasmic transport. *Nature* 525, 129–133.
- Frey, S., Gorlich, D., 2007. A saturated FG-repeat hydrogel can reproduce the permeability properties of nuclear pore complexes. *Cell* 130, 512–523.
- Gall, J.G., 2003. The centennial of the Cajal body. *Nat. Rev. Mol. Cell Biol.* 4, 975–980.
- Gan, W., Rhoads, R.E., 1996. Internal initiation of translation directed by the 5'-untranslated region of the mRNA for eIF4G, a factor involved in the picornavirus-induced switch from cap-dependent to internal initiation. *J. Biol. Chem.* 271, 623–626.
- Gao, F.B., Richter, J.D., Cleveland, D.W., 2017. Rethinking unconventional translation in neurodegeneration. *Cell* 171, 994–1000.
- Gemayel, R., Vines, M.D., Legendre, M., Verstrepen, K.J., 2010. Variable tandem repeats accelerate evolution of coding and regulatory sequences. *Annu. Rev. Genet.* 44, 445–477.
- Gendron, T.F., Bieniek, K.F., Zhang, Y.J., Jansen-West, K., Ash, P.E., Caulfield, T., Daugherty, L., Dunmore, J.H., Castanedes-Casey, M., Chew, J., Cosio, D.M., van Blitterswijk, M., Lee, W.C., Rademakers, R., Boylan, K.B., Dickson, D.W., Petrucelli, L., 2013. Antisense transcripts of the expanded C9ORF72 hexanucleotide repeat form nuclear RNA foci and undergo repeat-associated non-ATG translation in c9FTD/ALS. *Acta Neuropathol.* 126, 829–844.
- Gendron, T.F., van Blitterswijk, M., Bieniek, K.F., Daugherty, L.M., Jiang, J., Rush, B.K., Pedraza, O., Lucas, J.A., Murray, M.E., Desaro, P., Robertson, A., Overstreet, K., Thomas, C.S., Crook, J.E., Castanedes-Casey, M., Rousseau, L., Josephs, K.A., Parisi, J.E., Knopman, D.S., Petersen, R.C., Boeve, B.F., Graff-Radford, N.R., Rademakers, R., Lagier-Tourenne, C., Edbauer, D., Cleveland, D.W., Dickson, D.W., Petrucelli, L., Boylan, K.B., 2015. Cerebellar c9RAN proteins associate with clinical and neuropathological characteristics of C9ORF72 repeat expansion carriers. *Acta Neuropathol.* 130, 559–573.
- Girstmair, H., Saffert, P., Rode, S., Czech, A., Holland, G., Bannert, N., Ignatova, Z., 2013. Depletion of cognate charged transfer RNA causes translational frameshifting within the expanded CAG stretch in huntingtin. *Cell Rep.* 3, 148–159.
- Gomes, E., Shorter, J., 2018. The molecular language of membraneless organelles. *J. Biol. Chem.* 294, 115–127.
- Greco, C.M., Berman, R.F., Martin, R.M., Tassone, F., Schwartz, P.H., Chang, A., Trapp, B.D., Iwahashi, C., Brunberg, J., Grigsby, J., Hessel, D., Becker, E.J., Papazian, J., Leehey, M.A., Hagerman, R.J., Hagerman, P.J., 2006. Neuropathology of fragile X-associated tremor/ataxia syndrome (FXTAS). *Brain* 129, 243–255.
- Green, K.M., Linsalata, A.E., Todd, P.K., 2016. RAN translation-what makes it run? *Brain Res.* 1647, 30–42.
- Green, K.M., Glineburg, M.R., Kearse, M.G., Flores, B.N., Linsalata, A.E., Fedak, S.J., Goldstrohm, A.C., Barmada, S.J., Todd, P.K., 2017. RAN translation at C9orf72-associated repeat expansions is selectively enhanced by the integrated stress response. *Nat. Commun.* 8, 2005.
- Grima, J.C., Daigle, J.G., Arbez, N., Cunningham, K.C., Zhang, K., Ochaba, J., Geater, C., Morozko, E., Stocksdale, J., Glatzer, J.C., Pham, J.T., Ahmed, I., Peng, Q., Wadhwa, H., Pletnikova, O., Troncoso, J.C., Duan, W., Snyder, S.H., Ranum, L.P.W., Thompson, L.M., Lloyd, T.E., Ross, C.A., Rothstein, J.D., 2017. Mutant huntingtin disrupts the nuclear pore complex. *Neuron* 94, 93–107 (e6).
- Gross, C., Bassell, G.J., 2012. Excess protein synthesis in FXS patient lymphoblastoid cells can be rescued with a p110beta-selective inhibitor. *Mol. Med.* 18, 336–345.
- Gross, C., Yao, X., Pong, D.L., Jeromin, A., Bassell, G.J., 2011. Fragile X mental retardation protein regulates protein expression and mRNA translation of the potassium channel Kv4.2. *J. Neurosci.* 31, 5693–5698.
- Group, The Huntington's Disease Collaborative Research, 1993. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* 72, 971–983.



- Guo, L., Kim, H.J., Wang, H., Monaghan, J., Freyermuth, F., Sung, J.C., O'Donovan, K., Fare, C.M., Diaz, Z., Singh, N., Zhang, Z.C., Coughlin, M., Sweeny, E.A., DeSantis, M.E., Jackrel, M.E., Rodell, C.B., Burdick, J.A., King, O.D., Gitler, A.D., Lagier-Tourenne, C., Pandey, U.B., Chook, Y.M., Taylor, J.P., Shorter, J., 2018. Nuclear-import receptors reverse aberrant phase transitions of rna-binding proteins with prion-like domains. *Cell* 173, 677–692 (e20).
- Haeussler, A.R., Donnelly, C.J., Periz, G., Simko, E.A., Shaw, P.G., Kim, M.S., Maragakis, N.J., Troncoso, J.C., Pandey, A., Sattler, R., Rothstein, J.D., Wang, J., 2014. C9orf72 nucleotide repeat structures initiate molecular cascades of disease. *Nature* 507, 195–200.
- Hagerman, P., 2013. Fragile X-associated tremor/ataxia syndrome (FXTAS): pathology and mechanisms. *Acta Neuropathol.* 126, 1–19.
- Hannan, A.J., 2018a. Tandem repeats and repeatomes: delving deeper into the 'dark matter' of genomes. *EBioMedicine* 31, 3–4.
- Hannan, A.J., 2018b. Tandem repeats mediating genetic plasticity in health and disease. *Nat. Rev. Genet.* 19, 286–298.
- Haran, G., 2012. How, when and why proteins collapse: the relation to folding. *Curr. Opin. Struct. Biol.* 22, 14–20.
- Harding, H.P., Novoa, I., Zhang, Y., Zeng, H., Wek, R., Schapira, M., Ron, D., 2000. Regulated translation initiation controls stress-induced gene expression in mammalian cells. *Mol. Cell* 6, 1099–1108.
- Harding, H.P., Zhang, Y., Zeng, H., Novoa, I., Lu, P.D., Calton, M., Sadri, N., Yun, C., Popko, B., Paules, R., Stojdl, D.F., Bell, J.C., Hettmann, T., Leiden, J.M., Ron, D., 2003. An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol. Cell* 11, 619–633.
- Hart, M.P., Gitler, A.D., 2012. ALS-associated ataxin 2 polyQ expansions enhance stress-induced caspase 3 activation and increase TDP-43 pathological modifications. *J. Neurosci.* 32, 9133–9142.
- Hartmann, H., Hornburg, D., Czuppa, M., Bader, J., Michaelsen, M., Farny, D., Arzberger, T., Mann, M., Meissner, F., Edbauer, D., 2018. Proteomics and C9orf72 neuro-pathology identify ribosomes as poly-GR/PR interactors driving toxicity. *Life Sci. Alliance* 1, e201800070.
- Hashem, V., Galloway, J.N., Mori, M., Willemsen, R., Oostra, B.A., Paylor, R., Nelson, D.L., 2009. Ectopic expression of CGG containing mRNA is neurotoxic in mammals. *Hum. Mol. Genet.* 18, 2443–2451.
- Hilliker, A., Gao, Z., Jankowsky, E., Parker, R., 2011. The DEAD-box protein Ded1 modulates translation by the formation and resolution of an eIF4F-mRNA complex. *Mol. Cell* 43, 962–972.
- Hinnebusch, A.G., 2011. Molecular mechanism of scanning and start codon selection in eukaryotes. *Microbiol. Mol. Biol. Rev.* 75, 434–467 (first page of table of contents).
- Hinnebusch, A.G., Ivanov, I.P., Sonenberg, N., 2016. Translational control by 5'-untranslated regions of eukaryotic mRNAs. *Science* 352, 1413–1416.
- Hoelz, A., Debler, E.W., Blobel, G., 2011. The structure of the nuclear pore complex. *Annu. Rev. Biochem.* 80, 613–643.
- Holmquist, G.P., Dancis, B., 1979. Telomere replication, kinetochore organizers, and satellite DNA evolution. *Proc. Natl. Acad. Sci. U. S. A.* 76, 4566–4570.
- Huber, K.M., Gallagher, S.M., Warren, S.T., Bear, M.F., 2002. Altered synaptic plasticity in a mouse model of fragile X mental retardation. *Proc. Natl. Acad. Sci. U. S. A.* 99, 7746–7750.
- Hyman, A.A., Brangwynne, C.P., 2011. Beyond stereospecificity: liquids and mesoscale organization of cytoplasm. *Dev. Cell* 21, 14–16.
- Hyman, A.A., Weber, C.A., Julicher, F., 2014. Liquid-liquid phase separation in biology. *Annu. Rev. Cell Dev. Biol.* 30, 39–58.
- Ifrim, M.F., Williams, K.R., Bassell, G.J., 2015. Single-Molecule Imaging of PSD-95 mRNA Translation in Dendrites and Its Dysregulation in a Mouse Model of Fragile X Syndrome. *J. Neurosci.* 35, 7116–7130.
- Imbert, G., Saudou, F., Yvert, G., Devys, D., Trottier, Y., Garnier, J.M., Weber, C., Mandel, J.L., Cancel, G., Abbas, N., Durr, A., Didierjean, O., Stevanin, G., Agid, Y., Brice, A., 1996. Cloning of the gene for spinocerebellar ataxia 2 reveals a locus with high sensitivity to expanded CAG/glutamine repeats. *Nat. Genet.* 14, 285–291.
- Ishiguro, T., Sato, N., Ueyama, M., Fujikake, N., Sellier, C., Kanegami, A., Tokuda, E., Zamiri, B., Gall-Duncan, T., Mirceta, M., Furukawa, Y., Yokota, T., Wada, K., Taylor, J.P., Pearson, C.E., Charlet-Bergerand, N., Mizusawa, H., Nagai, Y., Ishikawa, K., 2017. Regulatory role of RNA chaperone TDP-43 for RNA misfolding and repeat-associated translation in SCA31. *Neuron* 94 (108–24), e7.
- Ishiura, H., Doi, K., Mitsui, J., Yoshimura, J., Matsukawa, M.K., Fujiyama, A., Toyoshima, Y., Kakita, A., Takahashi, H., Suzuki, Y., Sugano, S., Qu, W., Ichikawa, K., Yurino, H., Higasa, K., Shibata, S., Mitsue, A., Tanaka, M., Ichikawa, Y., Takahashi, Y., Date, H., Matsukawa, T., Kanda, J., Nakamoto, F.K., Higashihara, M., Abe, K., Koike, R., Sasagawa, M., Kuroha, Y., Hasegawa, N., Kanewasa, N., Kondo, T., Hitomi, T., Tada, M., Takano, H., Saito, Y., Sanpei, K., Onodera, O., Nishizawa, M., Nakamura, M., Yasuda, T., Sakiyama, Y., Otsuka, M., Ueki, A., Kaida, K.I., Shimizu, J., Hanajima, R., Hayashi, T., Terao, Y., Inomata-Terada, S., Hamada, M., Shiota, Y., Kubota, A., Ugawa, Y., Koh, K., Takiyama, Y., Ohsawa-Yoshida, N., Ishiura, S., Yamasaki, R., Tamaoka, A., Akiyama, H., Otsuki, T., Sano, A., Ikeda, A., Goto, J., Morishita, S., Tsuji, S., 2018. Expansions of intronic TTCTA and TTTTA repeats in benign adult familial myoclonic epilepsy. *Nat. Genet.* 50, 581–590.
- Jain, A., Vale, R.D., 2017. RNA phase transitions in repeat expansion disorders. *Nature* 546, 243–247.
- Jang, S.K., Krausslich, H.G., Nicklin, M.J., Duke, G.M., Palmenberg, A.C., Wimmer, E., 1988. A segment of the 5' nontranslated region of encephalomyocarditis virus RNA directs internal entry of ribosomes during in vitro translation. *J. Virol.* 62, 2636–2643.
- Jiang, H., Mankodi, A., Swanson, M.S., Moxley, R.T., Thornton, C.A., 2004. Myotonic dystrophy type 1 is associated with nuclear foci of mutant RNA, sequestration of muscleblind proteins and deregulated alternative splicing in neurons. *Hum. Mol. Genet.* 13, 3079–3088.
- Jin, P., Duan, R., Qurashi, A., Qin, Y., Tian, D., Rosser, T.C., Liu, H., Feng, Y., Warren, S.T., 2007. Pur alpha binds to rCGG repeats and modulates repeat-mediated neurodegeneration in a Drosophila model of fragile X tremor/ataxia syndrome. *Neuron* 55, 556–564.
- Jones, K.W., 1970. Chromosomal and nuclear location of mouse satellite DNA in individual cells. *Nature* 225, 912–915.
- Jones, K.W., 1973. Satellite DNA. *J. Med. Genet.* 10, 273–281.
- Jovicic, A., Mertens, J., Boeynaems, S., Bogaert, E., Chai, N., Yamada, S.B., Paul 3rd, J.W., Sun, S., Herdy, J.R., Bieri, G., Kramer, N.J., Gage, F.H., Van Den Bosch, L., Robberecht, W., Gitler, A.D., 2015. Modifiers of C9orf72 dipeptide repeat toxicity connect nucleocytoplasmic transport defects to FTD/ALS. *Nat. Neurosci.* 18, 1226–1229.
- Kamsteeg, E.J., Kress, W., Catalli, C., Hertz, J.M., Witsch-Baumgartner, M., Buckley, M.F., van Engelen, B.G., Schwartz, M., Scheffer, H., 2012. Best practice guidelines and recommendations on the molecular diagnosis of myotonic dystrophy types 1 and 2. *Eur. J. Hum. Genet.* 20, 1203–1208.
- Kanekura, K., Yagi, T., Cammack, A.J., Mahadevan, J., Kuroda, M., Harms, M.B., Miller, T.M., Urano, F., 2016. Poly-dipeptides encoded by the C9ORF72 repeats block global protein translation. *Hum. Mol. Genet.* 25, 1803–1813.
- Katsuno, M., Adachi, H., Kume, A., Li, M., Nakagomi, Y., Niwa, H., Sang, C., Kobayashi, Y., Doyu, M., Sobue, G., 2002. Testosterone reduction prevents phenotypic expression in a transgenic mouse model of spinal and bulbar muscular atrophy. *Neuron* 35, 843–854.
- Katsuno, M., Adachi, H., Doyu, M., Minamiyama, M., Sang, C., Kobayashi, Y., Inukai, A., Sobue, G., 2003. Leuporelin rescues polyglutamine-dependent phenotypes in a transgenic mouse model of spinal and bulbar muscular atrophy. *Nat. Med.* 9, 768–773.
- Kearse, M.G., Todd, P.K., 2014. Repeat-associated non-AUG translation and its impact in neurodegenerative disease. *Neurotherapeutics* 11, 721–731.
- Kearse, M.G., Green, K.M., Krans, A., Rodriguez, C.M., Linsalata, A.E., Goldstrohm, A.C., Todd, P.K., 2016. CGG Repeat-Associated Non-AUG Translation Utilizes a Cap-Dependent Scanning Mechanism of Initiation to Produce Toxic Proteins. *Mol. Cell* 62, 314–322.
- Kedersha, N.L., Gupta, M., Li, W., Miller, I., Anderson, P., 1999. RNA-binding proteins TIA-1 and TIAR link the phosphorylation of eIF-2 alpha to the assembly of mammalian stress granules. *J. Cell Biol.* 147, 1431–1442.
- Kim, H.J., Taylor, J.P., 2017. Lost in transportation: nucleocytoplasmic transport defects in ALS and other neurodegenerative diseases. *Neuron* 96, 285–297.
- Kim, H.J., Kim, N.C., Wang, Y.D., Scarborough, E.A., Moore, J., Diaz, Z., MacLea, K.S., Freibaum, B., Li, S., Molliex, A., Kanagaraj, A.P., Carter, R., Boylan, K.B., Wojtas, A.M., Rademakers, R., Pinkus, J.L., Greenberg, S.A., Trojanowski, J.Q., Traynor, B.J., Smith, B.N., Topp, S., Gkazi, A.S., Miller, J., Shaw, C.E., Kottlors, M., Kirschner, J., Pestronk, A., Li, Y.R., Ford, A.F., Gitler, A.D., Benatar, M., King, O.D., Kimonis, V.E., Ross, E.D., Weihl, C.C., Shorter, J., Taylor, J.P., 2013. Mutations in prion-like domains in hnRNPA2B1 and hnRNPA1 cause multisystem proteinopathy and ALS. *Nature* 495, 467–473.
- Kit, S., 1962. Species differences in animal deoxyribonucleic acid as revealed by equilibrium sedimentation in density gradients. *Nature* 193, 274–275.
- Kordasiewicz, H.B., Stanek, L.M., Wancewicz, E.V., Mazur, C., McAlonis, M.M., Pytel, K.A., Artates, J.W., Weiss, A., Cheng, S.H., Shihabuddin, L.S., Hung, G., Bennett, C.F., Cleveland, D.W., 2012. Sustained therapeutic reversal of Huntington's disease by transient repression of huntingtin synthesis. *Neuron* 74, 1031–1044.
- Kozak, M., 1984. Selection of initiation sites by eucaryotic ribosomes: effect of inserting AUG triplets upstream from the coding sequence for preproinsulin. *Nucleic Acids Res.* 12, 3873–3893.
- Kozlov, G., Safaei, N., Rosenauer, A., Gehring, K., 2010. Structural basis of binding of P-body-associated proteins GW182 and ataxin-2 by the Mille domain of poly(A)-binding protein. *J. Biol. Chem.* 285, 13599–13606.
- Krans, A., Kearse, M.G., Todd, P.K., 2016. Repeat-associated non-AUG translation from antisense CCG repeats in fragile X tremor/ataxia syndrome. *Ann. Neurol.* 80, 871–881.
- Kruglyak, S., Durrett, R.T., Schug, M.D., Aquadro, C.F., 1998. Equilibrium distributions of microsatellite repeat length resulting from a balance between slippage events and point mutations. *Proc. Natl. Acad. Sci. U. S. A.* 95, 10774–10778.
- Kwon, I., Xiang, S., Kato, M., Wu, L., Theodoropoulos, P., Wang, T., Kim, J., Yun, J., Xie, Y., McKnight, S.L., 2014. Poly-dipeptides encoded by the C9orf72 repeats bind nucleoli, impede RNA biogenesis, and kill cells. *Science* 345, 1139–1145.
- La Spada, A.R., Wilson, E.M., Lubahn, D.B., Harding, A.E., Fischbeck, K.H., 1991. Androgen receptor gene mutations in X-linked spinal and bulbar muscular atrophy. *Nature* 352, 77–79.
- LaCroix, A.J., Stables, D., Sahraoui, R., Adam, M.P., Mehaffey, M., Kernan, K., Myers, C.T., Fagerstrom, C., Anadiotis, G., Akkari, Y.M., Robbins, K.M., Gripp, K.W., Baratela, W.A.R., Bober, M.B., Duker, A.L., Doherty, D., Dempsey, J.C., Miller, D.G., Kircher, M., Bamshad, M.J., Nickerson, D.A., Mefford, H.C., Sol-Church, K., 2019. GGC repeat expansion and exon 1 methylation of XYLTI1 is a common pathogenic variant in baratela-scott syndrome. *Am. J. Hum. Genet.* 104, 35–44.
- Laggerbauer, B., Ostareck, D., Keidel, E.M., Ostareck-Lederer, A., Fischer, U., 2001. Evidence that fragile X mental retardation protein is a negative regulator of translation. *Hum. Mol. Genet.* 10, 329–338.
- Lagier-Tourenne, C., Baughn, M., Rigo, F., Sun, S., Liu, P., Li, H.R., Jiang, J., Watt, A.T., Chun, S., Katz, M., Qiu, J., Sun, Y., Ling, S.C., Zhu, Q., Polymeridou, M., Drenner, K., Artates, J.W., McAlonis-Downes, M., Markmiller, S., Hutt, K.R., Pizzo, D.P., Cady, J., Harms, M.B., Baloh, R.H., Vandenberg, S.R., Yeo, G.W., Fu, X.D., Bennett, C.F., Cleveland, D.W., Ravits, J., 2013. Targeted degradation of sense and antisense C9orf72 RNA foci as therapy for ALS and frontotemporal degeneration. *Proc. Natl.*

- Acad. Sci. U. S. A. 110, E4530–E4539.
- Lander, E.S., Linton, L.M., Birren, B., Nussbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., Funke, R., Gage, D., Harris, K., Heaford, A., Howland, J., Kann, L., LeHoczy, J., LeVine, R., McEwan, P., McKernan, K., Meldrum, J., Mesirov, J.P., Miranda, C., Morris, W., Naylor, J., Raymond, C., Rosetti, M., Santos, R., Sheridan, A., Sougnez, C., Stange-Thomann, Y., Stojanovic, N., Subramanian, A., Wyman, D., Rogers, J., Sulston, J., Ainscough, R., Beck, S., Bentley, D., Burton, J., Clee, C., Carter, N., Coulson, A., Deadman, R., Deloukas, P., Dunham, A., Dunham, I., Durbin, R., French, L., Grafham, D., Gregory, S., Hubbard, T., Humphray, S., Hunt, A., Jones, M., Lloyd, C., McMurray, A., Matthews, L., Mercer, S., Milne, S., Mullikin, J.C., Mungall, A., Plumb, R., Ross, M., Shownkeen, R., Sims, S., Waterston, R.H., Wilson, R.K., Hillier, L.W., McPherson, J.D., Marra, M.A., Mardis, E.R., Fulton, L.A., Chinwalla, A.T., Pepin, K.H., Gish, W.R., Chissoe, S.L., Wendl, M.C., Delehaanty, K.D., Miner, T.L., Delehaanty, A., Kramer, J.B., Cook, L.L., Fulton, R.S., Johnson, D.L., Minx, P.J., Clifton, S.W., Hawkins, T., Branscomb, E., Predki, P., Richardson, P., Wenning, S., Slezak, T., Doggett, N., Cheng, J.F., Olsen, A., Lucas, S., Elkin, C., Ueberbacher, E., Frazier, M., Gibbs, R.A., Muzny, D.M., Scherer, S.E., Bouck, J.B., Sodergren, E.J., Worley, K.C., Rives, C.M., Gorrell, J.H., Metzker, M.L., Naylor, S.L., Kucherlapati, R.S., Nelson, D.L., Weinstock, G.M., Sakaki, Y., Fujiyama, A., Hattori, M., Yada, T., Toyoda, A., Itoh, T., Kawagoe, C., Watanabe, H., Totoki, Y., Taylor, T., Weissenbach, J., Heilig, R., Saurin, W., Artiguenave, F., Brottier, P., Bruls, T., Pelletier, E., Robert, C., Wincker, P., Smith, D.R., Doucette-Stamm, L., Rubenfield, M., Weinstock, K., Lee, H.M., Dubois, J., Rosenthal, A., Platzer, M., Nyakatura, G., Taudien, S., Rump, A., Yang, H., Yu, J., Wang, J., Huang, G., Gu, J., Hood, L., Rowen, L., Madan, A., Qin, S., Davis, R.W., Federspiel, N.A., Abola, A.P., Proctor, M.J., Myers, R.M., Schmutz, J., Dickson, M., Grimwood, J., Cox, D.R., Olson, M.V., Kaul, R., Raymond, C., Shimizu, N., Kawasaki, K., Minoshima, S., Evans, G.A., Athanasiou, M., Schultz, R., Roe, B.A., Chen, F., Pan, H., Ramser, J., Lehrach, H., Reinhardt, R., McCombie, R., de la Bastide, M., Dedhia, N., Blocker, H., Hornischer, K., Nordsiek, G., Agarwala, R., Aravind, L., Bailey, J.A., Bateman, A., Batzoglou, S., Birney, E., Bork, P., Brown, D.G., Burge, C.B., Cerutti, L., Chen, H.C., Church, D., Clamp, M., Copley, R.R., Doerks, T., Eddy, S.R., Eichler, E.E., Furey, T.S., Galagan, J., Gilbert, J.G., Harmon, C., Hayashizaki, Y., Haussler, D., Hermjakob, H., Hokamp, K., Jang, W., Johnson, L.S., Jones, T.A., Kasif, S., Kasprzyk, A., Kennedy, S., Kent, W.J., Kitts, P., Koonin, E.V., Korf, I., Kulp, D., Lancet, D., Lowe, T.M., McLysaght, A., Mikkelsen, T., Moran, J.V., Mulder, N., Pollara, V.J., Ponting, C.P., Schuler, G., Schultz, J., Slater, G., Smit, A.F., Stupka, E., Szustakowski, J., Thierry-Mieg, D., Thierry-Mieg, J., Wagner, L., Wallis, J., Wheeler, R., Williams, A., Wolf, Y.I., Wolfe, K.H., Yang, S.P., Yeh, R.F., Collins, F., Guyer, M.S., Peterson, J., Felsenfeld, A., Wetterstrand, K.A., Patrino, A., Morgan, M.J., de Jong, P., Catanese, J.J., Osoegawa, K., Shizuya, H., Choi, S., Chen, Y.J., Szustakowski, J., 2001. Initial sequencing and analysis of the human genome. *Nature* 409, 860–921.
- Lastres-Becker, I., Nonis, D., Eich, F., Klinkenberg, M., Gorospe, M., Kotter, P., Klein, F.A., Kedersha, N., Auburger, G., 2016. Mammalian ataxin-2 modulates translation control at the pre-initiation complex via PI3K/mTOR and is induced by starvation. *Biochim. Biophys. Acta* 1862, 1558–1569.
- Lee, Y.B., Chen, H.J., Peres, J.N., Gomez-Deza, J., Attig, J., Stalekar, M., Troakes, C., Nishimura, A.L., Scotter, E.L., Vance, C., Adachi, Y., Sardone, V., Miller, J.W., Smith, B.N., Gallo, J.M., Ule, J., Hirth, F., Rogelj, B., Houart, C., Shaw, C.E., 2013. Hexanucleotide repeats in ALS/FTD form length-dependent RNA foci, sequester RNA binding proteins, and are neurotoxic. *Cell Rep.* 5, 1178–1186.
- Lee, K.H., Zhang, P., Kim, H.J., Mitrea, D.M., Sarkar, M., Freibaum, B.D., Cika, J., Coughlin, M., Messing, J., Molliex, A., Maxwell, B.A., Kim, N.C., Temirov, J., Moore, J., Kolaitis, R.M., Shaw, T.I., Bai, B., Peng, J., Kriwacki, R.W., Taylor, J.P., 2016. C9orf72 dipeptide repeats impair the assembly, dynamics, and function of membrane-less organelles. *Cell* 167, 774–788 (e17).
- Lee, Y.B., Baskaran, P., Gomez-Deza, J., Chen, H.J., Nishimura, A.L., Smith, B.N., Troakes, C., Adachi, Y., Stepto, A., Petrucelli, L., Gallo, J.M., Hirth, F., Rogelj, B., Guthrie, S., Shaw, C.E., 2017. C9orf72 poly GA RAN-translated protein plays a key role in amyotrophic lateral sclerosis via aggregation and toxicity. *Hum. Mol. Genet.* 26, 4765–4777.
- Lee, J.K., Conrad, A., Epping, E., Mathews, K., Magnotta, V., Dawson, J.D., Nopoulos, P., 2018. Effect of trinucleotide repeats in the huntington's gene on intelligence. *EBioMedicine* 31, 47–53.
- Li, Z., Zhang, Y., Ku, L., Wilkinson, K.D., Warren, S.T., Feng, Y., 2001. The fragile X mental retardation protein inhibits translation via interacting with mRNA. *Nucleic Acids Res.* 29, 2276–2283.
- Li, Y.R., King, O.D., Shorter, J., Gitler, A.D., 2013. Stress granules as crucibles of ALS pathogenesis. *J. Cell Biol.* 201, 361–372.
- Lieberman, A.P., Shakkottai, V.G., Albin, R.L., 2019. Polyglutamine repeats in neurodegenerative diseases. *Annu. Rev. Pathol.* 14, 1–27.
- Lin, Y., Protter, D.S., Rosen, M.K., Parker, R., 2015. Formation and maturation of phase-separated liquid droplets by RNA-binding proteins. *Mol. Cell* 60, 208–219.
- Lin, Y., Mori, E., Kato, M., Xiang, S., Wu, L., Kwon, I., McKnight, S.L., 2016. Toxic PR poly-dipeptides encoded by the C9orf72 repeat expansion target LC domain polymers. *Cell* 167 (789–802), e12.
- Liquori, C.L., Ricker, K., Moseley, M.L., Jacobsen, J.F., Kress, W., Naylor, S.L., Day, J.W., Ranum, L.P., 2001. Myotonic dystrophy type 2 caused by a CCTG expansion in intron 1 of ZNF9. *Science* 293, 864–867.
- Liu, B., Li, Y., Stackpole, E.E., Novak, A., Gao, Y., Zhao, Y., Zhao, X., Richter, J.D., 2018. Regulatory discrimination of mRNAs by FMRP controls mouse adult neural stem cell differentiation. *Proc. Natl. Acad. Sci. U. S. A.* 115 (E11397–e1405).
- Liu-Yesucevitz, L., Bassell, G.J., Gitler, A.D., Hart, A.C., Klann, E., Richter, J.D., Warren, S.T., Wolozin, B., 2011. Local RNA translation at the synapse and in disease. *J. Neurosci.* 31, 16086–16093.
- Lu, C., Lin, L., Tan, H., Wu, H., Sherman, S.L., Gao, F., Jin, P., Chen, D., 2012. Fragile X premutation RNA is sufficient to cause primary ovarian insufficiency in mice. *Hum. Mol. Genet.* 21, 5039–5047.
- Mackenzie, I.R., Frick, P., Grasser, F.A., Gendron, T.F., Petrucelli, L., Cashman, N.R., Edbauer, D., Kremmer, E., Prudlo, J., Troost, D., Neumann, M., 2015. Quantitative analysis and clinico-pathological correlations of different dipeptide repeat proteinopathies in C9ORF72 mutation carriers. *Acta Neuropathol.* 130, 845–861.
- Makise, M., Mackay, D.R., Elgort, S., Shankaran, S.S., Adam, S.A., Ullman, K.S., 2012. The nup153-nup50 protein interface and its role in nuclear import. *J. Biol. Chem.* 287, 38515–38522.
- Malinowska, L., Kroschwald, S., Alberti, S., 2013. Protein disorder, prion propensities, and self-organizing macromolecular collectives. *Biochim. Biophys. Acta* 1834, 918–931.
- Mankodi, A., Urbani, C.R., Yuan, Q.P., Moxley, R.T., Sansone, V., Krym, M., Henderson, D., Schalling, M., Swanson, M.S., Thornton, C.A., 2001. Muscleblind localizes to nuclear foci of aberrant RNA in myotonic dystrophy types 1 and 2. *Hum. Mol. Genet.* 10, 2165–2170.
- Mann, D.M., Rollinson, S., Robinson, A., Bennion Callister, J., Thompson, J.C., Snowden, J.S., Gendron, T., Petrucelli, L., Masuda-Suzukake, M., Hasegawa, M., Davidson, Y., Pickering-Brown, S., 2013. Dipeptide repeat proteins are present in the p62 positive inclusions in patients with frontotemporal lobar degeneration and motor neuron disease associated with expansions in C9ORF72. *Acta Neuropathol. Commun.* 1, 68.
- Margolis, J.M., Schoser, B.G., Moseley, M.L., Day, J.W., Ranum, L.P., 2006. DM2 intronic expansions: evidence for CCUG accumulation without flanking sequence or effects on ZNF9 mRNA processing or protein expression. *Hum. Mol. Genet.* 15, 1808–1815.
- McConnell, M.J., Moran, J.V., Abyzov, A., Akbarian, S., Bae, T., Cortes-Ciriano, I., Erwin, J.A., Fasching, L., Flasch, D.A., Freed, D., Ganz, J., Jaffe, A.E., Kwan, K.Y., Kwon, M., Lodato, M.A., Mills, R.E., Paquola, A.C.M., Rodin, R.E., Rosenbluh, C., Sestan, N., Sherman, M.A., Shin, J.H., Song, S., Straub, R.E., Thorpe, J., Weinberger, D.R., Urban, A.E., Zhou, B., Gage, F.H., Lehner, T., Senthil, G., Walsh, C.A., Chess, A., Courchesne, E., Gleeson, J.G., Kidd, J.M., Park, P.J., Pevsner, J., Vaccarino, F.M., 2017. Intersection of diverse neuronal genomes and neuropsychiatric disease: the brain somatic mosaicism network. *Science* 356.
- Meola, G., Cardani, R., 2015. Myotonic dystrophy type 2: an update on clinical aspects, genetic and pathomolecular mechanism. *J. Neuromuscul. Dis.* 2, S59–S71.
- Miller, J.W., Urbani, C.R., Teng-Ummay, P., Stenberg, M.G., Byrne, B.J., Thornton, C.A., Swanson, M.S., 2000. Recruitment of human muscleblind proteins to (CUG)(n) expansions associated with myotonic dystrophy. *EMBO J.* 19, 4439–4448.
- Mizielinska, S., Gronke, S., Niccoli, T., Ridler, C.E., Clayton, E.L., Devoy, A., Moens, T., Norona, F.E., Woollacott, I.O.C., Pietrzyk, J., Cleverley, K., Nicoll, A.J., Pickering-Brown, S., Dols, J., Cabecinha, M., Hendrich, O., Fratta, P., Fisher, E.M.C., Partridge, L., Isaacs, A.M., 2014. C9orf72 repeat expansions cause neurodegeneration in Drosophila through arginine-rich proteins. *Science* 345, 1192–1194.
- Mizielinska, S., Ridler, C.E., Balendra, R., Thoeng, A., Woodling, N.S., Grasser, F.A., Plagnol, V., Lashley, T., Partridge, L., Isaacs, A.M., 2017. Bidirectional nucleolar dysfunction in C9orf72 frontotemporal lobar degeneration. *Acta Neuropathol. Commun.* 5, 29.
- Moens, T.G., Mizielinska, S., Niccoli, T., Mitchell, J.S., Thoeng, A., Ridler, C.E., Gronke, S., Esser, J., Heslegrave, A., Zetterberg, H., Partridge, L., Isaacs, A.M., 2018. Sense and antisense RNA are not toxic in Drosophila models of C9orf72-associated ALS/FTD. *Acta Neuropathol.* 135, 445–457.
- Moens, T.G., Niccoli, T., Wilson, K.M., Atilano, M.L., Birsá, N., Gittings, L.M., Holbling, B.V., Dyson, M.C., Thoeng, A., Neeves, J., Glaria, I., Yu, L., Bussmann, J., Storkebaum, E., Pardo, M., Choudhary, J.S., Fratta, P., Partridge, L., Isaacs, A.M., 2019. C9orf72 arginine-rich dipeptide proteins interact with ribosomal proteins in vivo to induce a toxic translational arrest that is rescued by eIF1A. *Acta Neuropathol.* 137, 487–500.
- Mohr, D., Frey, S., Fischer, T., Guttler, T., Gorlich, D., 2009. Characterisation of the passive permeability barrier of nuclear pore complexes. *EMBO J.* 28, 2541–2553.
- Molliex, A., Temirov, J., Lee, J., Coughlin, M., Kanagaraj, A.P., Kim, H.J., Mittag, T., Taylor, J.P., 2015. Phase separation by low complexity domains promotes stress granule assembly and drives pathological fibrillization. *Cell* 163, 123–133.
- Mori, K., Weng, S.M., Arzberger, T., May, S., Rentzsch, K., Kremmer, E., Schmid, B., Kretschmar, H.A., Cruts, M., Van Broeckhoven, C., Haass, C., Edbauer, D., 2013. The C9orf72 GGGGCC repeat is translated into aggregating dipeptide-repeat proteins in FTLD/ALS. *Science* 339, 1335–1338.
- Morley, J.F., Brignull, H.R., Weyers, J.J., Morimoto, R.I., 2002. The threshold for polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and influenced by aging in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U. S. A.* 99, 10417–10422.
- Moseley, M.L., Zu, T., Ikeda, Y., Gao, W., Mosemiller, A.K., Daughters, R.S., Chen, G., Weatherspoon, M.R., Clark, H.B., Ebner, T.J., Day, J.W., Ranum, L.P., 2006. Bidirectional expression of CUG and CAG expansion transcripts and intranuclear polyglutamine inclusions in spinocerebellar ataxia type 8. *Nat. Genet.* 38, 758–769.
- Muslimov, I.A., Patel, M.V., Rose, A., Tiedge, H., 2011. Spatial code recognition in neuronal RNA targeting: role of RNA-hnRNP A2 interactions. *J. Cell Biol.* 194, 441–457.
- Nakamori, Masayuki, Sobczak, Krzysztof, Puwanant, Araya, Welle, Steve, Eichinger, Katy, Pandya, Shree, Dekdebrun, Jeannette, Heatwole, Chad R., McDermott, Michael P., Chen, Tian, Cline, Melissa, Tawil, Rabi, Osborne, Robert J., Wheeler, Thurman M., Swanson, Maurice S., Moxley 3rd, Richard T., Thornton, Charles A., 2013. Splicing biomarkers of disease severity in myotonic dystrophy. *Ann. Neurol.* 74, 862–872.
- Nonhoff, U., Ralser, M., Welzel, F., Piccini, I., Balzeret, D., Yaspo, M.L., Lehrach, H., Krobisch, S., 2007. Ataxin-2 interacts with the DEAD/H-box RNA helicase DDX6 and interferes with P-bodies and stress granules. *Mol. Biol. Cell* 18, 1385–1396.
- Oh, S.Y., He, F., Krans, A., Frazer, M., Taylor, J.P., Paulson, H.L., Todd, P.K., 2015. RAN translation at CGG repeats induces ubiquitin proteasome system impairment in models of fragile X-associated tremor ataxia syndrome. *Hum. Mol. Genet.* 24, 4317–4326.

- Ordway, J.M., Tallaksen-Greene, S., Gutekunst, C.A., Bernstein, E.M., Cearley, J.A., Wiener, H.W., Dure, L.S., Lindsey, R., Hersch, S.M., Jope, R.S., Albin, R.L., Detloff, P.J., 1997. Ectopically expressed CAG repeats cause intranuclear inclusions and a progressive late onset neurological phenotype in the mouse. *Cell* 91, 753–763.
- Orr, H.T., 2011. FTD and ALS: genetic ties that bind. *Neuron* 72, 189–190.
- Pakos-Zebrucka, K., Koryga, I., Mnich, K., Ljubic, M., Samali, A., 2016. The Integrated Stress Response. 17. pp. 1374–1395.
- Pardue, M.L., Gall, J.G., 1970. Chromosomal localization of mouse satellite DNA. *Science* 168, 1356–1358.
- Paulson, H., 2018a. Repeat expansion diseases. *Handb. Clin. Neurol.* 147, 105–123.
- Paulson, Henry, 2018b. Chapter 9 - repeat expansion diseases. In: Geschwind, Daniel H., Paulson, Henry L., Klein, Christine (Eds.), *Handb Clin Neurol* Elsevier.
- Paulson, H.L., Shakkottai, V.G., Clark, H.B., Orr, H.T., 2017. Polyglutamine spinocerebellar ataxias - from genes to potential treatments. *Nat. Rev. Neurosci.* 18, 613–626.
- Pelletier, J., Sonenberg, N., 1988. Internal initiation of translation of eukaryotic mRNA directed by a sequence derived from poliovirus RNA. *Nature* 334, 320–325.
- Pestova, T.V., Hellen, C.U., Shatsky, I.N., 1996a. Canonical eukaryotic initiation factors determine initiation of translation by internal ribosomal entry. *Mol. Cell. Biol.* 16, 6859–6869.
- Pestova, T.V., Shatsky, I.N., Hellen, C.U., 1996b. Functional dissection of eukaryotic initiation factor 4F: the 4A subunit and the central domain of the 4G subunit are sufficient to mediate internal entry of 43S preinitiation complexes. *Mol. Cell. Biol.* 16, 6870–6878.
- Pettersson, O.J., Aagaard, L., Jensen, T.G., Damgaard, C.K., 2015. Molecular mechanisms in DM1 - a focus on foci. *Nucleic Acids Res.* 43, 2433–2441.
- Protter, D.S.W., Parker, R., 2016. Principles and Properties of Stress Granules. *Trends Cell Biol.* 26, 668–679.
- Pulst, S.M., 1993. Spinocerebellar Ataxia Type 2. In: Adam, M.P., Ardinger, H.H., Pagon, R.A., Wallace, S.E., Bean, L.J.H., Stephens, K., Amemiya, A. (Eds.), *GeneReviews* (R). University of Washington, Seattle.
- Pulst, S.M., Nechiporuk, A., Starkman, S., 1993. Anticipation in spinocerebellar ataxia type 2. *Nat. Genet.* 5, 8–10.
- Pulst, S.M., Nechiporuk, A., Nechiporuk, T., Gispert, S., Chen, X.N., Lopes-Cendes, I., Pearlman, S., Starkman, S., Orozco-Diaz, G., Lunke, A., DeJong, P., Rouleau, G.A., Auburger, G., Korenberg, J.R., Figueroa, C., Sahba, S., 1996. Moderate expansion of a normally biallelic trinucleotide repeat in spinocerebellar ataxia type 2. *Nat. Genet.* 14, 269–276.
- Ray, P.S., Grover, R., Das, S., 2006. Two internal ribosome entry sites mediate the translation of p53 isoforms. *EMBO Rep.* 7, 404–410.
- Reddy, K., Zamiri, B., Stanley, S.Y., Macgregor Jr., R.B., Pearson, C.E., 2013. The disease-associated r(GGGGCC)<sub>n</sub> repeat from the C9orf72 gene forms tract length-dependent uni- and multimolecular RNA G-quadruplex structures. *J. Biol. Chem.* 288, 9860–9866.
- Reichelt, R., Holzenburg, A., Buhle Jr., E.L., Jarnik, M., Engel, A., Aebi, U., 1990. Correlation between structure and mass distribution of the nuclear pore complex and of distinct pore complex components. *J. Cell Biol.* 110, 883–894.
- Renoux, A.J., Todd, P.K., 2012. Neurodegeneration the RNA way. *Prog. Neurobiol.* 97, 173–189.
- Renton, A.E., Majounie, E., Waite, A., Simon-Sanchez, J., Rollinson, S., Gibbs, J.R., Schymick, J.C., Laaksovirta, H., van Swieten, J.C., Myllykangas, L., Kalimo, H., Paetau, A., Abramzon, Y., Remes, A.M., Kaganovich, A., Scholz, S.W., Duckworth, J., Ding, J., Harmer, D.W., Hernandez, D.G., Johnson, J.O., Mok, K., Ryten, M., Trabzuni, D., Guerreiro, R.J., Orrell, R.W., Neal, J., Murray, A., Pearson, J., Jansen, I.E., Sondervan, D., Seelaar, H., Blake, D., Young, K., Halliwell, N., Callister, J.B., Toulson, G., Richardson, A., Gerhard, A., Snowden, J., Mann, D., Neary, D., Nalls, M.A., Peuralinna, T., Jansson, L., Isoviita, V.M., Kaivorinne, A.L., Holtta-Vuori, M., Ikonen, E., Sulkava, R., Benatar, M., Wu, J., Chio, A., Restagno, G., Borghero, G., Sabatelli, M., Heckerman, D., Rogaeva, E., Zinman, L., Rothstein, J.D., Sendtner, M., Drepper, C., Eichler, E.E., Alkan, C., Abdullaev, Z., Pack, S.D., Dutra, A., Pak, E., Hardy, J., Singleton, A., Williams, N.M., Heutink, P., Pickering-Brown, S., Morris, H.R., Tienari, P.J., Traynor, B.J., 2011. A hexanucleotide repeat expansion in C9orf72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 72, 257–268.
- Ribbeck, K., Gorlich, D., 2001. Kinetic analysis of translocation through nuclear pore complexes. *EMBO J.* 20, 1320–1330.
- Richard, Guy-Franck, Kerrest, Alix, Dujon, Bernard, 2008. Comparative genomics and molecular dynamics of DNA repeats in eukaryotes. *Microbiol. Mol. Biol. Rev.* 72, 686–727.
- Rossi, S., Serrano, A., Gerbino, V., Giorgi, A., Di Francesco, L., Nencini, M., Bozzo, F., Schinina, M.E., Bagni, C., Cestra, G., Carri, M.T., Achsel, T., Cozzolino, M., 2015. Nuclear accumulation of mRNAs underlies G4C2-repeat-induced translational repression in a cellular model of C9orf72 ALS. *J. Cell Sci.* 128, 1787–1799.
- Saffert, P., Adamla, F., Schieweck, R., Atkins, J.F., Ignatova, Z., 2016. An expanded CAG repeat in huntingtin causes +1 frameshifting. *J. Biol. Chem.* 291, 18505–18513.
- Sanpei, K., Takano, H., Igarashi, S., Sato, T., Oyake, M., Sasaki, H., Wakisaka, A., Tashiro, K., Ishida, Y., Ikeuchi, T., Koide, R., Saito, M., Sato, A., Tanaka, T., Hanyu, S., Takiyama, Y., Nishizawa, M., Shimizu, N., Nomura, Y., Segawa, M., Iwabuchi, K., Eguchi, I., Tanaka, H., Takahashi, H., Tsuji, S., 1996. Identification of the spinocerebellar ataxia type 2 gene using a direct identification of repeat expansion and cloning technique, DIRECT. *Nat. Genet.* 14, 277–284.
- Sareen, D., O'Rourke, J.G., Meera, P., Muhammad, A.K., Grant, S., Simpkinson, M., Bell, S., Carmona, S., Ornelas, L., Sahabian, A., Gendron, T., Petrucelli, L., Baughn, M., Ravits, J., Harms, M.B., Rigo, F., Bennett, C.F., Otis, T.S., Svendsen, C.N., Baloh, R.H., 2013. Targeting RNA foci in iPSC-derived motor neurons from ALS patients with a C9orf72 repeat expansion. *Sci. Transl. Med.* 5 (208ra149).
- Sawaya, S., Bagshaw, A., Buschiazzo, E., Kumar, P., Chowdhury, S., Black, M.A., Gemmell, N., 2013. Microsatellite tandem repeats are abundant in human promoters and are associated with regulatory elements. *PLoS ONE* 8, e54710.
- Schmidt, H.B., Gorlich, D., 2015. Nup98 FG domains from diverse species spontaneously phase separate into particles with nuclear pore-like permselectivity. *Elife* 4.
- Schmidt, H.B., Gorlich, D., 2016. Transport selectivity of nuclear pores, phase separation, and membraneless organelles. *Trends Biochem. Sci.* 41, 46–61.
- Scoles, D.R., Ho, M.H., Dansithong, W., Pflieger, L.T., Petersen, L.W., Thai, K.K., Pulst, S.M., 2015. Repeat associated non-AUG translation (RAN translation) dependent on sequence downstream of the ATXN2 CAG repeat. *PLoS ONE* 10, e0128769.
- Scoles, D.R., Meera, P., Schneider, M.D., Paul, S., Dansithong, W., Figueroa, K.P., Hung, G., Rigo, F., Bennett, C.F., Otis, T.S., Pulst, S.M., 2017. Antisense oligonucleotide therapy for spinocerebellar ataxia type 2. *Nature* 544, 362–366.
- Sellier, C., Buijsen, R.A., He, F., Natla, S., Jung, L., Tropel, P., Gaucherot, A., Jacobs, H., Meziane, H., Vincent, A., Champy, M.F., Sorg, T., Pavlovic, G., Wattenhofer-Donze, M., Birling, M.C., Oulad-Abdelghani, M., Eberling, P., Ruffenach, F., Joint, M., Anheim, M., Martinez-Cerdeno, V., Tassone, F., Willemsen, R., Hukema, R.K., Viville, S., Martinat, C., Todd, P.K., Charlet-Berguerand, N., 2017. Translation of expanded CGG repeats into FMRpolyG is pathogenic and may contribute to fragile X tremor ataxia syndrome. *Neuron* 93, 331–347.
- Shakkottai, V.G., Fogel, B.L., 2013. 'Clinical neurogenetics: autosomal dominant spinocerebellar ataxia. *Neurol. Clin.* 31, 987–1007.
- Shi, K.Y., Mori, E., Nizami, Z.F., Lin, Y., Kato, M., Xiang, S., Wu, L.C., Ding, M., Yu, Y., Gall, J.G., McKnight, S.L., 2017. Toxic Prn poly-dipeptides encoded by the C9orf72 repeat expansion block nuclear import and export. *Proc. Natl. Acad. Sci. U. S. A.* 114, E1111–E1117.
- Siomi, H., Siomi, M.C., Nussbaum, R.L., Dreyfuss, G., 1993. The protein product of the fragile X gene, FMR1, has characteristics of an RNA-binding protein. *Cell* 74, 291–298.
- Siomi, H., Choi, M., Siomi, M.C., Nussbaum, R.L., Dreyfuss, G., 1994. Essential role for KH domains in RNA binding: impaired RNA binding by a mutation in the KH domain of FMR1 that causes fragile X syndrome. *Cell* 77, 33–39.
- Smit, A.F., 1999. Interspersed repeats and other mementos of transposable elements in mammalian genomes. *Curr. Opin. Genet. Dev.* 9, 657–663.
- Sone, Jun, Mitsushashi, Satomi, Fujita, Atsushi, Mizuguchi, Takeshi, Mori, Keiko, Koike, Haruki, Hashiguchi, Akihiro, Takashima, Hiroshi, Sugiyama, Hiroshi, Kohno, Yutaka, Takiyama, Yoshihisa, Maeda, Kengo, Doi, Hiroshi, Koyano, Shigeru, Takeuchi, Hideyuki, Kawamoto, Michi, Kohara, Nobuo, Ando, Tetsuo, Ieda, Toshiaki, Kita, Yasushi, Kokubun, Norito, Tsuboi, Yoshio, Katsuno, Masahisa, Iwasaki, Yasushi, Yoshida, Mari, Tanaka, Fumiaki, Suzuki, Ikko K., Frith, Martin C., Matsumoto, Naomichi, Sobue, Gen, 2019. Long-read sequencing identifies GGC repeat expansion in human-specific NOTCH2NLC associated with neuronal intranuclear inclusion disease. *BioRxiv* 515635.
- Sonobe, Y., Ghadge, G., Masaki, K., Sendoo, A., Fuchs, E., Roos, R.P., 2018. Translation of dipeptide repeat proteins from the C9orf72 expanded repeat is associated with cellular stress. *Neurobiol. Dis.* 116, 155–165.
- Soong, B.W., Paulson, H.L., 2007. Spinocerebellar ataxias: an update. *Curr. Opin. Neurol.* 20, 438–446.
- Soragni, E., Petrosyan, L., Rinkoski, T.A., Wieben, E.D., Baratz, K.H., Fautsch, M.P., Gottesfeld, J.M., 2018. Repeat-Associated Non-ATG (RAN) Translation in Fuchs' Endothelial Corneal Dystrophy. *Invest. Ophthalmol. Vis. Sci.* 59, 1888–1896.
- Spector, D.L., 2006. Snapshot: Cellular bodies. *Cell* 127, 1071.
- Starck, S.R., Jiang, V., Pavon-Eternod, M., Prasad, S., McCarthy, B., Pan, T., Shastri, N., 2012. Leucine-tRNA initiates at CUG start codons for protein synthesis and presentation by MHC class I. *Science* 336, 1719–1723.
- Stochmanki, S.J., Therrien, M., Laganier, J., Rochefort, D., Laurent, S., Karamera, L., Gaudet, R., Vyboh, K., Van Meyel, D.J., Di Cristo, G., Dion, P.A., Gaspar, C., Rouleau, G.A., 2012. Expanded ATXN3 frameshifting events are toxic in Drosophila and mammalian neuron models. *Hum. Mol. Genet.* 21, 2211–2218.
- Suzuki, H., Shibagaki, Y., Hattori, S., Matsuoka, M., 2018. The proline-arginine repeat protein linked to C9-ALS/FTD causes neuronal toxicity by inhibiting the DEAD-box RNA helicase-mediated ribosome biogenesis. *Cell Death Dis.* 9, 975.
- Swisher, K.D., Parker, R., 2010. Localization to, and effects of Pbp1, Pbp4, Lsm12, Dhh1, and Pab1 on stress granules in *Saccharomyces cerevisiae*. *PLoS ONE* 5, e10006.
- Tabet, R., Schaeffer, L., Freyermuth, F., Jambau, M., Workman, M., Lee, C.Z., Lin, C.C., Jiang, J., Jansen-West, K., Abou-Hamdan, H., Desaubry, L., Gendron, T., Petrucelli, L., Martin, F., Lagier-Tourenne, C., 2018. CUG initiation and frameshifting enable production of dipeptide repeat proteins from ALS/FTD C9orf72 transcripts. *Nat. Commun.* 9, 152.
- Tao, Z., Wang, H., Xia, Q., Li, K., Li, K., Jiang, X., Xu, G., Wang, G., Ying, Z., 2015. Nucleolar stress and impaired stress granule formation contribute to C9orf72 RAN translation-induced cytotoxicity. *Hum. Mol. Genet.* 24, 2426–2441.
- Tassone, F., Hagerman, R.J., Loesch, D.Z., Lachiewicz, A., Taylor, A.K., Hagerman, P.J., 2000a. Fragile X males with unmethylated, full mutation trinucleotide repeat expansions have elevated levels of FMR1 messenger RNA. *Am. J. Med. Genet.* 94, 232–236.
- Tassone, F., Hagerman, R.J., Taylor, A.K., Gane, L.W., Godfrey, T.E., Hagerman, P.J., 2000b. Elevated levels of FMR1 mRNA in carrier males: a new mechanism of involvement in the fragile-X syndrome. *Am. J. Hum. Genet.* 66, 6–15.
- Teixeira, D., Sheth, U., Valencia-Sanchez, M.A., Brenques, M., Parker, R., 2005. Processing bodies require RNA for assembly and contain nontranslating mRNAs. *Rna* 11, 371–382.
- Thomson, S.R., Seo, S.S., Barnes, S.A., Louros, S.R., Muscas, M., Dando, O., Kirby, C., Wyllie, D.J.A., Hardingham, G.E., Kind, P.C., Osterweil, E.K., 2017. Cell-type-specific translation profiling reveals a novel strategy for treating fragile X syndrome. *Neuron* 95, 550–563 (e5).
- Todd, P.K., Paulson, H.L., 2010. RNA-mediated neurodegeneration in repeat expansion disorders. *Ann. Neurol.* 67, 291–300.



- Todd, P.K., Mack, K.J., Malter, J.S., 2003. The fragile X mental retardation protein is required for type-I metabotropic glutamate receptor-dependent translation of PSD-95. *Proc. Natl. Acad. Sci. U. S. A.* 100, 14374–14378.
- Todd, P.K., Oh, S.Y., Krans, A., He, F., Sellier, C., Frazer, M., Renoux, A.J., Chen, K.C., Scaglione, K.M., Basur, V., Elenitoba-Johnson, K., Vonsattel, J.P., Louis, E.D., Sutton, M.A., Taylor, J.P., Mills, R.E., Charlet-Berguerand, N., Paulson, H.L., 2013. CGG repeat-associated translation mediates neurodegeneration in fragile X tremor ataxia syndrome. *Neuron* 78, 440–455.
- Toulouse, A., Au-Yeung, F., Gaspar, C., Roussel, J., Dion, P., Rouleau, G.A., 2005. Ribosomal frameshifting on MJD-1 transcripts with long CAG tracts. *Hum. Mol. Genet.* 14, 2649–2660.
- Treangen, T.J., Salzberg, S.L., 2011. Repetitive DNA and next-generation sequencing: computational challenges and solutions. *Nat. Rev. Genet.* 13, 36–46.
- Trottier, Y., Devys, D., Imbert, G., Saudou, F., An, I., Lutz, Y., Weber, C., Agid, Y., Hirsch, E.C., Mandel, J.L., 1995a. Cellular localization of the Huntington's disease protein and discrimination of the normal and mutated form. *Nat. Genet.* 10, 104–110.
- Trottier, Y., Lutz, Y., Stevanin, G., Imbert, G., Devys, D., Cancel, G., Saudou, F., Weber, C., David, G., Tora, L., et al., 1995b. Polyglutamine expansion as a pathological epitope in Huntington's disease and four dominant cerebellar ataxias. *Nature* 378, 403–406.
- Tsilifidis, C., MacKenzie, A.E., Mettler, G., Barcelo, J., Korneluk, R.G., 1992. Correlation between CTG trinucleotide repeat length and frequency of severe congenital myotonic dystrophy. *Nat. Genet.* 1, 192–195.
- Udagawa, T., Farny, N.G., Jakovcevski, M., Kaphzan, H., Alarcon, J.M., Anilkumar, S., Ivshina, M., Hurt, J.A., Nagaoka, K., Nalavadi, V.C., Lorenz, L.J., Bassell, G.J., Akbarian, S., Chattarji, S., Klann, E., Richter, J.D., 2013. Genetic and acute CPEB1 depletion ameliorate fragile X pathophysiology. *Nat. Med.* 19, 1473–1477.
- Usdin, K., 2008. The biological effects of simple tandem repeats: lessons from the repeat expansion diseases. *Genome Res.* 18, 1011–1019.
- Uversky, V.N., Dunker, A.K., 2010. Understanding protein non-folding. *Biochim. Biophys. Acta* 1804, 1231–1264.
- Vagner, S., Gensac, M.C., Maret, A., Bayard, F., Amalric, F., Prats, H., Prats, A.C., 1995. Alternative translation of human fibroblast growth factor 2 mRNA occurs by internal entry of ribosomes. *Mol. Cell. Biol.* 15, 35–44.
- Van Treeck, B., Protter, D.S.W., Matheny, T., Khong, A., Link, C.D., Parker, R., 2018. RNA self-assembly contributes to stress granule formation and defining the stress granule transcriptome. *Proc. Natl. Acad. Sci. U. S. A.* 115, 2734–2739.
- Vattem, K.M., Wek, R.C., 2004. Reinitiation involving upstream ORFs regulates ATF4 mRNA translation in mammalian cells. *Proc. Natl. Acad. Sci. U. S. A.* 101, 11269–11274.
- Verstrepen, K.J., Jansen, A., Lewitter, F., Fink, G.R., 2005. Intragenic tandem repeats generate functional variability. *Nat. Genet.* 37, 986–990.
- Vinces, M.D., Legendre, M., Caldara, M., Hagihara, M., Verstrepen, K.J., 2009. Unstable tandem repeats in promoters confer transcriptional evolvability. *Science* 324, 1213–1216.
- Viphakone, N., Hautbergue, G.M., Walsh, M., Chang, C.T., Holland, A., Folco, E.G., Reed, R., Wilson, S.A., 2012. TREX exposes the RNA-binding domain of Nxf1 to enable mRNA export. *Nat. Commun.* 3, 1006.
- Vitalis, A., Pappu, R.V., 2011. Assessing the contribution of heterogeneous distributions of oligomers to aggregation mechanisms of polyglutamine peptides. *Biophys. Chem.* 159, 14–23.
- Vitalis, A., Wang, X., Pappu, R.V., 2007. Quantitative characterization of intrinsic disorder in polyglutamine: insights from analysis based on polymer theories. *Biophys. J.* 93, 1923–1937.
- Walters, R.H., Murphy, R.M., 2009. Examining polyglutamine peptide length: a connection between collapsed conformations and increased aggregation. *J. Mol. Biol.* 393, 978–992.
- Wang, L.W., Berry-Kravis, E., Hagerman, R.J., 2010. Fragile X: leading the way for targeted treatments in autism. *Neurotherapeutics* 7, 264–274.
- Waring, M., Britten, R.J., 1966. Nucleotide sequence repetition: a rapidly reassociating fraction of mouse DNA. *Science* 154, 791–794.
- Warrick, J.M., Paulson, H.L., Gray-Board, G.L., Bui, Q.T., Fischbeck, K.H., Pittman, R.N., Bonini, N.M., 1998. Expanded polyglutamine protein forms nuclear inclusions and causes neural degeneration in *Drosophila*. *Cell* 93, 939–949.
- Wen, X., Tan, W., Westergard, T., Krishnamurthy, K., Markandaiah, S.S., Shi, Y., Lin, S., Schneider, N.A., Monaghan, J., Pandey, U.B., Pasinelli, P., Ichida, J.K., Trotti, D., 2014. Antisense proline-arginine RAN dipeptides linked to C9ORF72-ALS/FTD form toxic nuclear aggregates that initiate in vitro and in vivo neuronal death. *Neuron* 84, 1213–1225.
- Westergard, Thomas, McAvoy, Kevin, Russell, Katelyn, Wen, Xinmei, Pang, Yu, Morris, Brandie, Pasinelli, Piera, Trotti, Davide, Haeusler, Aaron, 2019. Repeat-associated non-AUG translation in C9orf72-ALS/FTD is driven by neuronal excitation and stress. *EMBO Mol. Med.* 0, e9423.
- Williams, A.J., Paulson, H.L., 2008. Polyglutamine neurodegeneration: protein misfolding revisited. *Trends Neurosci.* 31, 521–528.
- Wills, N.M., Atkins, J.F., 2006. The potential role of ribosomal frameshifting in generating aberrant proteins implicated in neurodegenerative diseases. *RNA* 12, 1149–1153.
- Wojciechowska, M., Olejniczak, M., Galka-Marciniak, P., Jazurek, M., Krzyzosiak, W.J., 2014. RAN translation and frameshifting as translational challenges at simple repeats of human neurodegenerative disorders. *Nucleic Acids Res.* 42, 11849–11864.
- Wolozin, B., 2012. Regulated protein aggregation: stress granules and neurodegeneration. *Mol. Neurodegener.* 7, 56.
- Xu, Z., Poidevin, M., Li, X., Li, Y., Shu, L., Nelson, D.L., Li, H., Hales, C.M., Gearing, M., Wingo, T.S., Jin, P., 2013. Expanded GGGGCC repeat RNA associated with amyotrophic lateral sclerosis and frontotemporal dementia causes neurodegeneration. *Proc. Natl. Acad. Sci. U. S. A.* 110, 7778–7783.
- Yamakawa, M., Ito, D., Honda, T., Kubo, K., Noda, M., Nakajima, K., Suzuki, N., 2015. Characterization of the dipeptide repeat protein in the molecular pathogenesis of c9FTD/ALS. *Hum. Mol. Genet.* 24, 1630–1645.
- Yang, D., Abdallah, A., Li, Z., Lu, Y., Almeida, S., Gao, F.B., 2015. FTD/ALS-associated poly(GR) protein impairs the Notch pathway and is recruited by poly(GA) into cytoplasmic inclusions. *Acta Neuropathol.* 130, 525–535.
- Young, S.K., Wek, R.C., 2016. Upstream open reading frames differentially regulate gene-specific translation in the integrated stress response. *J. Biol. Chem.* 291, 16927–16935.
- Yunis, J.J., Yasmin, W.G., 1971. Heterochromatin, satellite DNA, and cell function. Structural DNA of eucaryotes may support and protect genes and aid in speciation. *Science* 174, 1200–1209.
- Zalfa, F., Giorgi, M., Primerano, B., Moro, A., Di Penta, A., Reis, S., Oostra, B., Bagni, C., 2003. The fragile X syndrome protein FMRP associates with BC1 RNA and regulates the translation of specific mRNAs at synapses. *Cell* 112, 317–327.
- Zhang, Y.J., Jansen-West, K., Xu, Y.F., Gendron, T.F., Bieniek, K.F., Lin, W.L., Sasaguri, H., Caulfield, T., Hubbard, J., Daugherty, L., Chew, J., Belzil, V.V., Prudencio, M., Stankowski, J.N., Castaneda-Casey, M., Whitelaw, E., Ash, P.E., DeTure, M., Rademakers, R., Boylan, K.B., Dickson, D.W., Petrucelli, L., 2014. Aggregation-prone c9FTD/ALS poly(GA) RAN-translated proteins cause neurotoxicity by inducing ER stress. *Acta Neuropathol.* 128, 505–524.
- Zhang, K., Donnelly, C.J., Haeusler, A.R., Grima, J.C., Machamer, J.B., Steinwald, P., Daley, E.L., Miller, S.J., Cunningham, K.M., Vidensky, S., Gupta, S., Thomas, M.A., Hong, I., Chiu, S.L., Haganir, R.L., Ostrow, L.W., Matunis, M.J., Wang, J., Sattler, R., Lloyd, T.E., Rothstein, J.D., 2015. The C9orf72 repeat expansion disrupts nucleocytoplasmic transport. *Nature* 525, 56–61.
- Zhang, Y.J., Gendron, T.F., Grima, J.C., Sasaguri, H., Jansen-West, K., Xu, Y.F., Katzman, R.B., Gass, J., Murray, M.E., Shinohara, M., Lin, W.L., Garrett, A., Stankowski, J.N., Daugherty, L., Tong, J., Perkerson, E.A., Yue, M., Chew, J., Castaneda-Casey, M., Kurti, A., Wang, Z.S., Liesinger, A.M., Baker, J.D., Jiang, J., Lagier-Tourenne, C., Edbauer, D., Cleveland, D.W., Rademakers, R., Boylan, K.B., Bu, G., Link, C.D., Dickey, C.A., Rothstein, J.D., Dickson, D.W., Fryer, J.D., Petrucelli, L., 2016. C9ORF72 poly(GA) aggregates sequester and impair HR23 and nucleocytoplasmic transport proteins. *Nat. Neurosci.* 19, 668–677.
- Zhang, K., Daigle, J.G., Cunningham, K.M., Coyne, A.N., Ruan, K., Grima, J.C., Bowen, K.E., Wadhwa, H., Yang, P., Rigo, F., Taylor, J.P., Gitler, A.D., Rothstein, J.D., Lloyd, T.E., 2018a. Stress granule assembly disrupts nucleocytoplasmic transport. *Cell* 173, 958–971 (e17).
- Zhang, Y.J., Gendron, T.F., Ebbert, M.T.W., O'Raw, A.D., Yue, M., Jansen-West, K., Zhang, X., Prudencio, M., Chew, J., Cook, C.N., Daugherty, L.M., Tong, J., Song, Y., Pickles, S.R., Castaneda-Casey, M., Kurti, A., Rademakers, R., Oskarsson, B., Dickson, D.W., Hu, W., Gitler, A.D., Fryer, J.D., Petrucelli, L., 2018b. Poly(GR) impairs protein translation and stress granule dynamics in C9orf72-associated frontotemporal dementia and amyotrophic lateral sclerosis. *Nat. Med.* 24, 1136–1142.
- Zu, T., Gibbins, B., Doty, N.S., Gomes-Pereira, M., Huguette, A., Stone, M.D., Margolis, J., Peterson, M., Markowski, T.W., Ingram, M.A., Nan, Z., Forster, C., Low, W.C., Schoser, B., Somia, N.V., Clark, H.B., Schmechel, S., Bitterman, P.B., Gourdon, G., Swanson, M.S., Moseley, M., Ranum, L.P., 2011. Non-ATG-initiated translation directed by microsatellite expansions. *Proc. Natl. Acad. Sci. U. S. A.* 108, 260–265.
- Zu, T., Liu, Y., Bañez-Coronel, M., Reid, T., Pletnikova, O., Lewis, J., Miller, T.M., Harms, M.B., Falchuk, A.E., Subramony, S.H., Ostrow, L.W., Rothstein, J.D., Troncoso, J.C., Ranum, L.P., 2013. RAN proteins and RNA foci from antisense transcripts in C9ORF72 ALS and frontotemporal dementia. *Proc. Natl. Acad. Sci. U.S.A.* 110, E4968–E4977. <https://doi.org/10.1073/pnas.1315438110>.