

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

NeuroImage

journal homepage: www.elsevier.com/locate/neuroimage

The latency of a visual evoked potential tracks the onset of decision making

Michael D. Nunez^{a,b,*}, Aishwarya Gosai^{a,c}, Joachim Vandekerckhove^{a,d,e},
Ramesh Srinivasan^{a,b,e}

^a Department of Cognitive Sciences, University of California, Irvine, CA, USA

^b Department of Biomedical Engineering, University of California, Irvine, CA, USA

^c Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA

^d Department of Statistics, University of California, Irvine, CA, USA

^e Institute of Mathematical Behavioral Sciences, University of California, Irvine, CA, USA

ARTICLE INFO

Keywords:

Decision making
Figure-ground segregation
EEG
Visual evoked potentials (VEP)
Diffusion models
Hierarchical Bayesian methods

ABSTRACT

Encoding of a sensory stimulus is believed to be the first step in perceptual decision making. Previous research has shown that electrical signals recorded from the human brain track evidence accumulation during perceptual decision making (Gold and Shadlen, 2007; O'Connell et al., 2012; Philiastides et al., 2014). In this study we directly tested the hypothesis that the latency of the N200 recorded by EEG (a negative peak occurring between 150 and 275 ms after stimulus presentation in human participants) reflects the visual encoding time (VET) required for completion of figure-ground segregation before evidence accumulation. We show that N200 latencies vary across individuals, are modulated by external visual noise, and increase response time by x milliseconds when they increase by x milliseconds, reflecting a linear regression slope of 1. Simulations of cognitive decision-making theory show that variation in human response times not related to evidence accumulation (non-decision time; NDT), including VET, are tracked by the fastest response times. Evidence that VET is tracked by N200 latencies was found by fitting a linear model between trial-averaged N200 latencies and the 10th percentiles of response times, a model-independent estimate of NDT. Fitting a novel neuro-cognitive model of decision making also yielded a slope of 1 between N200 latency and model-estimated NDT in multiple visual noise conditions, indicating that N200 latencies track the completion of visual encoding and the onset of evidence accumulation. The N200 waveforms were localized to the cortical surface at distributed temporal and extrastriate locations, consistent with a distributed network engaged in figure-ground segregation of the target stimulus.

1. Introduction

When making a decision to stop or accelerate based on a traffic light, the driver must first recognize the presence of the traffic light. Decisions are often thought to be made based on a sequential sampling of evidence (Ratcliff and McKoon, 2008), in this scenario the driver samples visual information about the color, shape, and position of the traffic light while preparing the foot to depress one of two pedals. However the time between when the light of the traffic light first reaches the eye of the driver to the onset of decision making, i.e. the beginning of this sampling of visual information, is largely unknown. We define *visual encoding time* (VET) as the amount of time before these decision-related processes begin. Our goals in the following study were to 1) clarify estimation of VET in human participants using scalp-recorded electroencephalography (EEG) and 2) verify these time estimates using both non-parametric (i.e.

results depending upon a limited set of assumptions) and parametric approaches (i.e. results within the framework of a cognitive theory of quick human decision making).

While there has been abundant evidence of visual information appearing in the primary visual cortex approximately 60 milliseconds (ms) after stimulus onset (Schmolesky et al., 1998; Luck, 2012), further processing takes place within a network of brain areas comprising the visual system. The key objective of this further processing is the grouping together of elements of the scene coded in separate neurons into distinct objects, a process known as figure-ground segregation (Lamme, 1995). Masking, by presenting a distractor stimulus soon after the target stimulus onset, can interrupt figure-ground segregation up to 100 ms by degrading the encoding of the visual stimulus in cortical area V1 (Lamme et al., 2002). This work suggests that VET of the target stimulus is approximately 160 ms (100 ms masking time plus 60 ms for the time of

* Corresponding author. Department of Cognitive Sciences, University of California, Irvine, Irvine, CA, 92697, USA.
E-mail address: mdnunez1@uci.edu (M.D. Nunez).

<https://doi.org/10.1016/j.neuroimage.2019.04.052>

Received 5 September 2018; Received in revised form 23 January 2019; Accepted 18 April 2019

Available online 24 April 2019

1053-8119/Published by Elsevier Inc.

the distractor stimulus to reach primary visual cortex). Furthermore, on the basis of ERP studies in humans and physiological studies in animal models (Bach and Meigen, 1992; Roitman and Shadlen, 2002; Straube et al., 2010), VET is expected to be 150–200 ms in duration such that the estimated time depends upon the visual noise condition, target type, participant's attentional state, etc.

Human response times (time between stimulus onset and motor action execution) during decision making are theorized to be composed of VET, evidence accumulation time, response preparation time, and motor execution time. VET is separate from the decision-related processing, which has been experimentally found to reflect a sequential accumulation of evidence on level of the neuron (Gold and Shadlen, 2007; Shadlen and Kiani, 2013), the level of the scalp-recorded electroencephalograph (EEG; O'Connell et al., 2012; Philiastides et al., 2014; Kelly and O'Connell, 2015), and on the cognitive level as evidenced by cognitive models of human behavior (Voss et al., 2004; Ratcliff and McKoon, 2008). While evidence accumulation and motor planning may operate in parallel (Dmochowski and Norcia, 2015; Servant et al., 2015, 2016), it can be reasonably expected that figure-ground segregation is a sequential processing step that occurs before any evidence accumulation or motor planning. That is, the grouping of neural activity elicited by the stimulus across the visual system into a distinct “object” through figure-ground segregation is necessary before evidence accumulation and motor planning.

The figure-ground segregation speed of the human visual system has previously been estimated between 150 ms and 225 ms in humans with time-locked event related potentials (ERPs; Bach and Meigen, 1992; Thorpe et al., 1996; VanRullen and Thorpe, 2001) recorded by EEG using clever differences between experimental conditions. Bach and Meigen (1992) showed that texture integration of checkerboard targets formed from fields of oriented bars occurs between 161 and 225 ms. Thorpe et al. (1996) showed that decision making related to a go/no-go task based on similar visual scenes (containing animals or no animals) started about 150 ms after stimulus onset, and then VanRullen and Thorpe (2001) replicated their results using the same stimuli alternating as targets and non-targets. Loughnane et al. (2016) found initial evidence by looking at condition differences to suggest that stimulus-locked negative peaks in ERPs around 200 ms (N200s) influence the onset of a neural correlate of evidence accumulation in a random-dot motion task. In other studies, single-trial EEG potentials thought to be generative of N200 peaks were suspected to be related to a “pre-attentive” phase before evidence accumulation (Zhang et al., 2017, 2018). In random-dot motion experiments where participants (both humans and monkeys) are tasked to choose the direction of coherent motion of previously randomly moving dots, the feature-conjunction of multiple motion-selective neurons is realized by figure-ground segregation (i.e. the grouping together of elements of the scene coded in separate neurons into distinct objects). Single-neuron activity reflecting evidence accumulation in lateral intraparietal areas (LIP) typically begins at similar time periods (\approx 200 ms) after the experimental stimulus onsets in a random-dot motion paradigms (Roitman and Shadlen, 2002; Shadlen and Kiani, 2013).

In this study, our hypothesis was that N200 peak-latencies of ERPs would track VET no matter the external visual noise condition, participant, or participant's internal state that changed across EEG sessions. The experimental manipulation was important because we hypothesized that both the N200 peak-latency and VET (and thus NDT) would increase in conditions with more visual noise, but that the *relationship* between the two should not change in conditions with more visual noise. We used a visual target that was not used by the aforementioned studies (e.g. a visual Gabor stimulus) to add to growing evidence that VET takes about 200 ms across multiple visual target types. Using both direct linear regressions between EEG latency and fast response times and novel neuro-cognitive models of decision making, we verified that N200 peak-latencies tracked non-decision time (NDT), of which visual encoding time (VET) is an additive component, with regression slopes that imply an x millisecond increase in N200 peak-latency leads to an x millisecond

increase in NDT (i.e. a slope parameter of 1). These results helped us extend the theory of human perceptual decision making by incorporating the EEG-derived latency estimate of VET and evidence accumulation onset into the broader theory.

2. Methods

2.1. Experiments and participants

In order to evaluate the hypothesis that N200 latencies track VET, data was analyzed from two similar experiments. Data from Experiment 1 consisted of EEG recordings and behavioral observations from 12 unique participants (3 female, 9 male) with 2 sessions each. Data from Experiment 2 consisted of EEG recordings and behavioral observations from 4 unique participants (2 female, 2 male) with 7 sessions each. Two male participants participated in both experiments resulting in 14 unique participants across both experiments (5 female, 9 male) with 52 unique EEG sessions. Sessions of EEG collection and task performance for each participant in Experiment 1 were separated by at least 24 h. Sessions of EEG collection and task performance for each participant in Experiment 2 were separated by 1 week. All participants had normal or corrected-to-normal vision and had no history or family history of epilepsy. All participants gave written informed consent, and all data was collected at the University of California, Irvine with approval by the Institutional Review Board.

2.2. Experimental tasks

Data was included from two different experiments in this study in order to verify that any evidence for our hypothesis that we found (i.e. that N200 peak-latencies track VET) did not depend on the type of distracting visual stimuli. While we expect the distracting visual noise to yield different N200 peak-latencies and VETs across both experiments, we expect to find the same tracking of VET by the N200 peak-latency in both experiments across multiple visual contrast conditions. In both experiments, participants performed two-alternative forced choice tasks, specifically to determine whether a Gabor stimulus contained high or low spatial frequency content. Stimuli for Experiments 1 and 2 are built and displayed using the MATLAB Psychophysics toolbox (Psychtoolbox-3; www.psychtoolbox.org). Example stimuli are given in Fig. 1.

Participants performed perceptual decision making tasks in a dark room on a 61 cm LED monitor maintaining a distance of approximately 57 cm from retina to display. The monitor resolution was set as 1920*1280 pixels with a 120 Hz refresh rate. The participants were told to remain fixated on a small fixation spot while performing the task.

Gabors are sinusoidal grating patterns with a Gaussian falloff of contrast that produces maximal firing in certain neurons within the primary visual cortex (Webster et al., 1985). Participants in both experiments were tasked with identifying the spatial frequency (either “high” or “low” spatial frequency) of large Gabors (approximately 10 cm and 10° visual angle in diameter) that were randomly rotated and had a random phase on each trial (uniformly distributed draws across both rotation angle and phase). In Experiment 2, the high and low spatial frequencies of the target Gabors were 2.4 and 2.6 cycles per degree visual angle (cpd) respectively, while in Experiment 1, the targets were randomly drawn either from one of two shifted beta distributions, $Beta(13.5, 1.5)$ distribution plus 1.5 cpd or a $Beta(1.5, 13.5)$ distribution plus 2.5 cpd. These two target shifted-beta distributions had means 2.4 and 2.6 cpd respectively and were chosen so that there would exist small variability in the target difficulty (i.e. distance from 2.5 cpd) with two completely separable distributions. Both experiments contained three conditions of high, medium, and low visual noise contrast. Throughout each trial, samples of visual noise were displayed both before, and concurrent with the targets. A new sample was presented at regular intervals (25 ms, every 3rd video frame). In Experiment 1, a checkerboard noise pattern was displayed, while in Experiment 2, the




Fig. 1. Example stimuli of the *cue* and *response* intervals of medium noise conditions from Experiments 1 (top) and 2 (bottom). During the response interval, participants decided which spatial-frequency-target each Gabor represented, pressing a button with their left hand for a low spatial frequency target (2.4 cycles per degree visual angle, cpd) and pressing a button with their right hand for a high spatial frequency target (2.6 cpd). N200 waveforms were calculated time-locked to the onset of the Gabor stimulus during the response intervals. In both experiments a paradigm was used in which the visual noise changed at 40 Hz and the Gabor signal flickered at 30 Hz to evoke 40 Hz and 30 Hz responses in electrocortical activity that track attention to the noise and signal stimuli. While the attention analysis is beyond the scope of this paper, a similar analysis is presented by Nunez et al. (2015).

noise was spatially bandpass filtered to include only masking noise centered around 2 and 3 cpd (i.e. the noise was a mixture of two normal distributions centered at 2 and 3 cpd with .1 cpd standard deviations), equally masking both the high and low spatial frequency targets.

The time course of each trial was as follows: participants were asked to fixate on a small fixation spot in the center of the screen throughout the experiment, visual noise changing at 40 Hz was displayed a variable length of time between 500 ms and 1000 ms during the *cue* interval, then the Gabor signal stimulus was flickered at 30 Hz embedded in the noise stimuli for 1200 ms–2000 ms during the *response* interval. During the response interval, participants were asked to respond as accurately as possible in the time allowed with a button box, using their left hand to respond for low spatial frequency targets and their right hand for high spatial frequency targets. Auditory feedback was given after the response interval to indicate trial accuracy; this feedback was included in order to encourage participant vigilance. An entire session of task performance and EEG collection took participants approximately 1 h with breaks. Each session produced 8 blocks of 60 trials each for a total of 480 trials, with trials at the three levels of noise contrast intermixed within each block.

2.3. EEG recording

EEG was collected using Electrical Geodesic, Inc.'s 128 channel Geodesic sensor net and a Net Amps 200 series amplifier. Electrical activity from the scalp was recorded at a sampling rate of 1000 samples per second with a vertex reference and hardware bandpass filtered to either

1–50 Hz (EEG data from Experiment 1) or 1–100 Hz (EEG data from Experiment 2). The hardware band pass filter in Experiment 2 was chosen purposely to maintain high frequencies so that broadband noise was submitted to an Independent Component Analysis (ICA) in order to aid artifact correction. In data from both experiments, visual inspection of the raw time series and resulting ICA components was used to remove obvious artifact due to ocular movements, electrical artifact, electrode movement artifact, and some muscle (EMG) activity (see Makeig et al., 1996; Nunez et al., 2016, for further details), using the MATLAB repository artscreenEEG (Nunez et al., 2017a). Both data sets ultimately yielded artifact-resistant EEG data by first removing a small amount of artifactual time segments and then removing some linear electrode mixtures given by clear artifactual Independent Components. The EEG data was then common-averaged referenced and submitted to further processing as discussed below.

2.4. Estimation of N200 waveforms

N200 waveforms, calculated from the EEG signals following the onset of the Gabor patch during the *response* interval were obtained using five processing steps: 1) software bandpass filtering the EEG data from 1 to 10 Hz (forward-backward Butterworth IIR filter with stopbands at 0.25 and 20 Hz with 1 dB attenuation in the passband and 10 dB attenuation in the stopband) obtaining cleaner estimates of the early evoked potentials—note that slower, later evoked potentials were purposefully attenuated by using a 1 Hz highpass although some variance of the later

evoked potentials (such as the P300) may remain (see Acunzo et al., 2012; Karakaş et al., 2000) 2) subtracting the average baseline potential (i.e. the average amplitude in a 100 ms window before the stimulus), 3) averaging across trials to obtain a traditional ERP estimate at 128 electrodes for each participant (excluding those electrodes marked as artifact), 4) taking a singular value decomposition (SVD) of the ERPs for each experimental condition and EEG session, and then 5) using the first SVD component (the component that explained the most variance) as a channel weighting function in order to obtain a less-noisy estimate of the N200 waveforms for each experimental condition and EEG session as well as to obtain single-trial estimates (discussed below). To test robustness of filtering parameters (see VanRullen, 2011) we tested the effect of a 30 Hz lowpass filter instead of a 10 Hz lowpass filter to calculate the N200 waveforms. This filter change did not affect the conclusions reached in this paper because none of the statistics or parameters of the following analyses involving the N200 waveform changed significantly.

The goal of the singular value decomposition (SVD) was to improve the signal-to-noise ratio of both traditional ERP estimates and single-trial estimates by estimating an optimal spatial filter to detect the N200 (Kayser and Tenke, 2003; Parra et al., 2005). SVD is the algorithm used by most principal component analysis (PCA) algorithms and produces non-stochastic, deterministic results. N200 waveforms were obtained by finding the first principal component of the matrix of ERPs of 1000 ms post-stimulus and 100 ms pre-stimulus (samples T by channel C data, see Methods in Nunez et al. (2017b)). The first principal component consists of both a weight vector that produces a time series of the trial-averaged ERP and an associated vector of channel weights representing the spatial distribution of that component. Traditionally calculated ERPs (Luck, 2012) at specific parietal electrodes were also compared to the SVD estimated ERP and exhibited no difference in latency.

Single-trial estimates of the SVD-estimated ERP are then obtained by applying the weight vector as a spatial filter (i.e. weight summing over the vector of weights as show by Fig. 2). It should be noted that the first component of the SVD method yielded the N200 waveform in these experiments because 1) the neural response to the visual stimulus was large in the average ERP and 2) other waveform components of the ERP (such as the P300 response) were removed using the 1 Hz highpass. Had the experimental conditions and filtering parameters been different, the N200 waveform may have been placed in a SVD component that explained less of the variance.

The window to calculate minima in order to evaluate N200 peak-latency responses was found empirically. Out of the following windows: 101–299 ms post-stimulus, 101–249 ms post-stimulus, 151–224 ms post-stimulus, and 151–274 ms post-stimulus, the 151–274 ms window post-stimulus was found to capture the N200 waveform well with few false detections. Only 2 estimates out of 147

estimates of trial-averaged N200 peak-latency were detected at the 274 ms boundary and none at the 151 ms boundary. Single-trial N200 peak-latencies that were found with a minimum on the boundary were removed from the analysis, resulting in 14.2% of candidate trials removed (a maximum of 25.2% from the Exp 1. high noise condition and a minimum of 9.3% from the Exp 2. medium noise condition). A minimum at either 151 or 274 ms was indicative that single-trial N200 latencies were not well estimated on that trial, as the interval contains a ramp rather than a peak.

Deflection times were also calculated for the trial-averaged N200 waveforms by taking the window of 0–274 ms post-stimulus onset in the response interval and finding where the derivative of the ERP in that window first became negative before the N200 peak-latency. This reflects the point at which the signal begins to decrease before the N200 peak-latency. This particular time point has been explored in a previous study of visual encoding by Martin et al. (2010), albeit calculated using the point at which the signal reached two standard deviations below the baseline. Trial-averaged N200 waveforms and deflection time and peak-latency distributions are shown in Fig. 3. Deflection times are not separated by visual noise conditions because no evidence was found that deflection times changed based on visual noise (see Results).

2.5. Cognitive theory of human decision making

We tested the relationship between N200 peak-latency and VET directly by fitting response time and choice data from the participants to simple drift-diffusion models (DDM). In this theoretical framework, human visual encoding time (VET) is the first cognitive processing time after a visual stimulus is displayed, after which the participant accumulates evidence based on the visual input (see Fig. 4). The non-decision time (NDT) τ parameter is the sum of VET and other time-invariant processes across trials not associated with a sequential accumulation of evidence, such as motor execution time (Ratcliff, 1978; Ratcliff and McKoon, 2008). Fitting reaction time and choice behavior with DDM results only in NDT estimates, not direct estimates of the components of NDT, such as VET. We therefore tested whether the linear relationship between N200 peak-latency and NDT had a slope of 1, reflecting the hypothesis that N200 peak-latency is one additive component of NDT.

We assessed our ability to recover true non-decision time (NDT) estimates from response time (RT) percentiles and hierarchical Bayesian model fitting by simulating response time data. Response time data was simulated from drift-diffusion models with various amounts of trial-to-trial variability in NDT and evidence accumulation rate (i.e. drift rate) suggested by Ratcliff (1978) as well as specific amounts of contaminant processes, generating a random choice and reaction time (discussed below). These response times were simulated using DMAT, MATLAB software for simulating and estimating diffusion model parameters




Fig. 2. A visual representation of the singular value decomposition (SVD) method for finding single-trial estimates of evoked responses in EEG. The EEG presented here is time-locked to the signal onset during the response interval, such that the single-trial N200 waveform encoded the response to the signal onset. A single trial of EEG from one participant (Left) can be thought of as a time by channel ($T \times C$) matrix. SVD weights \mathbf{v} ($C \times 1$) are obtained from the ERP response (i.e. trial-averaged EEG; $T \times C$) and can be plotted on a cartoon representation of the human scalp with intermediate interpolated values (Middle). This specific trial's N200 waveform (Right) was obtained by multiplying the time series data from each channel on this trial by the associated weight in vector \mathbf{v} and then summing across all weighted channels. Weight averaging across channels is a common method to obtain larger signal-to-noise ratios in EEG (e.g. cleaner, more task-relevant neural signals, see Parra et al., 2005).




Fig. 3. All trial-averaged N200 waveforms across EEG sessions and the six visual noise experimental conditions ($N = 147$). Smoothed histograms for the N200 deflection times (top left distribution) as well as the N200 peak-latency times (all other distributions for each noise condition; Experiment 2 visual noise conditions denoted by star-texture) are displayed above the zero amplitude midline. There was a significant but weak correlation found between the N200 deflection times and the N200 peak-latency times ($\rho = .274$). N200 peak-latencies were dependent upon the noise condition of the experiment, while N200 deflection times were not found to vary dependent upon the noise condition.




Fig. 4. A graphical illustration of a Neural Drift Diffusion model in which the visual encoding time τ_e on single-trials describes the latency of the negative peaks of the N200 waveform on 146 experimental trials. Single-trial observations of the N200 peak-latency are found by using the SVD method (described previously) in one participant and experimental condition. Total NDT τ reflects both visual encoding time (VET) τ_e as well as motor execution time (MET) τ_m (e.g. motor response time of the participant after the decision is made) and can be estimated from response time distributions. Decision time is generated via a Brownian motion evidence accumulation process with evidence accumulation rate δ (drift rate), evidence accumulation noise ζ (diffusion coefficient), and evidence threshold (boundary parameter) α (for more information see Ratcliff, 1978; Ratcliff and McKoon, 2008; Nunez et al., 2015, 2017).

(Vandekerckhove and Tuerlinckx, 2008) using realistic ranges of the parameters with the following Uniform distribution draws: 0.35 to 0.5 s of NDT, -4 to 4 evidence units per second of drift rate, 0.8 to 1.4 evidence units required to make a decision (boundary parameter), 0 to 2 evidence units per second of trial-to-trial drift rate standard deviation, 0–0.1 s of additional possible trial-to-trial NDT around the mean. These simulations were generated for 30 EEG sessions (which could be assumed to be different and/or the same participants on different days) with 100 trials each, repeated 30 times with increasing amounts of contaminant trials (up to 10% of trials).

These simulations revealed that even with the presence of trial-to-trial variability in two cognitive parameters (NDT and evidence accumulation rate), means of non-decision time and evidence accumulation rate could still be recovered. In particular, mean NDT across trials was well estimated by 1) 10th percentiles of response time distributions (overestimating NDT with about 60 ms of consistent bias) and 2) hierarchical Bayesian fitting of a simple drift-diffusion model (DDM) accounting for between-session variability in the three free parameters (Wabersich and Vandekerckhove, 2014).

It should be noted that mean NDTs are not as well estimated in the presence of a large amount of contaminant trials without explicitly modeling contaminant trials (see Fig. 5). Contaminant trials are defined

as response time observations that are not due to a decision-making process and occur due to another random process (e.g. a “contaminant” process unrelated to the theorized cognitive process). Because these response times are not due to a decision process, the associated accuracy on those trials should be about 50% in a two-alternative forced choice task. Recently researchers have developed theory on the notion of “mind-wandering”, which is defined as cognition that departs from that which is useful to the task being completed (e.g. Hawkins et al., 2015; van Vugt et al., 2015; Fox et al., 2015; Lin et al., 2016). We assumed that each reaction time and choice observation was from a mixture model such that some percentage of trials was from a Diffusion Model process and some percentage of trials was from a uniform distribution of reaction times and a binomial distribution of accuracy with parameter $\theta = .5$ (that is, the sample mean of accuracy would be about 50%). Drugowitsch et al. (2012) previously used this method to account for so-called “lapse trials”, although we prefer the term “contaminant” trials in this paper.

2.6. N200 waveforms versus response times

To test our hypothesis using a simple method, we performed linear regressions of 1) trial-averaged N200 deflection times and peak-latencies versus 10th response time percentiles across noise conditions and EEG




Fig. 5. The robustness of non-decision time (NDT) estimation to contaminant processes was evaluated by calculating the slope parameter between true and estimated NDT for 30 simulated EEG sessions with 100 trials each, repeated 30 times with increasing amounts of contaminant trials. The dashed gold line represents a slope of 1 and thus would represent recovered NDTs for each participant. Without explicitly modeling contaminant trials, the addition of more trials from contaminant processes resulted in some non-decision time (NDT) estimates being negatively biased towards smaller values, resulting in smaller slope parameters (Green). This simulation also revealed that while 10th response time percentiles were positively biased as estimates of true NDT (e.g. true NDT were approximately 60 ms shorter than the 10th response time percentiles), the slope of 1 between estimated and true NDT relationship could be well estimated by the 10th percentiles (Purple). Explicitly modeling lapse trials (Orange) was shown to be a consistent procedure for finding NDT correlates in EEG data, although the estimated NDT are slightly underestimated.

sessions and 2) single-trial N200 peak-latencies versus response times. The fastest response times in each EEG session and noise condition are expected to be the linear sum of *visual encoding time* (VET) and other sequential cognitive processes not related to decision making. This is because in any response time process that contains a mix of stochastic evidence accumulation and fixed processes with smaller amounts of variability, NDT (time periods not associated with evidence accumulation during decision-making trials, see Voss et al., 2004; Ratcliff and McKoon, 2008) will be reflected in the fastest response times due to some trials having very fast (albeit possibly incorrect) evidence accumulation times but similar NDTs. We observed this in the simulations previously discussed. Thus, the fastest response times in each EEG session and noise condition were estimated by the 10th percentile. The 10th percentile was chosen because the smaller percentiles were more likely to include contaminants as discussed above.

Linear regression statistics reported are 95% confidence intervals related to the p values and t statistics in the Neyman and Pearson (1933) tradition as well as Bayes Factors (BF) which describe the amount of relative evidence (probability in the Bayesian definition) of a model which has a non-zero regression slope over a model which has a regression slope of zero (Jeffreys, 1961; Kass and Raftery, 1995; Rouder and Morey, 2012). Generally Bayes Factors over 3 are considered positive evidence for an effect (i.e. the effect is 3 times more likely under the alternative hypothesis than the null hypothesis) while over 20 is strong evidence (Kass and Raftery, 1995). Adjusted R^2 is also reported that describes the fraction of variance of the dependent variable (10th response time percentiles or single-trial reaction times) explained by the regressor variable (measures of neural processing speed). R_{adj}^2 ranges from 0 to 1. These statistics were generated by JASP, an open-source graphical software package for statistical analysis (JASP Team, 2017).

Also reported are Bayes Factors (BF1) of linear relationships between response time and N200 measures that describe the amount of relative evidence of the slope parameter being equal to 1 (e.g. an x ms increase in

N200 peak-latency corresponds to an x ms increase in visual encoding time) over a model which has a largely unknown regression slope. That is, the comparison density and prior probability of slope parameter was $\beta \sim \mathcal{N}(1, 3^2)$. These Bayes Factors compared the hypothesis of a “spike” at 1 with no uncertainty in possible slope values versus a large uncertainty with a standard deviation of 3 centered at 1. BF1 were calculated using the simple Savage-Dickey Density Ratio (Dickey and Lientz, 1970; Wagenmakers et al., 2010) which is the ratio of the height of the posterior distribution over the prior distribution at the tested value. These Bayes Factors were calculated by running simple JAGS code, a program that easily draws samples from marginal posterior distributions using Markov Chain Monte Carlo (MCMC; Plummer, 2003). These Bayes Factors, BF1, are typically much smaller than the Bayes Factors used to test the existence of an effect because less data is needed to prove the existence of an effect than to prove the effect is equal to a specific value (slope equal to 1, in this case).

2.7. Integrated neuro-cognitive model fitting

In order to test our N200-VET hypothesis directly, we fit hierarchical models to behavioral data and assumed linear influences of the N200 peak-latency on cognitive parameters. Thus all linear relationships between N200 peak-latency and NDT in every visual noise condition were estimated in a single-step in a hierarchical Bayesian framework.

Drift-diffusion modeling was applied to response time and accuracy data from Experiments 1 & 2 jointly, containing between session-differences in NDT τ , evidence accumulation rate δ (generally thought to reflect the average evidence accumulation ability of the participant across trials), and boundary parameter a (evidence threshold which is typically manipulated by a speed-accuracy trade-off condition) that were explained differences in N200 waveform peak-latencies z . All models were hierarchical, accounting for shared variance between session j , condition k , and experiment e variability which 1) ensured model fits

with small amounts of data (Lee, 2008; Lee and Newell, 2011; Vandekerckhove et al., 2011) and 2) provided the ability to describe data more accurately for observed and unobserved participants in future analyses (Wagenmakers, 2009; Nunez et al., 2017a,b). Bayesian hierarchical drift-diffusion models were fit using Wiener likelihood approximations in JAGS, a program that can easily sample from hierarchical models using Markov Chain Monte Carlo (MCMC) (Plummer, 2003; Wabersich and Vandekerckhove, 2014) using the pyjags Python package (Miąsko, 2017).

Hierarchical Model 1 tested a linear influence of trial-averaged N200 peak-latencies z_{jke} on each parameter. Hierarchical Model 2 tested the linear influence of single-trial estimated N200 peak-latencies on non-decision time. Notation “ \sim ” refers to a probability distribution of data or prior distribution of parameters (either from a normal distribution with notation \mathcal{N} , truncated normal distribution with truncation between a and b indicated by $\in (a, b)$, Gamma distribution with notation Γ , or Uniform distribution with notation \mathcal{U}).

Both models assumed that a certain proportion θ of response times (varying between 0% and 100% of trials) came from a contaminant lapse process (discussed above) that produced a random response (Bernoulli distribution with probability parameter $\rho = .5$) and a random (Uniformly distributed) response time between 0 and a large response time (bounded by the maximum observed response time for that EEG session in that visual noise condition). Note that notation $y \sim \mathcal{U}(-\max(\text{RT}), \max(\text{RT}))$ expresses this information when negative y encode error responses and positive y encode correct responses as in the package by Wabersich and Vandekerckhove (2014).

In Model 1, wide (uninformative) priors were given to the hierarchical effects μ_i of N200 peak-latency on drift-diffusion model parameters i , centered at 1 in order to calculate Bayes Factors (BF1) testing a slope-of-1 relationship using the Savage-Dickey density ratio (Verdinelli and Wasserman, 1995). As a sensitivity analysis, wide priors centered at 0 were used in previous fits of the model, which had inconsequential effects on the posterior distributions (i.e. parameter estimates). The final fit of Model 1 thus had the following embedded linear regression and prior structure:

$$\begin{aligned} \tau_{jke} &\sim \mathcal{N}(\eta_{(\tau)ke} + \gamma_{1ke} z_{jke}, \sigma_{(\tau)}^2) \in (0, 1), \eta_{(\tau)ke} \sim \mathcal{N}(0.3, 0.25^2), \sigma_{(\tau)} \sim \Gamma(0.2, 0.7) \end{aligned} \quad (1)$$

$$\delta_{jke} \sim \mathcal{N}(\eta_{(\delta)ke} + \gamma_{2ke} z_{jke}, \sigma_{(\delta)}^2) \in (-9, 9), \eta_{(\delta)ke} \sim \mathcal{N}(1, 2^2), \sigma_{(\delta)} \sim \Gamma(1, 1) \quad (2)$$

$$\alpha_{jke} \sim \mathcal{N}(\eta_{(\alpha)ke} + \gamma_{3ke} z_{jke}, \sigma_{(\alpha)}^2) \in (0.1, 3), \eta_{(\alpha)ke} \sim \mathcal{N}(1, 0.5^2), \sigma_{(\alpha)} \sim \Gamma(1, 1) \quad (3)$$

$$z_{jke} \sim \mathcal{N}(\zeta_{ke}, \sigma_{(z)}^2), \zeta_{ke} \sim \mathcal{N}(0.2, 0.1^2), \sigma_{(z)} \sim \Gamma(0.2, 1) \quad (4)$$

$$\gamma_{ike} \sim \mathcal{N}(\mu_i, \sigma_{(\gamma)}^2), \mu_i \sim \mathcal{N}(1, 3^2), \sigma_{(\gamma)} \sim \Gamma(1, 1) \quad (5)$$

Bayes Factors for each effect parameter γ within each visual noise condition and type were calculated using Python and R scripts with the Savage-Dickey density ratio (Verdinelli and Wasserman, 1995) of the posterior density over the prior density at $\gamma = 1$. These Bayes Factors provide the degree of evidence (defined as a probability ratio in Bayesian statistics) of a model where effect of trial-averaged N200 peak-latency on NDT is 1 versus a model where the effect is unknown (the prior model where $\gamma \sim \mathcal{N}(1, 3^2)$). The base likelihood that generates observed response time and accuracy data (both represented by vector \mathbf{y}) is given by the mixture of a DDM and a lapse process described above, with an uninformative prior on the proportion of lapse trials θ (uniformly distributed from 0% to 100%) for every session j , condition k , and

experiment e :

$$\begin{aligned} \mathbf{y}_{njke} &\sim (1 - \theta_{jke}) \text{DDM}(\tau_{jke}, \delta_{jke}, \alpha_{jke}) \\ &\quad + \theta_{jke} \mathcal{U}(-\max(\text{RT})_{jke}, \max(\text{RT})_{jke}) \end{aligned} \quad (6)$$

$$\theta_{jke} \sim \mathcal{U}(0, 1) \quad (7)$$

Model 2 was fit to find posterior distributions of decision-model parameters for *single-trial* N200 peak-latency linear effects on single-trial NDT (the sum of VET and motor response time as estimated by a hierarchical Bayesian account of a drift-diffusion model). The model was fit with linear relationships between N200 peak-latencies x on single-trials n and single-trial non-decision times, a similar framework presented in our previous work (see Nunez et al., 2017a,b). Also included were hierarchical linear effect parameters in order to summarize the linear effects for each noise condition and noise type (i.e. experiment). Model 2 had the following prior structure, a hierarchy very similar to Model 1:

$$\tau_{jke} \sim \mathcal{N}(\eta_{(\tau)ke}, \sigma_{(\tau)}^2) \in (0, 0.7), \eta_{(\tau)ke} \sim \mathcal{N}(0.3, 0.25^2), \sigma_{(\tau)} \sim \Gamma(0.2, 1) \quad (8)$$

$$\delta_{jke} \sim \mathcal{N}(\eta_{(\delta)ke}, \sigma_{(\delta)}^2) \in (-9, 9), \eta_{(\delta)ke} \sim \mathcal{N}(1, 2^2), \sigma_{(\delta)} \sim \Gamma(1, 1) \quad (9)$$

$$\alpha_{jke} \sim \mathcal{N}(\eta_{(\alpha)ke}, \sigma_{(\alpha)}^2) \in (0.1, 3), \eta_{(\alpha)ke} \sim \mathcal{N}(1, 0.7^2), \sigma_{(\alpha)} \sim \Gamma(1, 1) \quad (10)$$

$$z_{jke} \sim \mathcal{N}(\zeta_{ke}, \sigma_{(z)}^2), \zeta_{ke} \sim \mathcal{N}(0.2, 0.1^2), \sigma_{(z)} \sim \Gamma(0.2, 1) \quad (11)$$

$$\gamma_{ke} \sim \mathcal{N}(\mu_i, \sigma_{(\gamma)}^2), \mu_i \sim \mathcal{N}(1, 3^2), \sigma_{(\gamma)} \sim \Gamma(1, 1) \quad (12)$$

$$\lambda_{jke} \sim \mathcal{N}(\gamma_{ke}, \sigma_{(\lambda)}^2), \sigma_{(\lambda)} \sim \Gamma(1, 1), \sigma_{(x)} \sim \Gamma(0.2, 1) \quad (13)$$

Like Model 1, a lapse process was assumed in Model 2. Model 2 also had a simple normally distributed generative process for the N200 peak-latency x on trial n truncated between 0.151 and 0.274 s (see single-trial N200 peak-latency estimation discussion above). Model 2 thus had the following embedded linear regression structure on the likelihood level (where variable \mathbf{y} refers to the choice and response time on trial n from a drift-diffusion model; DDM):

$$\begin{aligned} \mathbf{y}_{njke} &\sim (1 - \theta_{jke}) \text{DDM}(\tau_{jke} + \lambda_{jke} x_{njke}, \delta_{jke}, \alpha_{jke}) \\ &\quad + \theta_{jke} \mathcal{U}(-\max(\text{RT})_{jke}, \max(\text{RT})_{jke}) \end{aligned} \quad (14)$$

$$x_{njke} \sim \mathcal{N}(z_{jke}, \sigma_{(x)}^2) \in (0.151, 0.274), \theta_{jke} \sim \mathcal{U}(0, 1) \quad (15)$$

2.8. Location of the N200 waveforms

We would expect VET to reflect a network-wide process of figure-ground segregation that involves multiple cortical regions. Therefore to further test the N200-VET hypothesis we wished to localize the N200 waveforms. Spline-interpolation on the scalp surface has been shown to be accurate for high-density EEG (approximately > 64 electrodes; Nunez and Srinivasan, 2006). To view the scalp representation of the observed N200 waveforms, we used spline-interpolation in each EEG session to estimate the potential between electrode contacts and to generate interpolated data for artifactual electrode contacts. This method yielded the best representation of the N200 peak-latency on the surface of a Montreal Neurological Institute (MNI) average human scalp (ICBM-152; Fonov et al., 2011, e.g. see the Top panel of Fig. 9).

EEG localization to a three-dimensional representation of the brain is an inexact process that is unsolvable without additional assumptions




Fig. 6. Top Left: A scatter plot of N200 deflection times and 10th response time percentiles (an estimate of non-decision time; NDT) Observations were generated per noise condition and per EEG collection session ($N = 147$). Top Right: A scatter plot of N200 peak-latencies and 10th response time percentiles. Bottom Left: A scatter plot of N200 peak-latencies and NDT estimated using the medians of posterior distributions from Model 1. Bottom Right: A scatter plot of single-trial N200 latencies versus single-trial response times. Observations were generated per trial ($N = 13,462$). Best-fit simple linear regression lines are shown in black with 95% confidence intervals for the intercept and slope parameters shown in gray. Overlaid on the linear regression lines are gold dashed lines representing the hypothesis with a slope of 1.

(Nunez and Srinivasan, 2006). However the surface spline Laplacian (i.e. “Current Source Density”, CSD) on a curved two-dimensional space does not require additional assumptions, has been shown to match closely to simulated cortical-surface activity using forward models (i.e the mapping of cortical activity to scalp potentials), are unaffected by reference electrode choices, and have shown consistent results when used with real EEG data (Nunez and Srinivasan, 2006; Kayser and Tenke, 2015). In addition, unlike 3D solutions, projections to the dura mater surface with CSD are theoretically solvable, and have been used with success in past

studies (see Nunez et al., 1994, for an example). Topographic maps of the surface spline Laplacian projection of the SVD-estimated, average N200 waveform across EEG sessions and noise conditions are shown in the Middle panel of Fig. 9. The minimum of the average true-ERP across all observations ($N = 147$) yielded a very similar Laplacian map.

Repeating a similar “source-localization” technique found in Nunez et al. (2017a,b), we localized the curved 2D surface spline Laplacians (Deng et al., 2012) to a folded cortical surface, which resulted with most amplitude existing on the gyral crowns (i.e. the “smooth” cortex). The




Fig. 7. Posterior distributions (Model 1) of effects of N200 peak-latency on non-decision time (NDT). Posterior distributions provide measures of parameter uncertainty in Bayesian model-fitting methods. If the posterior distributions had most of their density around the blue line, this would be strong evidence for the slope-of-1 relationship when the model assumptions are true. Posterior distributions with most of their density around the red line would be strong evidence for the null hypothesis (the parameter equals 0). Some evidence exists for the effects of N200 peak-latency on NDT to have a slope of 1 (x millisecond increase in N200 peak-latency corresponds to a x millisecond increase in NDT) as indicated by the Bayes Factors calculated with a Savage-Dickey density ratio (Verdinelli and Wasserman, 1995) of the posterior density over the prior distribution at $\gamma = 1$.

surface spline Laplacians were projected onto a cortical shape generated by an anatomical MRI of a single individual. This individual's brain image had cortical folds that better localized particular regions than the MNI brain shape. Each participant's N200 projection was performed using Tikhonov (L2) regularization to inverse the Finite Element (FE; Pommier and Renard, 2005) forward model based on the MNI ICBM-152 average head (Fonov et al., 2011), maintaining similar spatial distributions of activity of the surface Laplacians to the cortical surface. Cortical topographic maps of the average N200 waveform across EEG sessions and noise conditions are shown in the Bottom panel of Fig. 9. It should be noted that these projections to the cortical surface are only “Representative” sources (Nunez et al., 2019) and may not one-to-one relate to the genuine sources.

In the cortical maps each brain vertex was labeled using the Destrieux cortical atlas (Fischl et al., 2004). The locations described below were found in at least 70% of localizations across EEG sessions and experimental conditions using an empirically found cutoff (estimated at -10 units microamperes per mm^2). However we should note that the localization procedure must have some errors due to between-participant variance in both cortical and cranial shape and between-participant variance in tissue properties. These sources of variance could not easily be accounted for, and remains a source of error in this cortical projection technique.

2.9. Code accessibility

Pre-calculated EEG measures, raw behavioral data, MATLAB stimulus code, and MATLAB, Python, R, and JAGS analysis code are available on <https://osf.io/jg9u3/> and in the following repository <https://github.com/mdnunez/encodingN200> (as of March 2018).

3. Results

3.1. Data summaries: longer N200 peak-latencies with more difficult visual noise conditions

Data summaries of N200 deflection times, trial-averaged N200 peak-latencies, 10th response time percentiles (shown to be a stable but biased estimate of NDT in Fig. 5), median response times across trials, and mean accuracy across trials are given in Table 1, such that the total sample size of each measure was $N = 147$. A significant but weak correlation was found between the found N200 deflection times and the N200 peak-latencies ($\rho = .274, p < .001$). A significant correlation was found between the N200 peak-latencies and 10th response time percentiles (a less specific test of our main hypothesis; $\rho = .387, p < .001$). No correlation was found between the N200 deflection times and 10th response time percentiles ($\rho = -0.086, p = .30$). From the data summaries in Table 1 it is clear that N200 peak-latencies, 10th response time percentiles, and median reaction times generally become longer in higher-contrast noise conditions, an expected effect of visual noise condition on VET (due to more difficult figure-ground segregation). This was not true between the Low and Medium noise conditions in Experiment 1, although it should be noted that there was not much behavioral difference between Low and Medium noise contrast in Experiment 1 in response time nor accuracy. Decreasing accuracies with higher noise contrast are an expected effect of visual noise contrast on decision-making. The effects of visual noise condition on VET versus decision-making time (both components of response time) were separated by the integrated neuro-cognitive modeling results presented below.

Data summary across unique EEG sessions ($N = 49$) and conditions (total $N = 147$).

Data summary across unique subjects ($N = 14$)




Fig. 8. Posterior distributions (Model 1) of residual non-decision time (NDT) after accounting for the effect of N200 peak-latency. The N200 peak-latency explains most of the variance in non-decision time across visual noise conditions. The residual non-decision time is thought to be an approximate measure of motor execution time (MET) and is not expected to depend upon the visual noise condition. As expected, the posterior distributions are similar in each visual noise condition, yielding only minor evidence for a difference in motor execution times between the two experiments.




Fig. 9. Left and right sagittal and posterior views of localized trial-averaged N200 waveforms averaged across all EEG recordings and noise conditions ($N = 147$). (Top) MNI scalp potential topographies (averaged projections of the first SVD components at the N200 peak-latency) were generated using spline-interpolation between electrodes. (Middle) Current Source Density MNI scalp topographies were generated with a Laplacian transform of the aforementioned data (Nunez and Pilgreen, 1991; Deng et al., 2012; Kayser and Tenke, 2015). (Bottom) The “representative” cortical maps were obtained by projecting the MNI-scalp spline Laplacians onto one participant’s anatomical fMRI image via Tikhonov (L2) regularization, maintaining similar distributions of activity of the surface Laplacians on the cortical surface. Blue and orange regions correspond to areas that produce negative and positive potentials observed on the scalp respectively.

Table 1

Summary statistics for each data type used in this study across EEG sessions and visual noise conditions for a total sample size of 147. Experiment 1 had a sample size of 63 (21 unique EEG sessions per visual noise condition). Experiment 2 had a sample size of 84 (28 unique EEG sessions per visual noise condition). N200 peak-latencies, 10th percentile response times, and median response times become longer in more difficult (lower accuracy) noise conditions. This is not true of N200 deflection times in Experiment 2. VET is expected to become longer when segregation of the visual target is more difficult in conditions of high noise contrast.

		N200	N200	10 th Percentile	Median	Mean
		Deflection times	Peak-latencies	Response times	Response times	Accuracy
All observations	Overall Mean	95.5 ms	199.6 ms	575.0 ms	852.8 ms	78.76%
	5 th Percentile	48.3 ms	162.0 ms	443.6 ms	629.5 ms	58.36%
	Median	97.0 ms	192.0 ms	562.4 ms	845.8 ms	78.76%
	95 th Percentile	136.5 ms	259.8 ms	733.6 ms	1139.5 ms	95.32%
Experiment 1	Low Noise Mean	81.9 ms	166.3 ms	549.1 ms	833.0 ms	72.25%
	Medium Noise Mean	94.0 ms	193.0 ms	541.9 ms	821.3 ms	70.14%
	High Noise Mean	108.2 ms	244.5 ms	613.0 ms	896.8 ms	61.77%
Experiment 2	Low Noise Mean	100.1 ms	175.0 ms	545.3 ms	809.1 ms	90.07%
	Medium Noise Mean	93.6 ms	192.2 ms	565.3 ms	829.7 ms	89.59%
	High Noise Mean	94.5 ms	228.0 ms	629.9 ms	925.3 ms	80.72%

Table 2

Summary statistics for each data type averaged within each of the 14 unique participants. All measures varied across participants suggesting individual differences in VET and other decision-related cognitive processes.

	N200	N200	10 th Percentile	Median	Mean
	Deflection times	Peak-latencies	Response times	Response times	Accuracy
Minimum	57.7 ms	183.2 ms	482.3 ms	649.1 ms	62.67%
Mean	91.7 ms	199.5 ms	567.3 ms	839.7 ms	72.33%
Maximum	129.2 ms	225.3 ms	676.2 ms	1070.8 ms	90.14%

There were 14 unique participants across both experiments (with two participants participating in both experiments). In order to explore true individual differences, observations of the N200 deflection time, N200 peak-latency, 10th response time percentiles, median response times, and accuracies were averaged across the two experiments and all conditions for each participant. The range of these measures are given in Table 2. These measures varied across participants with a 18 ms standard deviation of N200 deflection time, 12 ms standard deviation of N200 peak-latency, 68 ms standard deviation of 10th response time percentiles, 125 ms standard deviation of median response times, and 9% standard deviation of accuracy across participants. These results suggest individual differences in both VET and decision-making time (e.g. response time related to an accumulation of evidence). Some of these individual differences account for the variation observed in N200 deflection times and N200 peak-latency distributions presented in Fig. 3. It should be noted that four of the participants were considered trained on the task, with two participants completing 9 sessions (both experiments) and two other participants completing 7 sessions of the second experiment. While not the focus of this particular study, future work should seek to discover if individual differences in VET are explained by individual differences in N200 peak-latencies (a hypothesis expected from the results of the research presented here). Assuming our hypothesis is true, individuals are expected to have larger differences in their decision-making time than VET.

3.2. Simple results: linear regressions suggest N200 peak-latencies shift reaction times

Testing the hypothesis that trial-averaged N200 peak-latency predicts session-level visual encoding time (VET), linear models between N200 peak-latencies and 10th response time percentiles (both correct and incorrect response times combined) were fit. In evidence accumulation models of quick decision making, the shifts of response time (RT) distributions (and thus the small RT percentiles) should correlate with NDT

(i.e. VET + motor execution time) while the shape of RT distributions (and thus the medium and larger RT percentiles) should correlate both with decision-making processes. Thus linear regressions can be explored without fitting decision-making models to data directly. Using simple linear regression, a slope close to 1 was found between trial-averaged N200 latencies and 10th RT percentiles as shown in the top right of Fig. 6 with a regression coefficient of $\beta_1 = 1.14$ ($N = 147$, 95% confidence interval: [0.65, 1.63], $p < .001$, $t = 4.56$, $BF = 1.58 \times 10^3$, $BF1 = 10.26$). Only 12% of the variance in 10th RT percentiles were explained by the variance across trial-averaged N200 latencies ($R^2_{adj} = .119$). This latter finding might be expected if other trial-invariant mechanisms besides visual encoding time (VET) contribute to the variance in NDT, such as motor execution time. Other regressions between trial-averaged N200 peak-latencies and 30th, 50th, 70th, and 90th RT percentiles and RT means were also significant with 95% confidence intervals that included 1. N200 peak-latencies explained similar or less variance in these RT measures (13%, 11%, 8%, 7% and 9% respectively), which is to be expected given that the larger RT percentiles contain more variance related to evidence accumulation. 1st RT percentiles were found to be explained by N200 peak-latencies only when 350 millisecond cutoff values were used to remove some contaminant trials, otherwise the regression was not significant.

The relationship is also expected to be observed on single-trials, albeit with more variance around the regression line. Single-trial estimates of N200 latencies were compared to response times across all data points. A slope near 1 was found between single-trial N200 peak-latencies and response times with a tight confidence interval as shown in the bottom right portion of Fig. 6 with a regression coefficient of $\beta_1 = 1.05$ ($N = 13,462$, 95% confidence interval: [0.93, 1.18], $p < .001$, $t = 16.65$, $BF = 6.30 \times 10^{57}$, $BF1 = 31.09$). However only about 2% of the variance of raw single-trial response times were explained ($R^2_{adj} = .020$). This small amount of variance explained is expected since evidence accumulation should contribute to most of the variance in response times on single trials rather than visual encoding time. Evidence accumulation will affect single-trial response times more than early RT percentiles (such as the 10th RT percentile) because the sequential sampling (Wiener) process yields increased variance for longer evidence accumulation windows.

No relationship was found between N200 deflection times and 10th RT percentiles ($N = 147$, $p = .251$, $t = -1.15$, $BF = 0.33$, $BF1 < .001$). The linear regression is shown in the top left of Fig. 6. In fact, the Bayes Factor $BF1$ displayed evidence strongly in favor of $\beta_1 \neq 1$ ($1/BF1 = 1.05 \times 10^4$). This suggests that while N200 deflection times may reflect an early visual processing stage, variance in N200 deflection times across noise conditions and EEG sessions does not correspond with variance in response time distribution shifts across noise conditions and EEG sessions.

3.3. Modeling results: direct evidence that N200 peak-latencies track visual encoding time

Each model was fit using JAGS with six Markov Chain Monte Carlo (MCMC) chains of 52,000 samples each run in parallel with 2,000 burn-in samples and a thinning parameter of 10 resulting in 5,000 posterior samples in each chain. The posterior samples from each chain were combined to form one posterior sample of 30,000 samples for each parameter. Model convergence was evaluated based on the Gelman-Rubin statistic \hat{R} , which compares the estimated between-chains and within-chain variances for each model parameter (Gelman and Rubin, 1992). Using this statistic both models were judged to have converged, with $\hat{R} \leq 1.08$ for all parameters in each model ($\hat{R} > 1.10$ for any parameter is usually deemed evidence for a lack of convergence of the entire model fit).

Model 1 ($\hat{R} \leq 1.02$ for all parameters) was used to evaluate the effect of trial-averaged N200 peak-latency on NDT across EEG sessions and noise conditions. The mean effect across all conditions of each experiment μ_γ provides a sense of how N200 peak-latency tracks NDT τ in general. The posterior distribution of this parameter was found to be around 1 in each condition and experiment as shown in Fig. 7 and the bottom left portion of Fig. 6, indicating that an x ms increase in N200 peak-latency corresponded to a roughly an x ms increase in NDT. The posterior distribution of the overall hierarchical effect of N200 peak-latency on NDT across sessions is plotted at the top of Fig. 7, indicating that the overall Bayes Factor for a slope-of-1 relationship between N1 latency and NDT is $BF_1 = 4.62$, reflecting moderate evidence (median = 1.39, 95% credible interval: [0.62, 2.26]). However all the posterior distributions are centered above 1 for each unique noise condition. This is expected if N200 peak-latencies reflect VET and estimated NDT are slightly underestimated when including a contaminant process as in Model 1, which is shown by the Orange regression line in Fig. 5. The effects of N200 peak-latency on two decision-making parameters was also explored in order to disentangle the three cognitive parameters. An undetermined effect of trial-averaged N200 peak-latency on evidence accumulation rate δ was found (median = -1.53 evidence units per second, 95% credible interval: [$-5.38, 2.27$]). An undetermined effect of trial-averaged N200 peak-latency on the evidence boundary α was found (median = 0.69 evidence units per second, 95% credible interval: [$-0.93, 2.44$]), assuming the evidence boundary is well estimated (which was not shown in simulation).

The residual non-decision time $\eta_{(\tau)}$ posterior distributions yield predictions about the motor execution time (MET), thought to be the second component of NDT that is not VET. Even though NDT changes based on

the visual noise condition, these changes should be reflected in the VET (and thus N200 peak-latencies) and not the motor execution time (MET). Thus the residual non-decision time should be similar in each visual noise condition. This is indeed the case, with all residual non-decision time posterior distributions overlapping significantly (see Fig. 8), reflecting a large probability that the residual NDT is equal in each noise condition. While the median posterior distributions range from 170 to 260 milliseconds, the best estimate of MET is given by the condition with the slope parameter posterior closest to 1, implied by the High Noise contrast condition in Experiment 1. This suggests that the best estimate for MET is 260 milliseconds, which matches estimates for MET we found in another study by Lui et al. (2018).

Model 2 ($\hat{R} \leq 1.08$ for all parameters) was fit in order to evaluate the effect of single-trial N200 peak-latency on non-decision time for each EEG session. In Model 2, the hierarchical effect parameter μ_γ provides a sense of how *single-trial* N200 peak-latency effects NDT. This parameter suggested that the overall single-trial effect was positive (median = 0.34, 95% credible interval: [0.22, 0.46]). This positive relationship on single-trials between N200 latency and NDT replicates our previous finding in different dataset (Nunez et al., 2017a,b). However no evidence was found that this single-trial effect was equal to 1 ($BF_1 < .001$). That is, there was strong evidence *against* a slope of 1 on single-trials between N200 peak-latency and NDT. Model 2 necessarily does not allow for measurement error in N200 peak-latencies on single trials. Model 2 thus may not accurately capture a true correspondence to VET when the single-trial neural measure contains measurement noise (see Hawkins et al., 2017, for a discussion on noisy neural measures embedded in cognitive models).

3.4. Brain networks that may generate N200 waveforms

N200 peak-latencies are distributed across the cortex and not localized to one particular area. This is reflective of a network that gives rise to the N200 peaks instead of local processing. The distributed network is evident from different focal regions of the average N200 waveform in the spline Laplacian (see Middle portion of Fig. 9) as opposed to locations only at the center of dual hemispheric posterior regions that might be inferred from the scalp potentials alone (see Top portion of Fig. 9). This is also evidenced by the deviating time course of all 128 electrode locations at the N200 peak latency in the spline Laplacian of each ERP, as shown in one example in Fig. 10 instead of only 1 or 2 electrode locations that show a peak around 181 ms. The spline Laplacian results leads us to reject the hypothesis of a single generator of N200 peak-latencies. This is also true of the localization presented at the Bottom of Fig. 9, however




Fig. 10. The time course of one indicative traditional ERP (Top) from Experiment 2 in the Medium noise condition for 128 electrode locations filtered from 1 to 10 Hz and spline Laplacian transform of the same ERP (Bottom) at the same locations. ERP potentials and spline Laplacians are shown on posterior views of the MNI scalp at 92 milliseconds (Gold, left) and 181 milliseconds (Pink, right) using spline interpolation between electrode locations. The variability across electrode locations of the spline Laplacian transform at 181 ms is indicative of a distributed network that gives rise to the N200 peak-latency. The topography of average SVD-localized N200 waveforms is shown in the middle portion of Fig. 9.

the sources estimated are only representative (see Nunez et al., 2019) and may not be that precise.

While the estimated cortical sources were distributed, the source estimation method described above suggested that N200 waveforms in this study could be found primarily in the following cortical locations (such that the locations listed here were present in at least 70% of the localizations across EEG sessions and noise conditions): the middle temporal gyri (both hemispheres), the postcentral gyri (both hemispheres), and the right angular gyrus. The right middle temporal gyrus was the location of the minimum amplitude (i.e. the greatest magnitude of the N200 peak) in 55.8% of observations across EEG sessions and noise conditions. While all other locations contained the largest negative amplitude in less than 10% of observations. Work by Mishkin et al. (1983) and Lamme and Roelfsema (2000), among others, suggest that the segregation of the target Gabor from visual noise would occur in the temporal “segregation” pathway of the primate cortex at about 150 ms (after a delay due to feedforward and feedback visual network behavior). The possible localization of the N200 waveforms in human extrastriate and temporal locations (especially the right middle temporal gyrus) is further evidence that the N200 peak-latency is the time point at which the target Gabor is “segregated” from visual noise in the human cortex, in order to accumulate visual evidence for a motor response.

4. Discussion

Evidence is presented both for and against the N200-VET hypothesis. Evidence against hypotheses are often left out of papers presenting initial results in both the Cognitive Psychology and Cognitive Neuroscience fields, often presented as confirmatory results. Such “censoring” of data can lead to biased statistical analyses (Sterling, 1959; Rosenthal, 1979; Guan and Vandekerckhove, 2016).

4.1. Evidence FOR N200 peak-latencies tracking visual encoding time

Both simple regression analyses yielded evidence that EEG measures of neural processing speed (N200 peak-latencies) track visual encoding times (VET). These regression fits indicate that there was 1) strong evidence for an x ms increase in response times when there was an x ms increase in single-trial N200 peak-latency and 2) evidence for an x ms increase in the 10th percentile of response times, an estimate of cognitive non-decision time (NDT), when there was an x ms increase in trial-averaged N200 peak-latencies. Furthermore, moderate evidence was found for a slope of 1 when fitting embedded linear regression of trial-averaged N200 peak-latency on NDT within the hierarchical drift-diffusion Model 1 for each visual noise condition. In addition, the residual NDT in Model 1 were similar across each visual noise types while N200 peak-latency described the differences in NDT across the different visual noise types, a result to be expected if residual NDT reflects motor execution time (MET).

Our findings in this study closely match findings discussed by Thorpe et al. (1996), Martin et al. (2010), Zhang et al. (2017), and Loughnane et al. (2016) as discussed in the Introduction of this paper. A phenomena of N200 peak-latencies reflecting VET would also explain why N200 latencies have been found to be affected by bottom-up and top-down attention (Vogel and Luck, 2000), which would be expected of VET because of the ability of the visual cortex to favor attended visual input (Luck et al., 1997). NDTs (of which VET is an additive component) have previously been shown to be affected by attention across participants and trials (Nunez et al., 2015, 2017). In this study we found significant noise-condition effects in N200 peak-latency (see Table 1) such that N200 peak-latencies become longer with larger amounts of visual noise contrast (this finding was not observed in the N200 deflection times), and it is expected the VET should become longer with larger amounts of visual noise due to a higher cognitive difficulty of target-noise segregation (e.g. a form of figure-ground segregation). Furthermore we found some individual differences in N200 peak-latencies across participants, which

matches expectations that individual differences in cognitive processing exist that explains individual differences in behavior (e.g. see Schubert et al., 2017, 2018).

4.2. Evidence AGAINST N200 peak-latencies tracking visual encoding time

Single-trial N200 latencies may be related to, but not be equal to, VET as indicated by the positive slopes not equal to 1 found between single-trial N200 peak-latencies and NDT in Model 2. This could be due to an overabundance of measurement noise in single-trial N200s. It may also be the case that the window (151–274 ms) used on single-trials introduced a “missing not at random” (MNAR) process that adversely affected the modeling. However, it is also possible that the single-trial N200 latencies do give a satisfactory account of VET with the specified window but fitting simple drift-diffusion parameter relationships fails on single-trials (see Hawkins et al., 2017), especially with the presence of contaminant processes. This would explain why we found a slope near 1 between single-trial N200 peak-latencies and response times (see bottom right panel of Fig. 6) but no evidence of this slope of 1 (albeit a significant and positive relationship) when fitting DDM parameters to N200 peak-latencies on single trials. Better theoretical models of behavior with trial-to-trial variability in NDT are probably needed (see Boehm et al., 2018), such as cognitive models that include both decision processes and mind-wandering processes (e.g. see van Vugt et al., 2015; Mittner et al., 2016). Known DDMs that account for intrinsic trial-to-trial variability in drift rate that are necessary to capture incorrect RT variance may also be helpful (see Heathcote et al., 2018) and is an interesting avenue of future work. Another alternative is a joint model of single-trial N200 latencies and human behavior that can be fit with Bayesian MCMC sampler programs, discussed below.

Some readers may be surprised that a slope-of-1 relationship was found between single-trial N200 peak-latencies and single-trial response times. We might expect single-trial attention effects on both VET and decision-related processes (e.g. see Nunez et al., 2017a,b) resulting in hypothesized slopes greater than 1 even when single-trial N200 peak-latencies have no effect on evidence accumulation directly. The initial results presented in this paper suggest that attention may affect VET and evidence accumulation independently. A further study must be conducted to determine whether single-trial attention effects confound these relationships when not accounting for the cognitive process using modeling.

Reviewing other studies in the literature, we found mixed evidence for the N200-VET hypothesis. However in many of these studies, cognitive models of decision making were not simulated to verify accurate measurements of NDT, as we performed in this study (see Fig. 5). This possible failure of parameter recovery is a known problem for some decision-model fitting procedures (Ratcliff and Tuerlinckx, 2002), and care should be taken to evaluate model recovery ability before hypothesis testing in future studies. As an example, Stanford et al. (2010) found NDT estimates of 100 ms by direct approximation using a SIMPLEX method, thus finding evidence for VET less than 100 ms that could not be tracked by an N200 ERP. However, the shift of the reaction time curves presented in Fig. 3e of the paper by Stanford et al. (2010) suggest that non-decision time (NDT) should have been estimated between 150 and 200 ms (see Fig. 2 of Lui et al., 2018, for an example of graphical NDT estimation). Considering that post-decisional delays in the study by Stanford et al. (2010) are approximated to only be fast saccades of about 30 ms, the 120–170 ms delay for VET would be in agreement with our results. Furthermore, Stanford et al. (2010) showed that when evidence accumulation was blocked for only 100 ms after target onset, monkeys were able to complete their decision, while when evidence accumulation was blocked for 200 ms after target onset, the monkeys were at chance accuracy, suggesting that visual encoding time must have a latency between 100 ms and 200 ms for a simple color discrimination task.

4.3. A neuro-cognitive theory of decision making

A neuro-cognitive model of encoding, evidence accumulation, and motor response is proposed that is identified with separable measures of VET and MET that are estimated from observed EEG, accuracy, and response time distributions. Fitting this type of model to data would allow direct hypothesis testing of observed trial-to-trial differences and individual differences in neural signals and human behavior during quick decision making.

Neuro-cognitive hierarchical models of VET, evidence accumulation, and motor response time become identifiable when treating EEG measures (e.g. N200 peak-latencies) as measures of VET. The theory is an extension of a specific class of decision-making models (e.g. “drift-diffusion” models, see Ratcliff and McKoon, 2008, for a review) which predicts that response time and accuracy are explained by a continuous accumulation of evidence towards certain pre-decided evidence thresholds (see Fig. 4).

A drift-diffusion model of accuracy ($w = 0$ or $w = 1$) and response time t on one trial is predicted to be a function of VET τ_e , MET τ_m , the drift rate δ which is the average rate of evidence accumulation within one trial, the diffusion coefficient ζ which is the variance around the average rate of evidence accumulation within a trial, and the amount of evidence α_t required to make a decision, which may be assumed to decrease with time. The bias towards correct or incorrect responses is equal to $z_0 = 0.5\alpha_t$. The joint density f of RT t and accuracy w of this simplified diffusion model is given in Equation (16). The density is derived from the solution given by Ratcliff (1978) and Tuerlinckx (2004).

$$\left\{ \begin{array}{l} f(t, w = 0 | \alpha, \zeta^2, \tau_e, \tau_m, \delta) = \frac{\pi \zeta^2}{\alpha^2} e^{-\frac{1}{2\zeta^2} [\delta \alpha_t + \delta^2 (t - \tau_e - \tau_m)]} \sum_{k=1}^{\infty} \left[k \sin\left(\frac{1}{2} \pi k\right) e^{-\frac{1}{2\alpha^2} \pi^2 \zeta^2 (t - \tau_e - \tau_m)} \right] \\ f(t, w = 1 | \alpha, \zeta^2, \tau_e, \tau_m, \delta) = f(t, w = 0 | \alpha, \zeta^2, \tau_e, \tau_m, -\delta) \end{array} \right. \quad (16)$$

Note that the density in Equation (16) is classically unidentifiable when estimating the model with only response time and accuracy observations for two reasons. 1) Visual encoding time τ_e and motor execution time τ_m both contribute to non-decision time $\tau = \tau_e + \tau_m$ which approximately equals the smallest observed response times that are not due to contaminant trials. 2) Only two of the three parameters related to evidence accumulation (i.e. drift rate δ , the diffusion coefficient ζ , and the evidence boundary α) can be found with behavioral data alone. For example, if α_t is constant with t , multiplying ζ by two and dividing both α and δ by two would result in the same fit of choice-RT (Wabersich and Vandekerckhove, 2014). Ideally, a version of the model given in Equation (16) with four free parameters (e.g. τ_e , τ_m , δ and a constant α in time t) would be estimated in JAGS (JASP Team, 2017) or Stan (Carpenter et al., 2016) to allow easy fitting of hierarchical models using the likelihood approximation by Navarro and Fuss (2009) (or Wiecki et al., 2013, for the likelihood approximation extended to include drift-rate variability). Alternatively, the model may be approximated by shortcuts in JAGS code with hierarchical models; research currently being conducted by the authors of this manuscript.

Fitting neuroscientific results to cognitive models that explain human behavior is a growing field of research (Mulder et al., 2014; Turner et al., 2015, 2017; de Hollander et al., 2016). The goal of future work is to further develop theory that reconciles observed brain function in the Cognitive Neuroscience literature with theoretical behavioral-cognitive relationships. This work could lead to the ability to fit the full density given in Equation (16) with identifiable parameters that explain both psychological phenomena and brain electrophysiology. This would allow

researchers and to directly test hypotheses in experimental data and provide researchers and clinicians with a method to explore the cognitive and brain-computational underpinnings behind observed human behavior.

Conflicts of interest

The authors declare no competing financial interests.

Author contributions

MDN, JV, and RS designed research. AG and MDN performed research. MDN analyzed data. MDN, AG, JV, and RS wrote the paper.

Significance statement

Encoding of a sensory stimulus is believed to be the first step in perceptual decision making. In this study, we report evidence that visual evoked potentials (EPs) around 200 ms after stimulus presentation track the time of visual figure-ground segregation before the onset of evidence accumulation during decision making. These EP latencies vary across individuals, are modulated by external visual noise, and increase response time by x milliseconds when they increase by x milliseconds. Hierarchical Bayesian model-fitting was also used to relate these EPs to a specific cognitive parameter that tracks time related to visual encoding in a decision-making model of response time. This work adds to the growing literature that suggests that EEG signals can track the component cognitive processes of decision making.

Acknowledgements

MDN, JV, and RS were supported by NSF grant #1658303. AG was supported by the Summer Undergraduate Research Program at the University of California, Irvine. Siyi Deng and Sam Thorpe are thanked for their contributions to the FEM solutions and anatomical fMRI image generation. Gabriel Weindel is thanked for his early comments and reanalysis of the public data paired with this paper. We would also like to thank Paul L. Nunez for his comments on the spline Laplacian-localization technique used in this paper. All reviewers are appreciated for their comments that lead to the improvement of this work. This study extends initial work presented in Chapter 4 of the PhD dissertation by Nunez (2017).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.neuroimage.2019.04.052>.

References

- Acunzo, D.J., MacKenzie, G., van Rossum, M.C., 2012. Systematic biases in early ERP and ERF components as a result of high-pass filtering. *J. Neurosci. Methods* 209, 212–218.
- Bach, M., Meigen, T., 1992. Electrophysiological correlates of texture segregation in the human visual evoked potential. *Vis. Res.* 32, 417–424.
- Boehm, U., Annis, J., Frank, M.J., Hawkins, G.E., Heathcote, A., Kellen, D., Krypotos, A.M., Lerche, V., Logan, G.D., Palmeri, T.J., et al., 2018. Estimating across-trial variability parameters of the diffusion decision model: expert advice and recommendations. *J. Math. Psychol.* 87, 46–75.

- Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M.A., Guo, J., Li, P., Riddell, A., 2016. Stan: a probabilistic programming language. *J. Stat. Softw.* 20, 1–37.
- de Hollander, G., Forstmann, B.U., Brown, S.D., 2016. Different ways of linking behavioral and neural data via computational cognitive models. *Biol. Psychiatry: Cogn. Neurosci. Neuroimag.* 1, 101–109.
- Deng, S., Winter, W., Thorpe, S., Srinivasan, R., 2012. Improved surface laplacian estimates of cortical potential using realistic models of head geometry. *IEEE (Inst. Electr. Electron. Eng.) Trans. Biomed. Eng.* 59, 2979–2985.
- Dickey, J.M., Lientz, B.P., 1970. The weighted likelihood ratio, sharp hypotheses about chances, the order of a Markov chain. *Ann. Math. Stat.* 41, 214–226.
- Dmochowski, J.P., Norcia, A.M., 2015. Cortical components of reaction-time during perceptual decisions in humans. *PLoS One* 10, e0143339.
- Drugowitsch, J., Moreno-Bote, R., Churchland, A.K., Shadlen, M.N., Pouget, A., 2012. The cost of accumulating evidence in perceptual decision making. *J. Neurosci.* 32, 3612–3628.
- Fischl, B., van der Kouwe, A., Destrieux, C., Halgren, E., Ségonne, F., Salat, D.H., Busa, E., Seidman, L.J., Goldstein, J., Kennedy, D., et al., 2004. Automatically parcellating the human cerebral cortex. *Cerebr. Cortex* 14, 11–22.
- Fonov, V., Evans, A.C., Botteron, K., Almli, C.R., McKinsty, R.C., Collins, D.L., Group, B.D.C., et al., 2011. Unbiased average age-appropriate atlases for pediatric studies. *Neuroimage* 54, 313–327.
- Fox, K.C.R., Spreng, R.N., Ellamil, M., Andrews-Hanna, J.R., Christoff, K., 2015. The wandering brain: meta-analysis of functional neuroimaging studies of mind-wandering and related spontaneous thought processes. *Neuroimage* 111, 611–621.
- Gelman, A., Rubin, D.B., 1992. Inference from iterative simulation using multiple sequences. *Stat. Sci.* 457–472.
- Gold, J.I., Shadlen, M.N., 2007. The neural basis of decision making. *Annu. Rev. Neurosci.* 30, 535–574.
- Guan, M., Vandekerckhove, J., 2016. A bayesian approach to mitigation of publication bias. *Psychon. Bull. Rev.* 23, 74–86.
- Hawkins, G.E., Mittner, M., Boekel, W., Heathcote, A., Forstmann, B.U., 2015. Toward a model-based cognitive neuroscience of mind wandering. *Neuroscience* 310, 290–305.
- Hawkins, G.E., Mittner, M., Forstmann, B.U., Heathcote, A., 2017. On the efficiency of neurally-informed cognitive models to identify latent cognitive states. *J. Math. Psychol.* 76, 142–155.
- Heathcote, A., Lin, Y.S., Reynolds, A., Strickland, L., Gretton, M., Matzke, D., 2018. Dynamic models of choice. *Behav. Res. Methods* 1–25.
- JASP Team, 2017. JASP (Computer software), Version 0.8.1.2.
- Jeffreys, H., 1961. *Theory of Probability*. Oxford University Press.
- Karakaş, S., Erzen, Ö.U., Başar, E., 2000. The genesis of human event-related responses explained through the theory of oscillatory neural assemblies. *Neurosci. Lett.* 285, 45–48.
- Kass, R.E., Raftery, A.E., 1995. Bayes factors. *J. Am. Stat. Assoc.* 90, 773–795.
- Kayser, J., Tenke, C.E., 2003. Optimizing PCA methodology for ERP component identification and measurement: theoretical rationale and empirical evaluation. *Clin. Neurophysiol.* 114, 2307–2325.
- Kayser, J., Tenke, C.E., 2015. Issues and considerations for using the scalp surface Laplacian in EEG/ERP research: a tutorial review. *Int. J. Psychophysiol.* 97, 189–209.
- Kelly, S.P., O'Connell, R.G., 2015. The neural processes underlying perceptual decision making in humans: recent progress and future directions. *J. Physiol. Paris* 109, 27–37.
- Lamme, V.A.F., 1995. The neurophysiology of figure-ground segregation in primary visual cortex. *J. Neurosci.* 15, 1605–1615.
- Lamme, V.A.F., Roelfsema, P.R., 2000. The distinct modes of vision offered by feedforward and recurrent processing. *Trends Neurosci.* 23, 571–579.
- Lamme, V.A.F., Zipser, K., Spekreijse, H., 2002. Masking interrupts figure-ground signals in v1. *J. Cogn. Neurosci.* 14, 1044–1053.
- Lee, M.D., 2008. Three case studies in the Bayesian analysis of cognitive models. *Psychon. Bull. Rev.* 15, 1–15.
- Lee, M.D., Newell, B.J., 2011. Using hierarchical Bayesian methods to examine the tools of decision-making. *Judgment Decis. Making* 6.
- Lin, C.T., Chuang, C.H., Kerick, S., Mullen, T., Jung, T.P., Ko, L.W., Chen, S.A., King, J.T., McDowell, K., 2016. Mind-wandering tends to occur under low perceptual demands during driving. *Sci. Rep.* 6.
- Loughnane, G.M., Newman, D.P., Bellgrove, M.A., Lalor, E.C., Kelly, S.P., O'Connell, R.G., 2016. Target selection signals influence perceptual decisions by modulating the onset and rate of evidence accumulation. *Curr. Biol.* 26, 496–502.
- Luck, S.J., 2012. Event-related potentials. In: Cooper, H., Camic, P.M., Long, D.L., Panter, A.T., Rindskopf, D., Sher, K.J. (Eds.), *APA Handbook of Research Methods in Psychology: Volume 1, Foundations, Planning, Measures, and Psychometrics*. American Psychological Association.
- Luck, S.J., Chelazzi, L., Hillyard, S.A., Desimone, R., 1997. Neural mechanisms of spatial selective attention in areas v1, v2, and v4 of macaque visual cortex. *J. Neurophysiol.* 77, 24–42.
- Lui, K.K., Nunez, M.D., Cassidy, J.M., Vandekerckhove, J., Cramer, S.C., Srinivasan, R., 2018. Timing of Readiness Potentials Reflect a Decision-Making Process in the Human Brain. 338806.
- Makeig, S., Bell, A.J., Jung, T.P., Sejnowski, T.J., 1996. Independent component analysis of electroencephalographic data. *Adv. Neural Inf. Process. Syst.* 8, 145–151.
- Martin, T., Huxlin, K.R., Kavcic, V., 2010. Motion-onset visual evoked potentials predict performance during a global direction discrimination task. *Neuropsychologia* 48, 3563–3572.
- Miąsko, T., 2017. Pyjags [computer software], version 1.2.2. <https://github.com/tmias/kopyjags>.
- Mishkin, M., Ungerleider, L.G., Macko, K.A., 1983. Object vision and spatial vision: two cortical pathways. *Trends Neurosci.* 6, 414–417.
- Mittner, M., Hawkins, G.E., Boekel, W., Forstmann, B.U., 2016. A neural model of mind wandering. *Trends Cognit. Sci.* 20, 570–578.
- Mulder, M.J., van Maanen, L., Forstmann, B.U., 2014. Perceptual decision neurosciences a model-based review. *Neuroscience* 277, 872–884.
- Navarro, D.J., Fuss, I.G., 2009. Fast and accurate calculations for first-passage times in Wiener diffusion models. *J. Math. Psychol.* 53, 222–230.
- Neyman, J., Pearson, E.S., 1933. On the problem of the most efficient tests of statistical hypotheses. *Philos. Trans. R. Soc. Lond. - Ser. A Contain. Pap. a Math. or Phys. Character* 231, 289–337.
- Nunez, M.D., 2017. Refining Understanding of Human Decision Making by Testing Integrated Neurocognitive Models of EEG, Choice and Reaction Time. Ph.D. diss. University of California, Irvine.
- Nunez, M.D., Horton, C., Deng, S., Winter, W., Srinivasan, R., 2017a. *artscreenEEG [computer software], version 0.3.0*. <https://github.com/mdnunez/artscreenEEG>.
- Nunez, M.D., Nunez, P.L., Srinivasan, R., 2016. Electroencephalography (EEG): neurophysics, experimental methods, and signal processing. In: Ombao, H., Linquist, M., Thompson, W., Aston, J. (Eds.), *Handbook of Neuroimaging Data Analysis*. Chapman & Hall/CRC, pp. 175–197.
- Nunez, M.D., Srinivasan, R., Vandekerckhove, J., 2015. Individual differences in attention influence perceptual decision making. *Front. Psychol.* 6, 18.
- Nunez, M.D., Vandekerckhove, J., Srinivasan, R., 2017b. How attention influences perceptual decision making: single-trial EEG correlates of drift-diffusion model parameters. *J. Math. Psychol.* 76, 117–130.
- Nunez, P.L., Nunez, M.D., Srinivasan, R., 2019. Multi-scale neural sources of EEG: genuine, equivalent, and representative. A tutorial review. *Brain Topogr.* 32 (2), 193–214.
- Nunez, P.L., Pilgreen, K.L., 1991. The spline-Laplacian in clinical neurophysiology: a method to improve EEG spatial resolution. *J. Clin. Neurophysiol.* 8, 397–413.
- Nunez, P.L., Silberstein, R.B., Cadushe, P.J., Wijesinghe, R.S., Westdorp, A.F., Srinivasan, R., 1994. A theoretical and experimental study of high resolution EEG based on surface Laplacians and cortical imaging. *Electroencephalogr. Clin. Neurophysiol.* 90, 40–57.
- Nunez, P.L., Srinivasan, R., 2006. *Electric Fields of the Brain: the Neurophysics of EEG*. Oxford University Press.
- O'Connell, R.G., Dockree, P.M., Kelly, S.P., 2012. A supramodal accumulation-to-bound signal that determines perceptual decisions in humans. *Nat. Neurosci.* 15, 1729–1735.
- Parra, L.C., Spence, C.D., Gerson, A.D., Sajda, P., 2005. Recipes for the linear analysis of EEG. *Neuroimage* 28, 326–341.
- Philiastides, M.G., Heekeren, H.R., Sajda, P., 2014. Human scalp potentials reflect a mixture of decision-related signals during perceptual choices. *J. Neurosci.* 34, 16877–16889.
- Plummer, M., 2003. JAGS: a program for analysis of Bayesian graphical models using Gibbs sampling. In: *Proceedings of the 3rd International Workshop on Distributed Statistical Computing*. DSC 2003, Vienna, Austria.
- Pommier, J., Renard, Y., 2005. Getfem++, an Open Source Generic C++ Library for Finite Element Methods.
- Ratcliff, R., 1978. A theory of memory retrieval. *Psychol. Rev.* 85, 59.
- Ratcliff, R., McKoon, G., 2008. The diffusion decision model: theory and data for two-choice decision tasks. *Neural Comput.* 20, 873–922.
- Ratcliff, R., Tuerlinckx, F., 2002. Estimating parameters of the diffusion model: approaches to dealing with contaminant reaction times and parameter variability. *Psychon. Bull. Rev.* 9, 438–481.
- Roitman, J.D., Shadlen, M.N., 2002. Response of neurons in the lateral intraparietal area during a combined visual discrimination reaction time task. *J. Neurosci.* 22, 9475–9489.
- Rosenthal, R., 1979. The file drawer problem and tolerance for null results. *Psychol. Bull.* 86, 638.
- Rouder, J.N., Morey, R.D., 2012. Default bayes factors for model selection in regression. *Multivariate Behav. Res.* 47, 877–903.
- Schmolesky, M.T., Wang, Y., Hanes, D.P., Thompson, K.G., Leutgeb, S., Schall, J.D., Leventhal, A.G., 1998. Signal timing across the macaque visual system. *J. Neurophysiol.* 79, 3272–3278.
- Schubert, A.L., Hagemann, D., Frischkorn, G.T., 2017. Is general intelligence little more than the speed of higher-order processing? *J. Exp. Psychol. Gen.* 146, 1498.
- Schubert, A.L., Nunez, M.D., Hagemann, D., Vandekerckhove, J., 2018. Individual differences in cortical processing speed predict cognitive abilities: a model-based cognitive neuroscience account. *Comput. Brain Behav.* 374827.
- Servant, M., White, C., Montagnini, A., Burle, B., 2015. Using covert response activation to test latent assumptions of formal decision-making models in humans. *J. Neurosci.* 35, 10371–10385.
- Servant, M., White, C., Montagnini, A., Burle, B., 2016. Linking theoretical decision-making mechanisms in the simon task with electrophysiological data: a model-based neuroscience study in humans. *J. Cogn. Neurosci.* 28, 1501–1521.
- Shadlen, M.N., Kiani, R., 2013. Decision making as a window on cognition. *Neuron* 80, 791–806.
- Stanford, T.R., Shankar, S., Massoglia, D.P., Costello, M.G., Salinas, E., 2010. Perceptual decision making in less than 30 milliseconds. *Nat. Neurosci.* 13, 379–385.
- Sterling, T.D., 1959. Publication decisions and their possible effects on inferences drawn from tests of significance or vice versa. *J. Am. Stat. Assoc.* 54, 30–34.
- Straube, S., Grimsen, C., Fahle, M., 2010. Electrophysiological correlates of figure-ground segregation directly reflect perceptual saliency. *Vis. Res.* 50, 509–521.
- Thorpe, S., Fize, D., Marlot, C., 1996. Speed of processing in the human visual system. *Nature* 381, 520.

- Tuerlinckx, F., 2004. The efficient computation of the cumulative distribution and probability density functions in the diffusion model. *Behav. Res. Methods Instrum. Comput.* 36, 702–716.
- Turner, B.M., Forstmann, B.U., Love, B.C., Palmeri, T.J., Van Maanen, L., 2017. Approaches to analysis in model-based cognitive neuroscience. *J. Math. Psychol.* 76, 65–79.
- Turner, B.M., van Maanen, L., Forstmann, B.U., 2015. Informing cognitive abstractions through neuroimaging: the neural drift diffusion model. *Psychol. Rev.* 122, 312–336.
- van Vugt, M.K., Taatgen, N.A., Sackur, J., Bastian, M., 2015. Modeling mind-wandering: a tool to better understand distraction. In: *Proceedings of the 13th International Conference on Cognitive Modeling*. University of Groningen, pp. 252–257.
- Vandekerckhove, J., Tuerlinckx, F., 2008. Diffusion model analysis with MATLAB: a DMAT primer. *Behav. Res. Methods* 40, 61–72.
- Vandekerckhove, J., Tuerlinckx, F., Lee, M.D., 2011. Hierarchical diffusion models for two-choice response times. *Psychol. Methods* 16, 44.
- VanRullen, R., 2011. Four common conceptual fallacies in mapping the time course of recognition. *Front. Psychol.* 2, 365.
- VanRullen, R., Thorpe, S.J., 2001. The time course of visual processing: from early perception to decision-making. *J. Cogn. Neurosci.* 13, 454–461.
- Verdinelli, I., Wasserman, L., 1995. Computing Bayes factors using a generalization of the savage-dickey density ratio. *J. Am. Stat. Assoc.* 90, 614–618, 00422.
- Vogel, E.K., Luck, S.J., 2000. The visual N1 component as an index of a discrimination process. *Psychophysiology* 37, 190–203.
- Voss, A., Rothermund, K., Voss, J., 2004. Interpreting the parameters of the diffusion model: an empirical validation. *Mem. Cognit.* 32, 1206–1220.
- Wabersich, D., Vandekerckhove, J., 2014. Extending JAGS: a tutorial on adding custom distributions to JAGS (with a diffusion model example). *Behav. Res. Methods* 46, 15–28.
- Wagenmakers, E.J., 2009. Methodological and empirical developments for the Ratcliff diffusion model of response times and accuracy. *Eur. J. Cogn. Psychol.* 21, 641–671.
- Wagenmakers, E.J., Lodewyckx, T., Kuriyal, H., Grasman, R., 2010. Bayesian hypothesis testing for psychologists: a tutorial on the savage–dickey method. *Cogn. Psychol.* 60, 158–189.
- Webster, M.A., De Valois, R.L., et al., 1985. Relationship between spatial-frequency and orientation tuning of striate-cortex cells. *JOSA A* 2, 1124–1132.
- Wiecki, T.V., Sofer, I., Frank, M.J., 2013. HDDM: hierarchical bayesian estimation of the drift-diffusion model in python. *Front. Neuroinf.* 7, 14.
- Zhang, Q., van Vugt, M., Borst, J.P., Anderson, J.R., 2018. Mapping working memory retrieval in space and in time: a combined electroencephalography and electrocorticography approach. *Neuroimage* 174, 472–484.
- Zhang, Q., Walsh, M.M., Anderson, J.R., 2017. The effects of probe similarity on retrieval and comparison processes in associative recognition. *J. Cogn. Neurosci.* 29 (2), 352–367.