



## Review

## Cellular senescence in cardiac diseases

Ippei Shimizu (MD, PhD)<sup>a,b,\*</sup>, Tohru Minamino (MD, PhD, FJCC)<sup>a,\*</sup><sup>a</sup> Department of Cardiovascular Biology and Medicine, Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan<sup>b</sup> Division of Molecular Aging and Cell Biology, Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan

## ARTICLE INFO

## Article history:

Received 20 May 2019

Accepted 20 May 2019

Available online 12 June 2019

## Keywords:

Cellular senescence

Heart failure

Senolysis

## ABSTRACT

Replicative capacity of somatic cells is limited. It indicates that aging also develops at the cellular level, and this is described as “cellular senescence”. Senescent cells become flattened, enlarged, and irreversibly lose capacity for proliferation. Lack of specific and conclusive markers for cellular senescence makes it difficult to comprehensively define and understand this biological process especially *in vivo*. Molecules including p53, p21, p16<sup>Ink4a</sup>, p38MAPK, and  $\gamma$ H2AX, telomere attrition, enhanced signals for SA- $\beta$ -gal, etc. are widely used to detect senescent cells, but these are indirect indicators of cellular senescence, and biological markers reflecting direct evidence need to be established. Genetic profiles are altered in senescent cells, letting these cells secrete pro-inflammatory molecules. Aging or age-related disorders including heart failure and atherosclerotic diseases link with an accumulation of cells undergoing cellular senescence in cardiovascular systems including heart and vessels. Senescent cells become pathogenic in most cases by mediating chronic sterile inflammation and tissue remodeling. A recent conceptual as well as technical breakthrough in this research area is “senolysis”, meaning the specific elimination of senescent cells. Genetic as well as pharmacological models with senolysis contributed to reverse aging phenotypes and ameliorated pathologies in age-related disorders without enhancing the risk of tumorigenesis, and opened a new avenue for aging research. Several compounds are identified as senolytics, and some are already tested in clinical settings. It was recently reported that senolysis reverses aging phenotype in cardiovascular disorders. Generating therapies targeting suppression or elimination of senescent cells would inhibit the progression of undesirable aspects of aging, and become promising therapies for cardiac diseases.

© 2019 The Authors. ISDN. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Contents

Introduction	314
Molecular mechanisms of cellular senescence	314
Markers of senescent cells	314
Roles of cellular senescence related molecules in cardiac disease	314
Cardiomyocytes	315
Endothelial cells	315
Fibroblasts	316
Progenitor cells	316
Senolytics targeting cardiac diseases	316
Conclusion and future directions	317
References	317

\* Corresponding authors at: Department of Cardiovascular Biology and Medicine, Niigata University Graduate School of Medical and Dental Sciences, 1-757 Asahimachidori, Chuo-ku, Niigata 951-8510, Japan.

E-mail addresses: [ippeishimizu@med.niigata-u.ac.jp](mailto:ippeishimizu@med.niigata-u.ac.jp), [ippeishimizu@yahoo.co.jp](mailto:ippeishimizu@yahoo.co.jp) (I. Shimizu), [t\\_minamino@yahoo.co.jp](mailto:t_minamino@yahoo.co.jp), [tminamino@med.niigata-u.ac.jp](mailto:tminamino@med.niigata-u.ac.jp) (T. Minamino).

## Introduction

Chronological aging is characterized by changes of intercellular communication, genomic instability, epigenetic alterations, telomere attrition, loss of proteostasis, mitochondrial dysfunction, stem cell exhaustion, deregulation of nutrient sensing, and cellular senescence [1]. All these contribute to deterioration at organ, cellular, and molecular levels, leading to a decline of physiological activity, leaving organisms prone to death. Aging *per se* is a physiological process, however, it associates with several undesirable aspects, and predisposes us to age-related diseases. Aging and age-related disorders progress through an integration of complex biological processes, and do not allow a simple approach to understand the whole picture, however, evidence indicates the central roles of cellular senescence in the pathogenesis of these conditions. Prevalence of age-associated diseases, including atherosclerotic disorders or heart failure increases with chronological aging, and cells positive for senescent markers are now well recognized to have causal roles for the progression of pathologies in these age-related diseases [2–4]. *In vitro* studies showed that exposure of young somatic cells to senescent cells promotes senescence of the young cells, and this is described as the “bystander effect” [5]. Pharmacological or genetic depletion of senescent cells contributed to reverse aging phenotype, and suppressed pathologies in chronological as well as age-related disease models [6–8]. In this review article, we would like to delineate the role of cellular senescence and the related molecules in cardiac diseases, and discuss the potential of therapies targeting senescent cells.

## Molecular mechanisms of cellular senescence

It was previously reported that replicative capacity of fibroblasts was limited [9]. This indicated that aging also develops at the cellular level, and now this is described as “cellular senescence”. Senescent cells become enlarged with a flattened morphology, and show irreversible loss of proliferative potential. Alteration in the expression of genetic profiles in these cells leads to secretion of pro-inflammatory molecules (known as the senescence-associated secretory phenotype (SASP)) [10]. Senescent cells accumulate in various organs of animal models or humans who are elderly and/or have age-related disorders, and thought to become pathogenic through introducing chronic sterile inflammation and tissue remodeling. There are two types of cellular senescence. As Hayflick et al. [9] previously reported, somatic cells have a limited proliferative capacity and eventually enter a state of irreversible growth arrest termed “replicative senescence.” Telomeres locate at each end of chromosome, and characterized as a region of repetitive nucleotide sequences, AGGGTT in vertebrates. Telomeres replicate incompletely during cell division, resulting in telomere attrition. Telomeres have critical roles for chromosomal stability and DNA replication. For this reason, when telomere shortening exceeds the physiological range, it is recognized as DNA damage and triggers replicative cellular senescence, mainly via the p53 or p16<sup>Ink4a</sup> signaling pathways. Another type of cellular senescence is described as “stress-induced premature senescence”. Premature senescence develops through various external as well as internal stress signals, and these include oxidative stress or irradiation, constitutive activation of mitogenic stimuli, oncogenic activation, and metabolic stress. The p53 or p16<sup>Ink4a</sup> signaling pathways also mediate this type of cellular senescence. Preference for p53 or p16<sup>Ink4a</sup> signaling pathway varies among species, and depends on the cell types [11,12]. In human cells, telomere dysfunction activates either p53 or p16<sup>Ink4a</sup> signaling, however, in rodent cells p53 signaling seems to be the only activated pathway [13]. p53 signaling is primarily triggered


for activation by DNA damage and telomere dysfunction, while p16<sup>Ink4a</sup> signaling is primarily linked to mitogenic stress, and general cellular stress [11,12,14]. It is now well recognized that the vast majority of stressors that induce cellular senescence activate either or both the p53/p21 or p16<sup>Ink4a</sup>/retinoblastoma protein pathways [4].

## Markers of senescent cells

As already described, most senescence-inducing stressors activate either the p53/p21 or p16<sup>Ink4a</sup>/retinoblastoma protein pathways, however, it needs to be noted that activation of these signaling pathways does not provide conclusive evidence that cells are senescent [15]. Some proliferating cancer cells express high levels of p16<sup>Ink4a</sup> [16]. p21 is also involved in “transient” cell cycle arrest in response to acute cell damage [17]. Phenotype of senescent cells has multiple variations, and cellular senescence needs to be defined through various combinations of markers. To date, several markers are used to detect senescent cells indirectly. Senescence-associated beta-galactosidase (SA-β-gal) activity at pH6 is broadly used to recognize cells as senescent. However, we here note that SA-β-gal activity is also known to increase in fibroblasts cultured under serum starvation [18]. In addition to p53, p21, and p16<sup>Ink4a</sup>, other well-known markers of cellular senescence are high levels of p38 mitogen-activated protein kinase (p38MAPK) or γH2AX, reflecting the activation of DNA damage responses. Senescence-associated heterochromatin foci (SAHF), and senescence-associated distention of satellites (SADs) are also considered to be indicators of cellular senescence [19,20]. It was previously shown that irreparable telomere damage develops in the absence of telomere attrition. Presence of DNA damage at telomeres were described as telomere-associated foci (TAF), and detected as colocalization of γH2AX, p53-binding protein 1 (53BP1) with telomeres. In fibroblasts, irradiation or H<sub>2</sub>O<sub>2</sub> administration increased TAF, and this was also shown to increase in liver, gut, and heart with aging [6,21]. Exploring biological markers reflecting direct evidence of cellular senescence continues to be an important research topic (Fig. 1).

## Roles of cellular senescence related molecules in cardiac disease

Accumulation of evidence indicates the close connection between pathways involved in cellular senescence with cardiovascular disorders. In humans with end-stage heart failure, p53 level and apoptosis were reported to increase in myocardium [22]. Patients with hypertrophic cardiomyopathy or dilated cardiomyopathy showed higher p53 expression in heart compared to non-failing heart [23,24]. In a murine left ventricular pressure overload model, cardiac-p53 level was increased, and this led to suppression of angiogenesis in the myocardium, promoting tissue hypoxia and cardiac dysfunction [25]. Shorter peripheral blood leukocyte telomere length (LTL) predicted adverse cardiovascular events in coronary artery disease patients [26,27], and recently it was shown that endurance training and interval training, but not resistance training, increased telomerase activity and telomere length in circulating leukocytes [28]. Insulin-like growth factor-binding protein-7 (IGFBP7) inhibits cell proliferation through G1 phase cell cycle arrest, and is considered as a protein member of senescence secretomes. IGFBP7 associates with tissue aging and obesity, and high circulating IGFBP7 was reported to link with poor prognosis in patients with heart failure with preserved ejection fraction (HFpEF) [29]. Interestingly, it was recently reported that neprilysin inhibition reduced circulating levels of this pro-senescent molecule [30]. Several representative cells including cardiomyocytes, endothelial cells, fibroblasts, and progenitor cells



**Fig. 1.** Markers of senescent cells. Indirect markers characterizing cellular senescence are as follows, and these vary among tissues and cells. (1) Enlarged and flat morphology, (2) cell cycle arrest (p53/p21, p16<sup>Ink4a</sup> signals become high), (3) DNA damage response ( $\gamma$ H2AX, p38 MAPK signals become high), (4) irreversible termination of proliferation, (5) high senescence-associated beta-galactosidase (SA- $\beta$ -gal) activity, (6) telomere attrition, (7) telomere dysfunction, (8) senescence-associated distention of satellites (SADS), and (9) senescence-associated heterochromatin foci (SAHF).

are involved in the maintenance of cardiac homeostasis, and roles of cellular senescence-related molecules in these cells are discussed next.

### Cardiomyocytes

Cellular senescence was traditionally recognized as a machinery to inhibit uncontrolled replication in proliferative cells, but nowadays it is thought that post-mitotic cells such as Purkinje neurons also develop a senescent-like phenotype. For example, murine cortical and Purkinje neurons were reported to exhibit an increase in  $\gamma$ H2AX, IL-6, 4-HNE, p38MAPK, and SA- $\beta$ -gal levels with aging [31]. Shortly after birth, cardiomyocytes exhibit cell cycle arrest due to the activation of DNA damage response introduced through exposure to higher concentration of oxygen in the postnatal environment [32]. Adult cardiomyocytes were long thought to be terminally differentiated post-mitotic cells, however, accumulating evidence indicates these cells retain proliferative capacity. It was previously reported that in humans, cardiomyocyte turnover was at a rate of <1% per year [33], and this was demonstrated to decline with aging both in humans and mice [34,35]. The underlying machinery of diminished cardiomyocyte turnover with aging is yet to be defined, and whether this is attributable to cardiomyocyte senescence is not clear due to the lack of specific senescence markers. In mice, it was reported that chronological aging links with an increase in cardiomyocyte size, together with reactive oxygen species (ROS) production, telomere attrition, and high level of p53 or p16<sup>Ink4a</sup> expression. In this study, compared to young mice (4 months of age), aged mice (20–22 months of age) exhibited increased left ventricular weight and cardiomyocyte volume, and showed reduction in cardiomyocyte number, together with reduced ventricular function, indicating the pathological roles of cardiomyocyte senescence in the aged heart [36]. In cardiomyocytes, toxic reagents such as doxorubicin administration was shown to upregulate p16<sup>Ink4a</sup>, p21 and SA- $\beta$ -gal level [37]. In humans, patients with heart failure demonstrated shorter cardiomyocyte telomeres compared with nonfailing donors. Hypertrophic cardiomyopathy (HCM) patients with reduced systolic function exhibited cardiomyocytes with shortest telomeres in association with resident DNA damage (as analyzed with  $\gamma$ H2AX or 53BP1 staining), and patients with ischemic cardiomyopathy (ICM) also showed shorter telomere length. Cardiomyocytes from idiopathic cardiomyopathy (IDCM) patients exhibited a trend to reduced telomere length, but this did not reach statistical significance. Interestingly, telomere attrition did not develop in vascular smooth muscle cells from failing hearts

of HCM, ICM, and IDCM patients, and this was comparable with non-failing donors. In this study, it was also shown that telomere shortening in cardiomyocytes did not develop with aging in non-failing donors [38]. Whether cardiomyocyte senescence exists *in vivo* was controversial, but recently, it was reported that length-independent telomere damage drives post-mitotic cardiomyocyte senescence [6]. In a chronologically aged murine model, TAF positive cardiomyocytes increased with aging (comparison of mice 3, 15, and 30 months of age). Interestingly,  $\gamma$ H2AX signal did not increase in cardiomyocytes, but increased only in telomeres. Irradiation increased TAF level in mouse embryonic cardiomyocytes, rat neonatal cardiomyocytes, and H9C2 myoblasts. In cardiomyocytes collected through Langendorff heart perfusion, transcripts p16<sup>Ink4a</sup> and p21 were increased, and staining studies showed p21 and SA- $\beta$ -gal signal increased with aging in cardiomyocytes in the heart. In right atrial appendage of humans, TAF signal also increased in cardiomyocytes in individuals aged >70 years, compared to <70 years of age. In this study, senescent cardiomyocytes were shown to contribute for non-canonical senescence-associated secretory phenotype, and promoted myofibroblast differentiation in fibroblasts and cardiomyocyte hypertrophy in mice [6]. Presence of senescence-like phenotype in cardiomyocytes was studied in atrium in humans, and whether this develops in human left ventricle remains an open question to be explored. Recently, an interesting paper showed the bidirectional role of p53 in cardiomyocytes. Mak et al. generated cardiomyocyte specific p53 knockout model by crossing MHC-MerCreMer<sup>+/-</sup> with p53<sup>fllox/fllox</sup> mice (here we describe as “CM-p53KO”). CM-p53 KO mice showed enhanced cardiac as well as cardiomyocyte hypertrophy, and this associated with reduced cardiac systolic dysfunction when these mice became 6 months-old. In contrast, CM-p53 KO exhibited improved cardiac function together with reduced cardiac as well as cardiomyocyte hypertrophy in left ventricular pressure overload model. These results indicated that p53 signal in cardiomyocytes contributed for inhibition of pathological cardiac hypertrophy in aging model, but augmented this in left ventricular pressure overload, showing that the roles of p53 in cardiomyocytes is context dependent [39].

### Endothelial cells

Endothelial cells in aged-sedentary individuals (around 60 years old) were reported to express higher protein level of p53, p21, and p16<sup>Ink4a</sup>, and this was also shown to reduce in aged exercising adults (around 57 years old) [40]. Preclinical studies indicated the pathological roles of endothelial cell-senescence in age-related

disorders including obesity, diabetes, and heart failure [41,42]. Heart failure can be characterized into two types depending on the level of systolic function. One is described as heart failure with reduced ejection fraction (HFrEF), another is classified as HFpEF, and both types of heart failure are prevalent among elderly persons [43]. In the failing heart, chronic sterile inflammation develops, and this is well recognized to promote cardiac remodeling [44]. In a murine left ventricular (LV) pressure overload model, activation of p53 signaling in vascular endothelial cells was reported to induce cardiac inflammation and remodeling [41]. In response to LV-pressure overload, p53 level increased in endothelial cells of capillaries in left ventricle, and had causal roles for the elevated expression of intercellular adhesion molecule (ICAM)-1 by these cells. This contributed to promoting inflammation in the failing heart in association with reduced systolic function. Another study showed depletion of p53 in endothelial cells ameliorated capillary rarefaction, improved cardiac function, and suppressed cardiac fibrosis and remodeling [45]. HFpEF occurs in approximately half of all patients with heart failure. Inflammation in coronary microvasculature is now thought to have central roles in the pathogenesis of HFpEF [46], and it was recently indicated that cellular senescence in endothelial cells may also be involved. When senescence-accelerated mice were fed a high-fat high-salt diet, endothelial cell senescence developed in cardiac tissues, and this coincided with the typical hemodynamic and structural changes of HFpEF [47]. Administration of palmitate, representing a metabolic stress, led to Sirt1 inactivation and increased senescent markers including p53, p16<sup>Ink4a</sup>, and SA- $\beta$ -gal in HUVECs [48]. Given that aged and/or obese population has higher prevalence for HFpEF, inhibition of endothelial cell senescence pathway may become a next generation therapy for this untreatable disorder.

#### Fibroblasts

Fibroblasts are considered to represent chief cardiac cell component, accounting for 27% of all cardiac cells in mice and 64% in rats. In humans, non-cardiomyocytes (mainly fibroblasts) are reported to account for 72% of total cells [49,50]. Studies indicate fibroblasts contribute for maintaining homeostasis in heart under physiological as well as stressed conditions. Aging was reported to associate with the presence of fibroblasts containing X-Gal crystals in pericardium [7]. During aging, visceral adipose tissue-derived osteopontin increased in plasma. Systemic genetic depletion of osteopontin, visceral adipose tissue depletion, or pharmacological osteopontin inhibition led to a significant increase in p16<sup>Ink4a</sup> level in cardiac fibroblasts, and contributed to the suppression of age-induced cardiac fibrosis. This indicated that osteopontin inhibited cellular senescence in cardiac fibroblasts, having causal roles for the augmentation of fibrosis in cardiac tissue. This showed fibroblast senescence associates with healthy cardiac aging [51]. Another study reported that senescence-associated genes including p53, p21, and ataxia telangiectasia mutated (ATM) increased after myocardial infarction in both mouse and human hearts. Heterozygous ATM systemic depletion resulted in reduced SA- $\beta$ -Gal or p53 signal in cardiac fibroblasts. Suppression of these senescence markers in fibroblasts contributed to enhance fibrosis in non-infarct area (this was comparable between genotypes in infarct area), and reduced systolic function in myocardial infarction model [52]. These preclinical studies indicate that inhibition of cellular senescence in fibroblasts becomes pathological in cardiac tissues. Whether this would be the case in humans is a critical question, and raises potential issues for therapies based on non-targeted delivery of senescence-suppressants which may be developed in the near future.


#### Progenitor cells

Cardiosphere-derived cells (CDCs) are progenitor cells in cardiac tissue and have a potential to differentiate into three types of cells present in this organ. These are cardiomyocytes, endothelial cells, and smooth muscle cells. CDCs were reported to prevent adverse cardiac remodeling, characterized as reduced fibrosis, ROS, and inflammation. This contributed to suppress the progression of systolic dysfunction, and improved survival in dilated cardiomyopathy mouse model [53]. CDCs were also reported to reverse HFpEF in Dahl salt-sensitive rats by suppression of fibrosis and inflammation in heart [54]. Direct injection of CDCs extracted from young Fisher rats (around 4 months old) into LV cavity of old Fisher rats (around 22 months old) suppressed cardiac fibrosis, improved diastolic function, ameliorated LV hypertrophy in aged rats. In this paper, *in vitro* studies showed young CDCs, extracted from human donors <2 years of age, were reported to secrete exosomes (CDC-XO), contributing to the elongation of telomeres, together with suppression of SA- $\beta$ -gal signal in CDCs obtained from >55-year old human donors [55]. Cardiac progenitor cells (CPCs) are another type of progenitor cell in the heart [56]. More than half of cardiac CPCs, defined as Sca-1<sup>pos</sup>/c-kit<sup>pos</sup>/CD31<sup>neg</sup>/CD45<sup>neg</sup>/Tryptase<sup>neg</sup>, isolated from aged (>70 years old) subjects showed higher levels of p16<sup>Ink4a</sup>, SA- $\beta$ -gal, and  $\gamma$ H2AX, had telomere attrition, and obtained SASP phenotype. Direct injection of senescent CPCs into left ventricle, led to reduced reparative potential in a murine myocardial infarction model compared to the introduction of non-senescent CPCs injection. This indicated aged human heart contains high proportion of senescent-CPCs, possibly contributing to the progression of cardiac dysfunction [57]. In castaneous (CAST) mice bearing short telomeres from birth, CPCs (defined as c-kit<sup>pos</sup>) were also shown to develop telomere attrition. CAST mice exhibited activation in p53 and autophagy, and promoted differentiation, senescence, and exhaustion of CPCs [58].

Bone marrow-derived progenitor cells are another type of cell reported to contribute to tissue repair in the failing heart [59]. In ischemic heart disease patients, bone marrow-derived progenitor cells had shorter telomere length, higher level of transcripts for p21 and p16<sup>Ink4a</sup>, and these associated with reduced myeloid differentiation capacity. This indicated the functional decline of BM-derived progenitor cells develops in ischemic heart disease [60]. Evidence shows that suppression of cellular senescence in progenitor cells is critical for maintaining homeostasis in cardiac tissues under stressed conditions.

#### Senolytics targeting cardiac diseases

Genetic or pharmacological models targeting the selective depletion of senescent cells (“senolysis”) were shown to reverse age-related pathological changes [7,61–64]. Aging associates with the accumulation of senescent cells in organs, and these are well accepted to mediate their pathogenic roles by promoting chronic sterile inflammation in obesity, diabetes, heart failure, or atherosclerotic models. INK-ATTAC mice were generated to enable researchers to produce the inducible elimination of p16<sup>Ink4a</sup>-expressing cells by the administration of a compound named AP20187. In a premature aging mouse model, elimination of p16<sup>Ink4a</sup>-positive senescent cells led to inhibition of aging phenotype in white adipose tissues, heart, and kidney. In cardiac tissue at 18 months, elimination of p16<sup>Ink4a</sup>-positive cells did not change heart rate, LV mass, thickness and diameter, ejection fraction, and fractional shortening compared to controls. However, cardiomyocyte cross-sectional area became significantly reduced with senolysis, suggesting that number of cardiomyocytes were increased in this group. Depletion of p16<sup>Ink4a</sup>-positive cells



**Fig. 2.** Emerging concept of therapies targeting failing hearts with senolysis. Rejuvenation of senescent-like cells by senolysis contributes to convert failing heart into healthy heart.

utilizing INK-ATTAC genetic system also contributed to enhanced capability of handling isoproterenol-induced stress, indicating that accumulation of p16<sup>Ink4a</sup>-positive cells has pathogenic roles to accelerate cardiac aging [7]. In another genetic model, p16<sup>Ink4a</sup>-3MR (3MR) mouse, a transgenic mouse that kills p16<sup>Ink4a</sup> positive cells upon administration of ganciclovir (GCV), elimination of p16<sup>Ink4a</sup> positive cells from plaques led to the regression of atherosclerotic lesions in low-density lipoprotein receptor-deficient mice [8]. Nowadays, several compounds are reported to mediate senolysis, and these are described as “senolytics” (reviewed in Katsuomi et al. [65]). Among them, dasatinib and quercetin (D + Q), and an anticancer agent (ABT263) are extensively tested. ABT263 was reported to have a senolytic effect, contributing to the selective clearance of p16<sup>Ink4a</sup>-positive senescent cells in bone marrow by inducing apoptosis in these cells. This led to rejuvenation of hematopoietic stem cells during aging [62]. D + Q administration improved vasomotor function in aged mice, and this was associated with reduced aortic calcification and osteogenic signaling in a murine model with hypercholesterolemia [66]. Recently, therapeutic potential of senolysis was also tested in cardiac tissues. D + Q administration significantly improved systolic cardiac function, together with reduced end-systolic left ventricular dimension in 24-month-old mice [64]. Global elimination of senescent cells in aged mice in pharmacological (D + Q) or genetic (INK-ATTAC) senolytic models led to the activation of resident CPCs in aged heart, and increased number of proliferative cardiomyocytes [57]. In another report, depletion of p16<sup>Ink4a</sup> positive cells in 27-month-old INK-ATTAC mice resulted in reduction of TAF positive cardiomyocytes, decline in cardiomyocyte size, and inhibited fibrotic area in heart. In a pharmacological senolytic model, ABT263 administration also reproduced these findings as observed in INK-ATTAC mice, and in both models, cardiomyocytes became positive for proliferative markers [6]. Another report also showed ABT263 administration eliminated senescent cardiomyocytes and attenuated profibrotic protein expression in aged mice. In this study, authors also tested roles of senolytic compound in myocardial infarction model, and showed clearance of senescent cells led to amelioration of myocardial remodeling, diastolic function, and improved survival rate in this ischemic heart failure model [67]. These indicated that elimination of senescent cells has a potential to reverse age-related dysfunction in cardiac tissues through promotion of regenerative capacity in this organ. Selective elimination of senescent cells with senolytics would become a next generation therapy for cardiac disorders [57,65,67–69] (Fig. 2).

## Conclusion and future directions

In this review article, we delineated the role of cellular senescence and related molecules in cardiac tissue. In addition to molecules including p53, p21, p16<sup>Ink4a</sup>, p38MAPK, and  $\gamma$ H2AX, telomere attrition, enhanced signals for SA- $\beta$ -gal, SAHF, TAF, SADs are also considered to be indicators of cellular senescence. Due to the lack of specific and conclusive senescence markers, it is especially challenging to conclude the presence of classically defined cellular senescent cells in organs. In many cases, if we follow traditional definitions, it would be appropriate to describe cells showing senescent markers as “senescent-like cells”, and studies exploring conclusive senescent markers continue to be an important research topic. We also need to note that pathological implications of cellular senescence are context dependent. Endothelial cell senescence seems to be always detrimental in age-related disorders including heart failure or obesity. However, suppression of cellular senescence in fibroblasts was reported to promote cardiac fibrosis and remodeling through the activation of myofibroblasts. Considering that endothelial cells and fibroblasts share similar pro-senescent stressors including saturated fatty acids, outcomes of therapies targeting the inhibition of cellular senescence would be obtained through the summation of complex biological processes. In addition to the consideration for the concern of tumorigenesis, it would be important to generate tissue and/or cell specific anti-senescence therapy. Senolysis, *i.e.* targeted elimination of senescent cells, is now highlighted worldwide, because this approach was shown to reverse pathological features of aging in genetic as well as pharmacological models. Importantly, so far, this therapy does not seem to promote tumorigenesis. In addition to careful observation to find possible side effects involved, it would also be critically important to establish cell-specific senolytic therapy for age-related cardiac diseases.

## References

- [1] Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell* 2013;153(6):1194–217.
- [2] Childs BG, Durik M, Baker DJ, van Deursen JM. Cellular senescence in aging and age-related disease: from mechanisms to therapy. *Nat Med* 2015;21(12):1424–35.
- [3] Shimizu I, Yoshida Y, Suda M, Minamino T. DNA damage response and metabolic disease. *Cell Metab* 2014;20(6):967–77.
- [4] van Deursen JM. The role of senescent cells in ageing. *Nature* 2014;509(7501):439–46.
- [5] Nelson G, Wordsworth J, Wang C, Jurk D, Lawless C, Martin-Ruiz C, et al. A senescent cell bystander effect: senescence-induced senescence. *Aging Cell* 2012;11(2):345–9.

- [6] Anderson R, Lagnado A, Maggiorani D, Walaszczyk A, Dookun E, Chapman J, et al. Length-independent telomere damage drives post-mitotic cardiomyocyte senescence. *EMBO J* 2019;38(5).
- [7] Baker DJ, Childs BG, Durik M, Wijers ME, Sieben CJ, Zhong J, et al. Naturally occurring p16(Ink4a)-positive cells shorten healthy lifespan. *Nature* 2016;530(7589):184–9.
- [8] Childs BG, Baker DJ, Wijshake T, Conover CA, Campisi J, van Deursen JM. Senescent intimal foam cells are deleterious at all stages of atherosclerosis. *Science* 2016;354(6311):472–7.
- [9] Hayflick L, Moorhead PS. The serial cultivation of human diploid cell strains. *Exp Cell Res* 1961;25:585–621.
- [10] Tchkonja T, Zhu Y, van Deursen J, Campisi J, Kirkland JL. Cellular senescence and the senescent secretory phenotype: therapeutic opportunities. *J Clin Invest* 2013;123(3):966–72.
- [11] Campisi J. Senescent cells, tumor suppression, and organismal aging: good citizens, bad neighbors. *Cell* 2005;120(4):513–22.
- [12] Beausejour CM, Krtolica A, Galimi F, Narita M, Lowe SW, Yaswen P, et al. Reversal of human cellular senescence: roles of the p53 and p16 pathways. *EMBO J* 2003;22(16):4212–22.
- [13] Jacobs JJ, de Lange T. Significant role for p16INK4a in p53-independent telomere-directed senescence. *Curr Biol* 2004;14(24):2302–8.
- [14] Donato AJ, Morgan RG, Walker AE, Lesniewski LA. Cellular and molecular biology of aging endothelial cells. *J Mol Cell Cardiol* 2015;89(Pt B):122–35.
- [15] Anderson R, Richardson GD, Passos JF. Mechanisms driving the ageing heart. *Exp Gerontol* 2018;109:5–15.
- [16] Shapiro GI, Edwards CD, Kobzik L, Godleski J, Richards W, Sugarbaker DJ, et al. Reciprocal Rb inactivation and P16(Ink4) expression in primary lung cancers and cell-lines. *Cancer Res* 1995;55(3):505–9.
- [17] Rodriguez R, Meuth M. Chk1 and p21 cooperate to prevent apoptosis during DNA replication fork stress. *Mol Biol Cell* 2006;17(1):402–12.
- [18] Severino J, Allen RG, Balin S, Balin A, Cristofalo VJ. Is beta-galactosidase staining a marker of senescence in vitro and in vivo? *Exp Cell Res* 2000;257(1):162–71.
- [19] Swanson EC, Manning B, Zhang H, Lawrence JB. Higher-order unfolding of satellite heterochromatin is a consistent and early event in cell senescence. *J Cell Biol* 2013;203(6):929–42.
- [20] Narita M, Nunez S, Heard E, Narita M, Lin AW, Hearn SA, et al. Rb-mediated heterochromatin formation and silencing of E2F target genes during cellular senescence. *Cell* 2003;113(6):703–16.
- [21] Hewitt G, Jurk D, Marques FDM, Correia-Melo C, Hardy T, Gackowska A, et al. Telomeres are favoured targets of a persistent DNA damage response in ageing and stress-induced senescence. *Nat Commun* 2012;3.
- [22] Song H, Conte JV, Foster AH, McLaughlin JS, Wei CM. Increased p53 protein expression in human failing myocardium. *J Heart Lung Transplant* 1999;18(8):744–9.
- [23] Predmore JM, Wang P, Davis F, Bartolone S, Westfall MV, Dyke DB, et al. Ubiquitin proteasome dysfunction in human hypertrophic and dilated cardiomyopathies. *Circulation* 2010;121(8):997–U33.
- [24] Birks EJ, Latif N, Enesa K, Folkvang T, Luong LA, Sarathchandra P, et al. Elevated p53 expression is associated with dysregulation of the ubiquitin-proteasome system in dilated cardiomyopathy. *Cardiovasc Res* 2008;79(3):472–80.
- [25] Sano M, Minamino T, Toko H, Miyauchi H, Orimo M, Qin Y, et al. p53-induced inhibition of Hif-1 causes cardiac dysfunction during pressure overload. *Nature* 2007;446(7134):444–8.
- [26] Hammadah M, Al Mheid I, Wilmot K, Ramadan R, Abdelhadi N, Alkhoder A, et al. Telomere shortening, regenerative capacity, and cardiovascular outcomes. *Circ Res* 2017;120(7):1130–8.
- [27] Margaritis M, Sanna F, Lazaros G, Akoumianakis I, Patel S, Antonopoulos AS, et al. Predictive value of telomere length on outcome following acute myocardial infarction: evidence for contrasting effects of vascular vs. blood oxidative stress. *Eur Heart J* 2017;38(41):3094–U29.
- [28] Werner CM, Hecksteden A, Morsch A, Zundler J, Wegmann M, Kratzsch J, et al. Differential effects of endurance, interval, and resistance training on telomerase activity and telomere length in a randomized, controlled study. *Eur Heart J* 2019;40(1):34–46.
- [29] Gandhi PU, Chow SL, Rector TS, Krum H, Gaggin HK, McMurray JJ, et al. Prognostic value of insulin-like growth factor-binding protein 7 in patients with heart failure and preserved ejection fraction. *J Card Fail* 2017;23(1):20–8.
- [30] Januzzi Jr JL, Packer M, Claggett B, Liu J, Shah AM, Zile MR, et al. IGFBP7 (insulin-like growth factor-binding protein-7) and neprilysin inhibition in patients with heart failure. *Circ Heart Fail* 2018;11(10):e005133.
- [31] Jurk D, Wang CF, Miwa S, Maddick M, Korolchuk V, Tzolou A, et al. Postmitotic neurons develop a p21-dependent senescence-like phenotype driven by a DNA damage response. *Aging Cell* 2012;11(6):996–1004.
- [32] Puente BN, Kimura W, Muralidhar SA, Moon J, Amatruda JF, Phelps KL, et al. The oxygen-rich postnatal environment induces cardiomyocyte cell-cycle arrest through DNA damage response (vol. 157, pg 741, 2014). *Cell* 2014;157(5):1243–50.
- [33] Bergmann O, Bhardwaj RD, Bernard S, Zdunek S, Barnabe-Heider F, Walsh S, et al. Evidence for cardiomyocyte renewal in humans. *Science* 2009;324(5923):98–102.
- [34] Bergmann O, Zdunek S, Felker A, Salehpour M, Alkass K, Bernard S, et al. Dynamics of cell generation and turnover in the human heart. *Cell* 2015;161(7):1566–75.
- [35] Richardson GD, Laval S, Owens WA. Cardiomyocyte regeneration in the mdx mouse model of nonischemic cardiomyopathy. *Stem Cells Dev* 2015;24(14):1672–9.
- [36] Torella D, Rota M, Nurzynska D, Musso E, Monsen A, Shiraishi I, et al. Cardiac stem cell and myocyte aging, heart failure, and insulin-like growth factor-1 overexpression. *Circ Res* 2004;94(4):514–24.
- [37] Spallarossa P, Altieri P, Aloï C, Garibaldi S, Barisione C, Ghigliotti G, et al. Doxorubicin induces senescence or apoptosis in rat neonatal cardiomyocytes by regulating the expression levels of the telomere binding factors 1 and 2. *Am J Physiol-Heart C* 2009;297(6):H2169–81.
- [38] Sharifi-Sanjani M, Oyster NM, Tichy ED, Bedi KC, Harel O, Margulies KB, et al. Cardiomyocyte-specific telomere shortening is a distinct signature of heart failure in humans. *J Am Heart Assoc* 2017;6(9).
- [39] Mac TW, Hauck L, Grothe D, Billia F. p53 regulates the cardiac transcriptome. *Proc Natl Acad Sci U S A* 2017;114(9):2331–6.
- [40] Rossman MJ, Kaplon RE, Hill SD, McNamara MN, Santos-Parker JR, Pierce GL, et al. Endothelial cell senescence with aging in healthy humans: prevention by habitual exercise and relation to vascular endothelial function. *Am J Physiol-Heart C* 2017;313(5):H890–5.
- [41] Yoshida Y, Shimizu I, Katsuomi G, Jiao S, Suda M, Hayashi Y, et al. p53-Induced inflammation exacerbates cardiac dysfunction during pressure overload. *J Mol Cell Cardiol* 2015;85:183–98.
- [42] Yokoyama M, Shimizu I, Nagasawa A, Yoshida Y, Katsuomi G, Wakasugi T, et al. p53 plays a crucial role in endothelial dysfunction associated with hyperglycemia and ischemia. *J Mol Cell Cardiol* 2019;129:105–17.
- [43] Borlaug BA. The pathophysiology of heart failure with preserved ejection fraction. *Nat Rev Cardiol* 2014;11(9):507–15.
- [44] Frieler RA, Mortensen RM. Immune cell and other noncardiomyocyte regulation of cardiac hypertrophy and remodeling. *Circulation* 2015;131(11):1019–30.
- [45] Gogiraju R, Xu X, Bochenek ML, Steinbrecher JH, Lehnart SE, Wenzel P, et al. Endothelial p53 deletion improves angiogenesis and prevents cardiac fibrosis and heart failure induced by pressure overload in mice. *J Am Heart Assoc* 2015;4(2).
- [46] Paulus WJ, Tschope C. A novel paradigm for heart failure with preserved ejection fraction comorbidities drive myocardial dysfunction and remodeling through coronary microvascular endothelial inflammation. *J Am Coll Cardiol* 2013;62(4):263–71.
- [47] Gevaert AB, Shakeri H, Leloup AJ, Van Hove CE, De Meyer GRY, Vrints CJ, et al. Endothelial senescence contributes to heart failure with preserved ejection fraction in an aging mouse model. *Circ Heart Fail* 2017;10(6).
- [48] Li Y, Peng Z, Wang C, Li L, Leng Y, Chen R, et al. Novel role of PKR in palmitate-induced Sirt1 inactivation and endothelial cell senescence. *Am J Physiol Heart Circ Physiol* 2018;315(3):H571–80.
- [49] Banerjee I, Fuseler JW, Price RL, Borg TK, Baudino TA. Determination of cell types and numbers during cardiac development in the neonatal and adult rat and mouse. *Am J Physiol Heart Circ Physiol* 2007;293(3):H1883–91.
- [50] Vliegen HW, van der Laarse A, Cornelisse CJ, Eulderink F. Myocardial changes in pressure overload-induced left ventricular hypertrophy. A study on tissue composition, polyploidization and multinucleation. *Eur Heart J* 1991;12(4):488–94.
- [51] Sawaki D, Czibik G, Pini M, Ternacle J, Suffee N, Mercedes R, et al. Visceral adipose tissue drives cardiac aging through modulation of fibroblast senescence by osteopontin production. *Circulation* 2018;138(8):809–22.
- [52] Jia L, Zhang W, Ma Y, Chen B, Liu Y, Piao C, et al. Haploinsufficiency of ataxia telangiectasia mutated accelerates heart failure after myocardial infarction. *J Am Heart Assoc* 2017;6(7).
- [53] Aminzadeh MA, Tseliou E, Sun B, Cheng K, Malliaras K, Makkar RR, et al. Therapeutic efficacy of cardiosphere-derived cells in a transgenic mouse model of non-ischaemic dilated cardiomyopathy. *Eur Heart J* 2015;36(12):751–62.
- [54] Gallet R, de Couto G, Simsoló E, Valle J, Sun B, Liu W, et al. Cardiosphere-derived cells reverse heart failure with preserved ejection fraction (HFpEF) in rats by decreasing fibrosis and inflammation. *JACC Basic Transl Sci* 2016;1(1–2):14–28.
- [55] Grigorian-Shamagian L, Liu WX, Fereydooni S, Middleton RC, Valle J, Cho JH, et al. Cardiac and systemic rejuvenation after cardiosphere-derived cell therapy in senescent rats. *Eur Heart J* 2017;38(39):2957–+.
- [56] Walravens AS, Vanhaverbeke M, Ottaviani L, Gillijns H, Trenson S, Vanden Driessche N, et al. Molecular signature of progenitor cells isolated from young and adult human hearts. *Sci Rep-UK* 2018;8.
- [57] Lewis-McDougall FC, Ruchaya PJ, Domenjo-Vila E, Shin Teoh T, Prata L, Cottle BJ, et al. Aged-senescent cells contribute to impaired heart regeneration. *Aging Cell* 2019;e12931.
- [58] Matsumoto C, Jiang Y, Emathingier J, Quijada P, Nguyen N, De La Torre A, et al. Short telomeres induce p53 and autophagy and modulate age-associated changes in cardiac progenitor cell fate. *Stem Cells* 2018;36(6):868–80.
- [59] Hatzistergos KE, Quevedo H, Oskouei BN, Hu QH, Feigenbaum GS, Margitich IS, et al. Bone marrow mesenchymal stem cells stimulate cardiac stem cell proliferation and differentiation. *Circ Res* 2010;107(7):913–+.
- [60] Nollet E, Hoymans VY, Rodrigus IR, De Bock D, Dom M, Van Hoof VOM, et al. Accelerated cellular senescence as underlying mechanism for functionally impaired bone marrow-derived progenitor cells in ischemic heart disease. *Atherosclerosis* 2017;260:138–46.
- [61] Baar MP, Brandt RM, Putavet DA, Klein JD, Derks KW, Bourgeois BR, et al. Targeted apoptosis of senescent cells restores tissue homeostasis in response to chemotoxicity and aging. *Cell* 2017;169(1):132–47.e16.

- [62] Chang JH, Wang YY, Shao LJ, Laberge RM, Demaria M, Campisi J, et al. Clearance of senescent cells by ABT263 rejuvenates aged hematopoietic stem cells in mice. *Nat Med* 2016;22(1):78-+.
- [63] Xu M, Palmer AK, Ding H, Weivoda MM, Pirtskhalava T, White TA, et al. Targeting senescent cells enhances adipogenesis and metabolic function in old age. *Elife* 2015;4.
- [64] Zhu Y, Tchkonina T, Pirtskhalava T, Gower AC, Ding HS, Giorgadze N, et al. The Achilles' heel of senescent cells: from transcriptome to senolytic drugs. *Aging Cell* 2015;14(4):644–58.
- [65] Katsuumi G, Shimizu I, Yoshida Y, Minamino T. Vascular senescence in cardiovascular and metabolic diseases. *Front Cardiovasc Med* 2018;5.
- [66] Roos CM, Zhang B, Palmer AK, Ogronnik MB, Pirtskhalava T, Thalji NM, et al. Chronic senolytic treatment alleviates established vasomotor dysfunction in aged or atherosclerotic mice. *Aging Cell* 2016;15(5):973–7.
- [67] Walaszczyk A, Dookun E, Redgrave R, Tual-Chalot S, Victorelli S, Spyridopoulos I, et al. Pharmacological clearance of senescent cells improves survival and recovery in aged mice following acute myocardial infarction. *Aging Cell* 2019; e12945.
- [68] Kirkland JL, Tchkonina T. Cellular senescence: a translational perspective. *Ebiomedicine* 2017;21:21–8.
- [69] Kirkland JL, Tchkonina T, Zhu Y, Niedernhofer LJ, Robbins PD. The clinical potential of senolytic drugs. *J Am Geriatr Soc* 2017;65(10):2297–301.