



Shear wave elastography investigation of multifidus stiffness in individuals with low back pain[☆]

Carlos Murillo, Deborah Falla, Alison Rushton, Andy Sanderson, Nicola R. Heneghan^{*}

Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, UK

ARTICLE INFO

Keywords:

Low back pain
Muscular stiffness
Lumbar multifidus
Shear wave elastography

ABSTRACT

The purpose of this study was to investigate differences in passive muscular stiffness between the superficial multifidus (SM) and deep multifidus (DM), and to compare their passive and active stiffness in individuals with low back pain (LBP) and asymptomatic individuals. Fifteen LBP individuals and 15 asymptomatic individuals were recruited. Passive stiffness of the SM and DM was measured bilaterally using shear wave elastography (SWE) with participants lying prone. Active stiffness was measured for the SM during trunk extension, and the contraction ratio was calculated. DM displayed higher passive muscular stiffness than SM in both the asymptomatic and LBP groups (14.41 ± 2.62 and 15.40 ± 2.77 kPa respectively; $p < 0.001$). Individuals with LBP exhibited higher passive muscular stiffness of SM (LBP: 10.15 ± 4.21 , asymptomatic: 6.84 ± 1.69 kPa; $p < 0.005$) and a lower contraction ratio (LBP: 1.54 ± 0.47 , asymptomatic: 2.65 ± 1.36 kPa; $p < 0.003$) compared to the asymptomatic group. The findings support a differentiation in passive muscular stiffness between SM and DM and provide evidence for an alteration in muscular stiffness at rest in individuals with LBP. The lower increase of muscular stiffness with contraction observed for those with LBP may reflect a deficit in activation of the multifidus.

1. Introduction

Research in the field of electromyography (EMG) has supported differences in function between the superficial (SM) and deep fibers of the multifidus (DM) and in addition, impaired function of this muscle in people with low back pain (LBP) (MacDonald et al., 2006). EMG research has supported first, differences in function between the superficial (SM) and deep fibers of the multifidus (DM) and second, impaired function of this muscle in people with LBP (Danneels et al., 2002; Moseley et al., 2002; MacDonald et al., 2009). It has been theorized that both, the differences in function between multifidus fibers and the functional impairment observed in people with LBP, may be related to the muscle structure, but research in this vein is inconclusive (Porterfield and DeRosa, 1998; Cagnie et al., 2015). However, investigating the mechanical properties of muscle, such as muscular stiffness, may offer a better understanding of variation within the multifidus fibers and the relationship between muscle structure and normal/altered function (Brandenburg et al., 2014; Roberts, 2016).

Shear wave elastography (SWE) provides a non-invasive quantitative measure of muscular stiffness (measured in shear elastic modulus)

at rest (passive) and during a contraction (active), which has shown to be positively related to the level of muscular activity and muscle force (Nordez and Hug, 2010; Brandenburg et al., 2014; Yoshitake et al., 2014; Ateş et al., 2015). SWE has previously been used to investigate the stiffness of the lumbar multifidus of asymptomatic individuals at rest and during contraction with good to excellent reliability (intra class correlation coefficients (ICC) values of between 0.77 and 0.94) (Moreau et al., 2016; Creze et al., 2017; Kopenhagen et al., 2018). However, no study has investigated whether or not differences in muscular stiffness exist between the SM and DM. Furthermore, only two studies have investigated passive muscular stiffness of multifidus in people with LBP, but the results are conflicting (Chan et al., 2012; Masaki et al., 2017).

In this study, we investigate (1) whether differences in muscular stiffness at rest exist between the SM and DM in asymptomatic and LBP individuals and (2) if differences in muscular stiffness at rest and with contraction exist in individuals with LBP compared to asymptomatic individuals. This study stands to provide novel insights into the normal mechanical properties of the multifidus muscle and how this is modified in individuals with LBP.

[☆] This study was approved by the University of Birmingham ethics committee and the procedures were conducted in agreement with the Declaration of Helsinki (ERN_17-0782).

^{*} Corresponding author.

E-mail address: n.heneghan@bham.ac.uk (N.R. Heneghan).

2. Methods

2.1. Participants

Fifteen individuals with LBP and 15 asymptomatic were recruited from staff and student communities at the University of Birmingham. All participants were eligible for this study if they were aged between 20 and 55 years, with 55 chosen as the maximum age to reduce the effect of age-related adipose infiltration within the muscle (Marcus et al., 2010). The LBP group included participants who had reported continuous LBP for more than 3 months or non-continuous pain for greater than 6 months with pain on at least half of the days (Krismer and Van Tulder, 2007). The asymptomatic group included participants without history of LBP. Exclusion criteria for both groups included neurological or respiratory disorders, pregnancy or previous spinal surgery. Individuals with LBP must not have been receiving treatment from a health care professional at the time of recruitment. Additional exclusion criteria for the LBP group included no known underlying pathology such as spinal stenosis, vertebral fracture, disc herniation, radicular low back pain with neurological deficit suggesting nerve root compression and/or ankylosing spondylitis (Krismer and Van Tulder, 2007). Ethical approval was granted by the University of Birmingham ethics committee (ERN_17-0782) and the procedures were conducted in agreement with the Declaration of Helsinki. Informed written consent was obtained from all participants.

2.2. Questionnaires

Participants with LBP completed the Numerical Rating Scale (NRS) to assess their pain intensity on the day of the measurement session and were also asked to rate their usual level of pain during the previous week. Additionally, the Oswestry Disability Index (ODI) and Tampa Scale for Kinesiophobia (TSK) were used to assess perceived disability and fear-avoidance behavior respectively (Vlaeyen et al., 1995; Fairbank and Pynsent, 2000).

2.3. Procedure

Stiffness of the SM and DM was measured bilaterally using an ultrasound imaging device with SWE (LOGIQ S8 GE Healthcare, Chicago USA) and a 9-linear array probe. All measurements were performed by the same experienced examiner trained in SWE measures. Participants were positioned in prone with a rolled towel placed under their abdomen to minimize the lumbar lordosis (Stokes et al., 2007). The ultrasound probe was placed 2 cm lateral to the level of the third lumbar spinous process (L3), which corresponds with the space between transverse process of L3 and L4; confirmed by the ultrasound image. The probe was placed on the skin with minimal pressure across all participants (Cortez et al., 2016). As muscle tissue is anisotropic, the ultrasound B-mode was used to identify the parallel orientation to the muscle fibers of SM; so the probe was positioned rotated towards the midline approximately 10° and also tilted approximately 10° from the sagittal plane (Cortez et al., 2016). Once the orientation of the muscle fibers was identified, the outline of the probe was marked on the participant's skin to ensure consistency in placement across measures. For the DM, it was not possible to identify the orientation of the fibers. The multifidus muscle was divided in two equal region of interest (ROI), which were located under the thoracolumbar fascia (TLF) (without including it) for the SM, and just below this position and above the articular processes of the vertebrae for the DM (Fig. 1). As the ROIs were defined to include the larger SM and DM area possible, these were different across participants.

To measure passive muscular stiffness of the SM and DM, participants remained five minutes lying down on the plinth before starting the acquisition to ensure that the muscle was at rest (Creze et al., 2017). The probe was placed on the area marked previously and was kept motionless for five seconds to obtain a well-defined elastography frame (Koo et al., 2013). Then, two acquisitions on each side allowed recording of nine continuous elastograms for SM and DM. Active

muscular stiffness measures of the SM were acquired during an isometric trunk extension akin to Ito test (1996), (~15° of trunk extension). The examiner visually monitored that participants did not drop the trunk extension position during the performance of the task (Ito et al., 1996). The SWE acquisition commenced when the participant reached a steady trunk extension position, and nine elastograms were acquired twice on each side with a 10-s rest between repetitions.

2.4. Image processing

After the SWE acquisition, an area was circled over the ROI for all saved elastograms. The few elastograms with artefacts caused by an attenuation effect were eliminated for the analysis to avoid under- or over-estimation of shear elastic values (MacDonald et al., 2016). Shear elastic modulus (μ) within each ROI were automatically calculated by the SWE software following the formula $\mu = \rho v^2$, where ρ is the density of the muscle tissue (assumed to be 1000 kg/m³) and v is the shear wave propagation velocity (Genisson et al., 2013). The mean of the two acquisitions was calculated to obtain representative values for each measure (Masaki et al., 2017). To quantify the increase of shear elastic modulus with contraction, the contraction ratio (Botanlioglu et al., 2013) was calculated for the SM by dividing shear modulus at rest from the mean of shear modulus with contraction (absolute values).

2.5. Statistical analysis

Descriptive statistics were used to analyze demographic data with inferential analysis including parametric and non-parametric tests used to compare groups. The Shapiro-Wilk normality test did not reveal significant deviation from normality for the measures of passive muscular stiffness and contraction ratio and paired-samples t-tests revealed no differences between sides for all measures, so the mean of the right and the left side was calculated for further analysis.

A two-way repeated measures analysis of variance (ANOVA) (with group as the between-subject independent variable and muscle fibers as within-subject factor) was performed to investigate if differences in shear elastic modulus at rest (passive muscular stiffness) of the SM and DM existed within and between groups. Pairwise comparisons with Bonferroni adjustment were used to determine significant differences. Independent samples t-tests were performed to compare the contraction ratio of the SM between groups. The intra-rater reliability of the SWE acquisitions (mean of 9 elastograms, right side asymptomatic group) was examined using two-way mixed-effects model [ICC (3.1)].

3. Results

3.1. Population characteristics

The characteristics of both groups are presented in Table 1. Both groups were comparable in age, gender, and BMI, with no significant differences seen between groups. The LBP group showed low disability and pain, with an average reported pain level at the time of data collection of 2.27 ± 1.62 out of 10.

3.2. Muscular stiffness

Figs. 2 and 3 show representative elastograms to determine passive muscular stiffness of the SM and DM, and active muscular stiffness of the SM for an asymptomatic individual and an individual with LBP. There was a significant difference between the shear elastic modulus at rest of the SM and DM as determined by the repeated measures ANOVA with Greenhouse-Geisser correction ($F(1,29) = 65.05, p < 0.001$). Post hoc comparisons revealed that shear elastic modulus at rest were higher in the DM than the SM in both groups ($p < 0.001$) (Table 2, Fig. 4). Moreover, shear elastic modulus of the SM at rest were greater for the LBP group relative to the asymptomatic group ($p = 0.005$). However, no significant differences in




Fig. 1. Representative elastograms recorded from an asymptomatic participant for the SM (A) and DM (B). The layers of tissue are marked from superficial to deep; Subcutaneous tissue (1), TLF and erector spinae aponeurosis (2), multifidus muscle (3) and transverse process of L4 (4). The white dashed line represents the junction between the multifidus and the vertebral processes. Blue colours signify lower muscular stiffness values measured in shear wave velocity (m/s) and red colours signify higher muscular stiffness values. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Table 1

Baseline characteristics of the participants, measured prior to the start of data collection.

Characteristic	LBP Mean ± SD	Asymptomatic Mean ± SD	P value
Age (years)	29.4 ± 10.80	26.71 ± 5.40	<i>p</i> = 0.90
Gender (% male)	46.70	53.30	<i>p</i> = 0.715
BMI (kg/m ²)	25.29 ± 3.18	24.01 ± 3.42	<i>p</i> = 0.54
NRS current pain (0–10)	2.27 ± 1.62		
NRS usual pain (0–10)	2.93 ± 1.98		
ODI (%)	12.7 ± 6.35		
TSK	34.4 ± 3.13		

LBP –Low Back Pain, BMI – Body Mass Index, NRS – Numeric Rating Scale for pain, ODI – Oswestry Disability Index, TSK – Tampa Scale for Kinesiophobia.

shear elastic modulus of the DM were found between groups (*p* = 0.181). An independent samples *t*-test revealed a lower contraction ratio of the SM for the LBP group compared to the asymptomatic controls (1.54 ± 0.47 and 2.65 ± 1.36 , *p* < 0.003) (Fig. 5). The ICC values (95% confidence interval) were 0.92 (0.79–0.97) and 0.90 (0.72–0.97) for shear elastic modulus at rest of the SM and DM respectively; and 0.81 (0.51–0.94) for shear elastic modulus of the SM with contraction.

4. Discussion

This is the first study to investigate whether differences in passive

muscular stiffness exist between the DM and SM both in asymptomatic participants and in people with LBP. The findings illustrate a difference in muscular stiffness between the SM and DM, supporting the existence of differences between the deep and superficial fibers of the multifidus (MacDonald et al., 2009; Moseley et al., 2002). In addition, individuals with LBP exhibited increased muscular stiffness of the SM at rest, and a reduced ability to stiffen this muscle with isometric trunk extension compared to asymptomatic individuals.

4.1. Passive muscular stiffness of SM and DM

Shear elastic modulus values at rest differed between the fibers of the multifidus, with the DM displaying greater shear elastic modulus values. Previous studies have evaluated stiffness of the multifidus but without differentiation between the DM and the SM or they have only examined the SM (Chan et al., 2012; Moreau et al., 2016; Masaki et al., 2017). In line with the current findings, higher shear elastic values at rest have been observed for the deep posterior cervical muscles relative to the superficial muscles using SWE (Dieterich et al., 2017).

In vitro animal studies have showed that type I fibers are stiffer than type II (Goubel and Marini, 1987; Petit et al., 1990); and therefore, the current findings may reflect differences in fiber type distribution between SM and DM. Histological research is inconclusive due to sample bias; but functional MRI have revealed differences in the relaxation time between SM and DM, suggesting that the DM has a higher percentage of type I fibers compared to the SM (Dickx et al., 2010; Cagnie




Fig. 2. Representative elastograms recorded from an asymptomatic participant; passive muscular stiffness of the SM (A), passive muscular stiffness of the DM (B) and active muscular stiffness of the SM (C).




Fig. 3. Representative elastograms recorded from an LBP participant; passive muscular stiffness of the SM (A), passive muscular stiffness of the DM (B) and active muscular stiffness of the SM (C).

Table 2

Values for muscular stiffness measurements and comparison of passive muscular stiffness of Superficial Multifidus (SM) and Deep Multifidus (DM) within and between groups.

	Passive muscular stiffness SM Shear elastic modulus (kPa)	Passive muscular stiffness DM Shear elastic modulus (kPa)	Post hoc comparisons between muscle fibers
LBP	10.15 ± 4.21	14.41 ± 2.62	$p < 0.001^*$, 95% CIs [1.797, 6.732]
Asymptomatic	6.84 ± 1.69	15.40 ± 2.77	$p < 0.001^*$, 95% CIs [6.797, 11.670]
Post hoc comparisons between groups	$p = 0.005^*$, 95% CIs [1175, 5.906]	$p = 0.181$, 95% CIs [-0.728, 5.906]	

LBP, low back pain; kPa, kilopascals; CIs, confidence intervals.




Fig. 4. Shear elastic modulus at rest (passive muscular stiffness) of the Superficial Multifidus (SM) and Deep Multifidus (DM) for LBP and asymptomatic groups. * $P < 0.05$.

et al., 2015). Type I fibers are more fatigue resistant than type I; and so, ideally suited to hold low load tonic activity contributing to the postural control (Porterfield and DeRosa, 1998). Thus, together with previous research, the current findings lend support to the existence of a structural differences between the SM and DM; which may have a functional implication in which the DM may provide spinal support (MacDonald et al., 2006).

4.2. Differences in multifidus stiffness in individuals with LBP

Greater shear elastic modulus values of the SM at rest were found for the LBP group when compared to asymptomatic participants. Masaki et al. (2017) previously reported significantly greater shear elastic modulus of multifidus at rest (measured at the level of L4) in individuals with LBP; however, Chan et al. (2012) did not observe group differences even if multifidus was examined at the same spinal level. In both studies, the ROI covered both the SM and DM and therefore, any potential differences between groups for SM muscular

stiffness may have been concealed by the DM values. Furthermore, Chan et al. (2012) utilized strain elastography, which is more operator dependent, potentially influencing their results (Brandenburg et al., 2014).

The differences in shear elastic modulus between LBP and asymptomatic individuals may reflect differences in muscle composition since passive stiffness is not only attributed to the contractile tissue within the muscle (Gillies and Lieber, 2011). Interestingly, Brown et al. (2011) induced lumbar disc degeneration in rabbits and found that, though the individual paravertebral muscle fibers became stiffer, the fiber bundles (composed of both muscle fibers and connective tissue) displayed a greater increase in stiffness. Thus, the increase of connective tissue due to a fibrotic proliferation may increase the shear elastic modulus values in LBP individuals (Brown et al., 2018), explaining the current findings and those reported by Masaki et al. (2017).

By contrast, the opposite findings reported by Chan et al. (2012) may be explained because of the higher adipose tissue infiltration found in their LBP group, which may have decreased the shear elastic




Fig. 5. Contraction ratio of the superficial multifidus (SM) for LBP and asymptomatic groups.

modulus values and concealed the between group differences (Roskopf et al., 2015). It has been found that the fat infiltration within multifidus may be caused by aging rather than by presence of pain (Lee et al., 2017). This may explain the higher adipose tissue infiltration reported by Chan et al. (2012) in the LBP group, which was older than the control group. In the same manner, the current findings of higher muscular stiffness may be result of a low level of adipose tissue infiltration in our LBP group, which was relatively young. In addition, though all participants had LBP for longer than 6 months, nearly all of them had non-continuous LBP and, therefore, may also exhibit a low amount of adipose tissue infiltration (Goubert et al., 2017).

4.3. Differences in contraction ratio

The participants with LBP presented a significantly lower contraction ratio; reflective of a smaller increase of muscular stiffness with contraction. The contraction ratio has previously been used to compare the increase of muscular stiffness with contraction between different conditions (pain/no pain) or between different muscles/muscle layers (Botanlioglu et al., 2013; Dieterich et al., 2017). As a normalized measurement for each participant, where muscular stiffness at rest differs between conditions, the contraction ratio allows for a more accurate estimation of differences in stiffness with contraction and force generation (Botanlioglu et al., 2013; Dieterich et al., 2017). Similar to the current findings, lower normalized active muscular stiffness was found in the deeper posterior neck muscles during isometric neck extension in individuals with neck pain (Dieterich et al., 2018).

As previous research has shown a positive linear relationship between shear elastic modulus, contraction and the level of muscular activity and muscle force, the current results may be compared in some extent to findings from EMG studies that investigated the activation of the SM during isometric contractions (Nordez and Hug, 2010; Yoshitake et al., 2014; Ateş et al., 2015). In agreement with the current findings, reduced activation of the multifidus has been observed during trunk extension in a prone position in individuals with acute and experimental LBP (Danneels et al., 2002; Dickx et al., 2008). It is speculated that this deficit in contraction found in individuals with LBP (reflected by a lower increase of muscular stiffness), may be explained in part by the proliferation of collagen content/connective tissue hypothesized above based on the finding of higher muscular stiffness at rest. These changes within the muscle would result in a decrease in the amount of contractile tissue and subsequently reduced ability to perform an efficient contraction (Goubert et al., 2017).

4.4. Methodological considerations

A limitation of SWE is the large inter-individual variability. Given that the SWE acquisitions were performed at a specific vertebral level and at a standardized distance from the spinous process, intra-muscular variations and regional differences likely explain a small extent of the variability with in the current data (Cortez et al., 2016; Stokes et al., 2007). The higher variability in shear modulus of the SM at rest in the LBP group likely reflects the large variability of individual neuromuscular adaptations due to LBP and/or an increase of the amount of non-contractile tissue (Hodges et al., 2013; Brown et al., 2018). Although elastograms with artefacts were removed from the analysis, the attenuation effect of the ultrasound push beam can be greater in the deep lumbar region due to the TLF, and might have generated artificial areas of very low/high stiffness, altering the muscular stiffness measurement and concealing the detection of significant differences between groups for the DM (MacDonald et al., 2016). Also, the assessment of the muscular stiffness of the DM with contraction was not included in the present study due to the poor-quality signal observed during the pilot sessions. Previous studies have reported poor quality signal during the evaluation of the deep abdominal muscles during contractions (MacDonald et al., 2016). Also, as trunk position was controlled visually as Ito et al. (1996) originally described, we cannot exclude small differences in trunk angle between groups, which could have affected measurements with contraction. Additionally, as LBP participants were not under treatment, the levels of pain and disability were fairly low; and so, different results may be obtained for individuals with more severe symptoms.

In conclusion, the present study provides new insights into the mechanical properties of the lumbar muscles. Specifically, the study demonstrates a difference in muscular stiffness between the DM and SM, with a greater shear elastic modulus values observed for the DM in both asymptomatic and LBP individuals. Greater shear elastic modulus values at rest of the SM was found in individuals with LBP. Finally, a deficit in the contraction of the SM during an isometric trunk extension task was observed for those with LBP, reflected by a lower increase of muscular stiffness with contraction.

Declaration of Competing Interest

This research did not receive any specific grant. The authors declare no conflict of interest.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jelekin.2019.05.004>.

References

- Ateş, F., Hug, F., Bouillard, K., Jubeau, M., Frappart, T., Couade, M., Bercoff, J., Nordez, A., 2015. Muscle shear elastic modulus is linearly related to muscle torque over the entire range of isometric contraction intensity. *J. Electromyogr. Kinesiol.* 25 (4), 703–708.
- Botanlioglu, H., Kantarci, F., Kaynak, G., Unal, Y., Ertan, S., Aydingoz, O., Erginer, R., Unlu, M.C., Mihmanli, I., Babacan, M., 2013. Shear wave elastography properties of vastus lateralis and vastus medialis obliquus muscles in normal subjects and female patients with patellofemoral pain syndrome. *Skeletal Radiol.* 42 (5), 659–666.
- Brandenburg, J.E., Eby, S.F., Song, P., Zhao, H., Brault, J.S., Chen, S., An, K.-N., 2014. Ultrasound elastography: the new frontier in direct measurement of muscle stiffness. *Arch. Phys. Med. Rehabil.* 95 (11), 2207–2219.
- Brown, E., Yoshitake, Y., Shinohara, M., Ueda, J., 2018. Automatic analysis of ultrasound shear-wave elastography in skeletal muscle without non-contractile tissue contamination. *Int. J. Intell. Robot. Appl.* 2 (2), 209–225.
- Brown, S.H., Gregory, D.E., Carr, A.J., Ward, S.R., Masuda, K., Lieber, R.L., 2011. Adaptations to the multifidus muscle in response to experimentally induced intervertebral disc degeneration. *Spine J. Meeting Abstr. LWW.*
- Cagnie, B., Dhooge, F., Schumacher, C., De Meulemeester, K., Petrovic, M., Van Oostervijck, J., Danneels, L., 2015. Fiber typing of the erector spinae and multifidus muscles in healthy controls and back pain patients: a systematic literature review. *J.*

- Manipulative Physiol. Ther. 38 (9), 653–663.
- Chan, S.-T., Fung, P.-K., Ng, N.-Y., Ngan, T.-L., Chong, M.-Y., Tang, C.-N., He, J.-F., Zheng, Y.-P., 2012. Dynamic changes of elasticity, cross-sectional area, and fat infiltration of multifidus at different postures in men with chronic low back pain. *Spine J.* 12 (5), 381–388.
- Cortez, C.D., Hermitte, L., Raiman, A., Mesmann, C., Lefort, T., Pialat, J., 2016. Ultrasound shear wave velocity in skeletal muscle: a reproducibility study. *Diagn. Intervent. Imaging* 97 (1), 71–79.
- Creze, M., Nyangoh Timoh, K., Gagey, O., Rocher, L., Bellin, M.F., Soubeyrand, M., 2017. Feasibility assessment of shear wave elastography to lumbar back muscles: a radio-anatomic study. *Clin. Anat.*
- Danneels, L., Coorevits, P., Cools, A., Vanderstraeten, G., Cambier, D., Witvrouw, E., De Cuyper, H., 2002. Differences in electromyographic activity in the multifidus muscle and the iliocostalis lumborum between healthy subjects and patients with sub-acute and chronic low back pain. *Eur. Spine J.* 11 (1), 13–19.
- Dickx, N., Cagnie, B., Achten, E., Vandemaele, P., Parlevliet, T., Danneels, L., 2008. Changes in lumbar muscle activity because of induced muscle pain evaluated by muscle functional magnetic resonance imaging. *Spine* 33 (26), E983–E989.
- Dickx, N., Cagnie, B., Achten, E., Vandemaele, P., Parlevliet, T., Danneels, L., 2010. Differentiation between deep and superficial fibers of the lumbar multifidus by magnetic resonance imaging. *Eur. Spine J.* 19 (1), 122–128.
- Dieterich, A., Yavuz, U., Nordez, A., Falla, D., Petzke, F., Hug, F., Nordez, A., 2018. Neck muscle stiffness is objectively not increased in chronic neck pain. In: XXII Congress of the International Society of Electrophysiology and Kinesiology, Dublin.
- Dieterich, A.V., Andrade, R.J., Le Sant, G., Falla, D., Petzke, F., Hug, F., Nordez, A., 2017. Shear wave elastography reveals different degrees of passive and active stiffness of the neck extensor muscles. *Eur. J. Appl. Physiol.* 117 (1), 171–178.
- Fairbank, J.C., Pynsent, P.B., 2000. The Oswestry disability index. *Spine* 25 (22), 2940–2953.
- Gennisson, J.-L., Deffieux, T., Fink, M., Tanter, M., 2013. Ultrasound elastography: principles and techniques. *Diagn. Intervent. Imaging* 94 (5), 487–495.
- Gillies, A.R., Lieber, R.L., 2011. Structure and function of the skeletal muscle extracellular matrix. *Muscle Nerve* 44 (3), 318–331.
- Goubel, F., Marini, J., 1987. Fibre type transition and stiffness modification of soleus muscle of trained rats. *Pflügers Archiv.* 410 (3), 321–325.
- Goubert, D., De Pauw, R., Meeus, M., Willems, T., Cagnie, B., Schoupe, S., Van Oosterwijck, J., Dhondt, E., Danneels, L., 2017. Lumbar muscle structure and function in chronic versus recurrent low back pain: a cross-sectional study. *Spine J.* 17 (9), 1285–1296.
- Hodges, P., Coppeters, M., MacDonald, D., Cholewicki, J., 2013. New insight into motor adaptation to pain revealed by a combination of modelling and empirical approaches. *Eur. J. Pain* 17 (8), 1138–1146.
- Ito, T., Shirado, O., Suzuki, H., Takahashi, M., Kaneda, K., Strax, T.E., 1996. Lumbar trunk muscle endurance testing: an inexpensive alternative to a machine for evaluation. *Arch. Phys. Med. Rehabil.* 77 (1), 75–79.
- Koo, T.K., Guo, J.-Y., Cohen, J.H., Parker, K.J., 2013. Relationship between shear elastic modulus and passive muscle force: an ex-vivo study. *J. Biomech.* 46 (12), 2053–2059.
- Koppenhaver, S., Kniss, J., Lilley, D., Oates, M., Fernández-de-las-Peñas, C., Maher, R., Croy, T., Shinohara, M., 2018. Reliability of ultrasound shear-wave elastography in assessing low back musculature elasticity in asymptomatic individuals. *J. Electromyogr. Kinesiol.* 39, 49–57.
- Krismser, M., Van Tulder, M., 2007. Low back pain (non-specific), Best practice & research clinical rheumatology 21(1), 77–91.
- Lee, S.H., Park, S.W., Kim, Y.B., Nam, T.K., Lee, Y.S.J.T.S.J. 2017. The fatty degeneration of lumbar paraspinal muscles on computed tomography scan according to age and disc level, 17(1), 81–87.
- MacDonald, D., Moseley, G.L., Hodges, P.W., 2009. Why do some patients keep hurting their back? Evidence of ongoing back muscle dysfunction during remission from recurrent back pain. *PAIN®* 142 (3), 183–188.
- MacDonald, D., Wan, A., McPhee, M., Tucker, K., Hug, F., 2016. Reliability of abdominal muscle stiffness measured using elastography during trunk rehabilitation exercises. *Ultrasound Med. Biol.* 42 (4), 1018–1025.
- MacDonald, D.A., Moseley, G.L., Hodges, P.W., 2006. The lumbar multifidus: does the evidence support clinical beliefs? *Manual Therapy* 11 (4), 254–263.
- Marcus, R.L., Addison, O., Kidde, J.P., Dibble, L.E., Lastayo, P.C., 2010. Skeletal muscle fat infiltration: impact of age, inactivity, and exercise. *J. Nutr. Health Aging* 14 (5), 362–366.
- Masaki, M., Aoyama, T., Murakami, T., Yanase, K., Ji, X., Tateuchi, H., Ichihashi, N., 2017. Association of low back pain with muscle stiffness and muscle mass of the lumbar back muscles, and sagittal spinal alignment in young and middle-aged medical workers. *Clin. Biomech.* 49, 128–133.
- Moreau, B., Vergari, C., Gad, H., Sandoz, B., Skalli, W., Laporte, S., 2016. Non-invasive assessment of human multifidus muscle stiffness using ultrasound shear wave elastography: a feasibility study. *Proc. Inst. Mech. Eng. [H]* 230 (8), 809–814.
- Moseley, G.L., Hodges, P.W., Gandevia, S.C., 2002. Deep and superficial fibers of the lumbar multifidus muscle are differentially active during voluntary arm movements. *Spine* 27 (2), E29–E36.
- Nordez, A., Hug, F., 2010. Muscle shear elastic modulus measured using supersonic shear imaging is highly related to muscle activity level. *J. Appl. Physiol.* 108 (5), 1389–1394.
- Petit, J., Filippi, G., Emonet-Denand, F., Hunt, C., Laporte, Y., 1990. Changes in muscle stiffness produced by motor units of different types in peroneus longus muscle of cat. *J. Neurophysiol.* 63 (1), 190–197.
- Porterfield, J., DeRosa, C., 1998. *Lumbopelvic Musculature: Structural and Functional Considerations, Mechanical Low Back Pain: Perspectives in Functional Anatomy*, second ed. WB Saunders, Philadelphia, PA, pp. 53–120.
- Roberts, T.J., 2016. Contribution of elastic tissues to the mechanics and energetics of muscle function during movement. *J. Exp. Biol.* 219 (2), 266–275.
- Roskopf, A.B., Ehrmann, C., Buck, F.M., Gerber, C., Flück, M., Pfirrmann, C.W.J.R., 2015. Quantitative shear-wave US elastography of the supraspinatus muscle: reliability of the method and relation to tendon integrity and muscle quality. *278(2)*, 465–474.
- Stokes, M., Hides, J., Elliott, J., Kiesel, K., Hodges, P., 2007. Rehabilitative ultrasound imaging of the posterior paraspinal muscles. *J. Orthop. Sports Phys. Ther.* 37 (10), 581–595.
- Vlaeyen, J.W., Kole-Snijders, A.M., Rottevel, A.M., Ruesink, R., Heuts, P.H., 1995. The role of fear of movement/(re) injury in pain disability. *J. Occup. Rehabil.* 5 (4), 235–252.
- Yoshitake, Y., Takai, Y., Kanehisa, H., Shinohara, M., 2014. Muscle shear modulus measured with ultrasound shear-wave elastography across a wide range of contraction intensity. *Muscle Nerve* 50 (1), 103–113.

Carlos Murillo is a physiotherapist who obtained his MRes in Spinal Pain from the University of Birmingham (United Kingdom) in 2018. He conducted his MRes thesis within the Centre of Precision Rehabilitation for Spinal Pain (CPR Spine); which was focused on investigating the alteration in muscular stiffness and activity of the lumbar muscles in people with low back pain. He is currently carrying out his PhD at Ghent University (Belgium) within a FWO funded research project.

Deborah Falla is Chair in Rehabilitation Science and Physiotherapy at the University of Birmingham, UK and is the Director of the Centre of Precision Rehabilitation for Spinal Pain (CPR Spine). Her research utilises state of the art electrophysiological measures to evaluate the control of human movement and how it is affected or adapted in response to various states (e.g. injury, fatigue, training, and pain). Her research aims to optimise the management of musculoskeletal disorders with a particular interest in spinal pain. She has published over 190 papers in international, peer-reviewed journals, and more than 300 conference papers/abstracts including over 30 invited/keynote lectures. Professor Falla has received several recognitions and awards for her work including the German Pain Research Prize, the George J. Davies - James A. Gould Excellence in Clinical Inquiry Award and the Delsys Prize for Electromyography Innovation. Professor Falla is an author of three books including the latest entitled “Management of neck pain disorders: a research informed approach” (Elsevier). Professor Falla acts as an Associate Editor for *Musculoskeletal Science & Practice*, the *Journal of Electromyography and Kinesiology* and *IEEE Transactions on Neural Systems and Rehabilitation Engineering*. She was President of the International Society of Electrophysiology and Kinesiology (ISEK) between 2016-2018.

Alison Rushton is Reader in Musculoskeletal Rehabilitation Sciences, at the University of Birmingham, UK. She is Deputy Director/Research Co-Lead of the Centre of Precision Rehabilitation for Spinal Pain and Programme Lead for the MRes Spinal Pain. Alison has a strong research profile, with > 100 publications in high impact journals including *BMJ*, *Spine* and *BJSM*. Her research is driven by patients and key issues in musculoskeletal clinical practice with a focus on spinal pain and dysfunction; personalising interventions to the individual patient, and developing evidence informed decision-making frameworks to inform safe/efficacious interventions. Alison is Chair of the Standards Committee of the International Federation of Orthopaedic Manipulative Physical therapy. And sits on the Editorial Board of *BMC Musculoskeletal*, *Plos One* and *Musculoskeletal Science and Practice* journals. Alison's contribution to musculoskeletal rehabilitation research has been recognised through the award of fellowships from the UK Chartered Society of Physiotherapy and the UK Musculoskeletal Association of Chartered Physiotherapists.

Andy Sanderson is a PhD student at the University of Birmingham's Centre of Precision Rehabilitation for Spinal Pain (CPR Spine). His research combines his prior anatomical training with cutting edge neurophysiological and kinematic techniques to assess muscle activity in the lumbar paraspinal musculature. In his PhD thesis, Andy is combining high-density electromyography and motion analysis to quantify changes in muscle activity movement caused by low back pain, with the aim to inform rehabilitation practice.

Nicola Heneghan is a Lecturer/Researcher in Musculoskeletal Rehabilitation Sciences in the Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), University of Birmingham, UK. Nicola's research investigates thoracic spine pain and dysfunction in different patient populations as well as collaborative research in patients with low back and neck pain, and following spinal surgery or trauma. She has published over 50 papers in international, peer-reviewed journals, and more than 75 conference papers/abstracts including invited/keynote lectures. She has supported > 120 physiotherapists to successful completion of postgraduate degrees (MSc, MRes and PhD) and currently supervises students across all programmes. External to her academic commitments she is currently the Chair of the MACP, the UK Member Organisation for IFOMPT, sits on the Internal Advisory Board for *Musculoskeletal Science & Practice* and is an Associate Editor for the *BMC Journal 'Systematic Reviews'*.