



Asymmetry of the frontal aslant tract is associated with lexical decision

Antonino Vallesi^{1,2} · Laura Babcock^{3,4}

Received: 27 May 2019 / Accepted: 27 February 2020 / Published online: 10 March 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Abstract

The frontal aslant tract (FAT) is a recently documented white matter tract that connects the inferior and superior frontal gyri with a tendency to be more pronounced in the left hemisphere. This tract has been found to play a role in language functions, particularly verbal fluency. However, it is not entirely clear to what extent FAT asymmetry is related to performance benefits in language-related tasks. In the present study, we aimed to fill this gap by examining the correlations between asymmetric micro- and macro-structural properties of the FAT and performance on verbal fluency and lexical decision tasks. The results showed no correlation between the FAT and verbal fluency; however, lexical decision was correlated with FAT laterality. Specifically, greater left lateralization in both micro- and macro-structural properties was related to faster lexical decision response times. The results were not due merely to motor or decision-making processes, as responses in a simple discrimination control task showed no correlation with laterality. These data are the first to suggest a role for the FAT in mediating processes underlying lexical decision.

Keywords Frontal aslant tract · Hemispheric asymmetries · Laterality · Verbal fluency · Lexical decision · Language functions · Structural connectivity

Introduction

The frontal aslant tract (FAT) is a recently identified white matter tract in the human brain (Catani et al. 2012). The FAT connects the inferior frontal gyrus (Brodmann areas 44/45) to superior medial frontal regions including anterior cingulate cortex, supplementary motor area (SMA), and

pre-supplementary motor area (pre-SMA), as corroborated by magnetic resonance imaging (MRI) functional connectivity between these regions (e.g., Crosson et al. 2001; see Dick et al. 2019, for a review). The FAT is typically left lateralized in terms of volume in right-handed individuals (Catani et al. 2012), suggesting a role in language functions, as the same lateralization has been observed also for other language-related tracts, including the long segment of the arcuate fasciculus (Thiebaut de Schotten et al. 2011b).

Indeed, a role in speech fluency has been confirmed for the left FAT by a number of clinical studies using intraoperative electrical stimulation during tumor resection (Fujii et al. 2015; Kemerdere et al. 2016; Kinoshita et al. 2015; Vassal et al. 2014). These studies showed that direct stimulation of the left FAT consistently resulted in speech disturbances (e.g., speech arrest, speech inhibition, stuttering). Further, Kinoshita (2015) found that patients with post-operative speech disturbances had smaller distances between the tumor resection site and the left FAT than those without this type of deficit; no right hemisphere patients showed language-related post-operative disturbances. Additionally, the average distance between tumor resection cavity and the left FAT was positively correlated with semantic and

Antonino Vallesi and Laura Babcock contributed equally to this work.

✉ Antonino Vallesi
antonino.vallesi@unipd.it

✉ Laura Babcock
lbabcock@gmail.com

¹ Department of Neuroscience & Padova Neuroscience Center, University of Padova, 35128 Padova, Italy

² Brain Imaging and Neural Dynamics Research Group, IRCCS San Camillo Hospital, Venice, Italy

³ Department of Neuroscience, D4, Biomedicum, Karolinska Institutet, 17177 Stockholm, Sweden

⁴ Department of Neuroscience, University of Padova, Padova 35128, Italy

phonemic fluency, but not naming ability (also see Chernoff et al. 2018 for analogous results).

Additional evidence that the left FAT is implicated in fluency comes from studies on patients with aphasia. In patients with primary progressive aphasia (PPA), the mean length of utterance and speech rate were correlated with proxy measures of white matter microstructural integrity of the left FAT such as fractional anisotropy (FA) and/or radial diffusivity (RD) (Catani et al. 2013; also see Mandelli et al. 2014). These measures did not correlate with grammatical processing or picture naming abilities. Correlations between verbal fluency abilities and the integrity of the left FAT (measured with lesion percentage and FA) were also seen in an examination of 51 post-stroke patients (Li et al. 2017). An additional study of post-stroke aphasic patients found that the white matter signal intensity (from a T1 scan) in the left FAT/anterior segment was the best predictor of speech fluency compared to other left hemisphere tracts involved in speech and language (Basilakos et al. 2014).

Beyond fluency, the left FAT seems well positioned to contribute to lexical retrieval processes given its end points in the inferior frontal gyrus (IFG) and pre-SMA, which have been linked to lexical processing. The former is assumed to be associated with effortful processes such as retrieval, manipulation, and selection of phonological representations (Fiez et al. 1999). In particular, left Brodmann area (BA) 45 is related to semantic aspects of language processing, and the more posterior left BA 44 is related to phonological and syntactic processing and speech programming (Amunts et al. 2004). Moreover, FA in left BA 45 has shown a negative association with lexical decision speed (Gold et al. 2007).

Left pre-SMA has been associated through fMRI studies with internally guided generation of lexical items, but only items with pre-existing representations in the brain (Crosson et al. 2001, 2003). This region is also activated during semantic decision tasks (Binder et al. 1997) and lexical decision tasks, especially for low-frequency words (Carreiras et al. 2006, 2009).

Finally, a neurosurgical study found that stimulation at the level of the pre-SMA and extending to the FAT in the left hemisphere resulted in morphological over-regularization in a noun-based verb-generation task (Sierpowska et al. 2015). The authors suggested that the stimulation acted by inhibiting more demanding lexical retrieval processes.

While the evidence above focuses on the left hemisphere, there is some indication that inter-hemispheric dynamics are important for lexical retrieval processes. For instance, healthy participants are faster in picture naming after inhibitory cathodal transcranial direct current stimulation (tDCS; vs. sham) of the right IFG, a result mostly due to fewer tip-of-the-tongue instances (Rosso et al. 2014). The interpretation of this finding was that, as the right hemisphere was inhibited by tDCS, inter-hemispheric rivalry toward the left

hemisphere was decreased. FMRI data were also collected during a word repetition task in the same participants. Effective connectivity analyses with dynamic causal modeling of these data showed hemispheric rivalry directly from right Broca's area to the left homologous area, and indirectly from left to right Broca's through the SMAs.

A similar role for inter-hemispheric dynamics has been found for fluency in relation to the FAT. In a study employing repetitive transcranial magnetic stimulation (rTMS) over left and right BA 47 (Smirni et al. 2017), performance on a phonemic fluency task was impaired after stimulation to left BA 47 (compared to sham), but improved after stimulation to right BA 47, although it must be noted that neuronavigation was performed with a probabilistic head model and not with the individual MRI. The authors suggested rTMS applied to the right hemisphere suppressed the inter-hemispheric inhibitory interactions.

Although task performance usually benefits from a division of labor between the two hemispheres (Banich and Belger 1990; Belger and Banich 1998), inter-hemispheric cooperation has been observed for complex, more demanding tasks, especially in the prefrontal cortex (e.g., Höller-Wallscheid et al. 2017), and may be beneficial for performance (Belger and Banich 1998; Berryman and Kennelly 1992).

Several studies have demonstrated that taking into account the inter-hemispheric asymmetries of intra-hemispheric white matter pathways is important to explain variability in performance across individuals for various cognitive functions, including language and attention systems (see Ocklenburg et al. 2016, for a review). As an example, the degree of structural asymmetry of the arcuate fasciculus, in terms of microstructure (FA) and/or macrostructure (volume, fiber density) is correlated with language-related functioning, with the direction of the advantage depending on the type of task. Catani (2007) found that a more symmetric arcuate fasciculus was associated with an advantage in verbal recall, while Lebel and Beaulieu (2009) found that children with leftward asymmetry in the arcuate fasciculus outperformed those with rightward asymmetry in vocabulary and phonological processing tasks. The arcuate fasciculus structural asymmetry also correlates with the degree of brain functional lateralization (Ocklenburg et al. 2013; James et al. 2015), which notably also predicts verbal abilities (Gotts et al. 2013). Similar relationships have been observed for other tracts also in non-language domains. For instance, a leftward asymmetry of the inferior corona radiata is positively correlated with the performance in Attention Network Test (ANT), in particular with the executive control sub-component (Yin et al. 2013), while the direction of structural asymmetry of the superior longitudinal fasciculus (SLF) II predicts the direction of behavioral asymmetry in visuospatial attention tasks (Thiebaut De Schotten et al. 2011a; also

see Budisavljevic et al. 2017b, for analogous results in visuomotor processing). Moreover, inter-individual differences in the asymmetry of the dorsal SLF tract predict the degree of lateralization in hand preference, visuospatial integration and fine motor control tasks (Howells et al. 2018).

However, little is known about whether and how inter-hemispheric white matter asymmetries in the FAT correlate with the ability to perform language tasks that have been associated with this tract. In this study, we tried to fill this gap by assessing correlations between asymmetric micro- and macrostructural properties of the FAT and performance on verbal fluency and lexical decision tasks. Given the previous findings that link these language abilities to the left FAT, we expected a correlation between leftward hemispheric FAT asymmetry and better performance on both of these linguistic tasks.

Methods

Participants

Twenty-nine healthy university students (19 females; mean age = 24.8, SD = 2.4, range 22–36) participated in the study. All participants were native Italian speakers with no known neurological or psychiatric problems. The Edinburgh Handedness Inventory (Oldfield 1971) was used to confirm that all participants were right handed (mean score = 85, SD = 18, range 30–100). The study was approved by the ethical committees of the Scuola Internazionale Superiore di Studi Avanzati (SISSA), “Istituto IRCCS E. Medea—La Nostra Famiglia”, and Azienda Ospedaliera di Padova. All participants gave written informed consent and were compensated for their time.

Verbal fluency task

Participants completed a verbal fluency task in which they were asked to name as many items as possible belonging to a category in 60 s.¹ Each participant completed six categories, three semantic categories and three letter categories, in their native language, Italian. The specific categories were drawn from a list of nine possible categories for each type, (adapted from Schwieter and Sunderman 2011; see Table 1 for all categories used).² Each category began

¹ Verbal fluency data were only available for 22 of the participants.

² In addition to their native Italian, participants also completed the task in two foreign languages. Only data from Italian are considered in the present analyses given the variability of these foreign languages (Dutch, English, French, German, and Spanish as well as the participants' abilities in them (scores ranging from 3 to 5 on a scale of 1–5, with 5 representing native-like).

Table 1 Categories used in the verbal fluency task

Set	Categories
Semantic set 1	Parti del corpo—body parts Professioni—occupations Strumenti musicali—musical instruments
Semantic set 2	Animali—animals Legume—vegetables Abbigliamento—clothing items
Semantic set 3	Mezzi di trasporto—methods of transportation Sport—sports Frutta—fruits
Letter set 1	A, M, F
Letter set 2	E, T, G
Letter set 3	O, D, L

with fixation cross for 500 ms, followed by a 100 ms beep (400 Hz) and then the category name was presented on the screen. Participants were given 60 s to name as many items belonging to the category as possible. The conclusion of the 60 s was signaled with another 100 ms beep. A blank screen appeared for 3000 ms before the next category. Participants were instructed that they would see two types of categories (semantic and letter) and have 60 s to name as many items as they could in the given category. They were also told that repetitions, words with the same root, and proper names would not be counted. Before completing the experimental categories, participants completed two practice categories (P and utensili da cucina—*kitchen items*) with an experimenter present to clarify any difficulties.

The entire task was recorded using a digital recorder for off-line transcription and coding. Transcription and coding were completed by a native Italian speaker. All responses initiated within the 60 s time limit for each category were transcribed and marked as either valid items or items to be excluded. For both category types, responses were excluded that were: not a word in the language, not part of the category, repetitions of a previous response in the same category, and proper names. Further for semantic categories, repetitions of a concept (e.g., eggplant and aubergine) and superordinates when subordinates were also named (e.g., bird, pigeon, bluejay) were excluded. Finally, for letter categories, only one word with a given root was counted (e.g., fast, faster, fastest). The total number of valid items produced across the six categories (three semantic and three letter) was used at the measure of interest.

Lexical decision task

Participants completed a lexical decision task in which they had to judge whether presented letter strings were real Italian words or non-words. The items in the present study were

taken from Paulesu et al. 2000 and personal communication. The real word items were disyllabic nouns that had stress on the first syllable and were among the 7500 most frequent Italian words (e.g., *collo-neck*, *mare-sea*, *porta-door*). The non-word items were also disyllabic and were formed by changing one or two phonemes in the real words (e.g., *coto*, *mave*, *borta*). Stimulus trials began with a fixation screen presented for 1000 ms. A letter string was then presented and remained onscreen until a response was given. Participants were asked to give a judgment on the letter string as quickly and accurately as possible. A ‘word’ response was made with the index finger of one hand and a ‘non-word’ response with the index finger of the other hand; hand assignments were counterbalanced across participants. The task began with 16 practice items followed by 160 experimental items, half of which were real words and half were non-words. Response times (RT) and accuracy were recorded; participants’ accuracy was very high (average 97%) and thus RT was used as the measure of interest.

Color and shape discrimination task

Participants completed a discrimination task in which they needed to indicate for each presented stimulus either the color or the shape, performed in separate blocks. This task was similar to the Lexical Decision Task, in that it required a forced-choice response from the participants; however, it did not require lexical processing. The stimuli consisted of red and blue hearts and stars presented individually on a white background. Trials began with a fixation cross presented for 1500 ms followed by a cue that indicated the present task. For color discrimination, the cue consisted of three colored rectangles (purple, orange, and yellow), while for shape discrimination three black shapes (triangle, circle, and square) were used. In both cases, the items were arranged linearly and appeared above the fixation cross location. The use of graphical cues had the benefit of limiting linguistic information. After either 100 or 1000 ms, distributed randomly and equally across the trials, the stimulus appeared in the center of the screen, below the cue, which remained onscreen. Participants were asked to indicate the color or shape as quickly and accurately as possible using the index fingers of their left and right hands. The four possible response-to-button mappings (left: red/heart, right: blue/star; left: red/star, right: blue/heart; left: blue/heart, right: red/star; left: blue/star, right: red/heart) were counterbalanced across participants. The trial ended when the participant made a response. Incorrect responses were followed by a 100 ms beep.

Participants completed two blocks of color discrimination and two blocks of shape discrimination; each block began with six practice trials, followed by 24 experimental trials. The blocks were presented using a sandwich design, meaning the first and last blocks required the same discrimination

task, with the other task presented in the second and third blocks.³ The specific assignment of tasks was counterbalanced across participants. Both response times and accuracy were recorded; RT was used as the measure of interest given the high accuracy rate of the participants (average 98%).

Diffusion imaging acquisition and processing

Images were acquired on a 3-T Philips Achieva whole-body scanner at the Santa Maria della Misericordia Hospital in Udine, Italy. Diffusion imaging was acquired with an eight-channel head coil using a single-shot, spin-echo, EPI sequence with the following parameters: TR = 9037 ms, TE = 70 ms, FOV = 240 mm × 223 mm, matrix size = 128 × 128, AP phase encoding, 57 contiguous axial slices (1.875 × 1.875 × 2.1 mm voxel resolution). For each slice, 64 diffusion-weighted images ($b = 1000$ s/mm²) and 1 non-weighted image ($b = 0$) were acquired. Data processing and tractography were performed using ExploreDTI (<https://www.exploredti.com>). First, subject motion and geometric distortions induced by eddy currents were corrected in a single step through rotation of the b-matrix. Then, a b-spline interpolated streamline algorithm was used to perform whole brain tractography (step size: 1 mm; FA threshold: 0.2; angle threshold: 35°). The output from ExploreDTI was imported to TrackVis (<https://www.trackvis.org>), where virtual in vivo dissections were performed, using software written by Natbrainlab (Thiebaut de Schotten et al. 2011a).

Tract dissection and measures

Dissection of the frontal aslant tract was completed manually (by LB) using a two regions of interest (ROI) method. Following previous tractography studies (e.g., Catani et al. 2012; Budisavljevic et al. 2017a), ROIs were placed in the inferior frontal gyrus and superior frontal gyrus on the fractional anisotropy images in the three planes. In both cases, the ROI encompassed the entirety of the gyrus. These two regions were used as ‘AND’ ROIs in tracking the FAT. Streamlines that did not follow the known anatomy of the FAT were removed. The FAT was tracked in both the left and right hemispheres (Fig. 1).

Additionally, the anterior segment of the arcuate fasciculus was dissected to serve as a control tract. The arcuate fasciculus is a white matter pathway well known for its role in language functions. This bundle can be divided into two pathways, a direct pathway connecting Broca’s area to Wernicke’s area and an indirect pathway mediated through the

³ Between the second and third block, the participants completed a mixed-task block requiring a different discrimination task on each trial; performance on that block is not considered in the present study.




Fig. 1 Right (in green) and left (in blue) frontal aslant tract in a single participant overlaid on their T1-weighted MR image. The image is generated looking posteriorly, thus the right hemisphere is represented on the left side of the image and vice versa (the cube on the bottom right depicts the brain orientation: A anterior, R right)

inferior parietal lobule (Catani et al. 2005). The anterior segment, which is considered part of the ventral branch of the superior longitudinal fasciculus (Martinos and Del Luca 2014), was selected due to its comparability to the FAT in terms of size and their shared endpoint in the inferior frontal gyrus. While the anterior segment is implicated in language functions (Catani and Mesulam 2008), it has not been linked to lexical retrieval making it an ideal choice to demonstrate the specificity of any effects to the FAT. Similar to the FAT, the anterior segment was tracked (by LB) bilaterally using two ‘AND’ ROIs. These were placed in Broca’s territory, encompassing the inferior frontal gyrus, middle frontal gyrus, and precentral gyrus, and in Geschwind’s territory composed of the supramarginal and angular gyri. Streamlines that did not follow the known anatomy of the anterior segment were removed.

Three measures of interest were extracted for each tract: number of streamlines (NoS), volume, and fractional anisotropy. NoS and volume provide information about the macrostructural properties of white matter tracts. Both measures were considered to avoid the shortcomings related to either measure (see Budisavljevic et al. 2017b, for further discussion). FA was used as it is the most widely used proxy of white matter microstructural properties. This measure indicates the degree of anisotropy in the diffusion process, ranging between 0 and 1. Values near 0 represent isotropic diffusion, while a value of 1 indicates diffusion along only one axis. In diffusion imaging, FA is taken as a proxy of fiber density, axonal diameter, and myelination of the white matter (Beaulieu 2002). To examine the asymmetry of the tracts, a lateralization index (LI) was computed for each of the three measures using the following formula: $LI = (\text{right} - \text{left}) / (\text{right} + \text{left})$ (Thiebaut de Schotten et al.

2011a). Thus, positive LI values indicate right lateralization and negative values left lateralization. The lateralization of both the FAT and the anterior segment of the arcuate fasciculus was neutral across all three measures in our sample (FAT: $ps \geq 0.241$; anterior segment: $ps \geq 0.134$), that is, the tracts showed no trend of left or right lateralization across the participants (cf. Thiebaut de Schotten et al. 2011b; 2011b; Budisavljevic et al. 2017b).

Statistical analyses

Statistical analyses were computed using IBM SPSS Statistics for Windows, Version 25.0. The two behavioral measures (verbal fluency, lexical decision RT) and three tract lateralization measures (NoS LI, volume LI, FA LI) were examined for normality with Kolmogorov–Smirnov tests. All measures were normally distributed ($p > 0.193$), except FA LI, for which there were three outliers. Removal of these participants resulted in a normal distribution for FA LI as well ($p > 0.200$). Pearson bivariate correlation analyses were used to examine the relationship between the behavioral measures and the lateralization measures. The six analyses were corrected for multiple comparisons using a false discovery rate (FDR) correction ($p < 0.05$). In addition, Pearson correlations were also computed between each behavioral measure and each white matter measure (i.e., NoS, volume, and FA) of the left and right FAT separately. Both corrected and uncorrected p values are provided.

Results

Verbal fluency was not correlated with any of the lateralization measures (NoS: $r = -0.258$, $p_{\text{uncorr}} = 0.247$, $p_{\text{corr}} = 0.346$; volume: $r = -0.237$, $p_{\text{uncorr}} = 0.288$, $p_{\text{corr}} = 0.346$; FA: $r = 0.183$, $p_{\text{uncorr}} = 0.439$, $p_{\text{corr}} = 0.439$). Lexical decision instead correlated with all three lateralization indices (NoS: $r = 0.424$, $p_{\text{uncorr}} = 0.022$, $p_{\text{corr}} = 0.044$; volume: $r = 0.475$, $p_{\text{uncorr}} = 0.027$, $p_{\text{corr}} = 0.072$; FA: $r = 0.539$, $p_{\text{uncorr}} = 0.004$, $p_{\text{corr}} = 0.024$; see Fig. 2). These correlations indicated that faster RTs on the lexical decision task were associated with left lateralization. The potential influence of lexical status on these relationships was examined with correlation analyses between the three lateralization indices and the RTs for words and non-words separately. Both words (NoS: $r = 0.367$, $p_{\text{uncorr}} = 0.050$, $p_{\text{corr}} = 0.050$; volume: $r = 0.392$, $p_{\text{uncorr}} = 0.036$, $p_{\text{corr}} = 0.043$; FA: $r = 0.540$, $p_{\text{uncorr}} = 0.004$, $p_{\text{corr}} = 0.021$) and non-words (NoS: $r = 0.431$, $p_{\text{uncorr}} = 0.020$, $p_{\text{corr}} = 0.030$; Volume: $r = 0.493$, $p_{\text{uncorr}} = 0.007$, $p_{\text{corr}} = 0.021$; FA: $r = 0.483$, $p_{\text{uncorr}} = 0.012$, $p_{\text{corr}} = 0.024$) correlated with all three lateralization measures. Finally, to confirm that the correlation was specific to lexical processes rather




Fig. 2 Correlations between the behavioral measures and tract lateralization measures. The top row shows correlations between verbal fluency score and the three lateralization measures, none of which were significant. The bottom row shows correlations between lexical

decision RT and the three lateralization measures; all three were significant; however, only the correlation with FA LI survived correction for multiple comparisons. This correlation indicated that faster RTs were associated with greater left lateralization

than motoric and/or decision-making processes, we examined the correlation between RT in the color and shape discrimination task and lateralization. This task was not correlated with any of the lateralization measures (NoS: $r = 0.316$, $p_{\text{uncorr}} = 0.095$, $p_{\text{corr}} = 0.095$; volume: $r = 0.318$, $p_{\text{uncorr}} = 0.093$, $p_{\text{corr}} = 0.095$; FA: $r = 0.341$, $p_{\text{uncorr}} = 0.089$, $p_{\text{corr}} = 0.095$).

Additionally, to understand whether verbal fluency or lexical decision may be related with the degree of laterality rather than with its (left vs. right) direction, we also assessed correlations between the behavioral tasks and the absolute laterality values; no significant correlations were found (all $ps > 0.217$).

Further, given that previous studies found a relation between language abilities and the left FAT, specifically, we also tested the correlations between the behavioral measures and the three tract measures in each hemisphere. Verbal fluency showed a correlation with the volume of the left FAT; however, this did not survive correction for multiple comparisons ($r = 0.426$, $p_{\text{uncorr}} = 0.048$, $p_{\text{corr}} = 0.162$). No other correlations reached significance before or after multiple comparison correction ($ps_{\text{uncorr}} \geq 0.054$, $ps_{\text{corr}} \geq 0.162$).

Finally, to confirm the specificity of the correlation to the FAT, we performed the same correlations using a control

tract, the anterior segment of the arcuate fasciculus. No significant correlations were found between performance on the lexical decision task and the lateralization of this tract (NoS: $r = -0.010$, $p_{\text{uncorr}} = 0.960$, $p_{\text{corr}} = 0.960$; volume: $r = 0.150$, $p_{\text{uncorr}} = 0.439$, $p_{\text{corr}} = 0.848$; FA: $r = 0.118$, $p_{\text{uncorr}} = 0.565$, $p_{\text{corr}} = 0.848$).

Discussion

In the present study, we examined the relationship between the FAT white matter structure and language functions, specifically verbal fluency and lexical decision, in healthy adults. Contrary to our hypotheses, which were mainly derived from findings with brain-damaged patients, our results showed no correlation between the lateralization indices of FAT micro- and macrostructure and verbal fluency; however, lexical decision was correlated with the laterality of the FAT. Specifically, faster lexical decision latency was associated with greater left lateralization of both the macrostructure (i.e., number of streamlines and volume) and the microstructure (i.e., fractional anisotropy) of the FAT. These results could not be explained as generally due to motor and/or decision-making processes, as responses in a simple discrimination task showed no correlation with laterality,

neither were these results explained by correlations with only the left FAT. Further, the results were specific to the FAT, and not found in another tract known to support language functions, namely, the anterior segment of the arcuate fasciculus. To the best of our knowledge, our results are the first to suggest a relation between FAT asymmetry and processes underlying lexical decision.

Verbal fluency was correlated with the volume of the left FAT only, extending the findings gathered primarily on clinical populations (Basilakos et al. 2014; Catani et al. 2013; Kinoshita et al. 2015) to healthy subjects, although this result did not survive multiple comparison correction. Importantly, the absence of correlation between FAT lateralization and performance in verbal fluency was unexpected, if one infers from results coming from clinical studies (e.g., Catani et al. 2013; Kinoshita et al. 2015). However, the present result is in line with previous work suggesting no correlation between speech fluency and the left FAT in healthy individuals (Kronfeld-Duenias et al. 2016). This finding could be tentatively interpreted as due to the fact that this type of task is clearly left lateralized in the prefrontal cortex (Gaillard et al. 2000; Schlösser et al. 1998), and thus it does not suffer much from interference from the right hemispheric homologous regions. On the other hand, there is evidence that word recognition, which is required in lexical decision tasks, involves inter-hemispheric exchange (Mohr et al. 1994; also see Ibrahim and Eviatar 2012) and is associated with inferior frontal activations extending bilaterally to the right hemisphere (e.g., Park et al. 2012). Although bilateral inferior frontal activation might be required when the task becomes more challenging, for instance when it requires extensive semantic processing (Edwards et al. 2005) and decision-making (Gan et al. 2013), inter-hemispheric collaboration could turn out to be detrimental for performance under less demanding contexts (cf. Banich and Belger 1990). Perhaps because of this, lexical decision processes may suffer more from inter-hemispheric interference. Having a strongly left-lateralized FAT may act as a safeguard against this interference from the right homologous region, which is more dedicated to complementary, possibly interfering processes, such as general action control, especially in the visuospatial domain (Dick et al. 2019).

A potential limitation of this study concerns the makeup of the participants, most of whom were multilingual. As multilingualism may affect the lateralization of some cortical regions and white matter tracts (e.g., Felton et al. 2017; Hämäläinen et al. 2017), the results may not generalize to a monolingual population and would need to be specifically investigated. However, bilinguals and multilinguals represent more than half of the world's population (Grosjean 2010). An additional limitation is the relatively low sample size. Despite this, we were still able to show positive results.

The final samples of 20 participants (for the verbal fluency tasks) or 26 (for the lexical decision task) did not allow us, however, to further investigate potentially interesting issues that would have required to split out our sample even further. Such issues include if and how sex differences or the degree of multilingualism modulate the relationship between the FAT and language functions. Future studies should address these and similar research questions.

In conclusion, the present study confirms the importance of investigating inter-hemispheric asymmetries for white matter pathways to understand inter-individual variability in cognitive task performance. Our findings clearly showed that a more left-lateralized frontal aslant tract is beneficial for performance in a verbal task such as lexical decision. Further studies are needed to more directly corroborate whether this pattern is due to a reduction of interference originating from non-language functions implemented in contralateral regions connected by the homologous tract.

Funding This project was funded by the European Research Council Starting Grant LEX-MEA no. 313692 (FP7/2007–2013) to AV.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Research involving human participants and/or animals All procedures performed in studies involving human participants were in accordance with the ethical standards of the Ethics Committee of Scuola Internazionale Superiore di Studi Avanzati—SISSA, Trieste (Prot. # 6772) and Ethics Committee of Azienda Ospedaliera di Padova (Prot. # 2758P), and with the 1964 Helsinki Declaration and its later amendments.

Informed consent Informed consent was obtained from all individual participants included in the study.

References

- Amunts K, Weiss PH, Mohlberg H et al (2004) Analysis of neural mechanisms underlying verbal fluency in cytoarchitectonically defined stereotaxic space—the roles of Brodmann areas 44 and 45. *Neuroimage* 22:42–56. <https://doi.org/10.1016/j.neuroimage.2003.12.031>
- Banich MT, Belger A (1990) Interhemispheric interaction: how do the hemispheres divide and conquer a task? *Cortex* 26:77–94
- Basilakos A, Fillmore PT, Rorden C et al (2014) Regional white matter damage predicts speech fluency in chronic post-stroke aphasia. *Front Hum Neurosci* 8:845. <https://doi.org/10.3389/fnhum.2014.00845>
- Beaulieu C (2002) The basis of anisotropic water diffusion in the nervous system—a technical review. *NMR Biomed* 15:435–455. <https://doi.org/10.1002/nbm.782>

- Belger A, Banich MT (1998) Costs and benefits of integrating information between the cerebral hemispheres: a computational perspective. *Neuropsychology* 12:380–398
- Berryman ML, Kennelly KJ (1992) Letter memory loads change more than visual-field advantage: interhemispheric coupling effects. *Brain Cogn* 18:152–168
- Binder JR, Frost JA, Hammeke TA et al (1997) Human brain language areas identified by functional magnetic resonance imaging. *J Neurosci* 17:353–362
- Budisavljevic S, Dell'Acqua F, Djordjilovic V et al (2017a) The role of the frontal aslant tract and premotor connections in visually guided hand movements. *Neuroimage* 146:419–428. <https://doi.org/10.1016/j.neuroimage.2016.10.051>
- Budisavljevic S, Dell'Acqua F, Zanatto D et al (2017b) Asymmetry and structure of the fronto-parietal networks underlie visuomotor processing in humans. *Cereb Cortex* 27:1532–1544. <https://doi.org/10.1093/cercor/bhv348>
- Carreiras M, Mechelli A, Price CJ (2006) Effect of word and syllable frequency on activation during lexical decision and reading aloud. *Hum Brain Mapp* 27:963–972. <https://doi.org/10.1002/hbm.20236>
- Carreiras M, Riba J, Vergara M et al (2009) Syllable congruency and word frequency effects on brain activation. *Hum Brain Mapp* 30:3079–3088. <https://doi.org/10.1002/hbm.20730>
- Catani M, Mesulan MM (2008) The arcuate fasciculus and the disconnection theme in language and aphasia: history and current state. *Cortex* 44(8):953–961. <https://doi.org/10.1016/j.cortex.2008.04.002>
- Catani M, Jones DK, Ffytche DH (2005) Perisylvian language networks of the human brain. *Ann Neurol* 57:8–16. <https://doi.org/10.1002/ana.20319>
- Catani M, Allin MPG, Husain M et al (2007) Symmetries in human brain language pathways correlate with verbal recall. *Proc Natl Acad Sci USA* 104:17163–17168. <https://doi.org/10.1073/pnas.0702116104>
- Catani M, Dell'Acqua F, Vergani F et al (2012) Short frontal lobe connections of the human brain. *Cortex* 48:273–291. <https://doi.org/10.1016/j.cortex.2011.12.001>
- Catani M, Mesulam MM, Jakobsen E et al (2013) A novel frontal pathway underlies verbal fluency in primary progressive aphasia. *Brain* 136:2619–2628. <https://doi.org/10.1093/brain/awt163>
- Chernoff BL, Teghipco A, Garcea FE et al (2018) A role for the frontal aslant tract in speech planning: a neurosurgical case study. *J Cog Neurosci* 30:752–769. https://doi.org/10.1162/jocn_a_01244
- Crosson B, Sadek JR, Maron L et al (2001) Relative shift in activity from medial to lateral frontal cortex during internally versus externally guided word generation. *J Cogn Neurosci* 13:272–283
- Crosson B, Benefield H, Cato MA et al (2003) Left and right basal ganglia and frontal activity during language generation: contributions to lexical, semantic, and phonological processes. *J Int Neuropsychol Soc* 9:1061–1077. <https://doi.org/10.1017/S135561770397010X>
- Dick AS, Garic D, Graziano P, Tremblay P (2019) The frontal aslant tract (FAT) and its role in speech, language and executive function. *Cortex* 111:148–163. <https://doi.org/10.1016/j.cortex.2018.10.015>
- Edwards JD, Pexman PM, Goodyear BG, Chambers CG (2005) An fMRI investigation of strategies for word recognition. *Brain Res Cogn Brain Res* 24:648–662. <https://doi.org/10.1016/j.cogbrainres.2005.03.016>
- Felton A, Vazquez D, Ramos-Nunez AJ, Greene MR, Macbeth A, Hernandez AE, Chiarello C (2017) Bilingualism influences structural indices of interhemispheric organization. *J Neuroling* 42:1–11. <https://doi.org/10.1016/j.jneuroling.2016.10.004>
- Fiez JA, Balota DA, Raichle ME, Petersen SE (1999) Effects of lexicality, frequency, and spelling-to-sound consistency on the functional anatomy of reading. *Neuron* 24:205–218
- Fujii M, Maesawa S, Motomura K et al (2015) Intraoperative subcortical mapping of a language-associated deep frontal tract connecting the superior frontal gyrus to Broca's area in the dominant hemisphere of patients with glioma. *J Neurosurg* 122:1390–1396. <https://doi.org/10.3171/2014.10.JNS14945>
- Gaillard WD, Hertz-Pannier L, Mott SH et al (2000) Functional anatomy of cognitive development: fMRI of verbal fluency in children and adults. *Neurology* 54:180–185
- Gan G, Büchel C, Isel F (2013) Effect of language task demands on the neural response during lexical access: a functional magnetic resonance imaging study. *Brain Behav* 3:402–416. <https://doi.org/10.1002/brb3.133>
- Gold BT, Powell DK, Xuan L et al (2007) Speed of lexical decision correlates with diffusion anisotropy in left parietal and frontal white matter: evidence from diffusion tensor imaging. *Neuropsychologia* 45:2439–2446. <https://doi.org/10.1016/j.neuropsychologia.2007.04.011>
- Gotts SJ, Jo HJ, Wallace GL et al (2013) Two distinct forms of functional lateralization in the human brain. *Proc Natl Acad Sci USA* 110:E3435–3444. <https://doi.org/10.1073/pnas.1302581110>
- Grosjean F (2010) *Bilingual: Life and reality*. Harvard University Press, Cambridge. <https://doi.org/10.4159/9780674056459>
- Hämäläinen S, Sairanen V, Leminen A, Lehtonen M (2017) Bilingualism modulates the white matter structure of language-related pathways. *NeuroImage* 152:249–257. <https://doi.org/10.1016/j.neuroimage.2017.02.081>
- Höller-Wallscheid MS, Thier P, Pomper JK, Lindner A (2017) Bilateral recruitment of prefrontal cortex in working memory is associated with task demand but not with age. *PNAS* 114:E830–E839. <https://doi.org/10.1073/pnas.1601983114>
- Howells H, Thiebaut de Schotten M, Dell'Acqua F, Beyh A, Zappalà G, Leslie A, Simmons A, Murphy DG, Catani M (2018) Frontoparietal tracts linked to lateralized hand preference and manual specialization. *Cereb Cortex* 28(7):2482–2494. <https://doi.org/10.1093/cercor/bhy040>
- Ibrahim R, Eviatar Z (2012) The contribution of the two hemispheres to lexical decision in different languages. *Behav Brain Funct* 8:3. <https://doi.org/10.1186/1744-9081-8-3>
- James JS, Kumari SR, Sreedharan RM et al (2015) Analyzing functional, structural, and anatomical correlation of hemispheric language lateralization in healthy subjects using functional MRI, diffusion tensor imaging, and voxel-based morphometry. *Neuro India* 63:49–57. <https://doi.org/10.4103/0028-3886.152634>
- Kemerdere R, de Champfleury NM, Deverdun J et al (2016) Role of the left frontal aslant tract in stuttering: a brain stimulation and tractographic study. *J Neurol* 263:157–167. <https://doi.org/10.1007/s00415-015-7949-3>
- Kinoshita M, de Champfleury NM, Deverdun J et al (2015) Role of fronto-striatal tract and frontal aslant tract in movement and speech: an axonal mapping study. *Brain Struct Funct* 220:3399–3412. <https://doi.org/10.1007/s00429-014-0863-0>
- Kronfeld-Duenias V, Amir O, Ezrati-Vinacour R et al (2016) The frontal aslant tract underlies speech fluency in persistent developmental stuttering. *Brain Struct Funct* 221:365–381. <https://doi.org/10.1007/s00429-014-0912-8>
- Lebel C, Beaulieu C (2009) Lateralization of the arcuate fasciculus from childhood to adulthood and its relation to cognitive abilities in children. *Hum Brain Mapp* 30:3563–3573. <https://doi.org/10.1002/hbm.20779>
- Li M, Zhang Y, Song L et al (2017) Structural connectivity subserving verbal fluency revealed by lesion-behavioral mapping in stroke patients. *Neuropsychologia* 101:85–96. <https://doi.org/10.1016/j.neuropsychologia.2017.05.008>
- Mandelli ML, Caverzasi E, Binney RJ et al (2014) Frontal white matter tracts sustaining speech production in primary progressive

- aphasia. *J Neurosci* 34:9754–9767. <https://doi.org/10.1523/JNEUROSCI.3464-13.2014>
- Martino J, De Lucas EM (2014) Subcortical anatomy of the lateral association fascicles of the brain: a review. *Clin Anat* 27(4):563–569. <https://doi.org/10.1002/ca.22321>
- Mohr B, Pulvermüller F, Zaidel E (1994) Lexical decision after left, right and bilateral presentation of function words, content words and non-words: evidence for interhemispheric interaction. *Neuropsychologia* 32:105–124
- Ocklenburg S, Hugdahl K, Westerhausen R (2013) Structural white matter asymmetries in relation to functional asymmetries during speech perception and production. *Neuroimage* 83:1088–1097. <https://doi.org/10.1016/j.neuroimage.2013.07.076>
- Ocklenburg S, Friedrich P, Güntürkün O, Genç E (2016) Intrahemispheric white matter asymmetries: the missing link between brain structure and functional lateralization? *Rev Neurosci* 27:465–480. <https://doi.org/10.1515/revneuro-2015-0052>
- Oldfield RC (1971) The assessment and analysis of handedness: the Edinburgh Inventory. *Neuropsychologia* 9:97–113
- Park HRP, Badzakova-Trajkov G, Waldie KE (2012) Brain activity in bilingual developmental dyslexia: an fMRI study. *Neurocase* 18:286–297. <https://doi.org/10.1080/13554794.2011.588182>
- Paulesu E, McCrory E, Fazio F et al (2000) A cultural effect on brain function. *Nat Neurosci* 3:91–96. <https://doi.org/10.1038/71163>
- Rosso C, Valabregue R, Arbizu C et al (2014) Connectivity between right inferior frontal gyrus and supplementary motor area predicts after-effects of right frontal cathodal tDCS on picture naming speed. *Brain Stimul* 7:122–129. <https://doi.org/10.1016/j.brs.2013.08.007>
- Schlösser R, Hutchinson M, Joseffer S et al (1998) Functional magnetic resonance imaging of human brain activity in a verbal fluency task. *J Neurol Neurosurg Psychiatry* 64:492–498. <https://doi.org/10.1136/jnnp.64.4.492>
- Schwieter JW, Sunderman G (2011) Inhibitory control processes and lexical access in trilingual speech production. *Linguist Approaches Bilingualism* 1:391–412. <https://doi.org/10.1075/lab.1.4.02sch>
- Sierpowska J, Gabarrós A, Fernandez-Coello A et al (2015) Morphological derivation overflow as a result of disruption of the left frontal aslant white matter tract. *Brain Lang* 142:54–64. <https://doi.org/10.1016/j.bandl.2015.01.005>
- Smirni D, Turriziani P, Mangano GR et al (2017) Modulating phonemic fluency performance in healthy subjects with transcranial magnetic stimulation over the left or right lateral frontal cortex. *Neuropsychologia* 102:109–115. <https://doi.org/10.1016/j.neuropsychologia.2017.06.006>
- Thiebaut de Schotten M, Dell’Acqua F, Forkel SJ, Simmons A, Vergani F, Murphy DG, Catani M (2011a) A lateralized brain network for visuospatial attention. *Nat Neurosci* 14(10):1245–1246. <https://doi.org/10.1038/nn.2905>
- Thiebaut de Schotten M, Ffytche DH, Bizzi A et al (2011b) Atlasing location, asymmetry and inter-subject variability of white matter tracts in the human brain with MR diffusion tractography. *Neuroimage* 54:49–59. <https://doi.org/10.1016/j.neuroimage.2010.07.055>
- Vassal F, Boutet C, Lemaire J-J, Nuti C (2014) New insights into the functional significance of the frontal aslant tract: an anatomic-functional study using intraoperative electrical stimulations combined with diffusion tensor imaging-based fiber tracking. *Br J Neurosurg* 28:685–687. <https://doi.org/10.3109/02688697.2014.889810>
- Yin X, Han Y, Ge H et al (2013) Inferior frontal white matter asymmetry correlates with executive control of attention. *Hum Brain Mapp* 34:796–813. <https://doi.org/10.1002/hbm.21477>

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.