



Correction to: Differences in prothrombotic response between the uninterrupted and interrupted apixaban therapies in patients undergoing cryoballoon ablation for paroxysmal atrial fibrillation: a randomized controlled study

Monami Ando^{1,2} · Yasuya Inden² · Yukihiro Yoshida¹ · Akinori Sairaku³ · Satoshi Yanagisawa² · Hirohiko Suzuki¹ · Ryo Watanabe¹ · Masaki Takenaka¹ · Mayuho Maeda¹ · Toyoaki Murohara²

Published online: 13 May 2019
© Springer Japan KK, part of Springer Nature 2019

Correction to: Heart and Vessels
<https://doi.org/10.1007/s00380-019-01370-9>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

In the original publication of the article, the Table 2 was published incorrectly. The correct Table 2 is as below.

Table 2 Procedural findings during ablation

	Group 1 (uninterrupted apixaban)	Group 2 (interrupted apixaban)	<i>p</i> value
Procedure time (min)	121.9 ± 18.2	128.5 ± 23.2	0.17
Fluoroscopic time (min)	24.5 ± 8.2	26.3 ± 6.3	0.12
Touch-up ablation	10 (31.3)	25 (38.5)	0.49

The original article can be found online at <https://doi.org/10.1007/s00380-019-01370-9>.

✉ Yasuya Inden
inden@med.nagoya-u.ac.jp

¹ Department of Cardiology, Cardiovascular Center, Nagoya Daini Red Cross Hospital, Nagoya, Japan

² Department of Cardiology, Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku, Nagoya, Aichi 466-8550, Japan

³ Department of Cardiology, Onomichi General Hospital, Hiroshima, Japan