



Neutropenia during frontline treatment of advanced Hodgkin lymphoma: Incidence, risk factors, and management

Anna Sureda^{a,*}, Eva Domingo-Domenech^a, Ashish Gautam^b

^a Clinical Hematology Department, Institut Català d'Oncologia – Hospitalet, Hospital Duran i Reynals, Gran Via de l'Hospitalet, 199 – 203, 08908, Barcelona, Spain

^b Millennium Pharmaceuticals, Inc. (a wholly owned subsidiary of Takeda Pharmaceutical Company Limited), 350 Massachusetts Avenue, Cambridge, MA, 02139, United States

ARTICLE INFO

Keywords:

Hodgkin lymphoma
Granulocyte colony-stimulating factor
Febrile neutropenia

ABSTRACT

Neutropenia, specifically febrile neutropenia (FN), can have profound sequelae (infection, hospitalization, mortality), and the risk of its development differs across chemotherapy regimens/according to patient characteristics. We conducted a comprehensive literature review regarding neutropenia in frontline treatment of adults with advanced Hodgkin lymphoma. Guidelines state primary prophylaxis (PP) with colony-stimulating factors (CSFs) should be implemented when the risk of FN is $\geq 20\%$; CSF PP is given with standard-of-care escalated BEACOPP, but the risk of FN with standard-of-care ABVD does not necessitate routine PP. Notably, the risk of neutropenia (including FN) is higher in clinical practice versus clinical studies, and physicians overestimate their adherence to CSF guidelines. ECHELON-1 demonstrated higher FN rates with brentuximab vedotin plus AVD (A + AVD) compared with ABVD (19% vs 8%) and led to the recommendation of PP with granulocyte-CSF (G-CSF) for all A + AVD patients, highlighting the importance of readjusting our risk-assessment thinking as standard backbone regimens are modified.

1. Introduction and methods

Advances in the multi-agent chemotherapy regimens used in advanced-stage Hodgkin lymphoma have dramatically changed disease prognosis. The most commonly used regimens now include ABVD (doxorubicin, bleomycin, vinblastine, and dacarbazine) and BEACOPP (bleomycin, etoposide, doxorubicin, cyclophosphamide, vincristine, procarbazine, and prednisone). Standard or 'baseline' BEACOPP is often 'escalated' utilizing increasing doses of etoposide, doxorubicin, and cyclophosphamide (Diehl et al., 2003; Engert, 2016). BEACOPP is generally considered to have a higher efficacy than ABVD, but is associated with a higher toxicity burden (Engert, 2016). Consequently, research is ongoing to incorporate new drugs into these standard ABVD and BEACOPP backbones to improve efficacy and/or reduce toxicity.

A common complication of myelosuppressive multi-agent chemotherapy regimens such as ABVD and BEACOPP is neutropenia. While neutropenia itself is not a troublesome adverse event (AE), severe neutropenia and, in particular, febrile neutropenia (FN) potentially have profound sequelae (Aapro et al., 2011; Klastersky et al., 2016; NCCN, 2019a, 2019c; Smith et al., 2015). There are two main strategies for reducing the occurrence and impact of neutropenia: (1) providing prophylactic myelopoietic support with growth factors (colony stimulating

factors [CSFs]) and/or (2) lowering myelotoxicity through initial choice of chemotherapeutic agent or modifying the chemotherapeutic dose, using dose delays and reductions, and missing doses. Furthermore, the risk of neutropenia is an important consideration in the context of the evolving treatment of Hodgkin lymphoma. We therefore conducted a comprehensive review of the literature regarding neutropenia and its management in the frontline treatment of adults with advanced Hodgkin lymphoma.

We used the Medline database (via PubMed) to search for English-language articles published since database inception through to August 1, 2017 using the following two search strategies: 1. ("neutropaenia" OR "neutropenia" OR "white blood cell*") AND ("Hodgkin lymphoma" OR "Hodgkin's lymphoma" OR "Hodgkin's disease" OR "Hodgkin disease") – 1300 search results; 2. ("colony-stimulating factor"[All Fields] OR "colony-stimulating factors"[All Fields]) AND ("guideline"[Publication Type] OR "guidelines as topic"[MeSH Terms] OR "guidelines"[All Fields]) OR "systematic review"[All Fields]) AND ("neoplasms"[MeSH Terms] OR "neoplasms"[All Fields] OR "cancer"[All Fields]) OR ("lymphoma"[MeSH Terms] OR "lymphoma"[All Fields]) OR ("neoplasms"[MeSH Terms] OR "neoplasms"[All Fields] OR "malignancies"[All Fields]) – 371 search results. We manually reviewed all retrieved titles and abstracts for relevance, and further assessed full papers that we judged appropriate for inclusion in this review.

* Corresponding author at: Head of the Hematology Department and Hematopoietic Stem Cell Transplant Programme, Institut Català d'Oncologia - Hospital Duran i Reynals, Gran Via de l'Hospitalet, 199 – 203, 08908, Barcelona, Spain.

E-mail addresses: asureda@iconcologia.net (A. Sureda), edomingo@iconcologia.net (E. Domingo-Domenech), Ashish.Gautam@Takeda.com (A. Gautam).

<https://doi.org/10.1016/j.critrevonc.2019.03.016>

Received 22 March 2018; Received in revised form 15 January 2019; Accepted 26 March 2019

1040-8428/ © 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1
Overall risk of severe and febrile neutropenia (or associated consequences) in prospective studies of patients with Hodgkin lymphoma.

| Study reference | Study population and characteristics | Reported incidence of neutropenia and associated complications | G-CSF permissible |
|---------------------------|---|--|--|
| Diehl et al. (2003) | Randomized controlled trial 1195 patients with advanced Hodgkin lymphoma (stage IIB, III, and IV) | Grade 4 leukopenia: COPP-ABVD (n = 260; 19%); BEACOPP (n = 469; 37%); BEACOPP _{escalated} (n = 466; 90%) Grade 4 infection: COPP-ABVD (n = 260; 1%); BEACOPP (n = 469; 3%); BEACOPP _{escalated} (n = 466; 8%) | Not specified |
| Federico et al. (2009) | Randomized controlled trial 317 patients with advanced Hodgkin lymphoma (stage IIB, III, and IV) | Grade 3/4 neutropenia: ABVD (n = 99; 34%); BEACOPP (n = 98; 54%); CEC (n = 98; 48%) Grade 3/4 infection: ABVD (n = 99; 2%); BEACOPP (n = 98; 14%); CEC (n = 98; 4%) | G-CSF given in addition to BEACOPP treatment (300 µg total, subcutaneously) |
| Cocorocchio et al. (2010) | Prospective single-arm study 82 patients with advanced Hodgkin lymphoma (stage IIA [bulky], IIB, III, and IV) | Grade 3/4 neutropenia: ChlVPP/ABVVP (n = 76; 32%) Grade 3/4 infection: ChlVPP/ABVVP (n = 76; 8%) FN: ChlVPP/ABVVP (n = 76; 8%) | Not specified |
| Younes et al. (2012a) | Prospective single-arm study 78 patients with advanced Hodgkin lymphoma (stage II [bulky], III, or IV) | Grade 3/4 granulocytopenia: Rituximab + ABVD (n = 78; 15% and 8%) | Not specified |
| Younes et al. (2013) | Randomized controlled trial 51 patients with advanced Hodgkin lymphoma (stage IIA [bulky], IIB, III, and IV) | Grade 3/4 neutropenia: A + ABVD (n = 25; 80%); A + AVD (n = 26; 77%) | Not specified |
| Gordon et al. (2013) | Randomized controlled trial 854 patients with advanced Hodgkin lymphoma (stage III and IV) | Grade 3/4 FN: A + ABVD (n = 25; 20%); A + AVD (n = 26; 8%) Grade 3 lymphopenia: Stanford V (n = 426; 78%); ABVD (n = 428; 42%) Grade 3 and 4 leukocytopenia: Stanford V (n = 426; 36% and 19%); ABVD (n = 428; 28% and 5%) | Not specified |
| Russo et al. (2014) | Prospective single-arm study 82 patients with advanced Hodgkin lymphoma (stage IIB, III, and IV) | Grade 4 neutropenia: ABVD _{DD-DI} (n = 82; 10%) Grade 3 febrile neutropenic infection: ABVD _{DD-DI} (n = 82; 6%) | All patients received primary G-CSF as support for ABVD. Pegylated G-CSF was not allowed in the study. |
| Carde et al. (2016) | Randomized controlled trial 549 patients with advanced Hodgkin lymphoma (stage III and IV) | Grade 4 neutropenia: ABVD ₈ (n = 275; 32%); BEACOPP ₄₊₄ (n = 274; 65%) Grade 4 FN: ABVD ₈ (n = 275; 6%); BEACOPP ₄₊₄ (n = 274; 34%) | Prophylactic G-CSF was mandatory with BEACOPP _{escalated} |
| Connors et al. (2018) | Randomized controlled trial 1334 patients with previously untreated advanced Hodgkin lymphoma (stage III and stage IV) | Grade 3/4 neutropenia: ABVD (n = 659; 39%); A + AVD (n = 662; 54%) FN: ABVD (n = 659; 8%); A + AVD (n = 662; 19%) | Prophylactic G-CSF was recommended for patients received A + AVD |

ABVD, doxorubicin, bleomycin, vinblastine, dacarbazine; ABVVP, etoposide, doxorubicin, bleomycin and vincristine; AVD, doxorubicin, vinblastine, dacarbazine; A + AVD, brentuximab vedotin + AVD; A + ABVD, brentuximab vedotin + ABVD; BEACOPP, bleomycin, etoposide, doxorubicin, cyclophosphamide, vincristine, procarbazine, and prednisone; CEC, cyclophosphamide, lomustine, vindesine, melphalan, prednisone, epidoxirubicin, vincristine, procarbazine, vinblastine, and bleomycin; ChlVPP, chlorambucil, vinblastine, procarbazine and prednisone; COPP, cyclophosphamide, vincristine, procarbazine, prednisone.

Here, we summarize the literature, focusing on the incidence of, risk factors for, and sequelae of neutropenia, in particular FN, and its management using CSF prophylaxis. We also evaluate how these aspects may potentially be affected within the context of the evolving treatment algorithm for Hodgkin lymphoma.

Incidences of neutropenic events in Hodgkin lymphoma (including FN if reported) are shown in Table 1. In general, based on the original studies, guidelines have classified BEACOPP as a chemotherapeutic regimen associated with a high risk of FN, and the risk of FN with ABVD is regarded as low-intermediate (depending on age) (Aapro et al., 2011; NCCN, 2019a).

1.1. Neutropenia, febrile neutropenia and its consequences

Neutropenia is a decrease in circulating neutrophils in the non-marginal pool and is defined in terms of the absolute neutrophil count (ANC), with a healthy person having an ANC of $1.5\text{--}8.0 \times 10^9$ cells/L. The severity of neutropenia is classified as mild ($1.0\text{--}1.5 \times 10^9$ cells/L), moderate ($0.5\text{--}1.0 \times 10^9$ cells/L), or severe ($< 0.5 \times 10^9$ cells/L). Due to the infection-fighting role of neutrophils, physicians are also concerned about the duration of neutropenia and the presence of fever. FN is defined as an ANC of [or expected to fall to] $< 0.5 \times 10^9$ cells/L in the presence of an oral temperature of $> 38.3^\circ\text{C}$ or $> 38.0^\circ\text{C}$ for 1 h (Klastersky et al., 2016; NCCN, 2019c).

The risk of developing neutropenia and/or FN is dependent on a variety of factors including the type of cancer, the chemotherapy regimen used, and patient characteristics, including a low neutrophil count at baseline (Lyman et al., 2014).

Certain chemotherapy agents and regimens are more myelosuppressive than others. For example, anthracyclines such as doxorubicin and alkylators such as cyclophosphamide, which are used in the ABVD

and BEACOPP regimens in the frontline treatment of Hodgkin lymphoma, are considered particularly myelosuppressive. Consequently, the risk of neutropenia and FN is dependent on the components used and the intensity (drug dose delivered per time unit) of the chemotherapeutic regimen (Lyman et al., 2014; Lyman, 2009; Lyman and Kuderer, 2003). In Hodgkin lymphoma, data from clinical studies demonstrate the difficulties in obtaining a comprehensive view of risk of neutropenia with specific regimens due to inconsistencies in reporting (grade 3/4 neutropenia vs FN vs febrile [neutropenic/non-neutropenic] infection) (Carde et al., 2016; Viviani et al., 2011; Fossa et al., 2012).

It is also important to recognize that our true understanding of the risk of FN is complicated by the administration of myelopoietic support per clinical study protocols and/or medical society and institutional guidelines. Additionally, studies have found differences in the rate of CSF use not only with different regimens, but also across different regions (Schwenkglens et al., 2010).

Myelosuppression increases with age (Repetto et al., 2003) and, consequently, the risk of neutropenia is increased in elderly patients (> 65 years) receiving chemotherapy (Aapro et al., 2011; Klastersky et al., 2016; NCCN, 2019a; Smith et al., 2015). The substantial contribution of age to therapeutic outcomes is such that several studies have investigated the effects of chemotherapy specifically in the elderly population with Hodgkin lymphoma (Ballova et al., 2005; Stamatoullas et al., 2015), and management guidelines have made specific recommendations regarding the elderly population and management of neutropenia (Repetto et al., 2003; NCCN, 2019b). For example, the rate of grade 4 or higher neutropenia in patients with Hodgkin lymphoma treated with ABVD or the Stanford V regimen (doxorubicin, vinblastine, mechlorethamine, vincristine, bleomycin, etoposide, and prednisone) was 64% in those aged ≥ 60 years versus 38% in patients aged < 60

years (Evens et al., 2013). Importantly, US and European guidelines for the management of disease recommend that escalated BEACOPP is either used with caution, or not at all, in individuals > 60 years of age (NCCN, 2019b; Eichenauer et al., 2014).

Other suggested risk factors for FN include advanced disease and comorbidities (both closely associated with age) (Smith et al., 2015; Lyman et al., 2014), bone marrow involvement (Moreau et al., 2009), poor performance status (NCCN, 2019a), and female gender (Scott et al., 2003).

In addition to risk of infection and fever, neutropenia also can increase the risk for hospitalizations (especially in patients with HL), and increases mortality (Dulisse et al., 2013). Patients with cancer receiving a reduced dose of chemotherapy due to neutropenic complications especially have a lower chance of remission and survival.

2. Management guidelines for primary prophylaxis with CSFs in Hodgkin lymphoma

2.1. CSFs

CSFs enhance the proliferation, maturation, and release of neutrophils into the bloodstream, resulting in a dose-dependent increase in circulating neutrophils. Granulocyte CSFs (G-CSFs) stimulate the growth of granulocyte colonies. G-CSFs available for the prevention of treatment-related FN are filgrastim, tbo-filgrastim (only available in the US), pegfilgrastim (a PEGylated form of G-CSF), lenograstim (a glycosylated form of G-CSF, only available in Europe), and filgrastim and pegfilgrastim biosimilars (Aapro et al., 2011; Klastersky et al., 2016; NCCN, 2019a; Smith et al., 2015). Short-acting G-CSFs filgrastim and lenograstim are administered daily (preferably subcutaneously) until post-nadir ANC recovery to normal or near-normal levels, and long-acting pegfilgrastim is administered as a single subcutaneous dose per chemotherapy cycle. It is generally understood that short-acting and long-acting G-CSFs are bioequivalent and clinical guidelines in both Europe and the US advocate the use of either (Aapro et al., 2011; Klastersky et al., 2016; NCCN, 2019a; Smith et al., 2015; Wang et al., 2015). A meta-analysis of 3493 patients in 17 randomized trials showed CSFs to be effective at reducing the incidence of FN, infection-related mortality, and all-cause mortality during chemotherapy treatment (Kuderer et al., 2007).

2.2. Key guidelines

The first guidelines on the use of CSFs were released in 2006 by ASCO (Smith et al., 2006); these were most recently updated in 2015 (Smith et al., 2015). Guidelines have also been released in the US by the National Comprehensive Cancer Network® (NCCN®), which releases regular updates (NCCN, 2019a, 2019b). In Europe, similar management guidelines have been released by both EORTC and ESMO (Aapro et al., 2011; Klastersky et al., 2016).

Guidelines are consistent in their recommendation for routine primary prophylaxis with G-CSF in patients receiving chemotherapy for whom there is a > 20% risk for FN based on patient-, disease-, and treatment-related factors (Aapro et al., 2011; Klastersky et al., 2016; NCCN, 2019a; Smith et al., 2015). As the risk of FN is greatest during the first course of chemotherapy, primary prophylaxis (use with first course of chemotherapy onwards) is recommended over secondary prophylaxis (use after an episode of severe or FN in the preceding course) (Klastersky et al., 2016). Additionally, since a previous episode of FN predisposes to further occurrence, it is important that the risk of FN and related complications is assessed at each cycle, and, if appropriate, secondary prophylaxis with CSF is initiated (Aapro et al., 2011; Klastersky et al., 2016; NCCN, 2019a, 2019c; Smith et al., 2015).

2.2.1. Use of CSFs with current standards of care in frontline advanced Hodgkin lymphoma

Based on the guidelines, standard current practice recommends routine primary prophylaxis with CSF is given when patients receive BEACOPP. For patients receiving chemotherapy regimens that are not considered to be high risk (e.g. ABVD), the decision regarding use of CSF prophylaxis can be confounded by the presence of additional risk factors that predispose patients to FN (Weycker et al., 2015). Prophylaxis with CSF is not recommended for routine use with ABVD, due to observed increased bleomycin-induced pulmonary toxicity (NCCN, 2019b; Martin et al., 2005). Two separate studies have confirmed that ABVD can be administered without safety concerns at the full-dose intensity without any CSF (Boleti and Mead, 2007; Evens et al., 2007).

In the early studies of BEACOPP versus ABVD (and cyclophosphamide, vincristine, procarbazine, and prednisone [COPP]), Diehl et al. reported G-CSF administration with 4% of COPP cycles, 14% of ABVD cycles, 10% of baseline BEACOPP courses, and 90% of escalated BEACOPP courses (Diehl et al., 1998). In 2007, Evens et al. published a retrospective review of all newly diagnosed patients with Hodgkin lymphoma treated at their institute in the US between 1996 and 2005. Among patients treated without empiric G-CSF, only 3/682 ABVD treatments were complicated by FN (Evens et al., 2007). The overall median normalized dose intensity for the 59 patients who received all ABVD cycles without G-CSF was 99.1%, leading the authors to conclude that ABVD may be fully and effectively administered without G-CSF support, regardless of a patient's ANC.

The influence of intensified treatment regimens on incidence of neutropenia and FN varies between regimens. With an intensified 6-cycle ChlVPP/ABVVP regimen (chlorambucil, vinblastine, procarbazine, doxorubicin, bleomycin, vincristine, and etoposide), Cocorocchio et al. reported grade 3/4 neutropenia in 32% of patients and FN in 8%, representing a considerable reduction in frequency compared with standard ChlVPP/ABVVP (Cocorocchio et al., 2010). In contrast, a retrospective study showed that the incidence of grade 3/4 neutropenia increased substantially with dose-dense ABVD compared with standard ABVD (58 vs 39%) (Tao et al., 2014). Despite routine CSF primary prophylaxis, Russo et al. reported grade 3 febrile neutropenic infection in 6% of patients treated with dose-dense and dose-intense ABVD (Russo et al., 2014).

2.2.2. Use of CSFs with new agents in frontline advanced Hodgkin lymphoma

Brentuximab vedotin is a new agent being investigated in the frontline treatment of advanced Hodgkin lymphoma. Brentuximab vedotin is a CD30-directed antibody-drug conjugate currently indicated for the treatment of patients with classical Hodgkin lymphoma after failure of autologous hematopoietic stem cell transplantation (auto-SCT), after failure of at least two prior multi-agent chemotherapy regimens in patients who are not auto-SCT candidates, and for treatment of patients with classical Hodgkin lymphoma at high risk of relapse or progression as post-auto-SCT consolidation. Thus, our experience in Hodgkin lymphoma in the clinic has largely been based on the use of brentuximab vedotin as a monotherapy in patients with prior treatments. In the pivotal phase 2 study of brentuximab vedotin in patients with relapsed or refractory Hodgkin lymphoma, 20% of patients had grade 3/4 neutropenia, but there were no cases of FN reported in the 102 patients enrolled (Younes et al., 2012b). All 102 patients had undergone auto-SCT, 66% had received prior radiation therapy, and the median number of prior chemotherapy regimens excluding auto-SCT was 3.5 (range, 1–13).

In a phase I study to assess the safety of frontline A + AVD and A + ABVD combinations in patients with advanced classical Hodgkin lymphoma, Q2W 1.2 mg/kg brentuximab vedotin combined with ABVD was associated with pulmonary toxicity; whereas A + AVD was considered tolerable (Younes et al., 2013; Connors et al., 2017). In the study, the use of CSF was permitted in accordance with institutional standard of care and if grade 3/4 neutropenia was observed CSF

support was used as prophylaxis in subsequent cycles. The authors reported that CSF was given to 84% of patients. The rate of FN was 20% with brentuximab vedotin in combination with ABVD and 8% with brentuximab vedotin in combination with AVD; none of the patients with FN had received previous CSF prophylaxis (Younes et al., 2013).

ECHELON-1, a large, international, open-label, multicenter, randomized phase 3 trial, compared brentuximab vedotin plus doxorubicin, vinblastine, and dacarbazine (A + AVD) with ABVD as frontline therapy in 1334 patients with stage III or IV classical Hodgkin lymphoma (Connors et al., 2018). FN was reported in 19% of patients receiving A + AVD and in 8% of patients receiving ABVD. Discussion with the independent data and safety monitoring committee (after 75% of enrollment was complete) led to the recommendation of primary prophylaxis with G-CSF for patients who were yet to be enrolled and who would receive the A + AVD regimen, owing to the higher incidence of FN in that group. In the A + AVD group, the incidence of FN was lower among the 83 patients who received primary prophylaxis with G-CSF (defined as use of G-CSF by day 5 of treatment) than among those who did not (11% vs 21%).

Further exploratory analyses of the ECHELON-1 clinical trial (Connors et al., 2018) showed that in patients receiving G-CSF primary prophylaxis, brentuximab vedotin dose reductions (20% vs 26%) and dose delays (35% vs 49%) were decreased, and hospitalization rates reduced (29% vs 38% patients with at least one hospitalization) compared with patients who received no G-CSF primary prophylaxis, thereby maintaining the dose density of the A + AVD regimen given to patients (Straus et al., 2018). These data suggest that the concomitant administration of G-CSF within the first five days of treatment may improve the efficacy of A + AVD, as shown by reduced risk of modified progression-free survival events by 25% with A + AVD with G-CSF compared with A + AVD without G-CSF and by 42% compared with ABVD (Straus et al., 2018).

3. Conclusions

Chemotherapy-induced FN is a potentially fatal complication of cancer treatment, and the prevention of FN reduces hospital admissions, antibiotic usage, and the need for dose reductions or delays in chemotherapy administration. According to guidelines, in patients with Hodgkin lymphoma who are treated with BEACOPP, primary prophylaxis with CSFs should routinely be administered. ABVD, however, is considered a low to medium-risk regimen, and primary prophylaxis is only appropriate in a subgroup of patients with patient-related risk factors. As new regimens are established for the frontline treatment of Hodgkin lymphoma, clinicians must reassess the risk of neutropenia and provide patients with CSF support accordingly. Results from ECHELON-1 demonstrate that primary prophylaxis with G-CSF appeared to mitigate the increased risk of FN and its sequelae in the subgroup of A + AVD patients who received primary prophylaxis, allowing delivery of the full dose density of A + AVD, and potentially improving the efficacy of A + AVD compared with that seen in patients who do not receive G-CSF prophylaxis.

Conflicts of interest

Anna Sureda reports consultancy, advisory boards, and travel grants from Takeda.

Eva Domingo-Domenech has no conflicts of interest to disclose.

Ashish Gautam is an employee of Millennium Pharmaceuticals Inc., a wholly owned subsidiary of Takeda Pharmaceutical Company Ltd.

Funding

The development of this publication was supported by Millennium Pharmaceuticals Inc., a wholly owned subsidiary of Takeda Pharmaceutical Company Limited.

Author contributions

AS and ED-D contributed to the conception and design of the article. AS and AG analyzed and interpreted data. All authors contributed to drafting and critical revision of the article, and approved the final version to be published.

Acknowledgments

The authors would like to acknowledge Sarah Feeny and Jenny Wilkinson of FireKite, an Ashfield company, part of UDG Healthcare plc, for writing support during the development of this manuscript, which was funded by Millennium Pharmaceuticals, Inc., and complied with Good Publication Practice 3 ethical guidelines (Battisti WP, et al. *Ann Intern Med* 2015;163:461–4).

References

- Aapro, M.S., Bohlius, J., Cameron, D.A., Dal Lago, L., Donnelly, J.P., Kearney, N., et al., 2011. 2010 update of EORTC guidelines for the use of granulocyte-colony stimulating factor to reduce the incidence of chemotherapy-induced febrile neutropenia in adult patients with lymphoproliferative disorders and solid tumours. *Eur. J. Cancer* 47, 8–32.
- Ballova, V., Ruffer, J.U., Haverkamp, H., Pfister, B., Muller-Hermelink, H.K., Duhmke, E., et al., 2005. A prospectively randomized trial carried out by the German Hodgkin Study Group (GHS) for elderly patients with advanced Hodgkin's disease comparing BEACOPP baseline and COPP-ABVD (study HD9elderly). *Ann. Oncol.* 16, 124–131.
- Boleti, E., Mead, G.M., 2007. ABVD for Hodgkin's lymphoma: full-dose chemotherapy without dose reductions or growth factors. *Ann. Oncol.* 18, 376–380.
- Carde, P., Karrasch, M., Fortpied, C., Brice, P., Khaled, H., Casasnovas, O., et al., 2016. Eight cycles of ABVD versus four cycles of BEACOPPescalated plus four cycles of BEACOPPbaseline in stage III to IV, international prognostic score ≥ 3 , high-risk Hodgkin lymphoma: first results of the phase III EORTC 20012 intergroup trial. *J. Clin. Oncol.* 34, 2028–2036.
- Cocorocchio, E., Vanazzi, A., Bassi, S., Peccatori, F., Antoniotti, P., Gigli, F., et al., 2010. Intensified CHVVP/ABVVP chemotherapy regimen and pegfilgrastim support in advanced Hodgkin lymphoma. *Ecancermedicallscience* 4, 184.
- Connors, J.M., Ansell, S.M., Fanale, M., Park, S.I., Younes, A., 2017. Five-year follow-up of brentuximab vedotin combined with ABVD or AVD for advanced-stage classical Hodgkin lymphoma. *Blood* 130, 1375–1377.
- Connors, J.M., Jurczak, W., Straus, D.J., Ansell, S.M., Kim, W.S., Gallamini, A., et al., 2018. Brentuximab vedotin with chemotherapy for stage III or IV Hodgkin's lymphoma. *N. Engl. J. Med.* 378, 331–344.
- Diehl, V., Franklin, J., Hasenclever, D., Tesch, H., Pfreundschuh, M., Lathan, B., et al., 1998. BEACOPP, a new dose-escalated and accelerated regimen, is at least as effective as COPP/ABVD in patients with advanced-stage Hodgkin's lymphoma: interim report from a trial of the German Hodgkin's Lymphoma Study Group. *J. Clin. Oncol.* 16, 3810–3821.
- Diehl, V., Franklin, J., Pfreundschuh, M., Lathan, B., Paulus, U., Hasenclever, D., et al., 2003. Standard and increased-dose BEACOPP chemotherapy compared with COPP-ABVD for advanced Hodgkin's disease. *N. Engl. J. Med.* 348, 2386–2395.
- Dulisse, B., Li, X., Gayle, J.A., Barron, R.L., Ernst, F.R., Rothman, K.J., et al., 2013. A retrospective study of the clinical and economic burden during hospitalizations among cancer patients with febrile neutropenia. *J. Med. Econ.* 16, 720–735.
- Eichenauer, D.A., Engert, A., Andre, M., Federico, M., Illidge, T., Hutchings, M., et al., 2014. Hodgkin's lymphoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann. Oncol.* 25 (Suppl. 3), iii70–5.
- Engert, A., 2016. ABVD or BEACOPP for advanced Hodgkin lymphoma. *J. Clin. Oncol.* 34, 1167–1169.
- Evens, A.M., Gilley, J., Ortiz, T., Gounder, M., Hou, N., Rademaker, A., et al., 2007. G-CSF is not necessary to maintain over 99% dose-intensity with ABVD in the treatment of Hodgkin lymphoma: low toxicity and excellent outcomes in a 10-year analysis. *Br. J. Haematol.* 137, 545–552.
- Evens, A.M., Hong, F., Gordon, L.I., Fisher, R.I., Bartlett, N.L., Connors, J.M., et al., 2013. The efficacy and tolerability of adriamycin, bleomycin, vinblastine, dacarbazine and Stanford V in older Hodgkin lymphoma patients: a comprehensive analysis from the North American intergroup trial E2496. *Br. J. Haematol.* 161, 76–86.
- Federico, M., Luminari, S., Iannitto, E., Polimeno, G., Marcheselli, L., Montanini, A., et al., 2009. ABVD compared with BEACOPP compared with CEC for the initial treatment of patients with advanced Hodgkin's lymphoma: results from the HD2000 Gruppo Italiano per lo Studio dei Linfomi Trial. *J. Clin. Oncol.* 27, 805–811.
- Fossa, A., Fiskvik, I.H., Kolstad, A., Lauritzen, G.F., Aurlien, E., Blystad, A.K., et al., 2012. Two escalated followed by six standard BEACOPP in advanced-stage high-risk classical Hodgkin lymphoma: high cure rates but increased risk of aseptic osteonecrosis. *Ann. Oncol.* 23, 1254–1259.
- Gordon, L.I., Hong, F., Fisher, R.I., Bartlett, N.L., Connors, J.M., Gascoyne, R.D., et al., 2013. Randomized phase III trial of ABVD versus Stanford V with or without radiation therapy in locally extensive and advanced-stage Hodgkin lymphoma: an intergroup study coordinated by the Eastern Cooperative Oncology Group (E2496). *J. Clin. Oncol.* 31, 684–691.

- Klastersky, J., de Naurois, J., Rolston, K., Rapoport, B., Maschmeyer, G., Aapro, M., et al., 2016. Management of febrile neutropenia: ESMO clinical practice guidelines. *Ann. Oncol.* 27, v111–v118.
- Kuderer, N.M., Dale, D.C., Crawford, J., Lyman, G.H., 2007. Impact of primary prophylaxis with granulocyte colony-stimulating factor on febrile neutropenia and mortality in adult cancer patients receiving chemotherapy: a systematic review. *J. Clin. Oncol.* 25, 3158–3167.
- Lyman, G.H., 2009. Impact of chemotherapy dose intensity on cancer patient outcomes. *J. Compr. Canc. Netw.* 7, 99–108.
- Lyman, G.H., Kuderer, N.M., 2003. Epidemiology of febrile neutropenia. *Support. Cancer Ther.* 1, 23–35.
- Lyman, G.H., Abella, E., Pettengell, R., 2014. Risk factors for febrile neutropenia among patients with cancer receiving chemotherapy: a systematic review. *Crit. Rev. Oncol. Hematol.* 90, 190–199.
- Martin, W.G., Ristow, K.M., Habermann, T.M., Colgan, J.P., Witzig, T.E., Ansell, S.M., 2005. Bleomycin pulmonary toxicity has a negative impact on the outcome of patients with Hodgkin's lymphoma. *J. Clin. Oncol.* 23, 7614–7620.
- Moreau, M., Klastersky, J., Schwarzbald, A., Muanza, F., Georgala, A., Aoun, M., et al., 2009. A general chemotherapy myelotoxicity score to predict febrile neutropenia in hematological malignancies. *Ann. Oncol.* 20, 513–519.
- Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Hematopoietic Growth Factors V.2.2019 © National Comprehensive Cancer Network, Inc. 2019a. All rights reserved. Accessed March 26, 2019. To view the most recent and complete version of the guideline, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.
- Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Hodgkin Lymphoma V.1.2019. © National Comprehensive Cancer Network, Inc. 2019b. All rights reserved. Accessed March 26, 2019. To view the most recent and complete version of the guideline, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.
- Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Prevention and Treatment of Cancer-Related Infections V.1.2019. © National Comprehensive Cancer Network, Inc. 2019c. All rights reserved. Accessed March 26, 2019. To view the most recent and complete version of the guideline, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.
- Repetto, L., Biganzoli, L., Koehne, C.H., Luebbe, A.S., Soubeyran, P., Tjan-Heijnen, V.C., et al., 2003. EORTC Cancer in the Elderly Task Force guidelines for the use of colony-stimulating factors in elderly patients with cancer. *Eur. J. Cancer* 39, 2264–2272.
- Russo, F., Corazzelli, G., Frigeri, F., Capobianco, G., Aloj, L., Volzone, F., et al., 2014. A phase II study of dose-dense and dose-intense ABVD (ABVDDD-D) without consolidation radiotherapy in patients with advanced Hodgkin lymphoma. *Br. J. Haematol.* 166, 118–129.
- Schwenkglenks, M., Pettengell, R., Szucs, T.D., Culakova, E., Lyman, G.H., 2010. Hodgkin lymphoma treatment with ABVD in the US and the EU: neutropenia occurrence and impaired chemotherapy delivery. *J. Hematol. Oncol.* 3, 27.
- Scott, S.D., Chrischilles, E.A., Link, B.K., Delgado, D.J., Fridman, M., Stolshek, B.S., 2003. Days of prophylactic filgrastim use to reduce febrile neutropenia in patients with non-Hodgkin's lymphoma treated with chemotherapy. *J. Manag. Care Pharm.* 9, 15–21.
- Smith, T.J., Khatcheressian, J., Lyman, G.H., Ozer, H., Armitage, J.O., Balducci, L., et al., 2006. Update of recommendations for the use of white blood cell growth factors: an evidence-based clinical practice guideline. *J. Clin. Oncol.* 24(24), 3187–3205.
- Smith, T.J., Bohlke, K., Lyman, G.H., Carson, K.R., Crawford, J., Cross, S.J., et al., 2015. Recommendations for the use of WBC growth factors: American Society of Clinical Oncology clinical practice guideline update. *J. Clin. Oncol.* 33, 3199–3212.
- Stamatoullas, A., Brice, P., Bouabdallah, R., Mareschal, S., Camus, V., Rahal, I., et al., 2015. Outcome of patients older than 60 years with classical Hodgkin lymphoma treated with front line ABVD chemotherapy: frequent pulmonary events suggest limiting the use of bleomycin in the elderly. *Br. J. Haematol.* 170, 179–184.
- Straus, D.J., Collins, G.P., Walewski, J.A., Zinzani, P.L., Grigg, A., Sureda, A.M., et al., 2018. Improving outcomes with brentuximab vedotin (BV) plus chemotherapy in patients with newly diagnosed advanced stage Hodgkin lymphoma. *J. Clin. Oncol.* 36.
- Tao, Y.X., Sun, S.Y., Kang, S.Y., Zhou, L.Q., Shi, Y.K., Li, Y.X., et al., 2014. Comparison of dose-dense ABVD and standard ABVD in the treatment of early unfavorable and advanced Hodgkin's lymphoma: a retrospective analysis. *J. Huazhong Univ. Sci. Technol. Med. Sci.* 34, 260–264.
- Viviani, S., Zinzani, P.L., Rambaldi, A., Brusamolino, E., Levis, A., Bonfante, V., et al., 2011. ABVD versus BEACOPP for Hodgkin's lymphoma when high-dose salvage is planned. *N. Engl. J. Med.* 365, 203–212.
- Wang, L., Baser, O., Kutikova, L., Page, J.H., Barron, R., 2015. The impact of primary prophylaxis with granulocyte colony-stimulating factors on febrile neutropenia during chemotherapy: a systematic review and meta-analysis of randomized controlled trials. *Support. Care Cancer* 23, 3131–3140.
- Weycker, D., Li, X., Barron, R., Wu, H., Morrow, P.K., Xu, H., et al., 2015. Importance of risk factors for febrile neutropenia among patients receiving chemotherapy regimens not classified as high-risk in guidelines for myeloid growth factor use. *J. Compr. Canc. Netw.* 13, 979–986.
- Younes, A., Oki, Y., McLaughlin, P., Copeland, A.R., Goy, A., Pro, B., et al., 2012a. Phase 2 study of rituximab plus ABVD in patients with newly diagnosed classical Hodgkin lymphoma. *Blood* 119, 4123–4128.
- Younes, A., Gopal, A.K., Smith, S.E., Ansell, S.M., Rosenblatt, J.D., Savage, K.J., et al., 2012b. Results of a pivotal phase II study of brentuximab vedotin for patients with relapsed or refractory Hodgkin's lymphoma. *J. Clin. Oncol.* 30, 2183–2189.
- Younes, A., Connors, J.M., Park, S.I., Fanale, M., O'Meara, M.M., Hunder, N.N., et al., 2013. Brentuximab vedotin combined with ABVD or AVD for patients with newly diagnosed Hodgkin's lymphoma: a phase 1, open-label, dose-escalation study. *Lancet Oncol.* 14, 1348–1356.

Anna Sureda, MD, PhD, is Head of the Haematology Department of Institut Català d'Oncologia–Hospital Duran I Reynals in Barcelona, Spain. Her primary focuses are on clinical investigations into the treatment of Hodgkin lymphoma, non-Hodgkin lymphoma, and multiple myeloma patients. She has participated in many phase 2 and phase 3 clinical trials for lymphoma patients. She has held membership of a number of national and international cancer and hematology societies, is a regular reviewer for several peer-reviewed hematology journals, and has co-authored more than 250 peer-reviewed journal articles.

Eva Domingo-Domenech, MD, graduated with a degree in Medicine from the University of Barcelona in 1995 and completed her residency in Haematology at the Hospital Universitario de Bellvitge, Barcelona in 2000. She is the Clinical Coordinator of the Hematology Department of Institut Català d'Oncologia – Hospital Duran i Reynals in Barcelona since June 2015. Eva Domingo Domenech has focused her career on clinical investigations into the treatment of Hodgkin's and non-Hodgkin's lymphomas patients. She is author or co-author of more than 46 indexed publications related with the epidemiology, treatment and transplantation of lymphoma patients.

Ashish Gautam, PhD, MBA, is Global Medical Lead in the Global Medical Affairs division of Millennium Pharmaceuticals Inc., a wholly owned subsidiary of Takeda Pharmaceutical Company Ltd. His research interests have been in advancing the understanding of oncologic pathways and clinical development of cancer drugs. He has published in several peer reviewed oncology focused journals.