



Equine assisted activities and therapies in children with autism spectrum disorder: A systematic review and a meta-analysis



Tomasz Trzmiel^{a,*}, Barbara Purandare^a, Michał Michalak^b, Ewa Zasadzka^a, Mariola Pawlaczyk^a

^a Department of Geriatric Medicine and Gerontology, Poznan University of Medical Sciences, Poland

^b Department of Computer Science and Statistics, Poznan University of Medical Sciences, Poland

ARTICLE INFO

Keywords:

Equine assisted activities and therapies
Autism spectrum disorder
Hippotherapy

ABSTRACT

Introduction: The multifactorial nature of Autism Spectrum Disorder (ASD) is the reason why complementary and alternative methods of treatment are sought in order to support the classic approach.

Objectives: The aim of the study was to assess the effectiveness of Equine-Assisted Activities and Therapies (EAAT) in ASD patients based on a review of the literature.

Methods: A review of the literature and a meta-analysis were conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. PUBMED, Cochrane Library, Web of Science, ClinicalTrials.gov and PEDro databases were searched until July 20, 2017. Only articles published in English, in a journal with a review process, after 1999, with a control group or presentation of comparative pre-/post-therapy results in ASD patients, and clear inclusion/exclusion criteria were considered. The methodological quality of the included studies was assessed using the Quality Assessment Tool for Quantitative Studies (QATQS). The meta-analysis of three studies was conducted.

Results: A total of 15 studies with 390 participants (aged: 3–16 years) were included. The interaction between psychosocial functioning and EAAT was investigated in most studies. Improvement was reported in the following domains: socialization, engagement, maladaptive behaviors, and shorter reaction time in problem-solving situations after EAAT. The meta-analysis revealed no statistically significant differences for the investigated effects.

Conclusions: Despite the need for further, more standardized research, the results of the studies included in this review allow us to conclude that EAAT may be a useful form of therapy in children with ASD.

1. Introduction

Modern medicine, with its continuous progress and breakthroughs, allows for the development of targeted therapies aiming at the pathomechanisms of various diseases, often based on the identification of individual therapeutic targets. Despite numerous medical advancements, the origins and causes of some diseases remain unknown and continue to pose a challenge to contemporary medicine. Autism Spectrum Disorder (ASD) is an example of a disease of unknown etiology and the number of the affected individuals is steadily increasing. In the US alone, ASD is diagnosed in as many as 16.8/1000 children aged 8 years, which significantly exceeded earlier predictions and makes it a public healthcare problem.¹

ASD is characterized by dysfunctional behaviors, communication and social-interaction difficulties. The affected individuals typically present stereotyped behaviors and struggle with establishing contact,

verbal and non-verbal communications, creating and understanding interpersonal relations.^{2–4} The etiology of ASD remains to be fully elucidated and the observational method is the only diagnostic tool available.⁵ A relationship between ASD and gastrointestinal symptoms, gut microbiota, nerve conduction, sleep, immune and mitochondrial disorders has been demonstrated.^{3,6} The multifactorial nature of ASD is the reasons why a variety of therapies, including educational, behavioral and pharmacotherapy, have been implemented. Despite extensive search, we lack consensus on the effective management and pharmacotherapy of ASD. The available therapeutic methods are not sufficient even in terms of symptomatic treatment.¹ Parents of children with ASD frequently seek complementary and alternative methods of treatment,⁷ e.g. equine-assisted activities and therapies (EAAT), to support the classic approach. As the number of individuals undergoing EAAT increases annually, it seems prudent to evaluate the effectiveness of EAAT for ASD treatment.

* Corresponding author at: Department of Geriatric Medicine and Gerontology, Poznan University of Medical Sciences, ul. Święcickiego 6, Poznan 60-781, Poland.
E-mail address: ttrzmiel@ump.edu.pl (T. Trzmiel).

<https://doi.org/10.1016/j.ctim.2018.11.004>

Received 23 August 2018; Received in revised form 1 November 2018; Accepted 2 November 2018

Available online 05 November 2018

0965-2299/ © 2018 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

EAAT incorporates hippotherapy (HPOT), i.e. an integrated therapeutic program, and therapeutic riding (TR), which originated from recreational activities. Both types of therapy use the contact with a horse and equine movement, which provides rhythmic movement to the patient's body. EAAT affects the entire body, regulating blood pressure and muscle tone, and leading to improved balance.^{8,9}

The literature offers numerous reports about the effectiveness of EAAT for the improvement of the physical functions of the body, although they mostly concern patients with cerebral palsy (CP).^{10–13} Still, the popularity of EAAT is growing and the number of publications rises accordingly.

The aim of the study was to conduct a systematic review of the literature and a meta-analysis of the published trials with comparison groups about the effectiveness of EAAT among ASD patients.

2. Methods

A review of the literature and a meta-analysis were conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.¹⁴ The procedures (search strategy, inclusion/exclusion criteria, and data extraction) were established and included in the protocol. An approval of an Ethics Committee is not required in such studies.

3. Search strategy

The following online data bases were searched: PUBMED, Cochrane Library, Web of Science, ClinicalTrials.gov and PEDro. An algorithm with key words (*Autism AND Hippotherapy OR Equine-Assisted Therapy*) was used to find suitable publications. Two of the authors (TT and EZ) conducted their independent searches between January 1, 2000 and July 20, 2017. Additionally, the reference sections of the included articles were manually inspected to identify unique records. Only articles published in English were taken into consideration.

4. Inclusion/exclusion criteria

The inclusion criteria for the reports were as follows: (a) published in English in a journal with a review process, after 1999, (b) research study with a control group or presentation of comparative pre- and post-therapy results in ASD patients, (c) clearly defined inclusion and exclusion criteria for the study groups and controls, (d) only EAAT was used in the study group as compared to controls.

The following articles were excluded: (a) study populations including other patients than with ASD, (b) use of a horseback riding stimulator instead of live horses, (c) use of additional therapies, or surgical interventions, in the study group as compared to controls.

5. Quality assessment

All studies included in the qualitative synthesis were evaluated using the Quality Assessment Tool for Quantitative Studies (QATQS),¹⁵ which allows to determine their methodological quality. QATQS assesses 8 sections: selection bias, study design, confounders, blinding, data collection methods, withdrawals and dropouts, intervention integrity, analysis. Each section may be classified as 'weak', 'moderate' or 'strong', according to a reviewer's dictionary. If one section is evaluated as 'weak', the entire study is deemed 'moderate', if more than one section is 'weak', the study is automatically 'weak', and if neither section is 'weak', the study is evaluated as 'strong'. The assessments were performed independently by two authors (TT and BP). If agreement on the quality assessment could not be reached by two authors, the third author was consulted (EZ).

6. Data extraction and analysis

The following data were extracted from each study: first author, year of publication, study population characteristics, study design, inclusion/exclusion criteria, intervention characteristics, assessment of the outcome, and results.

The meta-analysis was performed on the summary results of the Vineland Adaptive Behavior Scales (VABS)16 and their results in Social subscale and Communication subscale in the form of standardized mean differences (SMDs). Three studies,^{17–19} which used that tool to evaluate the effects of the therapy and presented the results in a way which facilitated a comparison were analyzed. A standard χ^2 test was used to test the heterogeneity between trial results. The I² parameter was used to measure inconsistency between the results. For the χ^2 statistic, the p-value of < 0.10 was considered to indicate significant heterogeneity. The I² statistic value of $\geq 50\%$ was considered to indicate significant heterogeneity. If significant heterogeneity existed between the studies, a random-effects model of analysis was used. Otherwise, a fixed-effects model was used. Publication bias was assessed visually by funnel plots and statistically by Egger's test.

7. Results

The search results and the flow diagram of the study selection are summarized in Fig. 1. A total of 15 studies were included. More than one test to assess the effects of HPOT was used in most (10 out of 15) studies. Domains associated with psychosocial functioning (socialization, communication, engagement, stereotypical behaviors) were evaluated in 14 studies,^{18–28,17,29,30} cortisol levels in saliva - 1,³¹ execution of daily activities - 5,^{18,19,26,28,17} gait parameters - 1,²⁰ and motor functions - 6.^{15–17,23,25,26} Using QATQS, 6 studies were deemed to be 'strong', 6 - 'moderate' and 3 - 'weak'. Detailed results are presented in Table 1. Data collection methods section and blinding section were awarded the highest and the lowest number of points, respectively. The results for individual sections are presented in Fig. 2.

8. Patient characteristics and study design

A total of 390 participants (aged: 3–16 years, mean: 5.14–10.2 years) from 15 articles were included in the study. The subjects were predominantly male (308 (79%) - males, 72 (21%) - females). All participants were diagnosed with ASD. In 7 studies^{18,19,25,26,28,17,30} they had to meet the DSM IV - TR (Diagnostic and Statistical Manual of Mental Disorders 4th edition, text revision) criteria,³² whereas in the remaining studies the criteria for ASD diagnosis were either not presented,^{20,22–24,27,29,31} or ICD-10 criteria for ASD were used.²¹ Patient characteristics for individual study groups are presented in Table 2.

TR was the intervention of choice in the clear majority of the studies,^{18–25,27,28,17,29–31} and HPOT was used in only one study.²⁶ The number of individual therapeutic sessions and their duration varied between the studies, with 30 min. for the shortest,^{20,30} and 180 min. for the longest¹⁸ session. Duration of the session was not disclosed in 1 study.³¹ Detailed characteristics of the studies are presented in Table 3.

9. Results of the therapeutic interventions

The interaction between psychosocial functioning and EAAT was investigated in most studies included in our analysis. Borgi et al¹⁹ who used TR in ASD patients, reported improved socialization scores of the VABS as compared to controls who did not attend TR (change between baseline and final scores, mean \pm SD, EAT: 0.72 ± 0.22 , controls: 0.23 ± 0.21 , ANOVA Time x Group interaction $F(1,18) = 5.30$, $p = 0.034$, Tukey test $p < 0.01$), and shorter reaction time in problem-solving situations (change between baseline and final scores using the Tower of London (TOL) test, mean \pm SD, EAT: -20.7 ± 6.6 , controls: -6.46 ± 5.2 , ANOVA Time x Group interaction, $F(1,19) = 5.85$,




Fig. 1. Study flow diagram.

p = 0.026; Tukey test p < 0.01).

Anderson et al¹⁸ reported a markedly decreased maladaptive behavior traits of the VABS maladaptive behavior score after using EAA (Equine Assisted Activities) ($F_{(1,11)} = 5.65$, $\eta^2 = 0.34$) ($F_{(1,11)} = 5.19$, $p = 0.04$). However, the results of other EQ/SQ parts and of the VABS correlated with EAA did not show statistical significance. Gabriels et al²⁸ who used the Autism Behavior Checklist (ABC-C),³⁵ found significantly reduced Hyperactivity (effect size =

0.50, $p = 0.02$) and Irritability (effect size 0.53, $p = 0.01$) in the TR intervention group and improved Social Cognition and Communication subscales with SRS (Social Response Scale)³⁶ as compared to the Barn Activities (BA) control group. The BA control group was instructed about horseback riding and caring for horses but had not contact with horses. Instead, a life-sized stuffed horse was used. In another study, Gabriels et al¹⁷ demonstrated a statistically significant improvement between the TR group and controls in the following ABC-C subscales:

Table 1
Evaluation of the methodological quality of each study using QATQS.

	Selection Bias	Design	Confounders	Blinding	Data collection methods	Withdrawals and drop-outs	Summary
Ajzenman 2013	?	?	+	+	+	+	?
Anderson 2016	+	?	+	+	+	+	+
Bass 2009	+	+	+	+	+	+	?
Borgi 2016	+	+	+	?	+	+	+
Gabriels 2012	+	?	+	+	+	+	+
Gabriels 2015	+	+	+	?	+	+	+
Garcia - Gomez 2017	+	+	+	?	+	+	+
Jenkins 2013	?	+	?	+	+	+	?
Kern 2011	+	?	+	+	+	+	?
Lanning 2014	?	?	+	?	+	?	+
Llambias 2016	?	?	+	?	?	?	?
Memishevskij 2010	+	+	+	?	+	+	+
Steiner 2015	+	+	+	?	+	+	?
Tabares 2012	+	?	+	?	+	+	+
Ward 2013	+	?	+	?	+	+	+




Fig. 2. Evaluation of the individual items within the quality sections presented as percentages across all included studies.

Irritability ($p = 0.004$), Lethargy ($p = 0.007$), Stereotypic Behavior ($p = 0.02$), and Hyperactivity ($p = 0.008$). Bass et al³⁰ found a statistically significantly improved SRS results (baseline vs. final) in the TR group ($r(10) = 2.87$, $p = 0.017$, $d = 0.66$) as compared to controls ($t(10) = 0.108$, $p = 0.916$, $d = 0.02$). Similar results were observed after comparing the Sensory Profile³⁷ results in both groups – a significant improvement was noted in the TR group and only a slight improvement in controls ($t(18) = -7.29$, $p < 0.01$, $d = -0.059$ and $t(13) = -1.77$, $p = 0.101$, respectively). Garcia-Gomez et al²⁴ used the BASC -I³⁸ questionnaire and found lower levels of aggressiveness, from 4.6 to 3.3 ($p = 0.039$, $d = 0.220$). Ward et al²⁵ used the GARS-2 (Gilliam Autism

Rating Scale: Second Edition) test^{39,40}, and noted an improvement in the TR group with regard to social interactions ($F(5.60) = 4.61$, $p < 0.05$ and individual items of the SPSC (Sensory Profile School Companion)⁴¹; Registration and Sensitivity subscales, respectively $F(5,100) = 2.29$, $p < 0.05$ and $F(5,100) = 2.99$, $p < 0.05$, School Factor Scores $F(20.322.6) = 1.89$, $p < 0.05$ and Section Scores $F(925.358.1) = 2.05$, $p < 0.05$. These authors observed a significant improvement after 6 weeks of therapy as compared to baseline, and a marked deterioration (back to baseline values) during a 6-week break, followed by another improvement after another 6 weeks of therapy.

Ajzenman et al²⁶ proved that a 12-week HPOT intervention improves the following areas: postural control, participation in daily activities, and adaptive behaviors. Means of all pre- and post-HPOT VABS-II items⁴² were 64.00 (± 15.30) and 70.67 (± 18.50 , $d = 0.393$ $p = 0.027$), respectively. As far as postural control was concerned, the improved areas were as follows: movement variability of sway area for COP (center of pressure) (change: 12%, $p = 0.028$, $d = 0.1999$), COM (center of mass) velocity, both, AP (anterior – posterior) axis and ML (medial – lateral) axis (changes: 102%, $p = 0.046$, $d = 1.316$ and 20%, $p = 0.046$, $d = 0.845$, respectively).

Llambias et al²⁹ investigated engagement in meaningful activities among children with ASD undergoing equine-assisted therapy (EAT). The authors coded the child’s response to requests and instructions of a therapist as ‘engaged’ or ‘not engaged’ for all of the evaluated behaviors. The percentage of time engaged during all segments was the outcome measure. The therapy significantly improved engagement in all subjects undergoing EAT (from 51.8%–77.8% at baseline to 95.5%–99.3% post-therapy).

Tabares et al³¹ measured progesterone and cortisol levels in salivary samples from 8 boys undergoing HPOT and found the cortisol levels to steadily decrease (from 2.79 ± 0.52 ng/mL at baseline to 2.23 ± 0.75 ng/mL after all sessions), and progesterone levels to increase (from 21.58 ± 12.0 pg/mL at baseline to 26.03 ± 11.98 pg/mL after all sessions ($p \leq 0.05$)). The authors explained that the elevated level of cortisol after the first session was the result of stress caused by a change in the daily routine, which is not well-tolerated by autistic children.

Steiner and Kertesz,²⁰ analyzed changes in gait cycle parameters in children with autism and found improved length of the gait cycle in the TR group and no improvement in the control group. A slight improvement of the gait cycle on the left side was accompanied by a slight weakening on the right side, resulting in increased asymmetry.

Kern et al²² observed an overall improvement in behaviors typical for autism during TR. They used the Childhood Autism Rating Score (CARS)^{43,44} to determine the intensity of the change and found lower CARS values after 3 ($t(21) = 2.73$, $p < 0.02$) and 6 ($t(16) = 3.33$, $p < 0.005$) months of therapy. No changes in the results were noted when two other tools were applied: the Timberlawn Parent – Child Interaction Scale^{45,46} and the Sensory Profile.

Lanning et al²⁷ noted an improved quality of life (QoL) evaluated with the use of two instruments: the Pediatric Quality of Life 4.0 (PedsQL 4.0) generic core scales⁴⁷ and Child Health Questionnaire (CHQ)⁴⁸ in autistic children undergoing EAA. Improved CHO scores for QoL were observed both, in the EAA group and controls (the latter were involved in social circles, which allowed for the children to take part in educational activities facilitated by students of psychology). The highest improvement for the controls and the EAA group was observed in the domains of self-esteem ($M1 = 56.25 \pm 18.77$, $M9 = 67.71$, ± 21.56) and general behavior ($M1 = 52.08$, ± 19.37 ; $M9 = 67.50$, ± 12.70), respectively. An increase in the psychosocial CHQ summary scores was reported for the control group ($M1 = 34.80$, \pm ; $M9 = 39.58$, ± 7.39), while higher physical and psychosocial summary scores were observed in the EAA group (respectively, $M1 = 40.80$, ± 13.43 ; $M9 = 44.51$, ± 16.72 and $M1 = 37.22$, ± 15.23 ; $M9 = 39.08$, ± 14.54). PedsQL scores of QoL were higher in the EAA group but a comparison of differences in QoL improvement in both groups, evaluated with PedsQL and CHQ, did not

Table 2
Characteristics of the participants of the included studies.

Study (first author and year)	N of participants	Age	Inclusion criteria	Exclusion criteria	Additional diseases
Aizenman 2013	6	5–12	ASD, age: 5–12 years, full term birth, ability to ambulate independently and follow a one - step direction	diagnosis of severe impairment, CP, epilepsy, any other neurological or psychiatric condition, severe behavioral issues, restricted ability to sit unaided, contraindications for HPOT according PATH list, previous exposure to EAAAT	nd ^a
Anderson 2016	15	5–16	ASD	riding experience with horses before the study	Attention Deficit Hyperactivity Disorder ADHA (20%), Hypersensitivity and Sensory Integration Disorder (53%)
Bass 2009	34	4–10	ASD	nd	nd
Borgi 2015	28	6–12	ASD, age: verbal, 6–12 years, IQ > 70 on the Wechsler Intelligence Scale for Children III, lack of previous experiences with THR	serious motor/neurological problem, allergies, fearful of horses	nd
Gabriels 2012	42	6–16	ASD age: 6–16 years, Social Communication Questionnaire > = 15, Aberrant Behavior Checklist - Community > = 11, NVIQ > = 40	dangerous behavior in past, animal abuse in past or animal phobia, more than 2 h of EAAAT in past 6 months, weight exceeding policies to ensure health and safety of staff	comorbid psychiatric conditions 38% of group
Gabriels 2015	127	6–16	ASD age: 6–16 years, Aberrant Behavior Checklist - Community > = 11, NVIQ > = 40	previous THR within past 3 years, history of animal abuse or phobia of horses	nd
Garcia - Gomez 2014	16	7–16	ASD	nd	nd
Jenkins 2013	7	6–14	ASD, no prior exposure to THR or HPOT, live within 30 miles of research site.	nd	Verbal and Motor Apraxia, Tuberos Sclerosis
Kern 2011	24	3–12	Age: 3–12 years, ASD, Childhood Autism Rating Scale (CARS) > = 30,	nd	nd
Lanning 2014	25	4–15	ASD, ability to communicate with staff, no previous EAA during last 6 months	nd	nd
Llambias 2016	7	4–7	ASD, age: 3–8 years, understand English, and have had no riding experiences for ≥ 3 months before the study, no aversion to horses	repeated aggressive behavior, comorbidities (e.g., deafness, blindness, epilepsy) that could interfere with the intervention	ADHD
Memisevic 2010	4	8–10	ASD	nd	mild intellectual disability, severe intellectual disability
Seiner 2015	26	10–13	ASD	nd	nd
Tabares 2012	8	8–16	ASD	nd	nd
Ward 2013	21	–	ASD	CAB-T tested critical behaviors	nd

^a nd – no data.

Table 3
Intervention and measure characteristics of the included studies.

Study (first author and year)	Type of therapeutic intervention	Number of sessions	Duration of 1 session	Period of therapeutic intervention	Measurement	Measured domains
Ajzenman 2013	HPOT	12	45 min.	12 weeks	Force plates and Video Motion Capture, Vineland Adaptive Behavior Scales II (Parent/caregiver rating form) VABS II, Child Activity Card Sort CACS	communication, daily living skills, socialization, motor skills, self-care, community mobility, high demand leisure, low demand leisure, social interaction, domestic and education
Anderson 2016	TR, Horsemanship, Stable Management	6	180 min.	6 weeks (1 - initial training and 5 - proper therapy)	Autism-Spectrum Quotient for Children(ASQ-child), Autism - Spectrum Quotient for Adolescents(ASQ-adolescent), Empathizing quotient/systemizing quotient Vineland Adaptive Behavior Scale (VABS) social responsiveness scale, the sensory profile	communication, daily living skills, socialization, motor skills, self-care, community mobility, high demand leisure, low demand leisure, social interaction, domestic and education
Bass 2009	mounting/dismounting, exercise, riding skills, mounted games, horsemanship activities	12	60 min.	12 weeks		social awareness, social cognition, social communication, social motivation, and autistic mannerisms
Borgi 2015	grooming, hand walking the horse, horseback riding, feeding and talking to horse	25	60–70 min.	6 months	Vineland Adaptive Behavior Scale (VABS), Tower of London (TOL)	communication, daily living skills, socialization, motor skills, executive functioning (mainly planning and problem-solving)
Gabriels 2012	putting riding helmet on, sitting and waiting on the bench, mounting horse, THR activities, dismounting, grooming horse, and putting away equipment	10	45 min.	10 weeks	The Aberrant Behavior Checklist-community, Vineland Adaptive Behavioral Scales – Interview Edition, Survey Form (VABS-II, Bruininks-Oseretsky Test of Motor Proficiency (BOT-2, Short Form, Sensory Integration and Praxis Test (SIPT)	irritability/agitation, lethargy/social withdrawal, stereotyped behavior, hyperactivity, and inappropriate speech; communication, daily living skills, and socialization skills: fine motor precision, fine motor integration, manual dexterity, balance, running speed and agility, upper-limb coordination and strength, sensory processes and praxis
Gabriels 2015	preparing to mount, mounting, riding activities, dismounting, grooming, putting equipment away	10	≥ 45 min.	10 weeks	Peabody Picture Vocabulary Test 4 ed PPVT-4, Systematic Analysis of Language Transcripts, Bruininks-Oseretsky Test of Motor Proficiency–2nd Ed (BOT-2), two subscales of the Sensory Integration and Praxis Test (SIPT)42: Praxis on Verbal Command and Postural Praxis, Vineland Adaptive Behavioral Scales – 2nd Edition, ABC-C, Social Responsiveness Scale (SRS),	irritability/agitation, lethargy/social withdrawal, stereotyped behavior, hyperactivity, and inappropriate speech, social awareness, social cognition, social communication, autistic mannerisms, social motivation, words/different words used, sensory integration praxis, postural praxis, praxis on verbal command, adaptive behaviors, communication, daily living activities, socialization
Garcia-Gomez 2014	preparing to mount, mounting, riding activities, dismounting, gathering equipment, saying goodbye to horses	24	45 min.	12 weeks	questionnaire based on Quality-of-life Model, Behavior Assessment System for Children - for Teachers BASC-T	behavior, emotional well-being, interpersonal relationship, personal development, physical well-being, self-determination, social inclusion, familial well-being.
Jenkins 2013	preparing horse, mounting and riding, dismounting, gathering and putting away equipment	9	60 min.	9 weeks	observation, CBCL child behavior checklist, Teacher rating form (TRF)	expression of happiness/unhappiness, initiations and responses to initiations, problem behavior and off-task behavior, posture
Kern 2011	riding, managing the horse in stable, feeding	24	60 min.	nd	Childhood Autism Rating Scale (CARS), Timberlawn Parent-Child Interaction Scale, Sensory Profile, Quality of Life Enjoyment and Satisfaction Questionnaire, Treatment Satisfaction survey	behavior, interaction between parent and child (empathy, expressiveness, responsiveness, positive regard, negative regard, mood and tone), frequency of response to various sensory experiences (processing, modulation, response), quality of life, satisfaction with treatment
Lanning 2014	safety lessons, grooming, riding activities	12	60 min.	12 weeks	Pediatric Quality of life 4.0 Generic Core Scales (PedsQL), Child Health Questionnaire (CHQ)	physical/ emotional/social/ school functioning. physical functioning, role/social emotional/ behavioral, role/social limitations-physical, bodily pain/discomfort, behavior, mental health, self-esteem, general health perception, parental impact –emotional, parental impact- time engagement
Llambias 2016	off-horse activities: grooming or feeding the horse, saddling and leading the horse, art activities (e.g., drawing the horse), and	9–12	45–60 min.	nd	video recording capture children response. When the therapist made a request, the child's response was coded.	

(continued on next page)

Table 3 (continued)

Study (first author and year)	Type of therapeutic intervention	Number of sessions	Duration of 1 session	Period of therapeutic intervention	Measurement	Measured domains
Memisevic 2010	games related to horses. On-horse activities: playing games such as carrying an object from one part of the arena to the other or grabbing rings. horsemanship activities, riding and exercises, equine-assisted psychotherapy	10	30 min.	10 weeks	The Autism Treatment Evaluation Checklist (ATEC)	speech/language communication, sociability, sensory/cognitive awareness, health/physical behavior
Steiner 2015	TR	4 × 2	30 min.	2 × 1 month	Pedagogical Analysis and Curriculum (PAC test), Gait analyzer - Ariel Performance Analysis System (APAS) Saliva Analysis - ELISA	communication, self-care, motor skills, socialization, gait cycle parameters cortisol level in saliva
Tabares 2012	contact and mounting the horse, horseback riding, dismount, interaction with horse	4	nd	1 month		
Ward 2013	orientation, mounting and riding, riding skills, closure	nd	≥ 40 min.	6 weeks of TR/ 6 weeks of break/4weeks of TR/ 6weeks of break/8weeks of TR	Gilliam Autism Rating Scale 2 (GARS2), Sensory Profile School Companion (SPSC)	stereotyped behavior, communication, social interaction, responsiveness to sensory experiences (how responsiveness is manifested in classroom)

reveal any statistical significance ($p \leq 0.05$).

Memishevijk and Hodzhikj²¹ measured the effects of EAT using the Autism Treatment Evaluation Checklist (ATEC).⁴⁹ ATEC scores in individual subscales were improved as compared to baseline (respectively, 0–25% for the speech/language/communication domains, 3–30% for sociability, 0–35% for sensory/cognitive awareness, and 0–11% for health/physical/behavior), although not in all subjects. Regardless, the summary scores were improved after therapy in all cases (by 1–25%).

Only Jenkins et al²³ found no evidence for statistically significant beneficial impact of TR on the affect, linguistic competence, spontaneous initiations, or problem behavior of autistic children. Given the results of their study, these authors believe that TR should be treated as a leisure activity rather than therapy. However, we are of the opinion that their study, being only a single report, should not belittle the significance of the evidence on the beneficial effects of EAAT.

10. VABS meta –results of the analysis

The meta-analysis revealed no statistical significance for the investigated effects, either for the VABS summary scores or the Socialization and Communication subscales (Tables 4–6). Egger's Test showed no publication bias for any of the cases.

11. Discussion

In our study, a systematic review of the studies with comparison groups which evaluated the effectiveness of EAAT in ASD therapy was conducted. The meta-analysis proved to be a challenge due to significant differences between research tools and result presentation in all of the analyzed texts. Most publications on the effectiveness of EAAT concern children with cerebral palsy, the most common neurodevelopmental disorder in that age group.^{50–52} In comparison, the number of studies on ASD is relatively small. ASD is less common, although the number of cases continues to rise, especially in the developed countries, and constitutes a challenge for the parents, caregivers, and health care system.^{53,54} Publications on EAAT effectiveness in ASD individuals focus on the behavioral aspects and interaction with the environment, as the impaired areas associated with social functioning, communication, response to stimuli, and engagement constitute the primary diagnostic criteria.^{55–57} Aggressive behavior is another typical feature of ASD and may affect as much as 68% of the ASD population.⁵⁸ Garcia-Gomez et al²⁴ reported lower levels of aggressiveness after TR, which is consistent with the findings of Nurenberg et al⁵⁹ who investigated chronic psychiatric inpatients with violent behavior undergoing Equine Assisted Psychotherapy or Animal Assisted Therapy. In another study among children with Attention –Deficit Hyperactivity Disorder (ADHD), Garcia-Gomez et al⁶⁰ observed a notable improvement in aggressive behavior variables after using EAT. The results of both abovementioned studies^{59,60} indicate that EAAT has a beneficial effect and reduces aggressiveness, which is consistent with the findings of Garcia-Gomez et al²⁴

Ajzenman et al²⁶ reported improved stability and postural parameters in ASD children after EAT. Other authors presented similar findings,^{61–64} although the studies were conducted among children with CP, and reported improved muscle symmetry,^{61,62} stability of the head and the trunk,⁶³ motor function, and stability.⁶⁴ Zadnikar and Kastri¹³ based on their analysis of 10 studies, confirmed improved stability and posture control in children with CP, whereas Vermohlen et al⁶⁵ reported improved balance, spasticity, and quality of life in 70 patients with Multiple Sclerosis undergoing HPOT. Other authors, who investigated children and adolescents with posture and motor impairment, did not find statistically significant improvements regarding these parameters after hippotherapy.⁶⁶

Improved social functioning (interaction, engagement, communication) was reported in 10 studies presented in our

Table 4
Data and Forest plot displaying VABS Total Score meta-analysis results.

Study	HPOT		Control			Standardized mean difference, random, 95% CI		Standardized mean difference, random, 95% CI	
	Mean	SD	N	Mean	SD	N			
Anderson 2016	214,86	121,63	15	210,13	124,32	15	0,038	-0,677	0,754
Borgi 2015	298,67	18,5	6	274,17	15,3	6	1,443	0,147	2,740
Gabriels 2012	458,8	24	40	434,6	24,1	40	0,985	0,521	1,450
Total			61			61	0,742	-0,010	1,494

Heterogeneity $\chi^2 = 5.87$, $df = 2$, $p = 0.053$
 $I^2 = 66.0\%$
 $z = 1.93$ $p = 0.053$

analysis^{18,19,21,25–2817,29,30}. Wilson et al⁶⁷ in their qualitative study, found that HPOT contributed to higher confidence, self-esteem, self-control, coping skills, and a decrease in undesirable behaviors in healthy adolescents. Their findings are supported by a study of Hession et al⁶⁸ who investigated the effects of TR on the mood, cognitive competence and gait parameters in children with dyspraxia. These authors emphasized not only the beneficial effects of a physical motion of riding a horse, but also the potential value of audiovisual stimulation during equine-assisted therapy. Beat-based rhythms which are experienced while horseback riding may stimulate the areas of the brain responsible for memory, perception of emotions, motor control, and learning.^{68–70} Notably, Hwang Jang et al⁷¹ conducted an analysis on the effects of hippotherapy on dexterity and psychosocial functioning of children with CP but found no significant improvements in psychosocial functioning.

The meta-analysis of 3 out of 15 studies, where the same tool – the VABS scale – was used to measure the effects of therapy, was conducted. Despite the lack of statistically significant effects of EAAT, it is important to bear in mind that the analysis was conducted in only 3 publications and 61 participants, which is proof of an insufficient number of studies on the matter rather than lack of EAAT effectiveness.

Beneficial effects of EAAT have been reported by most authors whose studies were discussed in the review and included better social functioning, reduced aggressiveness, and improved stability of the trunk. Regardless, data presentation, not to mention analysis, synthesis and generalization, present a considerable challenge due to lack of a standardized approach to the subject. No consensus on the effectiveness of the therapy, a great variety of the tools and scales used by the researchers and, most of all, of the investigated factors deem most of the results valuable but not easily measurable and comparable. Furthermore, there is a distinct lack of studies on EAAT which meet the

quality requirements.⁷² The available publications frequently display methodology lapses, mostly stemming from small sample size, lack of comparison group, and blinding. The use of different EAAT therapeutic protocols, and especially different methods of measuring their effectiveness, constitute yet another problem, which often lowers the quality of a study. Despite all of the abovementioned areas for improvement, the analysis left us in no doubt about the benefits of EAAT for ASD patients, chief among them social functioning. Even lack of comprehensive analysis is not able to obscure the evidence of great value and effectiveness of EAAT. Our attempt to analyze the findings of the available literature reports might encourage other researchers to conduct studies with a larger sample size, more standardized methods of therapy and outcome measurements, which in turn will allow for a comprehensive and objective evaluation of clinical usefulness of EAAT.

12. Conclusions

The overwhelming majority of the available reports demonstrated high effectiveness of EAAT, especially with regard to improved social functioning. Also, EAAT has been proven to significantly reduce aggressive behaviors and improve trunk stability. Nevertheless, it is impossible to draw universal conclusions due to the considerable discrepancies in therapeutic protocols and measurement instruments of the abovementioned studies. Further, longitudinal trials, with standardized EAAT protocols and representative large sample groups are necessary. Also, it is crucial to establish homogeneous tools to measure therapeutic progress and outcomes, especially with regard to social functioning. In consequence, a more precise assessment of the effectiveness of EAAT among ASD patients will be possible. Despite the need for further research, the results of the studies included in this review allow us to conclude that EAAT may be a useful form of therapy in

Table 5
Data and Forest plot displaying VABS communication score meta-analysis results.

Study	HPOT		Control			Standardized mean difference, fixed, 95% CI		Standardized mean difference, fixed, 95% CI	
	Mean	SD	N	Mean	SD	N			
Anderson 2016	44,6	30,39	15	44,2	31,19	15	0,013	-0,703	0,729
Borgi 2015	74	19,52	6	65	18,54	6	0,473	-0,678	1,623
Gabriels 2012	149	24,8	40	143,6	24,9	40	0,217	-0,222	0,657
Total			61			61	0,191	-0,165	0,547

Heterogeneity $\chi^2 = 0,48$, $df = 2$, $p = 0.786$
 $I^2 = 0,0\%$
 $z = 105$ $p = 0.293$

Table 6
Data and Forest plot displaying VABS Social Score meta-analysis results.

Study	HPOT			Control			Standardized mean difference, fixed, 95% CI			Standardized mean difference, fixed, 95% CI		
	Mean	SD	N	Mean	SD	N						
Anderson 2016	45,07	22,48	15	45,27	23,21	15	-0,009	-0,724	0,707			
Borgi 2015	68,17	21,52	6	61	17,62	6	0,364	-0,779	1,507			
Gabriels 2012	113,2	24,4	40	104,9	29,9	40	0,289	-0,151	0,730			
Total			61			61	0,220	-0,130	0,580			

Heterogeneity $\chi^2 = 0,55$, $df = 2$, $p = 0,760$
 $I^2 = 0,0\%$
 $z = 1,22$ $p = 0,221$

children with ASD.

13. Study limitations

The two main limitations of our review are as follows: a relatively small sample size, which increases the risk of a calculation error, and differences in research methodology, which greatly hinders the comparison of the results.

Conflicts of interest

The authors declare that they have no conflicts of interest.

References

- Marchezan J, Winkler Dos Santos EGA, Deckmann I, Riesgo RDS. Immunological dysfunction in autism spectrum disorder: a potential target for therapy. *Neuroimmunomodulation*. 2018;1–20. <https://doi.org/10.1159/000492225>.
- Rossignol DA, Frye RE. A review of research trends in physiological abnormalities in autism spectrum disorders: immune dysregulation, inflammation, oxidative stress, mitochondrial dysfunction and environmental toxicant exposures. *Mol Psychiatry*. 2012;17(4):389–401.
- Frye RE, Rossignol DA. Identification and treatment of pathophysiological comorbidities of autism spectrum disorder to achieve optimal outcomes. *Clinical medicine insights*. *Pediatrics*. 2016;10:43–56.
- Saghazadeh A, Ahangari N, Hendi K, Saleh F, Rezaei N. Status of essential elements in autism spectrum disorder: systematic review and meta-analysis. *Rev Neurosci*. 2017;28(7):783–809.
- Murphy CM, Wilson CE, Robertson DM, et al. Autism spectrum disorder in adults: diagnosis, management, and health services development. *Neuropsychiatr Dis Treat*. 2016;12:1669–1686.
- Shaaban SY, El Gendy YG, Mehanna NS, et al. The role of probiotics in children with autism spectrum disorder: a prospective, open-label study. *Nutr Neurosci*. 2017;1–6.
- Volkmar F, Siegel M, Woodbury-Smith M, King B, McCracken J, State M. Practice parameter for the assessment and treatment of children and adolescents with autism spectrum disorder. *J Am Acad Child Adolesc Psychiatry*. 2014;53(2):237–257.
- Stergiou A, Tzoufi M, Ntzani E, Varvarousis D, Beris A, Ploumis A. Therapeutic effects of horseback riding interventions: a systematic review and meta-analysis. *Am J Phys Med Rehabil*. 2017.
- Lee J, Yun C-K. Effects of hippotherapy on the thickness of deep abdominal muscles and activity of daily living in children with intellectual disabilities. *J Phys Ther Sci*. 2017;29(4):779–782.
- Rigby BR, Grandjean PW. The efficacy of equine-assisted activities and therapies on improving physical function. *J Altern Complement Med*. 2016;22(1):9–24.
- Tsang S-H, Chen H-C, Tam K-W. Systematic review and meta-analysis of the effect of equine assisted activities and therapies on gross motor outcome in children with cerebral palsy. *Disabil Rehabil*. 2013;35(2):89–99.
- Whalen CN, Case-Smith J. Therapeutic effects of horseback riding therapy on gross motor function in children with cerebral palsy: a systematic review. *Phys Occup Ther Pediatr*. 2012;32(3):229–242.
- Zadnikar M, Kastrin A. Effects of hippotherapy and therapeutic horseback riding on postural control or balance in children with cerebral palsy: a meta-analysis. *Dev Med Child Neurol*. 2011;53(8):684–691.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
- National Collaborating Centre for Methods and Tools. *Quality assessment tool for quantitative studies*. Hamilton, ON: McMaster University; 2008 Updated 03 October, 2017.
- Sparrow S, Balla D, Cicchetti D. *The Vineland adaptive behavior scales: Interview edition, survey form manual*. Circle Pines, Minnesota: American Guidance Service; 1984.
- Gabriels RL, Agnew JA, Holt KD, et al. Pilot study measuring the effects of therapeutic horseback riding on school-age children and adolescents with autism spectrum disorders. *Res Autism Spectr Disord*. 2012;6(2):578–588.
- Anderson S, Meints K. Brief report: the effects of equine-assisted activities on the social functioning in children and adolescents with autism spectrum disorder. *J Autism Dev Disord*. 2016;46(10):3344–3352.
- Borgi M, Loliva D, Cerino S, et al. Effectiveness of a standardized equine-assisted therapy program for children with autism spectrum disorder. *J Autism Dev Disord*. 2016;46(1):1–9.
- Steiner H, Kertesz Z. Effects of therapeutic horse riding on gait cycle parameters and some aspects of behavior of children with autism. *Acta Physiol Hung*. 2015;102(3):324–335.
- Haris Memishevikj SH. The effects of equine-assisted therapy in improving the psychosocial functioning of children with autism. *J Spec Educ Rehabil*. 2010;11(3):57–67.
- Kern JK, Fletcher CL, Garver CR, et al. Prospective trial of equine-assisted activities in autism spectrum disorder. *Altern Ther Health Med*. 2011;17(3):14–20.
- Jenkins SR, DiGennaro Reed FD. An experimental analysis of the effects of therapeutic horseback riding on the behavior of children with autism. *Res Autism Spectr Disord*. 2013;7(6):721–740.
- García-Gómez A, López Risco M, Rubio JC, Guerrero E, García-Peña IM. Effects of a program of adapted therapeutic horse-riding in a group of autism Spectrum disorder children. *EJREP*. 2017;12(32).
- Ward SC, Whalon K, Rusnak K, Wendell K, Paschall N. The association between therapeutic horseback riding and the social communication and sensory reactions of children with autism. *J Autism Dev Disord*. 2013;43(9):2190–2198.
- Ajzenman HF, Standeven JW, Shurtleff TL. Effect of hippotherapy on motor control, adaptive behaviors, and participation in children with autism spectrum disorder: a pilot study. *Am J Occup Ther*. 2013;67(6):653–663.
- Lanning BA, Baier MEM, Ivey-Hatz J, Krenek N, Tubbs JD. Effects of equine assisted activities on autism spectrum disorder. *J Autism Dev Disord*. 2014;44(8):1897–1907.
- Gabriels RL, Pan Z, Dechant B, Agnew JA, Brim N, Mesibov G. Randomized controlled trial of therapeutic horseback riding in children and adolescents with autism spectrum disorder. *J Am Acad Child Adolesc Psychiatry*. 2015;54(7):541–549.
- Llambias C, Magill-Evans J, Smith V, Warren S. Equine-assisted occupational therapy: increasing engagement for children with autism Spectrum disorder. *Am J Occup Ther*. 2016;70(6) 7006220040p1–7006220040p9.
- Bass MM, Duchowny CA, Llabre MM. The effect of therapeutic horseback riding on social functioning in children with autism. *J Autism Dev Disord*. 2009;39(9):1261–1267.
- Tabares C, Vicente F, Sánchez S, Aparicio A, Alejo S, Cubero J. Quantification of hormonal changes by effects of hippotherapy in the autistic population. *Neurochem J*. 2012;6(4):311–316.
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th ed. Washington, DC: American Psychiatric Association; 2000 text revision.
- Auyeung B, Wheelwright S, Allison C, Atkinson M, Samarawickrema N, Baron-Cohen S. The children's empathy quotient and systemizing quotient: sex differences in typical development and in autism spectrum conditions. *J Autism Dev Disord*. 2009;39(11):1509–1521.
- Tavassoli T, Miller LJ, Schoen SA, Jo Brout J, Sullivan J, Baron-Cohen S. Sensory reactivity, empathizing and systemizing in autism spectrum conditions and sensory processing disorder. *Dev Cogn Neurosci*. 2018;29:72–77.
- Krug D, Arick J, Almond P. *Autism behavior checklist – ABC*. Austin, Texas: PRO-ED; 1993.
- Bruni TP. Test review. Social responsiveness scale—Second edition (SRS-2). Constantino J. N. Gruber C. P. (2012). *Social Responsiveness Scale—Second Edition (SRS-2)*. Torrance, CA: Western Psychological Services. *J Psychoeduc Assess*. 2013;32(4):365–369.
- Dunn W. *The sensory profile: Examiner's manual*. San Antonio, Texas: Psychological

- Corporation; 1999.
38. Reynolds CR, Kamphaus RW. *BASC: Behavior assessment system for children*. Circle Pines, Minnesota: American Guidance Service; 1992.
 39. Montgomery JM, Newton B, Smith C. Test review. Gilliam, J. (2006). GARS-2: Gilliam Autism Rating Scale—Second Edition. Austin, TX: PRO-ED. *J Psychoeduc Assess*. 2008;26(4):395–401.
 40. South M, Williams BJ, McMahon WM, et al. Utility of the Gilliam autism rating scale in research and clinical populations. *J Autism Dev Disord*. 2002;32(6):593–599.
 41. Dunn W. *Sensory profile, school companion: User's manual*. San Antonio, Texas: Pearson Psych Corp.; 2006.
 42. Sparrow SS, Cicchetti DV, Balla DA. *Vineland adaptive behavior scales*. 2nd ed. Circle Pines, Minnesota: American Guidance Service; 2005.
 43. Schopler E, Reichler RJ, DeVellis RF, Daly K. Toward objective classification of childhood autism: Childhood Autism Rating Scale (CARS). *J Autism Dev Disord*. 1980;10(1):91–103.
 44. Schopler E, Reichler RJ, Renner BR. *The childhood autism rating scale*. Los Angeles, California: California Western Psychological Services; 1994.
 45. Lewis JM. *The birth of a family*. New York, New York: Brunner/Mazel; 1989.
 46. Lewis JM. *How's your family? A guide to identifying your family's strengths and weaknesses*. New York, New York: Brunner/Mazel; 1979.
 47. Varni JW, Seid M, Kurtin PS. PedsQL 4.0: reliability and validity of the pediatric quality of life inventory version 4.0 generic core scales in healthy and patient populations. *Med Care*. 2001;39(8):800–812.
 48. Langraf JM, Ware JE. *The CHQ user's manual*. Boston: HealthActCHQ Inc.; 1999.
 49. Rimland B ES. *Autism Treatment Evaluation Checklist (ATEC)*. <http://www.autism.com/ari/atec/ateonline.htm>. Updated 2016. (Accessed 8 May 2018).
 50. Stavsky M, Mor O, Mastrolia SA, Greenbaum S, Than NG, Erez O. Cerebral palsy-trends in epidemiology and recent development in prenatal mechanisms of disease, treatment, and prevention. *Front Pediatr*. 2017;5:21.
 51. Tseng S-H, Lee J-Y, Chou Y-L, Sheu M-L, Lee Y-W. Association between socio-economic status and cerebral palsy. *PLoS One*. 2018;13(1):e0191724.
 52. McIntyre S. The continually changing epidemiology of cerebral palsy. *Acta Paediatr (Oslo, Norway: 1992)*. 2018;107(3):374–375.
 53. Christensen DL, Baio J, van Naarden Braun K, et al. Prevalence and characteristics of autism spectrum disorder among children aged 8 years—autism and developmental disabilities monitoring network, 11 sites, United States, 2012. *Morb Mortal Wkly Rep*. 2016;65(3):1–23 Surveillance summaries (Washington, D.C.: 2002).
 54. Lyall K, Croen L, Daniels J, et al. The changing epidemiology of autism spectrum disorders. *Annu Rev Public Health*. 2017;38:81–102.
 55. Zwaigenbaum L, Penner M. Autism spectrum disorder: advances in diagnosis and evaluation. *BMJ*. 2018;361:k1674 Clinical research ed.
 56. Crehan ET, Baer J, Althoff RR, Constantino JN. Tracking the influence of autistic traits on competencies among school aged children with subthreshold autistic traits: a longitudinal study. *Child Psychiatry Hum Dev*. 2018 Epub 2018 May 11.
 57. Sharma SR, Gonda X, Tarazi FI. Autism spectrum disorder: classification, diagnosis and therapy. *Pharmacol Ther*. 2018 Epub 2018 May 12.
 58. Lai M-C, Lombardo MV, Baron-Cohen S. Autism. *Lancet*. 2014;383(9920):896–910.
 59. Nurenberg JR, Schleifer SJ, Shaffer TM, et al. Animal-assisted therapy with chronic psychiatric inpatients: equine-assisted psychotherapy and aggressive behavior. *Psychiatr Serv*. 2015;66(1):80–86.
 60. García-Gómez A, Rodríguez-Jiménez M, Guerrero-Barona E, Rubio-Jiménez JC, García-Peña I, Moreno-Manso JM. Benefits of an experimental program of equestrian therapy for children with ADHD. *Res Dev Disabil*. 2016;59:176–185.
 61. Benda W, McGibbon NH, Grant KL. Improvements in muscle symmetry in children with cerebral palsy after equine-assisted therapy (hippotherapy). *J Altern Complement Med*. 2003;9(6):817–825.
 62. McGibbon NH, Benda W, Duncan BR, Silkwood-Sherer D. Immediate and long-term effects of hippotherapy on symmetry of adductor muscle activity and functional ability in children with spastic cerebral palsy. *Arch Phys Med Rehabil*. 2009;90(6):966–974.
 63. Shurtleff TL, Standeven JW, Engsborg JR. Changes in dynamic trunk/head stability and functional reach after hippotherapy. *Arch Phys Med Rehabil*. 2009;90(7):1185–1195.
 64. McGibbon NH, Andrade CK, Widener G, Cintas HL. Effect of an equine-movement therapy program on gait, energy expenditure, and motor function in children with spastic cerebral palsy: a pilot study. *Dev Med Child Neurol*. 1998;40(11):754–762.
 65. Vermöhlen V, Schiller P, Schickendantz S, et al. Hippotherapy for patients with multiple sclerosis: a multicenter randomized controlled trial (MS-HIPPO). *Mult Scler*. 2018;24(September (10)):1375–1382.
 66. Schwesig R, Neumann S, Richter D, et al. Der Einfluss des Therapeutischen Reitens auf den Gang und die Haltungsverregulation. *Sportverletzung Sportschaden: Organ der Gesellschaft für Orthopädisch-Traumatologische Sportmedizin*. 2009;23(2):84–94.
 67. Wilson K, Buultjens M, Monfries M, Karimi L. Equine-assisted psychotherapy for adolescents experiencing depression and/or anxiety: a therapist's perspective. *Clin Child Psychol Psychiatry*. 2017;22(1):16–33.
 68. Hession CE, Eastwood B, Watterson D, Lehane CM, Oxley N, Murphy BA. Therapeutic horse riding improves cognition, mood arousal, and ambulation in children with dyspraxia. *J Altern Complement Med*. 2014;20(1):19–23.
 69. Grahn JA, Brett M. Rhythm and beat perception in motor areas of the brain. *J Cogn Neurosci*. 2007;19(5):893–906.
 70. Bengtsson SL, Ullén F, Ehrsson HH, et al. Listening to rhythms activates motor and premotor cortices. *Cortex*. 2009;45(1):62–71.
 71. Jang CH, Joo MC, Noh SE, et al. Effects of hippotherapy on psychosocial aspects in children with cerebral palsy and their caregivers: a pilot study. *Ann Rehabil Med*. 2016;40(2):230–236.
 72. Anestis MD, Anestis JC, Zawilinski LL, Hopkins TA, Lilienfeld SO. Equine-related treatments for mental disorders lack empirical support: a systematic review of empirical investigations. *J Clin Psychol*. 2014;70(12):1115–1132.