



## The influence of meibomian gland loss on ocular surface clinical parameters

Laura Rico-del-Viejo<sup>a,\*</sup>, José Manuel Benítez-del-Castillo<sup>b</sup>, Fernando Javier Gómez-Sanz<sup>a</sup>,  
María García-Montero<sup>a</sup>, Clara Llorens-Quintana<sup>c</sup>, David Madrid-Costa<sup>a</sup>

<sup>a</sup> Optometry and Vision Department, Faculty of Optics and Optometry, Complutense University of Madrid, Madrid, Spain

<sup>b</sup> Ophthalmology Department, San Carlos University Hospital, Complutense University, Madrid, Spain

<sup>c</sup> Department of Biomedical Engineering, Faculty of Fundamental Problems of Technology, Wrocław University of Science and Technology, Wrocław, Poland

### ARTICLE INFO

#### Keywords:

Meibomian gland loss  
Infrared Meibography  
Ocular surface  
Tear film  
Symptomatology  
Age

### ABSTRACT

**Purpose:** To assess the relationship between the meibomian gland loss (MGL) and relevant ocular surface clinical parameters as well as the influence of age in this relationship.

**Methods:** A total of 161 participants (mean age;  $42 \pm 17$  years) were enrolled in this study. Infrared meibography was performed using Keratograph 5M (K5M; Oculus GmbH, Wetzlar). Participants were divided into five groups according to total meiboscore and the ocular surface parameters of each MGL group were studied. In addition, the relationship between MGL and the ocular surface parameters was established including age as covariant.

**Results:** Both eyelids were taken into account since no association between the MGL from upper and lower eyelid was found ( $k$  value = 0.2;  $p = 0.3$ ) despite they were significantly correlated ( $r = 0.3$ ;  $p < 0.001$ ). No statistically significant differences were found in symptomatology among different MGL groups. Statistically significant differences were found among MGL groups in tear osmolarity ( $p = 0.02$ ), bulbar redness ( $p = 0.04$ ), corneal and conjunctival staining ( $p = 0.01$  and  $p = 0.004$ , respectively). Despite this, only corneal staining showed a significant correlation with MGL when age was covariant ( $r = 0.2$ ;  $p = 0.04$ ).

**Conclusions:** MGL higher than 50% seems to be accompanied by signs on the ocular surface. Furthermore, age demonstrated to be a relevant factor when assessing MGL. For this reason, future studies should compare age-matched groups in order to know the contribution of the MGL on the ocular surface and establish valid cut-off values for dry eye diagnosis.

### 1. Introduction

Dry eye disease (DED) understanding has grown over the last three decades. This multifactorial disorder is characterised by a loss of homeostasis of the tear film and accompanied by ocular symptoms, neurosensory abnormalities, ocular surface inflammation and damage [1]. DED prevalence ranges between 5% and 50% worldwide [2] and it increases linearly with age which makes DED a growing public health issue as the global population of older people is expected to be more than double its current amount by 2050 [3]. According to TFOS DEWS II DED classification, the aqueous deficiency (ADDE) and evaporative dry eye (EDE) are the two major types of DED and are considered to exist on a continuum rather than as separate entities [4]. It has been reported that the evaporative component is more common, making EDE the most common type of DED [2]. Meibomian gland dysfunction (MGD) has been considered the major cause of EDE [5]. Changes in the quantity and quality of the meibomian gland (MG) secretion lead to an

unstable lipid layer of the tear film, provoking an increase of the evaporation of the underlying aqueous layer [6]. Therefore, any change that occurs in the structure or function of MG could have an important clinical impact. Currently, non-contact infrared meibography (NIM) is widely used to assess MG non-invasively [7–9]. Several scoring systems for assessment of the meibomian gland loss (MGL) or dropout [7,10–14] have been proposed. The *meiboscore* proposed by Arita et al. [7] is one of the most commonly used scoring system for MGL evaluation. Previous studies found significant correlations between MGL and some tear film parameters (such as tear film break-up time (TBUT) [15–17], non-invasive tear film break-up time (NIBUT) [18], lipid layer thickness (LLT) [19], Schirmer test [20], MG secretion quality [16] and corneal staining [17]) as well as subjective symptomatology (Ocular-surface-disease-index (OSDI) [10] and McMonnies [21] questionnaires), suggesting its possible diagnostic value [22]. However, others studies concluded that assessing MGL alone as clinical parameter has not enough DED diagnostic value, and that it should be interpreted

\* Corresponding author at: Faculty of Optics and Optometry, Avda Arcos de Jalón, 118, 28037 Madrid, Complutense University of Madrid, Spain.  
E-mail address: [larico@ucm.es](mailto:larico@ucm.es) (L. Rico-del-Viejo).

<https://doi.org/10.1016/j.clae.2019.04.004>

Received 19 December 2018; Received in revised form 31 March 2019; Accepted 3 April 2019

1367-0484/ © 2019 British Contact Lens Association. Published by Elsevier Ltd. All rights reserved.

together with other clinical parameters [16,23,24]. Furthermore, it has been well-documented that MGL increases with age in both healthy subjects [7,15,25] and DED patients [26]. Indeed, a significant positive correlation between age and MGL was detected, indicating that the number of MG decreases with age [7]. On the other hand, in a previous study it has been observed that several ocular surface parameters such as ocular redness, corneal and conjunctival staining are highly influenced by ageing [27]. Additionally, TFOS DEWS II recently underlined that clinical DED signs increase in a higher amount by decade compared with symptoms [2]. These findings highlight the role of age in both MGL and ocular surface parameters. Thus, the aim of this prospective study is to assess the relationship between MGL and ocular surface parameters and the role of age on this relationship.

## 2. Material and methods

### 2.1. Participants

One-hundred sixty-one participants were included in this prospective study. Participants were recruited via email notices sent to the institutional e-mails from the academic community and advertisements placed on noticeboards at the university. All examinations were completed in the Faculty of Optics and Optometry at Complutense University of Madrid. This study was reviewed and approved by the Ethics Committee of San Carlos University Hospital (Madrid, Spain) and adhered to the tenets of the Declaration of Helsinki. Written informed consent was obtained from all included participants after explanation of the purpose and possible consequences of the study. Participants were required to be 18 years of age or older, be able to complete the questionnaires, understand the procedures and obtain an evaluable meibography image of the upper (UL) and lower (LL) eyelids. On the other hand, participants were excluded if they had a history of any active ocular disease different from DED and MGD (corneal ulcers, herpes simplex, keratitis, etc), any uncontrolled severe systemic disease that may affect the eye (Sjögren syndrome, diabetes type II, dermatological diseases, etc) or any ocular surgery or trauma that could affect the tear distribution and any eyelid margin abnormality. Contact lens wearers were accepted but they were required not to use their contact lenses within the week before performing the clinical examination.

### 2.2. Study protocol

All measurements were accomplished by the same examiner and performed from the least to the most invasive in order to minimize the effect of the previous measurement. Only the right eye (RE) of each participant was assessed (see Fig. 1).

#### 2.2.1. Symptomatology assessment

Participants were required to complete two of the most common DED Questionnaires during the examination: The OSDI questionnaire [28] and the Standard Patient Evaluation of Eye Dryness (SPEED) questionnaire [29].

#### 2.2.2. Tear film osmolarity

Tear film osmolarity (TFO) was measured using the TearLab Osmolarity System (TearLab Corp, San Diego, CA, USA) according to the manufacturer's instructions. It was conducted in first place in order to avoid reflex tearing or the instillation of any dye that could affect the results.

#### 2.2.3. Keratograph 5M automated measurements

Automated measurements were performed using the Keratograph 5 M (K5 M; Oculus GmbH, Wetzlar, Germany) equipped with the modified tear film scanning function. Three measurements of the tear meniscus height (TMhk), first tear film break-up time (NIK BUT-fr), the average time of all tear film break-up time incidents (NIK BUT-avg),

bulbar redness (BR) and limbal redness (LR) were obtained automatically with Oculus K5 M software according to the manufacturer's instructions. Each NIK BUT measurement was performed with at least 3 min of interval between them in order to allow tear film stability to recover. Additionally, these measurements were performed after TFO in order to avoid the influence of the tear produced by the eye during NIK BUT in the TFO values.

#### 2.2.4. Slit-lamp examination

Slit-lamp examination of the cornea and conjunctiva was performed under diffuse illumination using x10 – x16 magnification. Corneal integrity was assessed by instilling a fluorescein dye. Two minutes later, corneal staining was graded using the Oxford scoring scheme [30]. TBUT was measured three times with a stopwatch and was averaged for the subsequent analysis. Furthermore, bulbar conjunctival integrity was assessed using lissamine green and graded using the Oxford scoring scheme [30]. The meibum quality expression from the central 8 MG was assessed on a scale from 0 to 3: 0 = clear meibum readily expressed; 1 = cloudy meibum expressed with mild pressure; 2 = cloudy meibum expressed with more than moderate pressure; 3 = meibum could not be expressed even with strong pressure.

#### 2.2.5. Keratograph 5M infrared meibography

NIM was performed in order to assess MG morphology of the UL and LL of each participant. MGL of the UL and LL was graded subjectively by an experienced examiner using the *meiboscope* introduced by Arita et al. (grade 0, no gland loss; grade 1, area of gland loss < 33%; grade 2, area of gland loss 33%–67%; and grade 3, area of gland loss > 67%) [31]. The meiboscope for each eyelid was summed to give a total score between 0 and 6 (or from 0 to 6). This *total meiboscope* was used to divide participants into five groups according with the amount of MGL (see Fig. 2). In this regard, the group 1 was constituted by participants who presented a total meiboscope of 0; group 2 by participants who showed a total meiboscope of 1; group 3 by participants who showed a total meiboscope of 2; group 4 by participants who showed a total meiboscope of 3 and group 5 by participants who showed a total meiboscope of 4, 5 or 6. According the groups established, the percentage range of MGL was 0, 0–16.6, 16.5–33, 33–49.5 and > 50, for groups 1, 2, 3, 4 and 5 respectively.

#### 2.2.6. Tear film volume

Schirmer's test was performed with topical anaesthesia (Colirio Anestésico Doble®, Alcon Laboratories, Spain) and was the final test carried out in the examination protocol. One drop of topical anaesthesia was instilled on the conjunctival lower fornix of the RE, 5 min prior to do the test. Then, the Schirmer strip (35-mm Whatman filter paper; Tiedra Laboratories, Spain) was placed in the lower conjunctival sac at the junction of the lateral and middle thirds (avoiding contact with the cornea) and after 5 min the length of wetting was recorded. Participants were seated at rest and were asked to close the eyes during the test.

### 2.3. Data analysis

The values are expressed as means  $\pm$  standard deviation (SD) and the significance level was set  $p < 0.05$  with 95% of confidence level. Normality of the data distribution was tested using the Kolmogorov–Smirnov test. Accordingly, Cohen's kappa coefficient (weighted kappa value- 95% of confidence) was calculated and classified as follow: 0.00 (poor), 0.00–0.20 (slight), 0.21–0.40 (fair), 0.41–0.60 (moderate), 0.61–0.80 (substantial) and 0.80–1.0 (close to perfect) [32]. Kruskal–Wallis test was used for comparisons between different MGL groups. When statistically significant differences were found ( $p < 0.05$ ), post hoc tests were performed for multiple comparisons applying Bonferroni corrections. Correlation coefficients between MGL of both eyelids and ocular surface parameters were calculated with the Spearman correlation coefficient. In addition, partial




Fig. 1. Flow diagram of the study protocol. Numbers indicate the order of performance of each test (from less invasive towards to the most invasive).

correlation was performed including age as a covariant. These correlations were considered strong if they were > 0.80, moderately strong if they were between 0.5 and 0.8, fair if they were within the range of 0.3 and 0.5 and poor if they were < 0.30 [33].


### 3. Results

The demographic characteristics of the participants are summarized in Table 1. From the 161 participants, 91 were females and 70 males. Age range among participants was 19–88 years with a mean age of 42 ± 17 years.

#### 3.1. Association and correlation between UL and LL

A contingency table was used to compare MGL between UL and LL (Table 2) in order to know if it was necessary to evaluate the MGL in one eyelid or in both. Weighted kappa statistics showed no statistically significant agreement (weighted k value = 0.226; p = 0.372; range 0.099–0.353, 95% confidence limit) indicating no association between both eyelids. Despite it, a fair and statistically significant correlation was found between UL and LL (Spearman: r = 0.308; p < 0.001). For this reason, both eyelids should be assessed in order to know the overall condition of MG and its possible influence in the ocular surface state.

Table 3 shows the demographics of the participants as a function of total meiboscore.



**Fig. 2.** Different grades of MGL in the UL and LL obtained by non-contact infrared meibography (Oculus Keratograph K5M). **A.1** and **A.2** Participants without MGL in the UL and LL (total meiboscore 0; = 0%). **B.1** and **B.2** Participants with slight MGL in the UL and LL (total meiboscore 1; < 33%). **C.1** and **C.2** Participants with moderate MGL in the UL and LL (total meiboscore 2; 33–66%). **D.1** and **D.2** Participant with severe MGL in the UL and LL (total meiboscore 3; > 66%).

**Table 1**

Demographic data from the participants of the study. Results are expressed in mean ± standard deviation (SD) for each parameter.

N	161
Age (years)	42 ± 17
Age range (years)	19 to 88
Male/ Female (%)	43/57
OSDI (scores)	17.47 ± 15.65
SPEED (scores)	7 ± 5
TFO (mOsm/L)	309 ± 17
TMHk (mm)	0.26 ± 0.08
REDNESS (scores)	
Limb	0.70 ± 0.37
Bulbar	1.11 ± 0.49
NIK BUT (seconds)	
First	9.05 ± 5.41
Average	12.02 ± 5.26
TBUT (seconds)	4.36 ± 2.07
SCHIRMER TEST (mm)	12 ± 7

OSDI: The Ocular-surface-disease-index; SPEED: Standard Patient Evaluation of Eye Dryness; TFO: tear film osmolarity; TMHk: tear meniscus height; Bulbar and limbal redness were graded automatically by K5M software; NIK BUT-fr: first rupture non-invasive Keratograph tear film break-up time; NIK BUT-avg: average of non-invasive Keratograph tear film break-up time.

**3.2. Symptomatology assessment**

Results regarding subjective symptomatology are in Table 4. No statistically significant differences were found in OSDI (p = 0.385) and SPEED (p = 0.506) questionnaires among different MGL groups.

**3.3. Classical clinical parameters**

Classical clinical parameters results for each group are summarized in Table 5. Statistically significant differences were found among groups in TFO (p = 0.023), corneal (p = 0.015) and conjunctival staining (p = 0.004). For TFO, there were statistically significant differences between group 5 and groups 1,2 and 3 (p = 0.029; p = 0.014; p = 0.001, respectively). For corneal staining, there were statistically significant differences between group 5 and groups 1 and 2 (p = 0.045 and 0.003, respectively). Regarding conjunctival staining, there were also statistically significant differences between group 5 and groups 1 and 2 (p = 0.027 and 0.007, respectively) and between group 4 and groups 1 and 2 (p = 0.045 and 0.014, respectively).

**Table 2**

Contingency table of MGL for UL and LL.

		Upper Eyelid				
Lower Eyelid	Meiboscore*	0	1	2	3	Row Totals
0	0	21	22	4	0	47
		44.68%	46.81%	8.51%	0%	
1	0	36	36	12	2	83
		39.76%	43.37%	14.46%	2.41%	
2	0	57	53	48	28	233
		57.89%	53.73%	48.00%	28.51%	
3	0	3	8	9	3	23
		13.04%	34.78%	39.13%	13.04%	
3	0	5	11	36	42	94
		5.26%	11.94%	36.00%	42.86%	
3	0	0	1	0	2	3
		0%	33.33%	0%	66.67%	
3	0	0	1	0	2	3
		0%	1.49%	0%	28.57%	
Column Totals		57	67	25	7	156

UL: Upper eyelid; LL: Lower eyelid; MGL: Meibomian gland loss.

\* Meiboscore: Grade 0, no gland loss; grade 1, area of gland loss < 33% of the total gland area; grade 2, area of gland loss 33%–67%; and grade 3, area of gland loss > 67%.

**3.4. Keratograph 5M automated measurements**

Table 6 shows the automated measurements obtained with K5 M for each meiboscore group. Statistically significant differences in BR were found between groups 5 and 1 (p = 0.043) and groups 5 and 2 (p = 0.010).

**3.5. Relationship between MGL and ocular surface parameters considering age as covariant**

Partial correlations between MGL and ocular surface parameters are in Table 7. No relationship was found between MGL and any ocular surface parameter except for corneal staining (r = 0.208, p = 0.041) when age was included as covariant.

**4. Discussion**

NIM has been proven to be a useful technique for non-invasive observation of MG morphology in order to help physicians improve DED diagnosis and treatments [7,34,35].

The present study assesses the correlation between MGL and some other ocular surface parameters. Accordingly, first, it has to be

**Table 3**

Characteristics and distribution of the sample according to MGL grade (meiboscure). Results expressed in mean ± standard deviation (SD) and percentage (%).

Total Meiboscure <sup>a</sup>	Group 1 0 (reference)	Group 2 1	Group 3 2	Group 4 3	Group 5 4,5 and 6
MGL (% range)	0	[0-16.5]	[16.5-33]	[33-49.5]	> 50
N (%)	21 (13.0)	59 (36.7)	44 (27.3)	20 (12.4)	17 (10.6)
AGE (years)	34 ± 11	37 ± 15	46 ± 17	50 ± 18	60 ± 18
FEMALE n/N (%)	9/91 (9.9)	33/91 (36.3)	27/91 (29.7)	8/91 (8.8)	12/91 (13.2)
MALE n/N (%)	12/70 (17.1)	25/70 (35.7)	17/70 (24.3)	12/70 (17.1)	4/70 (5.7)

\* The total meiboscure is the sum of the meiboscure of both eyelids (0–6).

**Table 4**

Comparison of symptomatology scores among different MGL grades. Results expressed in mean ± standard deviation (SD) for each parameter (units).

TOTAL MEIBOSCORE GROUP (MGL%)	SYMPTOMATOLOGY	
	OSDI (scores)	SPEED (scores)
Group 1 (0)	14.30 ± 13.95	7 ± 5
Group 2 (0-16.5)	16.34 ± 13.90	7 ± 5
Group 3 (16.5-33)	18.30 ± 16.76	7 ± 5
Group 4 (33-49.5)	15.61 ± 15.21	6 ± 5
Group 5 (> 50)	23.28 ± 23.28	9 ± 5
p-value	0.385	0.506

\*statistically significant differences among groups; p < 0.05.

OSDI: The Ocular-surface-disease-index; SPEED: Standard Patient Evaluation of Eye Dryness.

determined if both (UL and LL) or only one eyelid should be included for MGL assessment. This matter is quite controversial since several studies have considered different criteria to choose both or only one eyelid. Routinely, LL is the most commonly assessed due to its accessibility which provokes less discomfort to the patient. It is believed that comparable outcome can be expected for MGL assessment choosing one or two eyelids, as UL and LL MGL seem to be significantly correlated [18]. Dogan et al. [36] assessed meibography images of 30 patients and proposed to evaluate only the UL for MGL assessment because of its correlation with TBUT and better inter-examiner agreement on MGL. On the other hand, Finis et al. [16] performed a retrospective analysis of 128 patients and found a strong correlation between meiboscure of the UL and the LL as well as with the total meiboscure. This suggests that MGL assessment based on the evaluation of the LL might be enough for the clinical routine. In the present study a positive and fair statistically significant correlation between MGL of UL and LL was found. However, there was no agreement as revealed by the Kappa statistic, suggesting that, despite there was a relationship between the MGL of both eyelids, exist a bias in their meiboscure. The findings of the present study could support those found by Pult et al. [18], who suggested the MGL assessment of both eyelids.

After concluding that both eyelids should be taken into account in order to assess MGL, participants were classified into five groups according to the total meiboscure [7].

**Table 5**

Comparison of classical clinical parameters among different MGL grades. Results expressed in mean ± standard deviation (SD) for each parameter (units).

TOTAL MEIBOSCORE GROUP (MGL%)	CLASSICAL CLINICAL PARAMETERS				
	TFO (mOsm/L)	TBUT (seconds)	SCHIRMER (mm)	CORNEAL STAINING (scores)	CONJUNCTIVAL STAINING (scores)
Group 1 (0)	308 ± 16	4.86 ± 2.52	13 ± 6	0.76 ± 0.70	1.19 ± 0.68
Group 2 (0-16.5)	309 ± 19	4.85 ± 2.57	13 ± 8	0.62 ± 0.85	1.20 ± 0.83
Group 3 (16.5-33)	304 ± 15	4.04 ± 1.46	11 ± 7	0.98 ± 0.86	1.67 ± 1.02
Group 4 (33-49.5)	312 ± 12	3.96 ± 1.22	10 ± 7	0.85 ± 0.67	1.75 ± 0.79
Group 5 (> 50)	326 ± 18	3.64 ± 1.51	11 ± 4	1.47 ± 1.23	1.88 ± 0.93
p-value	0.023*	0.249	0.160	0.015*	0.004*

TFO: tear film osmolarity; TBUT: tear film break-up time; Corneal and conjunctival staining were graded using Oxford Staining Score System.

\* statistically significant differences among groups; p < 0.05.

DED symptomatology was evaluated using OSDI and SPEED questionnaires in the current study. No statistically significant differences were found in DED questionnaires among different MGL groups. However, as Table 5 shows, the symptomatology score for group 5 was, in both questionnaires, clinically higher than other groups. It has been reported that a MGL of > 32% is likely to be accompanied with associated detectable clinical symptoms [10]. Similarly, several studies have found correlation between MGL and OSDI scores [25,37] and others did not [26,36], neither with SPEED questionnaire [38].

Regarding MGL and its correlation with other clinical parameters, statistically significant differences were found in TFO between group 5 and groups 1,2 and 3. Besides, statistically significant differences were found in corneal staining as well as conjunctival staining between group 5 and groups 1 and 2. These results are in agreement with those obtained by Feng et al. [26] who found a positive correlation between corneal staining and MGL in DED patients. Previous studies have suggested that when the amount of MG is reduced, the secretion of MG decreases. It might induce tear film homeostasis loss and greater tear film evaporation which could lead to surface epithelial damage as well as disturbance of the glycocalyx and goblet cell mucins [39].

In the present study no differences were found in TBUT and Schirmer test among different MGL groups. These results are in accordance with other studies that did not find any correlation between MGL and these tear film parameters [40] or, instead, a low correlation [26]. Nevertheless, Arita et al. [41] studied a population with MGD and found that Schirmer test was positively correlated with the meiboscure. Thus, patients with less amount or damaged MG would have an increase of fluid that may compensate the decreased function of the lipid layer. All these findings together suggest that these ocular surface parameters could be affected by MGL in patients who suffer from MGD since the glandular loss may exacerbate the signs of MGD in comparison with those who only present MGL without other ocular condition.

On the other hand, no statistically significant differences were found in any of the 5KM parameters among MGL groups except for BR. Recently, Ji et al. [42] found a correlation between MGL grade and NIKBUT-avg, NIKBUT and LLT in patients with DED and MGD. This discrepancy between studies could be due to the population studied. Indeed, the correlation found in this study might be because only subjects with DED and MGD were included.

The findings of the present study suggest that a MGL higher than

**Table 6**Comparison of K5M automated measurements among different MGL grades. Results expressed in mean  $\pm$  standard deviation (SD) for each parameter (units).

TOTAL MEIBOSCORE GROUP (MGL%)	K5M AUTOMATED MEASUREMENTS				
	TMHk (mm)	BR (scores)	LR (scores)	NIK BUT-first (seconds)	NIK BUT-avg (seconds)
<b>Group 1</b> (0)	0.23 $\pm$ 0.05	1.00 $\pm$ 0.46	0.68 $\pm$ 0.35	11.23 $\pm$ 6.82	13.81 $\pm$ 6.15
<b>Group 2</b> (0-16.5)	0.26 $\pm$ 0.71	0.99 $\pm$ 0.44	0.62 $\pm$ 0.32	8.79 $\pm$ 5.12	11.80 $\pm$ 5.02
<b>Group 3</b> (16.5-33)	0.27 $\pm$ 0.89	1.16 $\pm$ 0.55	0.70 $\pm$ 0.40	8.83 $\pm$ 4.95	12.32 $\pm$ 4.97
<b>Group 4</b> (33-49.5)	0.26 $\pm$ 0.80	1.25 $\pm$ 0.53	0.76 $\pm$ 0.37	10.19 $\pm$ 5.96	12.48 $\pm$ 5.52
<b>Group 5</b> (> 50)	0.28 $\pm$ 0.96	1.30 $\pm$ 0.41	0.92 $\pm$ 0.41	6.76 $\pm$ 4.08	10.07 $\pm$ 4.95
p-value	0.405	0.040*	0.063	0.213	0.427

TMHk: tear meniscus height; Bulbar (BR) and limbal redness (LR) were graded automatically by K5 M software; NIK BUT-fr: first rupture non-invasive. Keratograph tear film break-up time; NIK BUT-avg: average of non-invasive Keratograph tear film break-up time.

\* statistically significant differences among groups;  $p < 0.05$ .

**Table 7**

Relationship between MGL and the ocular surface parameters considering age as covariant.

	MGL (Covariant: Age)	
	Correlation coefficient	<i>p</i>
<b>DED Questionnaires</b>		
OSDI	0.011	0.914
SPEED	0.012	0.904
<b>K5 M Parameters</b>		
TMHk	0.059	0.563
BR	-0.003	0.969
LR	-0.044	0.668
NIK BUT-fr	-0.095	0.345
NIK BUT-avg	-0.054	0.597
<b>Classical Clinical Parameters</b>		
TFO	0.088	0.393
Corneal staining	0.208	0.041*
Conjunctival staining	-0.004	0.966
TBUT	-0.163	0.112
Schirmer test	-0.065	0.527
<b>MG features</b>		
MG quality secretion	0.044	0.587

(*r*, Spearman correlation coefficient).

OSDI: The Ocular-surface-disease-index (scores); SPEED: Standard Patient Evaluation of Eye Dryness (scores). TMHk: tear meniscus height (mm); Bulbar and limbal redness (BR and LR) were graded automatically by K5 M software; NIK BUT-first: first rupture non-invasive Keratograph tear film break-up time (seconds); NIK BUT-avg: average of non-invasive Keratograph tear film break-up time (seconds). TFO: tear film osmolarity(mOsm/L); TBUT: tear film break-up time (seconds); Schirmer test (mm); Corneal and conjunctival staining were graded using Oxford Staining Score System; MG: Meibomian glands.

\* statistically significant;  $p < 0.05$ .

50% is accompanied by signs of increased osmolarity, redness and staining of the ocular surface. However, it is important to highlight that the mean age of the participants in this study was higher in those groups with higher MGL. Therefore, it is not possible to ascertain if these signs in ocular surface parameters are due to MGL or to age. Indeed, the great influence of aging on MG morphology and function is well-known and documented in the literature [7,18,27,40]. In order to address this point, the relationship between MGL and the ocular surface parameters was assessed considering age as a covariant. However, when it was performed, only the corneal staining was correlated with MGL. These results emphasize the influence of ageing in the MG morphology but also in several ocular surface parameters such as corneal staining, LLT or tear volume among others [27,43]. It has been previously reported [27] and points out the importance of considering participants age when performing research studies focused on the ocular surface. Thus, in order to assess the real impact or influence of the MGL in ocular surface parameters (which are influenced by age), it would be necessary to compare age-matched groups. For example, MGL is positively correlated with meibum quality, suggesting an impaired

gland function when the amount of MGL increase [19,42,44]. Despite this, when age is covariant, the relationship between MGL and meibum quality is absent, indicating that meibum quality could be decreased either because of a higher amount of MGL or because of it naturally decreased with ageing. These findings suggest that different thresholds should be considered regarding abnormal ocular and tear film surface parameters according to age. The next step should be focused on determining the normal threshold of each parameter for each age range.

In summary, the current study suggests that a MGL higher than 50% is accompanied by signs in the ocular surface. Furthermore, in the light of these findings, future studies must consider age due to the great influence on the MG morphology and compare age-matched groups in order to comprehend the contribution of the MGL on the ocular surface as well as establish valid cut-off values for DED diagnosis.

## Funding

This research has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 642760.

## Disclosure

The authors have no proprietary interest in any of the devices mentioned in this article.

## Acknowledgements

This research has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 642760.

## References

- [1] J.P. Craig, J.D. Nelson, D.T. Azar, C. Belmonte, A.J. Bron, S.K. Chauhan, et al., The TFOS dry eye workshop II: executive summary, *Ocul Surf* 15 (4) (2017) 802–812.
- [2] F. Stapleton, M. Alves, V.Y. Bunya, I. Jalbert, K. Lekhanont, F. Malet, et al., TFOS DEWS II epidemiology report, *Ocul Surf* 15 (2017) 334–365.
- [3] United Nations, Department of economic and social affairs - population division, *World Population Ageing*, 2017.
- [4] J.P. Craig, K.K. Nichols, J.J. Nichols, B. Caffery, H.S. Dua, E.K. Akpek, et al., TFOS DEWS II definition and classification report, *Ocul Surf* 15 (2017) 276–283.
- [5] K.K. Nichols, G.N. Foulks, A.J. Bron, B.J. Glasgow, M. Dogru, K. Tsubota, et al., The international workshop on meibomian gland dysfunction: executive summary, *Investig Ophthalmol Vis Sci* 52 (2011) 1922–1929.
- [6] A.J. Bron, C.S. de Paiva, S.K. Chauhan, S. Bonini, E.E. Gabison, S. Jain, et al., TFOS DEWS II pathophysiology report, *Ocul Surf* 15 (2017) 438–510.
- [7] R. Arita, K. Itoh, K. Inoue, S. Amano, Noncontact infrared meibography to document age-related changes of the meibomian glands in a normal population, *Ophthalmology* 115 (2008) 911–915.
- [8] R. Arita, K. Itoh, S. Maeda, K. Maeda, S. Amano, A newly developed noninvasive and mobile pen-shaped meibography system, *Cornea* 32 (2013) 242–247.
- [9] H.S. Hwang, C.W. Park, C.K. Joo, Novel noncontact meibography with anterior segment optical coherence tomography: hosik meibography, *Cornea* 32 (2013) 40–43.

- [10] H. Pult, B.H. Riede-Pult, Non-contact meibography: keep it simple but effective, *Contact Lens Anterior Eye* 35 (2012) 77–80.
- [11] J.J. Nichols, D.A. Berntsen, G.L. Mitchell, K.K. Nichols, An assessment of grading scales for meibography images, *Cornea* 24 (2005) 382–388.
- [12] W.D. Mathers, M. Billborough, Meibomian gland function and giant papillary conjunctivitis, *Am J Ophthalmol* 114 (1992) 188–192.
- [13] J. Shimazaki, M. Sakata, K. Tsubota, Ocular surface changes and discomfort in patients with meibomian gland dysfunction, *Arch Ophthalmol* 113 (1995) 1266–1270.
- [14] S.C. Pflugfelder, S.C.G. Tseng, O. Sanabria, H. Kell, C.G. Garcia, C. Felix, et al., Evaluation of subjective assessments and objective diagnostic tests for diagnosing tear-film disorders known to cause ocular irritation, *Cornea* 17 (1998) 38–56.
- [15] Y. Ban, S. Shimazaki-Den, K. Tsubota, J. Shimazaki, Morphological evaluation of meibomian glands using noncontact infrared meibography, *Ocul Surf* 11 (2013) 47–53.
- [16] D. Finis, P. Ackermann, N. Pischel, C. König, J. Hayajneh, M. Borrelli, et al., Evaluation of meibomian gland dysfunction and local distribution of meibomian gland atrophy by non-contact infrared meibography, *Curr Eye Res* 40 (2015) 982–989.
- [17] Y. Yin, L. Gong, Uneven meibomian gland dropout over the tarsal plate and its correlation with meibomian gland dysfunction, *Cornea* 34 (2015) 1200–1205.
- [18] H. Pult, B.H. Riede-Pult, J.J. Nichols, Relation between upper and lower lids' meibomian gland morphology, tear film, and dry eye, *Optom Vis Sci* 89 (2012) E310–5.
- [19] Y. Eom, J.S. Lee, S.Y. Kang, H.M. Kim, J.S. Song, Correlation between quantitative measurements of tear film lipid layer thickness and meibomian gland loss in patients with obstructive meibomian gland dysfunction and normal controls, *Am J Ophthalmol* 155 (2013) 1104–1110.
- [20] T. Mizoguchi, R. Arita, S. Fukuoka, N. Morishige, Morphology and function of meibomian glands and other tear film parameters in junior high school students, *Cornea* 36 (2017) 922–926.
- [21] P.R.K. Turnbull, S.L. Misra, J.P. Craig, Comparison of treatment effect across varying severities of meibomian gland dropout, *Cont Lens Anterior Eye* 41 (2017) 88–92.
- [22] G. Geerling, C. Baudouin, P. Aragona, M. Rolando, K.G. Boboridis, J.M. Benítez-del-Castillo, et al., Emerging strategies for the diagnosis and treatment of meibomian gland dysfunction: proceedings of the OCEAN group meeting, *Ocul Surf* 15 (2017) 179–192.
- [23] R. Arita, K. Itoh, S. Maeda, K. Maeda, A. Furuta, S. Fukuoka, et al., Proposed diagnostic criteria for obstructive meibomian gland dysfunction, *Ophthalmology* 116 (2009) 2058–2063.
- [24] M.A. Lemp, L.A. Crews, A.J. Bron, G.N. Foulks, B.D. Sullivan, Distribution of aqueous-deficient and evaporative dry eye in a clinic-based patient cohort: a retrospective study, *Cornea* 31 (2012) 472–478.
- [25] N.S. Yeotikar, H. Zhu, M. Markoulli, K.K. Nichols, T. Naduvilath, E.B. Papas, Functional and morphologic changes of meibomian glands in an asymptomatic adult population, *Investig Ophthalmol Vis Sci* 57 (2016) 3996–4007.
- [26] Y. Feng, Z. Gao, K. Feng, H. Qu, J. Hong, Meibomian gland dropout in patients with dry eye disease in China, *Curr Eye Res* 39 (2014) 965–972.
- [27] L. Rico-del-Viejo, A. Lorente-Velázquez, J.L. Hernández-Verdejo, R. García-Mata, J.M. Benítez-del-Castillo, D. Madrid-Costa, The effect of ageing on the ocular surface parameters, *Cont Lens Anterior Eye* 41 (2018) 5–12.
- [28] R.M. Schiffman, M.D. Christianson, G. Jacobsen, J.D. Hirsch, B.L. Reis, Reliability and validity of the ocular surface disease index, *Arch Ophthalmol* 118 (2000) 615–621.
- [29] W. Ngo, P. Situ, N. Keir, D. Korb, C. Blackie, T. Simpson, Psychometric properties and validation of the standard patient evaluation of eye dryness questionnaire, *Cornea* 32 (2013) 1204–1210.
- [30] A.J. Bron, V.E. Evans, J.A. Smith, Grading of corneal and conjunctival staining in the context of other dry eye tests, *Cornea* 22 (2003) 640–650.
- [31] R. Arita, K. Itoh, K.A.S. Inoue, Noncontact infrared meibography to document age-related changes of the meibomian glands in a normal population, *Ophthalmology* 15 (2008) 911–915.
- [32] J.R.K.G. Landis, The measurement of observer agreement for categorical data, *Biometrics* 33 (1977) 159–174.
- [33] Y.H. Chang, *Biostatistics 104: correlational analysis*, Singapore Med J (44) (2003) 614–619.
- [34] H. Pult, J.J. Nichols, A review of meibography, *Optom Vis Sci* 89 (2012) E760–E769.
- [35] Arita R. Meibography: a japanese perspective, *Invest Ophthalmol Vis Sci* 59 (2018) DES48–DES55.
- [36] A.S. Dogan, M. Kosker, N. Arslan, C. Gurdal, Interexaminer reliability of meibography: upper or lower eyelid? *Eye Cont Lens* 44 (2018) 113–117.
- [37] H. Pult, Relationships between meibomian gland loss and age, sex, and dry eye, *Eye Cont Lens* 44 (2018) S318–S324.
- [38] P.K. Gupta, M.N. Stevens, N. Kashyap, Y. Priestley, Prevalence of meibomian gland atrophy in a pediatric population, *Cornea* 37 (2018) 426–430.
- [39] A. Lemp, The definition and classification of dry eye disease: report of the definition and classification of the Dry Eye WorkShop (2007), *Ocul Surf* 5 (2007).
- [40] A. Machalińska, A. Zakrzewska, K. Safranow, B. Wiszniewska, B. Machaliński, Risk factors and symptoms of meibomian gland loss in a healthy population, *J Ophthalmol* 2016 (2016) 1–8.
- [41] R. Arita, N. Morishige, S. Koh, R. Shirakawa, M. Kawashima, T. Sakimoto, et al., Increased tear fluid production as a compensatory response to meibomian gland loss: a multicenter cross-sectional study, *Ophthalmology* 122 (2015) 925–933.
- [42] Y.W. Ji, J. Lee, H. Lee, K.Y. Seo, E.K. Kim, T.I. Kim, Automated measurement of tear film dynamics and lipid layer thickness for assessment of non-sjögren dry eye syndrome with meibomian gland dysfunction, *Cornea* 36 (2017) 176–182.
- [43] C. Maïssa, M. Guillon, Tear film dynamics and lipid layer characteristics-Effect of age and gender, *Cont Lens Anterior Eye* 33 (2010) 176–182.
- [44] Y. Eom, K.E. Choi, S.Y. Kang, H.K. Lee, H.M. Kim, J.S. Song, Comparison of meibomian gland loss and expressed meibum grade between the upper and lower eyelids in patients with obstructive meibomian gland dysfunction, *Cornea* 33 (2014) 448–452.