


HOSTED BY



Contents lists available at ScienceDirect

# International Journal of Pediatrics and Adolescent Medicine

journal homepage: <http://www.elsevier.com/locate/ijpam>

## Stressful life events, gender and obesity: A prospective, population-based study of adolescents in British Columbia

A.I. Conklin<sup>a, b, \*</sup>, S.X.R. Guo<sup>c</sup>, C.A. Yao<sup>b</sup>, A.C.T. Tam<sup>d</sup>, C.G. Richardson<sup>b, e</sup><sup>a</sup> Collaboration for Outcomes Research and Evaluation (CORE), Faculty of Pharmaceutical Sciences, University of British Columbia, Vancouver, BC, Canada<sup>b</sup> Centre for Health Evaluation and Outcome Sciences (CHEOS), Providence Healthcare Research Institute, St. Paul's Hospital, Vancouver, BC, Canada<sup>c</sup> Behavioural Neuroscience, Faculty of Science, University of British Columbia, Vancouver, BC, Canada<sup>d</sup> Psychology, Faculty of Arts, University of British Columbia, Vancouver, Canada<sup>e</sup> School of Population and Public Health, Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada

### ARTICLE INFO

#### Article history:

Received 26 December 2018

Accepted 3 March 2019

Available online 11 March 2019

#### Keywords:

Life change events

Obesity

Adolescents

Gender

Cohort studies

### ABSTRACT

**Objective:** To determine whether stressful life events are related to levels of obesity in a group of ethnically diverse Canadian youth and the extent to which the relationship differs by gender.

**Methods:** This study of 905 adolescents (age 13–17 years) from a BC population-based cohort (BASUS) used self-reported data from Wave 5 (2011 fall) on stressful life events and socio-demographic factors and from Wave 6 (2012 spring) on weight and height. Multivariable logistic regression models conditioned on known confounders and used a cross-product term for effect modification by gender. Post-estimation analysis calculated gender-specific predicted mean probabilities of having obesity associated with greater frequency of stressful life events.

**Results:** Compared to young men reporting no stressful life events in the previous year, young men reporting one event were nearly 50% more likely to have obesity at 6-month follow-up (OR 1.47 [95% CI: 0.63, 3.41]) and those reporting multiple stressful life events were twice as likely to have obesity at 6-month follow-up (OR 2.07 [95% CI: 0.79–5.43]). Only young women reporting multiple events showed a higher likelihood of having obesity at the end of the study (OR 1.32 [95% CI: 0.41–4.18]) than their counterparts reporting no life events.

**Conclusions:** Results suggest that the frequency of major life events may be an important social stressor associated with obesity in adolescents, particularly for young men. However, findings should be replicated in larger samples using measured anthropometry to inform future obesity prevention strategies.

© 2019 Publishing services provided by Elsevier B.V. on behalf of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 1. Introduction

Adolescent obesity is a global health concern despite serious efforts towards its mitigation [1]. A third of young people in Canada have excess weight [2], which is a substantial public health challenge, as obesity is a strong risk factor for numerous chronic conditions [3,4], and 80% of adolescents with obesity have it in adulthood also [5]. Although weight gain is common during adolescence, stress-related weight gain is also likely to increase

during this developmental period, as adolescents are particularly vulnerable to disturbances of the stress response pathways [6,7]. Understanding the role of social stressors such as adverse life events is critical to design effective obesity interventions for young people who are greatly need it, as shown in a recent Cochrane review [8].

Young people experience a unique period of developmental vulnerability so that major life events like social stressors can permanently impact their brain development as well as their metabolically relevant systems, which, in turn, could negatively affect proper physical functioning and emotional regulation during adulthood [9]. Unlike other age groups, young people are distinct insofar, as they are more vulnerable to the health impact of stress while also being more likely to have little control over the stressful life events that they experience [10]. The independent stressors

\* Corresponding author. University of British Columbia, 2405 Westbrook Mall, Vancouver, BC, V6T 1Z3, Canada.

E-mail address: [annalijn.conklin@ubc.ca](mailto:annalijn.conklin@ubc.ca) (A.I. Conklin).

Peer review under responsibility of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia.

reported by young people range in magnitude from more common everyday life events (25%) such as separating from a romantic partner or parental divorce to more extreme but less common life events (6.2%), including death of a parent or surviving a violent act [11]. By the age of 16 years, it is estimated that a quarter of young people will have experienced at least one high-magnitude stressor [11]. As compared to adults, young people will be more emotionally vulnerable to social stressors because their development-specific brain alterations mean they have higher basal levels of hypothalamic–pituitary–adrenal (HPA) activity and heightened biological stress reactivity [12]. Higher HPA activation and cortisol levels have known effects on central adiposity [13].

There is some evidence to show that adolescents' differential brain development is associated with greater responsiveness to emotion-related social cues and incentives, which occurs at the time when they have underdeveloped neurological pathways that are responsible for rational decision-making and resisting temptations [14]. In this development context, social stressors that induce changes to the brain in young people could have negative metabolic consequences by making them more susceptible to the hedonic effects of highly palatable foods (i.e. rich in sugar and fat) [15]. The consumption of energy-dense foods triggers the brain's reward pathway and becomes a reinforcing behaviour in the same way as that of drug use in young people, underpinned by the principles of reward and reinforcement [16]. It is also important to consider how stressful life events affect weight-related outcomes differently for young women and young men, given sex hormone differences in HPA activation and stress reactivity [17].

Thus, behavioural alterations that occur from physiological adaptations or from coping responses to stressful life events constitute another pathway to poor metabolic outcomes [18]. More specifically, social stressors could lead to young people not only eating poor-quality diets but also sitting more, exercising and sleeping less as well as consuming illegal substances such as tobacco and alcohol—all behaviours that have been associated with adiposity.

Indeed, sleep deprivation is associated with obesity, and the interaction between stressful life events and altered sleep patterns could create a potentially damaging nexus of negative metabolic effects [19]. Social stressors can induce changes to sleep–wake cycles that amplify the negative effects of development-specific sleep deprivation on satiety hormone (leptin) regulation [20] and emotional regulation [21] and thus lead to higher intakes of palatable, energy-dense foods and excess weight increase [22]. Such behavioural changes and lifestyle habits formed during adolescence may endure into adulthood due to the underlying physiological development [23] and could help explain why adolescent obesity predicts adult risk more than the risk in young age groups [5].

Again, the impact of social stressors on obesity prevalence may differ between young women and young men because sex chromosomes are known to regulate habit formation [24], and both the prevalence of sleep problems [25] and the amount of sleep deprivation required to affect metabolic outcomes is greater in young women than in young men [26]. However, few studies apply gender-based analysis to address the extent to which women and men differ in their exposures to stressful life events is associated with differential vulnerability to poor health [27], and current evidence on the metabolic effects of stressful life events is limited to cross-sectional designs and populations of children or adults only [28,29]. This study aimed to investigate the relationship between stressful life events and obesity in a population-based cohort of young people using a gender perspective [30]. In applying a gender perspective similar to that in our previous work [31], we hypothesized that greater levels of stressful life events would be associated

with higher obesity at follow-up and that the associations would show gender-specific patterning.

## 2. Subjects and methods

### 2.1. Study population

This study used data from participants in the population-based *British Columbia Adolescent Substance Use Survey (BASUS)* cohort study that comprised 3170 students aged 13–18 years from 86 secondary schools in British Columbia, Canada [31], who completed biannual online surveys from October 2009 (Wave 1) to December 2012 (Wave 7). We used self-reported data from Wave 5 (2011 fall) on stressful life events ( $n = 1543$ ), sleep and wake times ( $n = 1469$ ) and socio-demographics ( $n = 973$ ) and from Wave 6 (2012 spring) on weight and height ( $n = 1575$ ). Written informed consent was obtained from all volunteer students following school district procedures; ethics approval for the BASUS study (no. H08–02841) was given by the Behavioural Research Ethics Board of the University of British Columbia [31,32].

### 2.2. Measures

#### 2.2.1. Stressful life events

We used the junior high life experiences survey [33] listing 13 events specific to assess exposure to stressful life events, similar to another work [34]. The survey asks participants to report retrospectively over 12 months whether or not they experienced any of the following events: *family moved to new house, close friend committed suicide, family member committed suicide, parents separated/divorced, one parent died, participant found out about their being adopted, first-time job loss, victim of violence, broke up with girlfriend/boyfriend, first intercourse, had sexually transmitted disease, expelled from school, first-time police involvement* [33,34]. Participants reported whether or not they experienced a given event in the previous 12 months. The total number of events was calculated, and students were classified into a three-level categorical variable for their exposure to zero, one and multiple ( $\geq 2$ ) recent stressful life events.

#### 2.2.2. Obesity status

We measured obesity status at study follow-up based on body mass index (BMI), which we calculated from self-reported data on height and weight. Gender- and age-specific thresholds for BMI were used to classify participants as either having obesity ( $>97$ th percentile) or not (3rd to 97th percentile), based on the Dietitians of Canada WHO Growth Charts for Canada [35]. We therefore excluded participants with BMI-for-age percentiles falling below the normal weight category ( $n = 43$ ) and those with BMI-for-age values exceeding the range in the charts ( $n = 18$ ).

#### 2.2.3. Covariates

Concurrent socio-demographic variables included ethnicity (Caucasian [reference], Asian, Aboriginal and Other [Latino, Black, and others]), maternal education (university or higher [reference], some vocation or college, secondary school and below secondary school) and pubertal stage (post-pubertal (reference), late pubertal, mid-pubertal, early pubertal and pre-pubertal) [31]. As chronological age is a less sensitive measure of development-related weight change [36], we used pubertal stage, which we derived from sex-specific questions related to pubertal development following an established method [37].

### 2.3. Statistical analysis

Descriptive statistics summarised baseline sociodemographic characteristics and obesity across levels of stressful life events. Chi-square test was used to assess the significance of gender differences in the proportion reporting different levels of stressful life events. Our main analysis had the *a priori* strategy to examine the gender-specific relationship between stressful life events and obesity; thus, we performed a series of logistic regression models including an interaction term between the exposure and the variable for self-reported male/female and sequentially adjusted for pubertal stage, ethnicity and maternal education. The final sample analyzed was 905 and appeared similar to the full cohort in terms of baseline measures of sociodemographic characteristics and health outcomes (Supplementary Table 1).

Sensitivity analyses tested the robustness of the main findings to the use of alternative socioeconomic indicator, as family income as perceived relative to peers may be more reliably reported than mother's education level. Thus, models were separately adjusted for being far above average/slightly average (reference), slightly above average, average, slightly below average and slightly below average/far below average ( $n = 943$ ) [31]. Separate models also tested for additional confounding by depressive symptomatology (CESD score  $\geq 24$ ) [38] ( $n = 796$ ). All statistical analyses were conducted using Stata 14.1 (StataCorp. 2013). We report our results using odds ratios (OR) and 95% confidence intervals (95% CI).

### 3. Results

Our sample of adolescents (13–17 years) had an average age of 14.8 years (SD 0.7), with 58% women and 51% white ethnicity. Most (65%) rated their health as very good/excellent, many reported having a university-educated mother (41%) and a relatively above-

average family income (44%) and few (7%) were classified as having obesity at follow-up. Notably, a higher proportion of young men (11%) than young women (5%) had obesity at follow-up. Nearly a third ( $n = 297$ ) of young people reported experiencing one or more stressful life events: 20% reported one recent event and 11% reported multiple recent events. A higher proportion of young women reported one and multiple stressful life events, which showed significant difference from the proportions of young men reporting exposure to stressful life events ( $P = .005$ ). The most common stressful life events for young people in BC were moving to a new home and splitting up from a boyfriend or girlfriend (Table 1).

Table 2 shows the distribution of social and health characteristics across levels of stressful life events. Exposure to multiple stressful life events was more prevalent among females than males, post-pubertal adolescents, those with depression, lower self-rated health and lower socio-economic status. There was a higher prevalence of obesity at follow-up among participants reporting multiple stressful life events (11%) than those not exposed (6%) (see Table 2).

#### 3.1. Gender differences in the relationship between stressful life events and obesity

Multivariable-adjusted regression models showed a monotonic association between greater frequency of stressful life events and obesity at study follow-up in young men only (Table 3, Model D). Compared to young men reporting no stressful life events, those reporting one event were 47% more likely to have obesity (OR 1.47 [95% CI: 0.63–3.41]) and those reporting multiple events two times more likely to have obesity (2.07 [0.79–5.43]). Only young women reporting multiple stressful life events showed higher odds of subsequent obesity (1.32 [0.41–4.18]) than those reporting none. Findings were attenuated for young men and women reporting multiple events when maternal education was replaced with household income (1.69 [0.28–4.35] and 1.07 [0.92–3.90], respectively) but were amplified when additionally adjusting for depression (2.54 [0.90–7.18] and 1.55 [0.44–5.40], respectively).

### 4. Discussion

This population-based study of young people aged 13–17 years found a nonsignificant association between two or more stressful life events and obesity status a year later, independent of known confounders. The pattern of association appeared monotonic for young men only. However, findings need to be replicated in a larger sample with more prevalent and clinically measured obesity to provide stronger empirical support for our initial hypotheses that exposure to greater amount of stressful life events would be linked to higher obesity levels in a gender-specific manner.

**Table 1**  
Frequency of self-reported stressful life events among young people in the BASUS cohort.

|                                    | Total, n | Males, n | Females, n |
|------------------------------------|----------|----------|------------|
| None                               | 1018     | 440      | 578        |
| Move to new house                  | 201      | 79       | 122        |
| Split up with girlfriend/boyfriend | 184      | 65       | 119        |
| Parents divorced                   | 79       | 27       | 52         |
| First time having intercourse      | 72       | 24       | 48         |
| Involved with police               | 65       | 31       | 34         |
| Victim of violence                 | 61       | 25       | 36         |
| Suicide of friend                  | 38       | 11       | 27         |
| Lost job                           | 31       | 12       | 19         |
| Suicide of family                  | 17       | 6        | 11         |
| Found out you have an STD          | 15       | 6        | 9          |
| Expelled from school               | 15       | 7        | 8          |
| Death of a parent                  | 14       | 7        | 7          |
| First told you are adopted         | 7        | 5        | 2          |

**Table 2**  
Sociodemographic and health characteristics of young people in the BASUS cohort across levels of stressful life events.

|                              | Mean (SD)<br>Age | Female | Highest maternal<br>education <sup>a</sup> | High family<br>income <sup>b</sup> | White<br>ethnicity | Depressed <sup>c</sup> | Excellent/very good<br>health | Post-pubertal<br>stage | Obesity<br>status <sup>d</sup> |
|------------------------------|------------------|--------|--------------------------------------------|------------------------------------|--------------------|------------------------|-------------------------------|------------------------|--------------------------------|
| Stressful Life Events        |                  |        |                                            |                                    |                    |                        |                               |                        |                                |
| None ( $n = 609$ )           | 14.7 (0.7)       | 57%    | 45%                                        | 20%                                | 48%                | 12%                    | 70%                           | 13%                    | 6%                             |
| One ( $n = 196$ )            | 14.8 (0.6)       | 65%    | 48%                                        | 14%                                | 61%                | 15%                    | 66%                           | 18%                    | 8%                             |
| Two or more<br>( $n = 100$ ) | 15.0 (0.7)       | 63%    | 42%                                        | 17%                                | 55%                | 47%                    | 59%                           | 21%                    | 11%                            |

BMI, body mass index ( $\text{kg}/\text{m}^2$ ). Measurement time points were age, gender, ethnicity, maternal education, family income, depression and general health (Fall 2011); height and weight (2012 spring). <sup>a</sup> University or higher. <sup>b</sup> far above average or slightly above average relative to peers. <sup>c</sup> Depression score  $\geq 24$ , assessed using Centre for Epidemiologic Studies Depression Scale. <sup>d</sup> BMI  $>97$ th percentile based on age- and gender-specific WHO growth charts.

**Table 3**

Odds ratios of obesity at follow-up associated with recent stressful life events in young women and young men in the BASUS cohort study.

|                       | Young Women |           |         |           |         |           |         |           |
|-----------------------|-------------|-----------|---------|-----------|---------|-----------|---------|-----------|
|                       | Model A     |           | Model B |           | Model C |           | Model D |           |
|                       | OR          | 95% CI    | OR      | 95% CI    | OR      | 95% CI    | OR      | 95% CI    |
| Stressful life events |             |           |         |           |         |           |         |           |
| None                  | 1.00        |           | 1.00    |           | 1.00    |           | 1.00    |           |
| One                   | 1.27        | .54, 2.99 | .80     | .29, 2.23 | .79     | .28, 2.21 | .88     | .31, 2.48 |
| Two or more           | 2.16        | .82, 5.68 | 2.13    | .80, 5.70 | 1.62    | .56, 4.69 | 1.32    | .41, 4.18 |
| Young Men             |             |           |         |           |         |           |         |           |
| Stressful life events |             |           |         |           |         |           |         |           |
| None                  | 1.00        |           | 1.00    |           | 1.00    |           | 1.00    |           |
| One                   | 1.44        | .67, 3.09 | 1.31    | .59, 2.92 | 1.31    | .58, 2.95 | 1.47    | .63, 3.41 |
| Two or more           | 1.80        | .74, 4.38 | 1.95    | .78, 4.89 | 1.79    | .70, 4.56 | 2.07    | .79, 5.43 |

Gender-specific odds ratios (95% CI) obtained by multivariable linear regression analysis using an interaction term between sex and stressful life events (Model A,  $n = 1117$ ) and adjusting for pubertal stage (Model B,  $n = 908$ ), ethnicity (Model C,  $n = 975$ ), and mother's education ( $n = 905$ ).

#### 4.1. Relevance to previous work

Overall, review evidence of longitudinal studies indicates that stressful life events are positively associated with obesity in young and adult populations [39,40]. Our results pointed to similar results, although not statistically significant. We also found potential gender differences, as higher odds of obesity were seen only in young women reporting two or more events but was monotonically patterned in young men. In adults, the prospective association seems to differ by gender whereby men have a higher obesity risk from stressful life events than women [40]. By contrast, cross-sectional evidence from two large epidemiological studies with U.S. adults reported female-specific effects of stressful life events on BMI [41,42]. Notably, Barry and Petry (2008) observed a strong positive association only in women when they examined exposure over the previous year to at least 12 stressful life events [42], which is a higher threshold than this study used. Overall estimates in children indicate that greater exposure to stressful life events are associated with a 12% greater likelihood of overweight status (OR 1.12) [43], and this association shows some gender differences in young children [44]. It is difficult to directly compare our results with those of previous works, as the literature uses heterogeneous nomenclature and operationalization of both stressful life events and obesity status, lacks a focus on this vulnerable age group of mid-to late-puberty and is silent on potential gender differences.

Several potential mechanisms could explain our finding of a relationship between increasing stressful life events and greater odds of having obesity, including physiological and behavioural responses. Chronic stress is known to increase overall activation of the HPA axis which can negatively affect hormonal regulation of metabolic activities. In particular, elevated cortisol levels in the body are associated with increased leptin levels and central adiposity [13]. Moreover, this disruption to leptin levels can lead to an inhibited sense of satiety, which, in turn, increases food consumption [13]. Concomitant with these physiological responses, stressful life events can lead to unhealthy behavioural responses. Poor coping mechanisms for stressful life events can further contribute to obesity risk through psychosocial effects that increase sedentary behaviour [45] and higher energy intake [46] and reduce physical activity [47].

The observed monotonic relationship between stressful life events and obesity in the short term among adolescent males could be explained by a combination of metabolic dysregulation and unhealthy behaviours that have known sex- and gender-based differences. Studies show that young males have a higher tendency for cortisol and autonomic nervous system reactivity than young females when exposed to stress; these works also find that

the sex-based difference in HPA response patterns in mid-adolescence is similar to those found in adulthood [17]. Notably, female sex hormone cycles exert modulating effects on HPA responsiveness to perceived stress [48], with the post-ovulation luteal phase dampening the magnitude of other circadian rhythms such as cortisol [49]. In addition, there may be gender-based differences that explain the null association in young women. It is possible that a higher number of stressful life events are required to show negative influences on their weight given that women typically experience a higher burden of certain high-magnitude events and their negative consequences. In Canada, for example, the likelihood that women over the age of 15 years will be sexually victimized 11 times greater than that for men or stalked 3 times higher than men [50]. Alternatively, young women's social roles and norms mean they tend to have a bigger and wider social network that is a potential support system for coping with stressful life events and may mitigate the negative metabolic consequences of recent stressful life events. Other gender-specific norms about overeating behaviour and thin body ideals could also explain why stressful life events were not associated with obesity in young women.

#### 4.2. Methodological considerations

Recall or social desirability bias could affect our self-reported exposures and outcomes [31,51]. Nevertheless, experiences of specific life events are important exposures to investigate, which provide concrete targets for intervention compared to global perceptions of psychological stress that are typically studied [39,40,43]. While social desirability bias in self-reported height and weight underestimates true obesity prevalence in a population, questions about the contextual influences of obesity risk can be answered by analyzing participants' relative rank in weight status based on self-reported BMI. Results may be biased by measurement error in the limited range of pre-specified response options and categories constructed for the exposure variables. This study may also be limited by bias from residual confounding from unmeasured factors such as young women's parity status, which could affect the outcome, but was not collected in the BASUS study.

Missing data predominantly on covariables may have introduced selection bias in our study, although our analyzed sample appeared to be representative of the full cohort apart from a slightly higher proportion of young men from high-income families and young women with university-educated mothers. Nevertheless, loss of sample size likely limited the power to detect a statistically significant result. Finally, our measure of stressful life events did not include assessment of event severity, which varied across the different types listed and may lead to differential gender-specific

effects on obesity risk.

Notwithstanding these limitations, a key strength of our study is the population-based cohort that is representative of the broader adolescent population in Canada's west coast [52], and inclusion of ethnicity, pubertal stage and socioeconomic status as known confounders. Importantly, this work advances the evidence base in several ways. First, we examined independent stressful life events rather than a generic measure of perceived psychological stress, and thereby provide novel evidence on distinct risk factors that serve as direct targets for future interventions. Second, we focused on young people aged 13–17 years who have a unique developmental period putting them at particularly high risk of negative metabolic consequences from major life events. Third, we contributed a critical gender perspective on the relationship and demonstrated a differential patterning of the relationship between stressful life events and obesity in this age group.

## 5. Conclusion

Healthy young people in BC reporting multiple stressful life events were more likely to have obesity in the short term, more so for young men than young women. Findings add to previous work supporting the need to consider addressing young people's experiences of stressful life events in future obesity prevention strategies. Additional gender-sensitive health research should investigate the behavioural and biological mechanisms linking stressful life events with obesity risk.

## Funding

This work received partial funding support from the University of British Columbia Work Learn Program summer 2017 (Projects 170278 & 170279). The BASUS cohort study was generously supported by the Canadian Institutes of Health Research (Grant #86729).

## Disclosure

We declare no conflicts of interest.

## Ethical statement

All volunteers gave written informed consent following school district procedures, and the study was approved by Behavioural Research Ethics Board of the University of British Columbia (No. H08–02841).

## Acknowledgments

The authors wish to thank the BASUS cohort participants for their contribution. Data from the BASUS Study cannot be deposited publicly due to ethics and consent requirements of the sensitive information from this vulnerable age group. Data requests can be made directly to the BASUS study team ([www.basus.ca](http://www.basus.ca)).

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijpam.2019.03.001>.

## References

- [1] Abarca-Gómez L, Abdeen ZA, Hamid ZA, Abu-Rmeileh NM, Acosta-Cazares B, Acuin C, et al. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet* 2017;390(10113):2627–42.
- [2] Rao DP, Kropac E, Do MT, Roberts KC, Jayaraman GC. Childhood overweight and obesity trends in Canada. *Health Promot Chronic Dis Prev Can Res Policy Pract* 2016;36(9):194.
- [3] Segula D. Complications of obesity in adults: a short review of the literature. *Malawi Med J* 2014 Mar;26(1):20–4.
- [4] Reilly JJ, Kelly J. Long-term impact of overweight and obesity in childhood and adolescence on morbidity and premature mortality in adulthood: systematic review. *Int J Obes* 2011;35(7):891–8.
- [5] Simmonds M, Llewellyn A, Owen CG, Woolacott N. Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev* 2016;17(2):95–107.
- [6] Spear LP. The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev* 2000 Jun;24(4):417–63.
- [7] Pervanidou P, Chrousos GP. Stress and obesity/metabolic syndrome in childhood and adolescence. *Int J Pediatr Obes IJPO Off J Int Assoc Study Obes* 2011 Sep;6(Suppl 1):21–8.
- [8] Waters E, de Silva-Sanigorski A, Hall BJ, Brown T, Campbell KJ, Gao Y, et al. Interventions for preventing obesity in children. *Cochrane Database Syst Rev* 2011 Dec;(12):CD001871.
- [9] Conklin AI, Guo SX, Tam AC, Richardson CG. Gender, stressful life events and interactions with sleep: a systematic review of determinants of adiposity in young people. *BMJ Open* 2018 Jul;8(7): e019982.
- [10] Harkness KL, Alavi N, Monroe SM, Slavich GM, Gotlib IH, Bagby RM. Gender differences in life events prior to onset of major depressive disorder: the moderating effect of age. *J Abnorm Psychol* 2010;119(4):791.
- [11] Costello EJ, Erkanli A, Fairbank JA, Angold A. The prevalence of potentially traumatic events in childhood and adolescence. *J Trauma Stress* 2002;15(2): 99–112.
- [12] Gunnar MR, Talge NM, Herrera A. Stressor paradigms in developmental studies: what does and does not work to produce mean increases in salivary cortisol. *Psychoneuroendocrinology* 2009 Aug;34(7):953–67.
- [13] Pervanidou P, Chrousos GP. Metabolic consequences of stress during childhood and adolescence. *Metabolism* 2012 May;61(5):611–9.
- [14] Casey B, Jones RM, Hare TA. The adolescent brain. *Ann N Y Acad Sci* 2008;1124(1):111–26.
- [15] Pervanidou P, Chrousos GP. Metabolic consequences of stress during childhood and adolescence. *Metabolism* 2012;61(5):611–9.
- [16] Volkow ND, Wang G-J, Fowler JS, Telang F. Overlapping neuronal circuits in addiction and obesity: evidence of systems pathology. *Philos Trans R Soc B Biol Sci* 2008;363(1507):3191–200.
- [17] Ordaz S, Luna B. Sex differences in physiological reactivity to acute psychosocial stress in adolescence. *Psychoneuroendocrinology* 2012 Aug;37(8): 1135–57.
- [18] Laws M. Examining the effects of psychosocial stress on the hypertension self-management behaviors of African American women. Richmond, Virginia: Virginia Commonwealth University; 2016. 137 pp.
- [19] Hirotsu C, Tufik S, Andersen ML. Interactions between sleep, stress, and metabolism: from physiological to pathological conditions. *Sleep Sci* 2015 Nov;8(3):143–52.
- [20] Al Khatib HK, Harding SV, Pot GK. The effects of partial sleep deprivation on energy balance: a systematic review and meta-analysis. *Eur J Clin Nutr* 2017;71:614–24.
- [21] Chaput J-P, Gray CE, Poitras VJ, Carson V, Gruber R, Olds T, et al. Systematic review of the relationships between sleep duration and health indicators in school-aged children and youth. *Appl Physiol Nutr Metabol* 2016 Jun;41: S266–82. 6 (Suppl. 3).
- [22] Pejovic S, Vgontzas AN, Basta M, Tsaoussoglou M, Zoumakis E, Vgontzas A, et al. Leptin and hunger levels in young healthy adults after one night of sleep loss. *J Sleep Res* 2010 Dec 1;19(4):552–8.
- [23] Bull NL. Dietary habits, food consumption, and nutrient intake during adolescence. *J Adolesc Health* 1992 Jul;13(5):384–8.
- [24] Quinn JJ, Hitchcott PK, Umeda EA, Arnold AP, Taylor JR. Sex chromosome complement regulates habit formation. *Nat Neurosci* 2007 Nov;10(11): 1398–400.
- [25] Vignau J, Bailly D, Duhamel A, Vervaecke P, Beuscart R, Collinet C. Epidemiologic study of sleep quality and troubles in French secondary school adolescents. *J Adolesc Health* 1997;21(5):343–50.
- [26] Eisenmann J, Ekkakakis P, Holmes M. Sleep duration and overweight among Australian children and adolescents. *Acta Paediatr* 2006 Aug 1;95(8):956–63.
- [27] Cohen S, Murphy MLM, Prather AA. Ten surprising facts about stressful life events and disease risk [cited 2018 Dec 7]. *Annu Rev Psychol* 2019 Jan 4;70(1). Available from: <https://www.annualreviews.org/doi/10.1146/annurev-psych-010418-102857>.
- [28] Wardle J, Chida Y, Gibson EL, Whitaker KL, Steptoe A. Stress and adiposity: a meta-analysis of longitudinal studies. *Obesity* 2011;19(4):771–8.
- [29] Elsenburg LK, van Wijk KJE, Liefbroer AC, Smidt N. Accumulation of adverse childhood events and overweight in children: a systematic review and meta-analysis. *Obesity* 2017 May;25(5):820–32.
- [30] Conklin AI, Guo SX, Tam AC, Richardson CG. Gender, stressful life events and interactions with sleep: a systematic review of determinants of adiposity in young people. *BMJ Open* 2018;8:e019982. <https://doi.org/10.1136/bmjopen-2017-019982>.
- [31] Conklin AI, Yao CA, Richardson CG. Chronic sleep deprivation and gender-

- specific risk of depression in adolescents: a prospective population-based study [cited 2019 Jan 12] BMC Public Health [Internet]. 2018 Dec;18(1). Available from: <https://bmcpublihealth.biomedcentral.com/articles/10.1186/s12889-018-5656-6>.
- [32] Richardson CG, Kwon J-Y, Ratner PA. Self-esteem and the initiation of substance use among adolescents. *Can J Public Health Rev Can Sante Publique* 2012 Nov 8;104(1):e60–3.
  - [33] Swearingen EM, Cohen LH. Measurement of adolescents' life events: the junior high life experiences survey. *Am J Community Psychol* 1985 Feb;13(1): 69–85.
  - [34] Meadows SO, Brown JS, Elder GH. Depressive symptoms, stress, and support: gendered trajectories from adolescence to young adulthood. *J Youth Adolesc* 2006 Feb;35(1):89–99.
  - [35] Dietitians of Canada. WHO Growth Charts for Canada: body mass index-for-age percentiles. 2014.
  - [36] Lee JM, Appugliese D, Kaciroti N, Corwyn RF, Bradley RH, Lumeng JC. Weight status in young girls and the onset of puberty. *Pediatrics* 2007 Mar 1;119(3): e624–30.
  - [37] Carskadon MA, Acebo C. A self-administered rating scale for pubertal development. *J Adolesc Health* 1993;14(3):190–5.
  - [38] Radloff LS. The use of the center for epidemiologic studies depression scale in adolescents and young adults. *J Youth Adolesc* 1991;20(2):149–66.
  - [39] Gundersen C, Mahatmya D, Garasky S, Lohman B. Linking psychosocial stressors and childhood obesity: stress and obesity. *Obes Rev* 2011 May;12(5): e54–63.
  - [40] Wardle J, Chida Y, Gibson EL, Whitaker KL, Steptoe A. Stress and adiposity: a meta-analysis of longitudinal studies. *Obesity* 2011;19(4):771–8.
  - [41] Udo T, Grilo CM, McKee SA. Gender differences in the impact of stressful life events on changes in body mass index. *Prev Med* 2014 Dec;69:49–53.
  - [42] Barry D, Petry N. Gender differences in associations between stressful life events and body mass index. *Prev Med* 2008 Nov;47(5):498–503.
  - [43] Elsenburg LK, van Wijk KJE, Liefbroer AC, Smidt N. Accumulation of adverse childhood events and overweight in children: a systematic review and meta-analysis: adverse Events and Childhood Overweight. *Obesity* 2017 May;25(5): 820–32.
  - [44] Suglia SF, Duarte CS, Chambers EC, Boynton-Jarrett R. Cumulative social risk and obesity in early childhood. *Pediatrics* 2012 May 1;129(5):e1173–9.
  - [45] Hoare E, Milton K, Foster C, Allender S. The associations between sedentary behaviour and mental health among adolescents: a systematic review. *Int J Behav Nutr Phys Act* 2016 Dec;13(1).
  - [46] Goossens L, Braet C, Van Vlierberghe L, Mels S. Loss of control over eating in overweight youngsters: the role of anxiety, depression and emotional eating. *Eur Eat Disord Rev* 2009 Jan;17(1):68–78.
  - [47] Stults-Kolehmainen MA, Sinha R. The effects of stress on physical activity and exercise. *Sports Med* 2014 Jan;44(1):81–121.
  - [48] Kirschbaum C, Kudielka BM, Gaab J, Schommer NC, Hellhammer DH. Impact of gender, menstrual cycle phase, and oral contraceptives on the activity of the hypothalamus-pituitary-adrenal axis. *Psychosom Med* 1999;61(2):154–62.
  - [49] Baker FC, Driver HS. Circadian rhythms, sleep, and the menstrual cycle. *Sleep Med* 2007 Sep;8(6):613–22.
  - [50] Sinha M. Measuring violence against women: statistical trends. *Juristat Can Cent Justice Statistics* 2013;1.
  - [51] Conklin AI. Gender, diet quality and obesity: economic and social determinants, and their interactions. older adults; 2014.
  - [52] Smith A, Stewart D, Poon C, Peled M, Saewyc E. McCreary Centre society. From hastings street to Haida Gwaii: provincial results of the 2013 BC adolescent health survey. Vancouver, BC: McCreary Centre Society; 2014. p. 1–88.