



Sleep duration and risk of breast cancer: The JACC Study

Jinhong Cao¹ · Ehab S. Eshak^{1,2} · Keyang Liu¹ · Isao Muraki¹ · Renzhe Cui¹ · Hiroyasu Iso¹ · Akiko Tamakoshi³ · JACC Study Group

Received: 1 August 2018 / Accepted: 4 October 2018 / Published online: 20 November 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract

Purpose The evidence on beneficial or adverse effects of sleep duration on risk of breast cancer remains controversial and limited, especially in Asia.

Methods A prospective study of 34,350 women aged 40–79 years in whom sleep duration, and menstrual and reproductive histories were determined by a self-administered questionnaire. The follow-up period was from 1988 to 2009, and hazard ratios (HRs) with 95% confidence intervals (CIs) of breast cancer incidence were calculated for shorter sleep duration in reference to sleep duration of ≥ 8 h/day by Cox proportional hazard models.

Results During 19.2-year median follow-up (236 cases), we found a significant inverse association between sleep duration and risk of breast cancer, especially among postmenopausal women and women with low parity (nulliparous and women with < 3 children); the multivariable HRs (95% CIs) among postmenopausal women who reported 7 h/day and ≤ 6 h/day of sleep in reference to ≥ 8 h/day were 1.49 (0.81–2.76) and 1.98 (1.08–3.70) (P for trend = 0.028), respectively, and the corresponding values among women with low parity were 1.50 (0.96–2.35) and 1.76 (1.01–2.79) (P for trend = 0.018).

Conclusions Short sleep duration was associated with increased risk of incident breast cancer, especially among postmenopausal women and women with low parity.

Keywords Sleep duration · Breast cancer · Incidence · Cohort study · Postmenopausal · Parity · Japan

Introduction

Breast cancer is the most common cancer among Japanese women followed by colon and rectum cancer [1]. The association between sleep duration and risk of breast cancer among women has a complex nature and remains controversial [2]. Previous case-control and cohort studies have reported no association between sleep duration and risk of breast cancer

[2–7]. However, several other studies showed lower risk of breast cancer with longer sleep duration (≥ 9 h/day) [8–10], while some other studies indicated that longer sleep duration (≥ 9 h/day) was associated with increased risk of breast cancer [11, 12].

Melatonin is suggested as an agent in the association between sleep duration and breast cancer [13, 14]. Melatonin (5-methoxytryptamine) is synthesized and secreted by the pineal gland in the brain and controls the body's circadian rhythm [13]. Darkness during sleep stimulates the release of melatonin [7, 9, 15], and melatonin may inhibit breast tumorigenesis directly by inhibiting mammary cell proliferation and invasiveness, and indirectly by decreasing estrogen levels via a down-regulation of the hypothalamic-pituitary reproductive axis and regulating the activity of the aromatases, the enzymes responsible for the local estrogens synthesis [16–19].

Some studies showed menopausal status to be associated with risk of breast cancer because of aging [20], higher levels of adiposity [21, 22], or endogenous estrogen [23]. Furthermore, short sleep duration, especially among

The JACC study group investigators are listed in Acknowledgements section.

✉ Hiroyasu Iso
iso@pbhel.med.osaka-u.ac.jp

¹ Public Health, Department of Social Medicine, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita-shi, 565-0871 Osaka, Japan

² Department of Public Health, Community and Preventive Medicine, Faculty of Medicine, Minia University, Minia, Egypt

³ Department of Public Health, Faculty of Medicine, Hokkaido University, Sapporo, Japan

postmenopausal women, was associated with high risk of breast cancer in several cohort studies [8–10, 24]. Abundant previous findings have also shown parity as an indicator for breast cancer risk [25–27]. Meanwhile, multiparous women have reported longer sleep duration than nulliparous women [28].

Thus, we thought to examine the associations between sleep duration and risk of breast cancer among premenopausal and postmenopausal women, and among low and high parous women in a large population-based Japanese study, the Japan Collaborative Cohort study (JACC).

Materials and methods

Study population and ascertainment of breast cancer

Details of the study design and subjects have been described elsewhere [29]. Briefly, the baseline data of the JACC Study were collected from 1988 to 1990, and 110,585 individuals (46,395 men and 64,190 women) aged 40 to 79 years in 45 study areas throughout Japan participated in the study. The follow-up survey for cancer incidence was conducted from the baseline, and finalized at the end of the 2009. In 24 areas out of the 45 study areas, data on cancer incidence such as date of diagnosis and primary site were collected simultaneously through population-based cancer registries or by reviewing the records of local and major hospitals. After excluding male subjects, we confined the analysis to women from these 24 areas where cancer incidence information are available ($n = 36,266$). Excluding data of women with previous diagnosis of breast cancer ($n = 11$), and women with missing data on sleep duration ($n = 1905$) left a total of 34,350 (19,529 premenopausal, and 14,821 postmenopausal) women for the analysis. This study was sponsored by the Ministry of Education, Sports and Science. Informed consent was obtained from participants to participate in the JACC study for the baseline questionnaire. The ethics committees of Nagoya University School of Medicine and Osaka University approved the protocol of this study.

Exposure and other covariates assessment

Participants completed a self-administered questionnaire including sleep duration, information on age, family history of diseases, history of hypertension, diabetes mellitus, cardiovascular diseases, cancer, height, weight, education background, smoking status, alcohol drinking habit, physical activity, mental stress, dietary habits, reproductive and menstrual history, menopause, and hormone use. Body mass index was calculated by dividing reported weight in kilograms by the square of reported height in meters.

Assessment of sleep duration

We obtained information about the average sleep duration on weekdays during the preceding year. The average sleep duration per day was classified into three categories: ≤ 6 , 7, and ≥ 8 h. Fractions hours were rounded off (e.g., 7 h represented responses from 7.0 to 7.9 h).

Statistical analysis

Mean values (standard deviations) and proportions of baseline risk characteristics were calculated, and the linear trends in those variables according to sleep duration were tested by the linear regression analysis for continuous variables and the logistic regression analysis for proportional variables. Person-years of follow-up were calculated from the responding date to the baseline questionnaire until the attainment of one out of four possible endpoints as follows: (1) incidence of breast cancer event, (2) relocation from the study area, (3) the end of the study on December 31, 2009, or (4) death, because some study areas discontinued the follow-up survey regarding cancer before 2009 (1994 in one study area, 1997 in two areas, 1999 in one area, 2000 in one area, 2002 in one area, 2003 in one area, 2006 in two areas, and 2008 in two areas).

Cox proportional hazard regression age- and multi-variable-adjusted models were used to estimate the hazard ratios (HRs) with 95% confidence intervals (CIs) for breast cancer incidence according to sleep duration (≤ 6 h, 7 h, and ≥ 8 h/day) as the reference to ≥ 8 h/day, and in relation to 1-SD decrement (1.07 h/d) of sleep duration. The confounding factors included age (continuous), age of menarche (< 14 , 14–15 and > 15 years), age of menopause (< 45 , 45–50 and > 50 years), age at first child birth (< 25 and ≥ 25 y), type of menopause (nature or operation), body mass index (continuous), sport time per week (never, < 1 , 1–2, 3–4 and ≥ 5 h/week), walking time per day (never, < 30 , 30–60 and ≥ 60 min/day), currently married (yes or no), smoking status (never, ex-smoker and current smoker), alcohol intake (never, ex-drinker and current drinker of 0.1–22.9, 23.0–45.9, and ≥ 46.0 g ethanol/day), parity (0, 1, 2 and ≥ 3), use of sex hormone (yes or no), family history of breast cancer (yes or no), and history of diabetes (yes or no). The stratification analyses were performed by potential effect modifiers such as menopausal status and number of children. Values for *P*-interaction were calculated for cross-product terms of menopausal status (dichotomous) or number of children (continuous) with sleep duration categories (1 to 3 corresponding to ≤ 6 , 7, and ≥ 8 h/day of sleep duration) for the categorical analysis and sleep duration (h/d) for the continuous

analysis. We used SAS Version 9.4 software (SAS Institute Inc, Cary, NC) for statistical analysis. All statistical tests were 2-tailed and values of $P < 0.05$ were regarded as significant.

Results

Table 1 shows the baseline characteristics of women according to sleep duration. Women who reported ≤ 6 h/day of sleep were more likely to have a family history of breast cancer and to have used sex hormones. On the other hand, women who reported ≥ 8 h/day of sleep were of older age

at menarche, with higher BMI and were more likely to have natural menopause and ≥ 3 children.

With reference to women with sleep duration ≥ 8 h/day, there was higher risk of breast cancer among women with shorter sleep duration in total women (Table 2). The multivariable HRs (95% CIs) of breast cancer were 1.36 (0.98–1.90) for 7 h and 1.31 (0.92–1.86) for ≤ 6 h sleep per day.

Table 3 shows the stratification analyses by menopausal status and parity (number of children). The inverse associations between sleep duration and risk of breast cancer were confined to postmenopausal women, the multivariable HRs (95% CIs) of breast cancer were 1.49 (0.81–2.76) for 7 h and 1.98

Table 1 Distributions of potential risk factors according to sleep duration in a cohort of 34,350 Japanese women

Parameters	Sleep duration			<i>P</i> for trend
	≤ 6 h	7 h	≥ 8 h	
No. at risk	10,064	12,979	11,307	
Age, years (SD)	60.7 (7.7)	60.2 (7.2)	62.6 (7.6)	<0.001
Age at menarche, years (SD)	15.0 (1.8)	15.0 (1.7)	15.3 (1.8)	<0.001
Age at menopause, years (SD)	48.6 (4.6)	48.6 (4.7)	48.7 (4.5)	0.318
Natural menopause, %	86.6	87.2	88.1	0.005
Age at first child birth, years (SD)	24.7 (3.3)	24.5 (3.1)	24.2 (3.0)	<0.001
Parity (number of children)				
0	3.0	3.4	3.2	<0.001
1	8.0	7.9	6.7	
2	34.5	35.8	28.7	
≥ 3	54.5	52.9	61.3	
Family history of breast cancer, %	1.5	1.4	1.2	0.023
Body mass index, kg/m ² (SD)	22.9 (3.2)	22.9 (3.1)	23.1 (3.3)	<0.001
Currently married, %	78.1	81.9	77.6	0.065
Sports ≥ 3 h/week, %	10.9	10.9	11.5	0.500
Walking time ≥ 60 min/day, %	73.1	74.1	74.6	0.013
Current smoking, %	2.6	2.1	2.7	0.250
Alcohol intake, g ethanol/day (SD)	8.6 (10.6)	8.4 (10.6)	9.1 (12.7)	0.838
Sex hormone use, %	5.1	4.6	3.7	<0.001
History of diabetes, %	5.7	4.1	6.1	0.403

Table 2 Age-adjusted and multivariable hazard ratios (95% confidence intervals) of incident breast cancer according to sleep duration for total women

	Sleep duration			<i>P</i> for trend	^a 1SD decrement (1.07 h/day) of sleep duration
	≤ 6 h	7 h	≥ 8 h		
Total women					
Person-year	165,359	221,198	194,205		
Breast cancer, <i>n</i>	74	101	61		
Age-adjusted HR (95% CI)	1.31 (0.93–1.85)	1.35 (1.98–1.86)	1.00	0.133	1.11 (0.97–1.27)
Multivariable HR (95%CI) ^b	1.31 (0.92–1.86)	1.36 (0.98–1.90)	1.00	0.149	1.13 (0.98–1.29)

^a1SD decrement in sleep duration = 1.07 h/day

^bAdjusted for age, age at menarche, age at first child birth, body mass index, parity (number of children), family history of breast cancer, marital status, sport time, walking time, alcohol intake, smoking status, hormone use, history of diabetes, and age and type of menopause

Table 3 Multivariable hazard ratios (95% confidence intervals) of incident breast cancer according to sleep duration, stratified by menopausal status, and number of children

	Sleep duration			<i>P</i> for trend	^a 1SD decrement (1.07 h/day) of sleep duration
	≤ 6 h	7 h	≥ 8 h		
Menopausal status					
Premenopause					
Person-year	102,201	138,570	114,858		
Breast cancer, <i>n</i>	46	74	43		
Age-adjusted HR (95% CI)	1.10 (0.72–1.69)	1.32 (0.90–1.94)	1.00	0.689	1.05 (0.89–1.24)
Multivariable HR (95% CI) ^b	1.11 (0.72–1.71)	1.34 (0.90–1.98)	1.00	0.687	1.07 (0.90–1.27)
Postmenopause					
Person-year	63,158	82,628	79,347		
Breast cancer, <i>n</i>	28	27	18		
Age-adjusted HR (95% CI)	1.92 (1.06–3.49)	1.43 (0.78–2.60)	1.00	0.031	1.26 (1.00–1.58)
Multivariable HR (95% CI) ^c	1.98 (1.08–3.70)	1.49 (0.81–2.76)	1.00	0.028	1.28 (1.01–1.61)
<i>P</i> _{interaction}				0.264	0.400
Number of children					
0–2					
Person-year	89,345	122,823	94,275		
Breast cancer, <i>n</i>	52	60	32		
Age-adjusted HR (95% CI)	1.66 (1.06–2.59)	1.40 (0.91–2.16)	1.00	0.027	1.21 (1.01–1.44)
Multivariable HR (95% CI) ^d	1.76 (1.01–2.79)	1.50 (0.96–2.35)	1.00	0.018	1.26 (1.05–1.51)
≥ 3					
Person-year	76,014	98,375	99,929		
Breast cancer, <i>n</i>	22	41	29		
Age-adjusted HR (95% CI)	0.88 (0.50–1.55)	1.27 (0.78–2.08)	1.00	0.692	0.98 (0.80–1.21)
Multivariable HR (95% CI) ^d	0.83 (0.46–1.48)	1.24 (0.75–2.04)	1.00	0.562	0.96 (0.77–1.20)
<i>P</i> _{interaction}				0.002	0.014

^a1SD decrement in sleep duration = 1.07 h/day

^bAdjusted for age, age at menarche, age at first child birth, body mass index, parity (number of children), family history of breast cancer, marital status, sport time, walking time, alcohol intake, smoking status, hormone use, history of diabetes

^cAdjusted further for age and type of menopause

^dAdjusted for age, age at menarche, age at first child birth, body mass index, family history of breast cancer, marital status, sport time, walking time, alcohol intake, smoking status, hormone use, history of diabetes, and age and type of menopause

(1.08–3.70) for ≤ 6 h sleep per day (*P* for trend = 0.028); however, the interaction by menopausal status was not statistically significant (*P*_{interaction} = 0.264). The inverse association was also evident among low parous women including nulliparous and women with < 3 children. The multivariable HRs (95% CIs) for breast cancer risk among low parous women who have reported 7 h and ≤ 6 h sleep per day compared with those reported ≥ 8 h/day were 1.50 (0.96–2.35) and 1.76 (1.01–2.79), respectively (*P* for trend = 0.018, *P*_{interaction} = 0.002).

Discussion

During 19.2-years median follow-up for 34,350 women aged ≥ 40 years, we observed that short sleep duration was associated with increased risk of incident breast cancer

among Japanese women. This positive association was more evident for postmenopausal women and women with number of children < 3, although the interaction with parity but not menopausal status was statistically significant.

The high risk of incident breast cancer with short sleep duration found in our study is consistent with findings from previous prospective cohort studies [8, 24]. Among 42,840 women of the Southern Community Cohort Study, shorter sleep was associated with increased risk of breast cancer; odds ratios (95% CIs) were 2.13 (1.15–3.93) for < 6 h/day, 1.66 (0.92–3.02) for 6 h/day, and 2.22 (1.19–4.12) for 7 h/day compared with ≥ 8 h/day (*P* for trend = 0.04) [24]. Similar results were found among 7 396 Finnish women [8]. However, those studies did not examine the association by menopausal status or parity. In Japan, Kakizaki et al. examined the association between sleep duration and

risk of incident breast cancer among 23,995 women in the Ohsaki National Health Insurance (NHI) Cohort Study, and showed that women who reported ≥ 9 h/day sleep in reference to those with ≤ 6 h/day had lower breast cancer risk: the multivariable HR (95% CI) was 0.29 (0.09–0.98, P for trend = 0.002). The reduced risk was observed among postmenopausal women; 0.74 (0.35–1.59, P for trend = 0.09) but not among premenopausal; 1.48 (0.56–3.93, P for trend = 0.27) (P for interaction = 0.70) [10]. Among 33 528 women participated in the Singapore Chinese Health Study, Wu et al. reported inverse trends in breast cancer risk across sleep duration categories among postmenopausal women, but not among total or premenopausal women. In reference to sleep duration for ≤ 6 h/d, the multivariable HRs (95% CIs) among postmenopausal women in the categories 7, 8, and ≥ 9 h/day were 0.94 (0.70–1.20), 0.81 (0.60–1.10), and 0.67 (0.40–1.10) (P for trend = 0.047) [9]. Previous studies have shown high risk of breast cancer in nulliparous or women with low parity than that in multiparous women [25]. Again, the inverse association between sleep duration and risk of breast cancer was observed among women with < 3 children including nulliparous women more than that among those with ≥ 3 children.

Shorter sleep duration was associated with lower levels of urinary melatonin; 42% lower in Chinese women reported ≤ 6 h of sleep duration than levels in women reported ≥ 9 h of sleep duration in the Singapore Chinese Health Study [9]. Because sleep stimulates the release of melatonin [9], melatonin is suggested a biological mediator for the sleep/breast cancer association. Higher levels of melatonin may associate with reduced risk of breast cancer by the following mechanisms; (a) melatonin interacts with estrogen receptors (ER) on the epithelial mammary cells, leading to direct inhibition of mammary cell proliferation and invasiveness [23]; (b) melatonin interacts with the neuroendocrine reproductive axis and the hypothalamic-pituitary reproductive axis, leading to a down-regulation of some hormones which promote tumor growth, especially gonadal estrogens and prolactin [30]; (c) melatonin inhibits telomerase enzymes activity, responsible for estrogen synthesis in tumor cells and adjacent peritumoral fat tissues [31]; (d) melatonin has antioxidant properties of melatonin which can suppress oncogenesis [32].

Postmenopausal women are at higher risk of breast cancer than premenopausal women because of higher levels of adiposity among postmenopausal women [21, 22] which serve as the primary source of endogenous estrogen transformed from androgen by enhanced aromatase expression and activity [22]. On the other hand, nulliparous and women with low parity showed lower urinary excretion of melatonin [15] and higher levels of estrogens and prolactin, but lower levels of sex hormone-binding globulins [33–35]. These factors were associated with increased risk of estrogen-receptor-positive

carcinogenic tumors. Thus, the sleep-induced melatonin secretion could, at least partially, explain the inverse association of sleep duration with risk of breast cancer in high risk group of postmenopausal women and women with low parity in our study.

The strengths of our study were its prospective design, which avoided recall bias and the availability of information on potential confounding factors. Our subjects were recruited from the general population, the sample was large and the response rate to the questionnaire was high [29, 36]. In addition, the cancer registry of the study had sufficient quality to reduce the possibility of misclassification of outcomes [37].

There are several limitations in this study. First, we did not obtain information about the quality of sleep, such as the presence or absence of sleep apnea or other sleep disorders, which were associated with increased risk of breast cancer via intermittent hypoxia and suppression of the immune system [38]. Second, we used self-reported information on sleep duration obtained only at baseline, we cannot rule out the possibility of change in sleep duration during the long follow-up, and the self-report may lead to some misclassification. However, self-reported sleep duration was shown to yield valid results in comparison with quantitative sleep assessment with actigraphy; $r = 0.57$ for nighttime sleep duration [39]. Third, we could not examine a potential adverse effect of long sleep duration, because of the low proportion of participants and the few number of cases in the long sleep duration of ≥ 9 h/day category (5.6% of participants and 6 breast cancer cases). Fourth, nulliparous women are at high risk for breast cancer [25]. However, only 6.7% of our studied women were nulliparous and there was no breast cancer case in the sleep duration category of ≥ 8 h/day. Therefore, we could not treat them as a separate category in the stratification analysis; but, we added them to women with low parity < 3 children. Finally, we did not collect the data on blood melatonin levels or urinary excretions; therefore, the exact contribution of melatonin in the observed associations between sleep duration and risk of incident breast cancer cannot be certified.

In summary, short sleep duration was associated with increased risk of incident breast cancer, especially among postmenopausal women and women with low parity. Health education to women about the need for proper sleep duration is suggested and further research is needed to confirm the observed associations.

Acknowledgements The authors thank all staff members involved in this study for their valuable help in conducting the baseline survey and follow-up. Study investigators: Members of the JACC Study Group: Dr. Akiko Tamakoshi (present chairperson of the study group), Hokkaido University Graduate School of Medicine; Dr. Mitsuru Mori, Sapporo Medical University School of Medicine; Dr. Yoshihiro Kaneko, Akita University Graduate School of Medicine; Dr. Ichiro Tsuji, Tohoku

University Graduate School of Medicine; Dr. Yosikazu Nakamura, Jichi Medical School; Dr. Hiroyasu Iso, Osaka University School of Medicine; Dr. Kazumasa Yamagishi, Faculty of Medicine, University of Tsukuba; Dr. Haruo Mikami, Chiba Cancer Center; Dr. Michiko Kurosawa, Juntendo University School of Medicine; Dr. Yoshiharu Hoshiyama, Yokohama Soei University; Dr. Naohito Tanabe, University of Niigata Prefecture; Dr. Koji Tamakoshi, Nagoya University Graduate School of Health Science; Dr. Kenji Wakai, Nagoya University Graduate School of Medicine; Dr. Shinkan Tokudome, National Institute of Health and Nutrition; Dr. Koji Suzuki, Fujita Health University School of Health Sciences; Drs. Shuji Hashimoto and Hiroshi Yatsuya, Fujita Health University School of Medicine; Dr. Shogo Kikuchi, Aichi Medical University School of Medicine; Dr. Yasuhiko Wada, Faculty of Nutrition, University of Kochi; Dr. Takashi Kawamura, Kyoto University Health Service; Dr. Yoshiyuki Watanabe, Kyoto Prefectural University of Medicine Graduate School of Medical Science; Dr. Kotaro Ozasa, Radiation Effects Research Foundation; Dr. Kazuya Mikami, Kyoto Prefectural University of Medicine Graduate School of Medical Science; Dr. Chigusa Date, School of Human Science and Environment, University of Hyogo; Dr. Kiyomi Sakata, Iwate Medical University; Dr. Yoichi Kurozawa, Tottori University Faculty of Medicine; Drs. Takesumi Yoshimura and Yoshihisa Fujino, University of Occupational and Environmental Health; Dr. Akira Shibata, Kurume University; Dr. Naoyuki Okamoto, Kanagawa Cancer Center; and Dr. Hideo Shio, Long-Term Care Health Facility Caretown Minamikusatsu, Shiga.

Funding This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT) (Monbusho); Grants-in-Aid for Scientific Research on Priority Areas of Cancer; and Grants-in-Aid for Scientific Research on Priority Areas of Cancer Epidemiology from MEXT (MonbuKagaku-sho) (Nos. 61010076, 62010074, 63010074, 1010068, 2151065, 3151064, 4151063, 5151069, 6279102, 11181101, 17015022, 18014011, 20014026, 20390156, and 26293138); Comprehensive Research on Cardiovascular and Life-Style Related Diseases (H26-Junkankitou [Seisaku]-Ippan-001 and H29-Junkankitou [Seishuu]-Ippan-003), JSPS KAKENHI Grant Number JP 16H06277, and Grants-in-Aid for China Scholarship Council (CSC file No. 201608050113).

Compliance with ethical standards

Conflict of interest All authors declare that they have no conflicts of interest.

Ethical approval All procedures performed in this study were in accordance with the Helsinki declaration and was approved by Osaka and Nagoya Universities research ethics committees.

Informed consent Informed consent was obtained from all participants included in the study at individual or community leader level.

References

- Matsuda A, Matsuda T, Shibata A, Katanoda K, Sobue T, Nishimoto H, Japan Cancer Surveillance Research Group (2013) Cancer incidence and incidence rates in Japan in 2008: A Study of 25 population-based cancer registries for the Monitoring of Cancer Incidence in Japan (MCIJ) Project. *Jpn J Clin Oncol* 44:388–396
- Qian X, Brinton LA, Schairer C, Matthews CE (2015) Sleep duration and breast cancer risk in the Breast Cancer Detection Demonstration Project follow-up cohort. *Br J Cancer* 112:567–571
- Vogtmann E, Levitan EB, Hale L, Shikany JM, Shah NA, Endeshaw Y, Lewis CE, Manson JE, Chlebowski RT (2013) Association between sleep and breast cancer incidence among postmenopausal women in the women's health initiative. *Sleep* 36:1437–1444
- Girschik J, Heyworth J, Fritschi L (2013) Self-reported Sleep Duration, Sleep Quality, and Breast Cancer Risk in a Population-based Case-Control Study. *Am J Epidemiol* 177:316–327
- Wu AH, Stanczyk FZ, Wang R, Koh WP, Yuan JM, Yu MC (2013) Sleep duration, spot urinary 6-sulfatoxymelatonin levels and risk of breast cancer among Chinese women in Singapore. *Int J Cancer* 132:891–896
- Qin Y, Zhou Y, Zhang X, Wei X, He J (2013) Sleep duration and breast cancer risk: a meta-analysis of observational studies. *Int J Cancer* 134:1166–1173
- Pinheiro SP, Schernhammer ES, Tworoger SS, Michels KB (2006) A prospective study on habitual duration of sleep and incidence of breast cancer in a large cohort of women. *Cancer Res* 66(10):5521–5525
- Verkasalo PK, Lillberg K, Stevens RG, Hublin C, Partinen M, Koskenvuo M, Kaprio J (2005) Sleep duration and breast cancer: a prospective cohort study. *Cancer Res* 65(20):9595–9601
- Wu AH, Wang R, Koh W, Stanczyk Z, Lee H, Yu MC (2008) Sleep duration, melatonin and breast cancer among Chinese women in Singapore. *Carcinogenesis* 29:1244–1248
- Kakizaki M, Kuriyama S, Sone T, Ohmori-Matsuda K, Hozawa A, Nakaya N, Fukudo S, Tsuji I (2008) Sleep duration and the risk of breast cancer: the Ohsaki Cohort Study. *Br J Cancer* 99:1502–1505
- Mcelroy JA, Newcomb PA, Titus-L, Egan KM (2006) Duration of sleep and breast cancer risk in a large population-based case-control study. *J Sleep Res* 15(3):241–249
- Hurley S, Goldberg D, Bernstein L, Reynolds P, Hurley S (2015) Sleep duration and cancer risk in women. *Cancer Causes Control* 26(7):1037–1045
- Brzezinski A (1997) Melatonin in humans. *N Engl J Med* 336:186–195
- Schernhammer ES, Schulmeister K (2004) Melatonin and cancer risk: does light at night compromise physiologic cancer protection by lowering serum melatonin levels? *Br J Cancer* 90:941–943
- Schernhammer ES, Kroenke CH, Dowsett M, Folkert E, Hankinson SE (2006) Urinary 6-sulfatoxymelatonin levels and their correlations with lifestyle factors and steroid hormone levels. *J Pineal Res* 40:116–124
- Anisimov VN (2003) The role of pineal gland in breast cancer development. *Crit Rev Oncol Hematol* 46:221–234
- Cohen M, Lippman M, Chabner B (1978) Role of the pineal gland in the aetiology and treatment of breast cancer. *Lancet* 2:814–881
- Blask DE, Wilson ST, Zalatan F (1997) Physiological melatonin inhibition of human breast cancer cell growth in vitro: evidence for a glutathione-mediated pathway. *Cancer Res* 57:1909–1914
- Cos S, Gonzalez A, Martinez-Campa C, Mediavilla MD, Alonso-Gonzalez C, Sanchez-Barcelo EJ (2006) Estrogen-signaling pathway: a link between breast cancer and melatonin oncostatic actions. *Cancer Detect Prev* 30(2):118–128
- Surakasula A, Nagarjunapu GC, Raghavaiah KV (2014) A comparative study of pre- and post-menopausal breast cancer: Risk factors, presentation, characteristics and management. *J Res Pharm Pract* 3(1):12–18
- Lovejoy JC, Champagne CM, de Jonge L, Xie H, Smith SR (2008) Increased visceral fat and decreased energy expenditure during the menopausal transition. *Int J Obes* 32(6):949–958

22. Cleary MP, Grossmann ME (2009) Obesity and breast cancer: the estrogen connection. *Endocrinology* 150(6):2537–2542
23. Molis TM, Spriggs LL, Hill SM (1994) Modulation of estrogen receptor mRNA expression by melatonin in MCF-7 human breast cancer cells. *Mol Endocrinol* 8:1681–1690
24. Xiao Q, Signorello LB, Brinton LA, Cohen SS, Blot WJ, Matthews CE (2016) Sleep duration and breast cancer risk among black and white women. *Sleep Med* 20:25–29
25. Lambe M, Hsieh CC, Chan HW, Ekblom A, Trichopoulos D, Adami HO (1996) Parity, age at first and last birth, and risk of breast cancer: a population-based study in Sweden. *Breast Cancer Res Treat* 38(3):305–311
26. Palmer JR, Boggs DA, Wise LA, Ambrosone CB, Adams-Campbell LL, Rosenberg L (2011) Parity and lactation in relation to estrogen receptor negative breast cancer in African American women. *Cancer Epidemiol Biomark Prev* 20:1883–1891
27. Ambrosone CB, Zirpoli G, Ruszczyc M, Shankar J, Hong CC, McIlwain D, Roberts M, Yao S, McCann SE, Ciupak G, Hwang H, Khoury T, Jandorf L, Bovbjerg DH, Pawlish K, Bandera E (2014) Parity and breastfeeding among African-American Women: differential effects on breast cancer risk by estrogen receptor status in the Women's Circle of Health Study. *Cancer Causes Control* 25(2):259–265
28. Signal TL, Gander PH, Sangalli MR, Travier N, Firestone RT, Tuohy JF (2007) Sleep duration and quality in healthy nulliparous and multiparous women across pregnancy and post-partum. *Aust N. Z J Obstet Gynaecol* 47(1):16–22
29. Tamakoshi A, Ozasa K, Fujino Y, Suzuki K, Sakata K, Mori M, Kikuchi S, Iso H, The JACC Study Group (2013) Cohort profile of the Japan Collaborative Cohort Study. *J Epidemiol* 23:227–232
30. Reiter RJ (1980) The pineal and its hormones in the control of reproduction in mammals. *Endocr Rev* 1:109–131
31. Leon-Blanco MM, Guerrero JM, Reiter RJ, Calvo JR, Pozo D (2003) Melatonin inhibits telomerase activity in the MCF-7 tumor cell line both in vivo and in vitro. *J Pineal Res* 35(3):204–211
32. Allegra M, Reiter RJ, Tan DX, Gentile C, Tesoriere L, Livrea MA (2003) The chemistry of melatonin's interaction with reactive species. *J Pineal Res* 34:1–10
33. Longcope C, Speizer FE (1995) Reproductive factors and family history of breast cancer in relation to plasma estrogen and prolactin levels in postmenopausal women in the Nurses' Health Study (United States). *Cancer Causes Control* 6:217–224
34. Kwa HG, Cleton F, Bulbrook RD, Wang DY, Hayward JL (1981) Plasma prolactin levels and breast cancer: relation to parity, weight and height, and age at first birth. *Int J Cancer* 28:31–34
35. Bernstein L, Pike MC, Ross RK, Judd HL, Brown JB, Henderson BE (1985) Estrogen and sex hormone-binding globulin levels in nulliparous and parous women. *J Natl Cancer Inst* 74:741–745
36. Tamakoshi K, Yatsuya H, Wakai K, Suzuki S, Nishio K, Lin Y, Niwa Y, Kondo T, Yamamoto A, Tokudome S, Toyoshima H, Tamakoshi A, JACC Study Group (2005) Impact of menstrual and reproductive factors on breast cancer risk in Japan: results of the JACC study. *Cancer Sci* 96(1):57–62
37. Ukawa S, Tamakoshi A, Mori M, Ikehara S, Shirakawa T, Yatsuya H, Iso H, JACC study group (2018) Association between average daily television viewing time and the incidence of ovarian cancer: findings from the Japan Collaborative Cohort Study. *Cancer Causes Control* 29(2):213–219
38. Fang HF, Miao NF, Chen CD, Sithole T, Chung MH (2015) Risk of cancer in patients with insomnia, parasomnia, and obstructive sleep apnea: a nationwide nested case-control study. *J Cancer* 6(11):1140–1147
39. Lockley SW, Skene DJ, Arendt J (1999) Comparison between subjective and actigraphic measurement of sleep and sleep rhythms. *J Sleep Res* 8:175–183