



Correction to: Kinesiophobia modulates lumbar movements in people with chronic low back pain: a kinematic analysis of lumbar bending and returning movement

Michihiro Osumi¹ · Masahiko Sumitani² · Yuko Otake² · Tomohiko Nishigami³ · Akira Mibu³ · Yuki Nishi¹ · Ryota Imai¹ · Gosuke Sato¹ · Yusuke Nagakura¹ · Shu Morioka¹


Published online: 13 June 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: European Spine Journal
<https://doi.org/10.1007/s00586-019-06010-4>

In the figure 2, “CLBP Low fear” which located right end of Time of Phase 1 is wrong. The correct statement is “CLBP High fear”. The complete correct figure 2 is given below.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Fig. 2 Comparison of phase duration among groups.
#Significant difference between CLBP high-fear and HC group ($p < 0.017$). §Significant difference between CLBP high-fear and CLBP low-fear groups ($p < 0.017$)



The original article can be found online at <https://doi.org/10.1007/s00586-019-06010-4>.

✉ Michihiro Osumi
m.ohsumi@kio.ac.jp

¹ Graduate School of Health Science, Kio University, 4-2-2 Umaminaka, Koryo-cho, Kitakatsuragi-gun, Nara 635-0832, Japan

² Department of Pain and Palliative Medicine, The University of Tokyo Hospital, Tokyo, Japan

³ Department of Nursing and Physical Therapy, Konan Woman’s University, Kobe, Hyogo, Japan