



Contents lists available at ScienceDirect

Indian Pacing and Electrophysiology Journal

journal homepage: www.elsevier.com/locate/IPEJ

Predictors of recurrence of paroxysmal atrial fibrillation following cryoablation based pulmonary vein isolation

Keywords:

Editorial
Atrial fibrillation
Recurrence
Predictors

Cryoablation based pulmonary vein isolation (PVI) is a safe and effective method for treatment of paroxysmal atrial fibrillation (PAF). Recurrence after cryoablation for PAF is not uncommon [1,2], ranging from 25 to 50%. Celik [3] and coworkers have attempted to identify factors associated with recurrence of PAF after cryoablation of PVs. Recurrence of AF after successful PVI (using radiofrequency catheter ablation) is often due to PV reconnection [4], other factors include non PV triggers, and other focal mechanisms. There are conflicting results about the predictors of AF recurrence after cryoablation based PVI. Present report by Celik et al. [3] published in this issue of Indian Pacing and Electrophysiology Journal addresses this problem. They performed cryoablation based PVI in 50 patients with PAF and followed them for 12 months. Left atrial volume (LAV), left atrial volume index (LAVi), serum galactin-3 (Gal-3) level and neutrophil to lymphocyte ratio (N/L ratio) were measured before ablation and at 6 and 12 months of follow-up post ablation. Atrial fibrillation recurred in 14 patients and 36 were free of recurrence. Increased LAV and LAVi measured at baseline were associated with AF recurrence after cryoablation based PVI. However, markers such as Gal-3 and N/L ratio were not helpful to predict AF recurrence in this cohort.

If we are able to identify clinical predictors of AF recurrence after cryoablation, it will help us to avoid unnecessary procedures, and complications related to it. It will help patient selection and improve clinical outcome following cryoablation based PVI. Although cryoablation based PVI eliminates the triggers, the effect of atrial substrate is often overlooked. Till date, several predictors of outcome of AF ablation have been identified which includes: age, duration of AF, LA size, number of antiarrhythmic drugs used pre-ablation, early recurrence of AF post ablation and co-morbidities including: obesity, sleep apnoea, hypertension, LA fibrosis [5–8].

The mechanism of AF includes interplay between electrical, contractile and structural remodeling, which are important synergistic components of the AF substrate. Further, the role of metabolic

milieu which includes pro-inflammatory, and pro-oxidant environment in the development of AF recurrence after ablation therapy should not be overlooked. Factors related to ongoing inflammation and oxidative stress including: red cell distribution width (RDW), N/L ratio, HsCRP (high sensitivity C-reactive protein) have shown some association with AF [9–14]. In a study by Aksu et al., [15] patients with AF recurrence after cryoablation had higher baseline RDW which is a measure of size and variability of red blood cells (RBCs). It is an inexpensive and easily available marker of inflammation and oxidative stress in AF. Higher RDW quartiles are related with left atrial enlargement (LAE), which is known to be an indicator of advanced LA structural remodeling. We can't determine if LAE is the cause or effect of PAF and elevated plasma level of RDW. Thus, further studies are needed to establish exact pathophysiologic and prognostic role of RDW in AF.

Epicardial adipose tissue (EAT) thickness has also been found to be associated with AF recurrence after catheter ablation [16–18]. It has been shown that EAT contains numerous autonomic ganglionated plexi, which take part in initiation and continuation of AF [19]. As shown by Kocyigit et al. [20] total periatrial EAT measured with cardiac computed tomography is an independent predictor of AF recurrence in patients undergoing cryoablation. Canpolat et al. [21] investigated the association of transforming growth factor (TGF)- β 1 (a marker of atrial fibrosis) with the extent of LA fibrosis using delayed enhancement cardiac magnetic resonance imaging (DEMRI), and effects of LA fibrosis on the success of cryoablation for lone AF. Increase in LA fibrosis of >20% predicted AF recurrence after catheter ablation with a sensitivity of 100% and specificity of 93.75%. Plasma TGF- β 1 level >15,894 pg/mL predicted the presence of LA fibrosis with sensitivity of 70.37% and specificity of 100%.

Next in line are the autoantibodies, which may have a role in structural atrial remodelling [22]. Anti-M2-R (antimuscarinic acetylcholine receptor antibody) level is a marker of LA fibrosis detected by DE-MRI in paroxysmal lone AF patients. Gurses et al. [23] demonstrated elevated anti-M2-R levels in patients with late recurrence of AF following cryoablation. Thus, by predicting the severity of LA fibrosis in DE-MRI, anti-M2-R levels may improve patient selection for successful AF ablation, avoiding procedures in patients likely to have recurrence of AF. Yalcin et al., [24] determined the preprocedural predictive value of anti- β 1-R (autoantibody against β 1 adrenergic receptor) levels for AF recurrence. Elevated anti- β 1-R levels (>159 ng/mL) was an independent predictor of AF recurrence following cryoablation in this study.

Isolation of PVs with cryoablation addresses the triggers, but substrate for perpetuation of AF may continue to remain untreated and progress over time.

Conclusion: There is a potential role of inflammatory indices to

Peer review under responsibility of Indian Heart Rhythm Society.

<https://doi.org/10.1016/j.ipej.2018.11.009>

0972-6292/Copyright © 2018, Indian Heart Rhythm Society. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

predict recurrence of AF after cryoablation. Various indices like-RDW, autoantibodies and periatrial fat have been associated with recurrence of AF after cryoablation, but their causal relation is yet to be proven. More work is needed to understand about the predictors of recurrence of AF following cryoablation based PVI.

Disclosure

None to declare.

References

- [1] Andrade JG, Khairy P, Guerra PG, Deyell MW, Rivard L, Macle L, Thibault B, Talajic M, Roy D, Dubuc M. Efficacy and safety of cryoballoon ablation for atrial fibrillation: a systematic review of published studies. *Heart Rhythm* 2011;8: 1444–51.
- [2] Oto A, Aytemir K, Canpolat U, Karakulak U, Evranos B, Sahiner L, Okutucu S, Kaya EB, Tokgozoglu L, Kabakci G. Pulmonary vein isolation with the cryoballoon technique in atrial fibrillation treatment: single centre experience. *Turk Kardiyol Dernegi Arsivi* 2013;41:299–309.
- [3] Celik AI, et al. Predictors of the paroxysmal atrial fibrillation recurrence following cryoballoon-based pulmonary vein isolation: assessment of left atrial volume, left atrial volume index, galectin-3 level and neutrophil-to-lymphocyte ratio. *Indian Pacing Electrophysiol J* 2018. <https://doi.org/10.1016/j.ipej.2018.07.003>.
- [4] Nery PB, Belliveau D, Nair GM, Bernick J, Redpath CJ, Szczotka A, Sadek MM, Green MS, Wells G, Birnie DH. Relationship between pulmonary vein reconnection and atrial fibrillation recurrence: a systematic review and meta-analysis. *JACC Clin Electrophysiol* 2016 Aug;2(4):474–83.
- [5] Akoum N, Daccarett M, McGann C, Segerson N, Vergara G, Kuppahally S, et al. Atrial fibrosis helps select the appropriate patient and strategy in catheter ablation of atrial fibrillation: a DE-MRI guided approach. *J Cardiovasc Electrophysiol* 2011;22:16–22.
- [6] Arya A, Hindricks G, Sommer P, Huo Y, Bollmann A, Gaspar T, Bode K, Huser D, Kottkamp H, Piorkowski C. Long-term results and the predictors of outcome of catheter ablation of atrial fibrillation using steerable sheath catheter navigation after single procedure in 674 patients. *Europace* 2010;12(2):173–80.
- [7] Aytemir K, Oto A, Canpolat U, Sunman H, Yorgun H, Sahiner L, Kaya EB. Immediate and medium-term outcomes of cryoballoon-based pulmonary vein isolation in patients with paroxysmal and persistent atrial fibrillation: single-centre experience. *J Intervent Card Electrophysiol* 2013;38(3):187–95.
- [8] Lin YJ, Tsao HM, Chang SL, et al. Prognostic implications of the high-sensitive C-reactive protein in the catheter ablation of atrial fibrillation. *Am J Cardiol* 2010;105(4):495–501.
- [9] Trivedi C, Di Biase L. Baseline neutrophil/lymphocyte ratio predicts recurrences after radiofrequency catheter ablation: results from prospective study on paroxysmal atrial fibrillation. *Circulation* 2013;128:A18588.
- [10] Guo Y, Lip GY, Apostolakis S. Inflammation in atrial fibrillation. *J Am Coll Cardiol* 2012;60(22):2263–70.
- [11] Aviles RJ, Martin DO, Apperson-Hansen C, Houghtaling PL, Rautaharju P, Kronmal RA, Tracy RP, Van Wagoner DR, Psaty BM, Lauer MS, Chung MK. Inflammation as a risk factor for atrial fibrillation. *Circulation* 2003;108(24): 3006–10.
- [12] Issac TT, Dokainish H, Lakkis NM. Role of inflammation in initiation and perpetuation of atrial fibrillation: a systematic review of the published data. *J Am Coll Cardiol* 2007;50(21):2021–8.
- [13] Koroglu S, Tuncer C, Acar G, Akcay A, Sokmen G, Yalcintas S, Nacar A, Altun B, Sokmen A. Relation of inflammatory and oxidative markers to the occurrence and recurrence of persistent atrial fibrillation. *Turk Kardiyol Dernegi Arsivi : Turk Kardiyoloji Derneginin yayin organidir* 2012;40(6):499–504.
- [14] Gurses KM, Yalcin MU, Kocyigit D, Evranos B, Ates AH, Yorgun H, Sahiner ML, Kaya EB, Ozer N, Oto MA, Aytemir K. Red blood cell distribution width predicts outcome of cryoballoon-based atrial fibrillation ablation. *J Intervent Card Electrophysiol* 2015 Jan;42(1):51–8.
- [15] Aksu T, Baysal E, Guler TE, Golcuk SE, Erden I, Ozcan KS. Predictors of atrial fibrillation recurrence after cryoballoon ablation. *Hematol Res Rev* 2015 Jun 29;6:211–7.
- [16] Wong CX, Abed HS, Molaee P, et al. Pericardial fat is associated with atrial fibrillation severity and ablation outcome. *J Am Coll Cardiol* 2011;57: 1745–51.
- [17] Tsao HM, Hu WC, Wu MH, et al. Quantitative analysis of quantity and distribution of epicardial adipose tissue surrounding the left atrium in patients with atrial fibrillation and effect of recurrence after ablation. *Am J Cardiol* 2011;107:1498–503.
- [18] Nagashima K, Okumura Y, Watanabe I, et al. Association between epicardial adipose tissue volumes on 3-dimensional reconstructed CT images and recurrence of atrial fibrillation after catheter ablation. *Circ J* 2011;75:2559–65.
- [19] Po SS, Nakagawa H, Jackman WM. Localization of left atrial ganglionated plexi in patients with atrial fibrillation. *J Cardiovasc Electrophysiol* 2009;20: 1186–9.
- [20] Kocyigit D, Gurses KM, Yalcin MU, Turk G, Evranos B, Yorgun H, Sahiner ML, Kaya EB, Hazirolan T, Tokgozoglu L, Oto MA, Ozer N, Aytemir K. Periatrial epicardial adipose tissue thickness is an independent predictor of atrial fibrillation recurrence after cryoballoon-based pulmonary vein isolation. *J Cardiovasc Comput Tomogr* 2015 Jul-Aug;9(4):295–302.
- [21] Canpolat U, Oto A, Hazirolan T, Sunman H, Yorgun H, Sahiner L, Kaya EB, Aytemir K. A prospective DE-MRI study evaluating the role of TGF- β 1 in left atrial fibrosis and implications for outcomes of cryoballoon-based catheter ablation: new insights into primary fibrotic atricardiomyopathy. *J Cardiovasc Electrophysiol* 2015 Mar;26(3):251–9.
- [22] Baba A, Fu M. Autoantibodies in atrial fibrillation: actor, biomarker or bystander? *Autoimmunity* 2008;41:470–2.
- [23] Gurses KM, Yalcin MU, Kocyigit D, Kesikli SA, Canpolat U, Yorgun H, Sahiner ML, Kaya EB, Hazirolan T, Ozer N, Oto MA, Guc D, Aytemir K. M2-muscarinic acetylcholine receptor autoantibody levels predict left atrial fibrosis severity in paroxysmal lone atrial fibrillation patients undergoing cryoablation. *Europace* 2015 Feb;17(2):239–46.
- [24] Yalcin MU, Gurses KM, Kocyigit D, Kesikli SA, Dural M, Evranos B, Yorgun H, Sahiner L, Kaya EB, Oto MA, Guc D, Aytemir K, Ozer N. Cardiac autoantibody levels predict recurrence following cryoballoon-based pulmonary vein isolation in paroxysmal atrial fibrillation patients. *J Cardiovasc Electrophysiol* 2015 Jun;26(6):615–21.

Daljeet Kaur, C. Narasimhan*

CARE Hospital, Banjara Hills, Hyderabad, Telangana, India

* Corresponding author. Department of Cardiology and Arrhythmia division, CARE Hospital and CARE Foundation, Banjara Hills, Hyderabad, Telangana, India.

E-mail address: calambur1@gmail.com (C. Narasimhan).

Available online 14 November 2018