



Knockout of Nr2e3 prevents rod photoreceptor differentiation and leads to selective L-/M-cone photoreceptor degeneration in zebrafish

Shanglun Xie^{a,1}, Shanshan Han^{a,1}, Zhen Qu^{a,1}, Fei Liu^a, Jingzhen Li^a, Shanshan Yu^a, James Reilly^b, Jiayi Tu^a, Xiliang Liu^a, Zhaojing Lu^a, Xuebin Hu^a, Tinsae Assefa Yimer^a, Yayun Qin^a, Yuwen Huang^a, Yuexia Lv^a, Tao Jiang^a, Xinhua Shu^b, Zhaohui Tang^a, Haibo Jia^{a,*}, Fulton Wong^c, Mugen Liu^{a,**}

^a Key Laboratory of Molecular Biophysics of Ministry of Education, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, Hubei 430074, PR China

^b Department of Life Sciences, Glasgow Caledonian University, Glasgow G4 0BA, Scotland, United Kingdom of Great Britain and Northern Ireland

^c Department of Ophthalmology, Duke University School of Medicine, Durham, NC 27710, USA

ARTICLE INFO

Keywords:

Nr2e3
CRISPR
Zebrafish
Photoreceptor
Differentiation
Degeneration

ABSTRACT

Mutations in the photoreceptor cell-specific nuclear receptor gene *Nr2e3* increased the number of S-cone photoreceptors in human and murine retinas and led to retinal degeneration that involved photoreceptor and non-photoreceptor cells. The mechanisms underlying these complex phenotypes remain unclear. In the hope of understanding the precise role of *Nr2e3* in photoreceptor cell fate determination and differentiation, we generated a line of *Nr2e3* knockout zebrafish using CRISPR technology. In these *Nr2e3*-null animals, rod precursors undergo terminal mitoses but fail to differentiate as rods. Rod-specific genes are not expressed and the outer segment (OS) fails to form. Formation and differentiation of cone photoreceptors is normal. Specifically, there is no increase in the number of UV-cone or S-cone photoreceptors. Laminated retinal structure is maintained. After normal development, L-/M-cones selectively degenerate, with progressive shortening of OS that starts at age 1 month. The amount of cone phototransduction proteins is concomitantly reduced, whereas UV- and S-cones have normal OS lengths even at age 10 months. In vitro studies show *Nr2e3* synergizes with *Crx* and *Nrl* to enhance *rhodopsin* gene expression. *Nr2e3* does not affect cone opsin expression. Our results extend the knowledge of *Nr2e3*'s roles and have specific implications for the interpretation of the phenotypes observed in human and murine retinas. Furthermore, our model may offer new opportunities in finding treatments for enhanced S-cone syndrome (ESCS) and other retinal degenerative diseases.

1. Introduction

To cope with the enormous range of ambient light levels encountered in daily life, the vertebrate retina utilizes two types of photoreceptors: rods that function in dim light and cones that function in bright light and mediate color vision. Although the signaling pathway for phototransduction is common for rods and cones, the signaling proteins are mostly coded by distinct sets of rod and cone specific genes. The human retina has one type of rod and three types of cone. Rods

express the visual pigment rhodopsin, whereas the three types of cones express the short (S, blue), medium (M, green) or long (L, red) wavelength cone opsin. The murine retina has one type of rod and only S- (blue) and M- (green) cones; some cones express both S and M opsins. The zebrafish has one type of rod and four types of cone: L-, M-, S- and UV-cone. The fourth type expresses a cone opsin that is maximally sensitive to UV light [1–6]. During development, the rods, cones and five other major classes of retinal cells are produced from a common pool of pluripotent progenitor cells. The progenitors undergo mitosis

Abbreviations: Nr2e3, nuclear receptor subfamily 2, group E, member 3; ESCS, enhanced S-cone syndrome; CRISPR, clustered regularly interspaced short palindromic repeat; hpf, hours post-fertilization; dpf, days post-fertilization; mpf, months post-fertilization; WT, wild type

* Correspondence to: H. Jia, Key Laboratory of Molecular Biophysics of Ministry of Education, Center for Human Genome Research, Huazhong University of Science and Technology, Wuhan, Hubei 430074, PR China.

** Corresponding author.

E-mail addresses: haibo.jia@hust.edu.cn (H. Jia), lium@mail.hust.edu.cn (M. Liu).

¹ These authors contributed equally to this work.

<https://doi.org/10.1016/j.bbadis.2019.01.022>

Received 22 August 2018; Received in revised form 20 January 2019; Accepted 21 January 2019

Available online 23 January 2019

0925-4439/ © 2019 Elsevier B.V. All rights reserved.

and the post-mitotic cells then undergo differentiation into specific types of retinal cells. Differentiation is conceptualized as a process of selective expression of sets of specific genes [7–12]. For rods and cones, several transcription factors that control photoreceptor-specific gene expression have been identified. Cone-rod homeobox transcription factor (Crx) is central to photoreceptor determination [13–15] and required for the development of photoreceptor outer segment (OS) and expression of photoreceptor-specific genes [14–17]. Crx is expressed both in rod and cone photoreceptors. Mutations in *Crx* cause photoreceptor degeneration in humans (autosomal dominant cone-rod dystrophy (CRD) [13,18], autosomal dominant retinitis pigmentosa (adRP) [19] and autosomal recessive Leber congenital amaurosis (LCA) [20]) and in mice [14]. Neural retina leucine zipper protein (Nrl) is a leucine zipper transcription factor that is required for rod development. Nrl is preferentially expressed in rods. It acts synergistically with Crx to regulate *rhodopsin* transcription. Mutations in human *Nrl* cause retinal degeneration: autosomal dominant retinitis pigmentosa (adRP) [21], autosomal recessive retinitis pigmentosa (arRP) [22] and enhanced S-cone syndrome (ESCS) [23]. More details about ESCS and RP will be presented later. Deletion of *Nrl* in mice caused the complete loss of rod function and super-normal S-cone function. In the absence of *Nrl*, the rods acquired some of the characteristics of S-cones. Accordingly, *Nrl* seems to act as a molecular switch during rod-cell development by directly modulating rod-specific genes while simultaneously inhibiting the S-cone pathway through the activation of *Nr2e3* [24].

The nuclear receptor transcription factor *Nr2e3* is expressed in photoreceptors. It forms a complex with Crx that enhances the expression of rod-specific genes and represses the expression of cone-specific genes in rods [25,26]. Mutations in human *Nr2e3* cause ESCS, an autosomal recessive disorder characterized by reduced rod, L- and M-cone responses and enhanced sensitivity to short-wave light due to supernumerary S-cones. In addition, patients with ESCS have dysmorphic retinas, showing an early disorganization of the laminated retinal structure with rosette formation in the outer nuclear layer (ONL) where photoreceptor cell bodies are located, and a slow, progressive retinal degeneration [27–30]. Mutations in *Nr2e3* have also been identified in patients with Goldmann-Favre syndrome, clumped pigmentary retinal degeneration [31], adRP [32], and arRP [33]. A spontaneous recessive null mutation in the murine *Nr2e3* has been identified in the *retinal degeneration 7 (rd7)* mice. The phenotype of this mutant includes progressive photoreceptor degeneration and attenuation of the electroretinogram (ERG) with time, abnormal retinal lamination with rosette formation in the ONL and a 2-fold increase in the number of S-cones [34,35]; however, the majority of the photoreceptors in the *rd7* retina are a hybrid cell type that expresses both rod- and cone-specific genes [36].

The roles played by *Nrl* and *Nr2e3* in rod/cone cell fate determination and differentiation have been exploited in a strategy of therapeutic cellular reprogramming to treat RP, a retinal degeneration characterized by primary rod defects and degeneration followed by secondary cone loss. Many RP-causing mutations occur in rod-specific genes. The rationale for such a strategy is that knocking down rod determinants such as *Nrl* or *Nr2e3* reprograms rods to a cone fate, rendering rod-specific gene mutations irrelevant, with consequent preservation of retinal structural integrity and function [37–40].

Although much is known about the function of *Nr2e3*, major questions remain unanswered. For example, the role of *Nr2e3* in retinal progenitors for cell fate determination is unclear [5]. In *Nr2e3*^{-/-} retinas, the aberrant S-cones could have arisen from an alteration of cell fate or dedifferentiation of postmitotic premature rods [25].

Given the strategic importance of these transcription factors in potential therapeutic intervention, it is important to better understand their function. We have decided to study *Nr2e3* in zebrafish by knocking out the gene using clustered regularly interspaced short palindromic repeat (CRISPR) technology. Architecture and cell types in zebrafish retina are conserved with that in human; furthermore, the

zebrafish is diurnal and possesses a cone dominant retina similar to the macula in human retina [41–43]. Importantly, genomic structure and amino acid sequence of zebrafish *Nr2e3*, especially the DNA-binding domain (DBD) and ligand binding domain (LBD), is highly conserved with the human ortholog [25,44]. For these reasons, zebrafish could be a suitable organism to study the roles of *Nr2e3* underlying photoreceptor development and differentiation. Here, we report on findings from the study of such *Nr2e3*-null zebrafish.

2. Materials and methods

2.1. Zebrafish maintenance and breeding

Wild-type AB strain of zebrafish were raised in recirculating water system (pH 6.6–7.4, 26–28.5 °C) and fed with brine shrimps three times a day under the condition of 14/10 h light/dark cycle. Embryos were kept in E3 medium (5 mM NaCl, 0.17 mM KCl, 0.33 mM CaCl₂, 0.33 mM MgSO₄) at 30 °C. The medium was changed once a day, after 96 h incubation, the larvae were transferred into tanks and fed with live paramecia three times a day. Our study was approved by the Ethics Committee of Huazhong University of Science and Technology.

2.2. Cas9 mRNA and gRNA synthesis

Plasmids that contain the template DNA for in vitro transcription of Cas9 mRNA were linearized by *NotI* and purified by ethanol precipitation. Capped Cas9 mRNA was synthesized using the mMACHINE Mmachine SP6 kit (Ambion, Austin, Texas, USA) and purified by LiCl precipitation. Plasmids that contain the gRNA scaffold sequence were linearized by *BamHI* and purified by ethanol precipitation. gRNA was synthesized using TranscriptAid T7 High Yield Transcription Kit (Thermo scientific, Wilmington, Delaware, USA) and purified by phenol:chloroform extraction and isopropanol precipitation.

2.3. Microinjection and genotyping

The compounds of gRNA and Cas9 mRNA were injected into one-cell stage WT embryos according to previously reported protocol [45]. Effective groups were raised to sexual maturity and outcrossed with WT fish to obtain F₁ zebrafish. F₁ males and F₁ females carrying the same mutations were crossed; homozygous fish were generated. A pair of primers (5'-TAGTTTCTGCCCTTATGTCTT-3', 5'-CTGTCTGATTGGAATGTTGT-3') that flanked the target site was used to amplify the 480 bp DNA fragments; these amplifying fragments were then identified by agarose electrophoresis and DNA sequencing. The homozygotes were also confirmed by quantitative Real-time PCR and Western blot.

2.4. Hematoxylin and eosin (H&E) staining

Whole mount zebrafish larvae or isolated eyes were fixed in the 4% paraformaldehyde (PFA) overnight at 4 °C. Next, larvae or eyes were dehydrated in 30% sucrose (diluted in PBS). After that, the tissues were embedded in OCT compound; embedded tissues were positioned parallel to the blade and were sectioned sagittally. For larvae each slice was 8 μm thick, and for eyes each slice was 12 μm thick. Operations were followed as in previous protocol [46,47]. We chose the slices containing optic nerve and stained with hematoxylin and eosin. Images were captured by Olympus BX53 microscope (Olympus Imaging, Tokyo, Japan).

2.5. Immunofluorescence

Retinal cryosections from WT and *Nr2e3*^{-/-} zebrafish were put at room temperature for 20 min for drying. Then the sections were permeabilized in PDT (PBS containing 1% DMSO and 0.1% TritonX-100) for 10 min. After being blocked with 10% goat serum in PBDT (PDT

containing 1% BSA) for 1 h, tissues on the slide were incubated with primary antibody overnight at 4 °C. The primary antibodies used in this study and the dilution ratio are shown in Supplementary Table 1. For the next step, slides were washed in PDT 3 times and then incubated with Alexa Fluor 488 nm secondary antibody (Life Technologies, Carlsbad, California, USA) at 37 °C for 1 h. DAPI (Sigma, St. Louis, Missouri, USA) was diluted with PBS and used to label nuclei at a concentration of 5 µg/ml. Images were acquired using a confocal laser-scanning microscope (FluoView FV1000 confocal microscope, Olympus Imaging). All operations were followed as previous protocol [48,49].

2.6. Transmission electron microscopy

Adult zebrafish eyes or whole body of larval zebrafish were fixed in 2.5% glutaraldehyde in 0.1 M PBS. After washing 3 times, the eyes were further fixed in 1% osmium tetroxide in 0.1 M PBS buffer at room temperature for 2 h. For the next step, the eyes were dehydrated in gradient ethanol and then incubated in acetone for 20 min at room temperature. The eyes were treated with 50% (1 h), 75% (3 h) and 100% (overnight) epoxy resin (mixed with acetone, v/v) successively, and then heated at 70 °C overnight. Ultrathin sections were produced using a Reichert-Jung ultramicrotome (Leica, Wetzlar, Germany). Sections were stained with 3% uranyl acetate and 3% lead citrate. Transmission electron microscopy (HT7700, Hitachi, Tokyo, Japan) was used to obtain images.

2.7. RT-PCR and Western blotting

Total RNAs were extracted from embryos or larvae using RNAiso Plus reagent (Takara, Kusatsu, Japan), cDNA was generated by MMLV reverse transcriptase (Invitrogen, Carlsbad, California, USA). The cDNA was used as templates to examine the mRNA level of targeted genes. These experiments were performed using AceQ™ qPCR SYBR® Green Master Mix (Vazyme, Nanjing, China) on the StepOnePlus™ real-time PCR System (ABI, Carlsbad, California, USA). The primers used for specific gene are listed in Supplementary Table 2. Proteins were extracted from adult fish eyes and homogenized in RIPA buffer (Beyotime, Shanghai, China) with protease inhibitor cocktail (Roche, Basel, Switzerland). Concentrations of proteins were determined using the BCA Protein Assay Kit (Beyotime). Lysates were mixed with loading buffer and heated in boiling water for 10 min, then stored immediately at –20 °C. The same amounts of protein were separated on SDS-PAGE and transferred to nitrocellulose membranes (Millipore, Danvers, Minnesota, USA). The membranes were then orderly blocked and incubated in primary antibodies overnight. Antibodies used for WB are listed in Supplementary Table 1. The membranes were washed and incubated in HRP-conjugated secondary antibodies (Thermo Fisher). Finally, the membranes were visualized using ChemiDoc™ XRS + system (Bio-Rad Laboratories, Berkeley, California, USA) to acquire the images. All operations were followed as in previous protocol [50–52].

2.8. EdU and TUNEL assays

A Cell-light EdU Apollo567 in Vitro Kit (RiboBio, Guangzhou, China) and a BrightRed Apoptosis Detection Kit (Vazyme) were used to detect the proliferous and apoptotic cells according to the manufacturer's instructions. Images were captured using a confocal microscope (FV1000, Olympus).

2.9. Cell culture, plasmids and transfection

Dulbecco's modified Eagle's medium (DMEM, Gibco, Grand Island, New York, USA) containing 10% fetal bovine serum (FBS, Gibco) was used to raise HEK293 cells. The full length of zebrafish *Nr2e3* coding sequence was cloned into the p3XFLAG-CMV-7.1 vector; the full length

of zebrafish *Crx* and *Nrl* coding sequences were cloned into pEGFP-C1 vector. Cell-type specific promoter sequences were amplified using specific primers and cloned into the pGL3 Luciferase Reporter Vectors. Primers used for vector construction are listed in Supplementary Table 3. Plasmids were transfected into HEK293 cells with Lipofectamine 3000 Reagent (Thermo Fisher) in Opti-MEM Reduced Serum Medium (Gibco). After 6 hour incubation, the medium was changed with DMEM containing FBS; after 36 hour incubation, cells were harvested and activity of luciferase was examined using Dual-Luciferase® Reporter Assay System (Promega, Madison, Wisconsin, USA).

2.10. Statistical analysis

All experiments were repeated at least three times. Significance was determined by two-tailed Student's *t*-test.

3. Results

3.1. Generation of *Nr2e3* mutant zebrafish strain by CRISPR/Cas9 technology

In order to generate the *Nr2e3* knockout zebrafish strain, cDNA (NM_001007368.1) sequence was screened on the special website (<http://www.crisprscan.org/>) for a high score oligo that was suitable to design as a target site: a 23-bp sequence located in exon 4 was thus selected (Fig. 1A). Cas9 mRNA and gRNA transcribed in vitro were microinjected into one-cell-stage embryos and mutagenic efficiency was evaluated after 48 hpf. All procedures followed previous protocol [45]. We identified a mutation that carries a 37 bp deletion c.485_521del in *Nr2e3* mRNA. This deletion caused a frame shift resulting in an abnormal p.Leu162Glnfs*30 *Nr2e3* protein (Fig. 1B). The mRNA level of *Nr2e3* was reduced to nearly 40% compared with those of WT control, an effect probably explained by nonsense-mediated RNA decay (Fig. 1C). Western blot assay was performed to evaluate the protein level of *Nr2e3*. A 46 kD protein was detected in WT extracts but was lost in the mutant samples (Fig. 1D). Thus, our knockout zebrafish are authentic *Nr2e3*-null mutants. The mutant animals appeared smaller during the larval period, but there was no difference in body size at juvenile and adult stages. No other macroscopic phenotypes were observed.

3.2. Differentiation of rod photoreceptors is arrested in *Nr2e3*-null zebrafish

Because *Nr2e3* was reported to express primarily in rods [25], the phenotype of rods was first analyzed in *Nr2e3* mutant zebrafish. Immunofluorescent staining was conducted using anti-rhodopsin antibody to label rod OSs. At 10 dpf, rhodopsin immunolabelling was observed clearly at the ventral and dorsal regions in WT retinas. The labelled pattern of dense cells in ventral retina, sporadic cells in dorsal retina and few cells in central retina is consistent with the depiction of origination and gradient distribution of rods [53]. However, rhodopsin immunolabelling was not found in *Nr2e3* mutant retinas (Fig. 2A). We examined this aspect of the phenotype in adult retinas at age 6 months and 10 months. Similarly, no rhodopsin labelled signals were detected in *Nr2e3* mutant retinas (Fig. 2A). The mRNA levels of rod genes *rhodopsin*, *gnat1*, *grk1a* and *pde6b* were evaluated by the real-time quantification PCR method. Results showed that these genes were not expressed (Fig. 2B). Consistent with the mRNA results, protein levels of rhodopsin, GNAT1, PDE6B and GNB1 were not expressed (Fig. 2C). These data indicate that rods were not developed in *Nr2e3* mutant zebrafish. However, expression of rod genes were rescued when *Nr2e3* mRNA was injected into the mutant embryos, this suggested that the phenotypes of *Nr2e3*^{-/-} zebrafish are due to loss of *Nr2e3* (Fig. S1).

To verify the above observations directly, we performed transmission electron microscopic (TEM) analysis to examine the ultrastructure




Fig. 1. Generation of *Nr2e3* knockout zebrafish. (A) The eight exons of zebrafish *Nr2e3* and the genomic target site are shown. The target sequence is highlighted. (B) Sanger sequencing of the target site in wildtype (WT) and *Nr2e3*^{-/-} zebrafish to confirm the genotype of c.485_521del (del37). The deleted sequence is indicated in the box. (C) *Nr2e3* mRNA levels in WT and *Nr2e3* mutant were examined by quantitative PCR at 36 hpf. Five independent experiments were performed. β -Actin served as endogenous control. The results are shown as mean \pm SD. ****P* < 0.001. (D) *Nr2e3* protein levels in WT and *Nr2e3* mutant were detected at 1 month old larvae by western blotting.

of the mutant retinas. At 10 dpf, unambiguous rod OSs developed well in WT retinas, but only cone OSs were found in *Nr2e3* mutant retinas (Fig. 3A, B). At 1 month, by contrast to the regularly organized and well-formed rod OSs in WT fish, no rod OSs were found in *Nr2e3* mutants (Fig. 3C–D').

Consistent with the lost expression of rod genes and lack of rod OSs, the number of rod nuclei was reduced. Based on the locations and morphological features of rod and cone nuclei [36], we identified small and round rod nuclei arranged regularly in WT ONL, but only sporadic rod nuclei were found in *Nr2e3* mutant retinas (Fig. S2). This feature could be seen clearly by DAPI staining (Fig. 2A).

Nr2e3 has been reported to express in rod progenitors and precursors in zebrafish [53]. Thus, we wondered if generation of rod lineage was implicated. The mRNA level of *pax6*, a marker of Müller glial cells that were identified as rod progenitors [53,54], was not changed (Fig. 4A). The mitotic cells in the central retina that possessed a rounded or fusiform shape were considered as rod progenitors or precursors [54,55]. Therefore, EdU incorporation assays were performed on our zebrafish. Consistent with the previous report, most EdU labelled cells were located at the dorsal and ventral ciliary marginal zone (CMZ), the persistent proliferative regions in the zebrafish retina [53,56]. We identified EdU positive cells that possessed a rounded or fusiform shape in the central ONL or INL of the mutant (arrowheads in Fig. 4B). Furthermore, the labelled pattern of *zpr-3*, another rod marker, at 72 hpf when rod lineage had emerged but before rhodopsin could be detected, was identical between WT and *Nr2e3* mutant retinas (Fig. 4C). These results show that generation of rod lineage is not affected.

To determine if these immature rods would undergo apoptosis before expressing the mature markers and developing OSs, we examined the retinal sections from WT and *Nr2e3*^{-/-} mutants at 10 dpf using terminal deoxynucleotidyl transferase dUTP nickend labeling method (TUNEL). In 18 WT retinal sections we detected 3 TUNEL positive cells in ONL and 12 cells in INL or GCL; by contrast, in 18 mutant sections no labelled cells were found in ONL and only 6 positive cells in INL or GCL (Fig. S3). So, the loss of rods in *Nr2e3* mutants could not be explained by apoptosis.

In conclusion, these results indicate that *Nr2e3* is required for rod differentiation but does not act in rod fate determination.

3.3. Cell fate determination and differentiation of cone photoreceptors are not affected in *Nr2e3* mutant zebrafish

Since in ESCS patients and *rd7* mice, *Nr2e3* mutations caused an overproduction of S-cones, thus we wondered if the immature rods were transformed into UV-cones, the closest homolog of mammalian S-cones [43]. Anti-SWS1 antibody was used to immunolabel the UV-cone OSs. Unlike the phenotypes in ESCS and *rd7*, UV-cones did not increase (Fig. 5A). We then examined the number of S-cones using anti-SWS2 antibody. Similarly, S-cones were not increased (Fig. 5B). The mRNA levels of cone genes *opn1lws1*, *opn1lws2*, *pde6c* and a transcription factor *tbx2b*, which is required for fate determination of UV-cones [10], were not changed (Fig. S4A). The observation that rods were lost but UV- and S-cones did not increase prompted us to analyze the phenotype of L-/M-cones. Anti-LWS and anti-RH2 antibodies were used to immunolabel L-/M-cone OSs, respectively. Again, the number of L-/M-




Fig. 2. *Nr2e3*^{-/-} zebrafish present a no-rod retina. (A) Retinal cryosections from wildtype (WT) and *Nr2e3* mutant were immunostained with anti-rhodopsin antibody at 10 dpf (dorsal is up), 6 months and 10 months (images come from the ventral retina). Confocal images showed there are no rods at all stages in mutant retinas. Scale bars: 20 μ m. (B) Quantitative PCR assays at 5 dpf demonstrated rod genes *rhodopsin*, *gnat1*, *grk1a* and *pde6b* were not expressed in *Nr2e3* mutants. Three independent experiments were performed. β -Actin served as endogenous control. The results are shown as mean \pm SD. *** P < 0.001. (C) Western blotting analysis at 2 months showed the proteins that participant in the rod phototransduction cascade were not expressed in *Nr2e3* mutants. α -Tubulin was used as endogenous control.

cones was not altered (Fig. 5C, D). Consistent with these results, the labelled patterns of PNA (peanut lectin) and GNAT2 (cone transducin subunit), proteins that are located in cone OSs, were identical to those in WT sections (Fig. S4B, C). These results suggest that formation and differentiation of cone photoreceptors proceed as normal in the mutant zebrafish. Specifically, there is no increase in the number of S-cone or UV-cone photoreceptors.

A previous study showed that loss of *Nr2e3* resulted in superabundant proliferation of cone progenitors in *rd7* mouse [5]. Therefore, we counted the number of EdU positive cells at the peripheral ONL, which are considered as nascent cone precursors [57], and found no difference between WT and mutant (arrows in Figs. 4B, S5). These data suggest that proliferation of cone progenitors does not increase in *Nr2e3* mutant zebrafish.

3.4. Laminated retinal structure is maintained in *Nr2e3* mutant zebrafish

Histological examinations were implemented to analyze the whole structure of mutant retinas. Retinal cryosections from WT and mutants at different ages were stained with hematoxylin and eosin. The laminated architecture was maintained in *Nr2e3* mutant retinas. Unlike the phenotype of *rd7* mouse [34], there were no whorls or rosettes in the ONL. Notably, from 1 month of age the lengths of photoreceptor OSs were gradually reduced. Consider that there were no rod OSs in mutant retinas, we surmised that cone OSs were shortened. At all stages that we

analyzed, the thickness of the inner retina was normal (Figs. 6A–D, S6).

3.5. Selective degeneration of L-/M-cone photoreceptors was observed in *Nr2e3* mutant zebrafish

Our histological data indicated that degeneration of cone photoreceptors was occurring in *Nr2e3* mutant zebrafish. To determine which cone subtypes were involved, immunostaining assays were implemented to analyze the phenotypes of the four types of cones in several adult stages. We found that the lengths of UV-cone and S-cone photoreceptor OSs were normal at 4 months and 10 months but OSs of L-/M-cones were shortened gradually, starting at age 1 month (Figs. 7A–D, S7A–D). In 6 months old retinas that were immunolabelled using anti-GNAT2 antibody, the labelled OSs were observed to be dividing into three layers in WT retinas. According to their spatial orientation and shape, these were OSs of UV-cones, S-cones and L-/M-cones. Only the OSs of L-/M-cones were shortened significantly (Fig. S8A). Consistent with the shortened OSs, the protein levels of GNB3 and GNAT2 were reduced dramatically at 6 months and 12 months (Fig. S8B). These results suggest that L-/M-cones selectively degenerate in *Nr2e3* mutant zebrafish.




Fig. 3. Ultrastructural analysis of photoreceptor OSs in wildtype (WT) and *Nr2e3*^{-/-} retinas. (A) At 10 dpf, recognizable rod OSs developed well in WT retinas, (B) whereas only cone OSs were observed in *Nr2e3* mutant retinas. (C) At 1 month of age, well-formed rod OSs were regularly organized in WT retinas; (D) similarly, no rod OSs were observed in *Nr2e3* mutant retinas. (C', D') Enlarged regions of the boxes in C and D. Arrowheads indicate the rod OSs, whereas arrows indicate the cone OSs. Scale bars: 1 μ m in (A, B); 5 μ m in (C, D and C', D').

3.6. *Nr2e3* enhances transcription of rhodopsin synergized with *Crx* and *Nrl*, but does not repress that of UV- or S-cone opsins

It is widely accepted that *Nr2e3* promotes rod differentiation by strengthening the expression of rod genes and repressing the expression of cone genes in rods. These opposing roles are achieved by cooperation with other transcription factors, especially *Crx* and *Nrl* [24–26,58]. From the phenotype of our mutant, which showed different effects the loss of *Nr2e3* had on rods and cones, we suspected that zebrafish *Nr2e3* would strengthen expression of rod genes synergized with *Crx* and *Nrl*, but would not affect that of cone genes. To confirm this hypothesis, reporter assays were conducted in HEK293 cells. The activity of

rhodopsin promoter (2.5 kb) was examined. We showed that *Nr2e3* alone did not increase rhodopsin promoter activity. Co-expressing *Crx* and *Nrl* increased the activity by 2.8-fold. When *Nr2e3* was expressed at the maximum concentration that we have assayed, rhodopsin promoter activity was increased by 4-fold, which was significantly higher than that expressing *Nr2e3* alone or co-expressing *Crx* and *Nrl* (Fig. 8A). This property of zebrafish *Nr2e3* on regulation of rhodopsin expression is conserved with that in mammalian retina [26].

The activity of rhodopsin promoter in this study is different from previous reports in that the mammalian rhodopsin promoter can be activated 70- or 175-fold [26,59]. The length of the 2.5 kb promoter sequence ought to be sufficient because in RHO-EGFP zebrafish a 1 kb




Fig. 4. Generation of rod lineage is not affected in *Nr2e3*^{-/-} zebrafish. (A) *Pax6a* mRNA levels were examined at 36 hpf and 48 hpf. Three independent experiments were performed. β -Actin served as endogenous control. The results are shown as mean \pm SD. NS, no significance. (B) Confocal images of the EdU labelled cells are shown at 60 hpf. Arrowheads indicate the rod progenitors or precursors. (a, b) Magnified images of the boxes in B. (C) Confocal images stained for *zpr-3* at 72 hpf. WT, wildtype. Scale bars: 20 μ m in (B, C); 10 μ m in (a, b).

sequence of rhodopsin promoter could drive EGFP expression in rod photoreceptors. It is possible that there are enhancers far from the promoter, or there are unidentified factors playing a positive role for expression of rhodopsin that were not included in the complexes, so

that the promoter did not reach the maximal activity.

To explain the phenotypes of UV- and S-cones, we tested the role of *Nr2e3* on expression of UV and blue opsin genes. We found that *Nr2e3* did not repress transcription of UV- or S-opsin (Fig. 8B, C). This




Fig. 5. Development of cone photoreceptors is normal in *Nr2e3*^{-/-} zebrafish. Confocal images stained for UV opsin (A), red opsin (C) and green opsin (D) at 5 dpf, as well as blue opsin at 7 dpf (B). The number of cone photoreceptors and distribution of cone opsins were normal. WT, wildtype. Scale bars: 20 μ m.




Fig. 6. Retinal lamination is retained in *Nr2e3*^{-/-} zebrafish despite OSs degeneration. Retinal cryosections from wildtype (WT) and *Nr2e3* mutant at 10 dpf (A), 1 month (B), 4 months (C) and 12 months (D) were stained with hematoxylin and eosin. RPE, retinal pigment epithelium; OS, outer segment; IS, inner segment; ONL, outer nuclear layer; OPL, outer plexiform layer; INL, inner nuclear layer; IPL, inner plexiform layer; GCL, ganglion cell layer. Scale bars: 40 μm.

property of zebrafish *Nr2e3* is different from that of the mammalian ortholog [26]. Because zebrafish *Nr2e3* did not repress expression of UV or S cone opsins, expression of these genes would not undergo expansion when *Nr2e3* was lost, thus the number of UV or S cones would not increase in *Nr2e3* mutant zebrafish.

Together, our reporter assays indicate that in zebrafish, *Nr2e3* may promote rod differentiation by reinforcing expression of rod genes. The fact that *Nr2e3* does not repress expression of UV or S cone genes may explain the phenotype of UV- and S-cones.

4. Discussion

In the hope of understanding the precise role of *Nr2e3* and providing interpretation of the phenotypes of ESCS and *rd7*, an *Nr2e3*-null zebrafish model was created in this study. We show that *Nr2e3* has no role in photoreceptor cell fate determination. It is necessary for rod but not cone photoreceptor differentiation. Lack of *Nr2e3* leads to selective L-/M-cone photoreceptor degeneration.

Loss of *Nr2e3* function leads to very different outcomes for rod development in mouse and in zebrafish. In the *rd7* mouse, rod genes were expressed normally at the mRNA and protein levels [5,25,36,60]. The physiological function of rods was maintained initially [61]. These observations are consistent with normal development of rods in the *rd7* mouse. By contrast, the post-mitotic premature rods of *Nr2e3*^{-/-} zebrafish fail to develop at all. The phenotype of *Nr2e3*^{-/-} zebrafish resembles that of *Nrl*^{-/-} mouse in that deletion of the gene ablated expression of rod genes and prevented formation of rod OSs. The physiological function of rods was not detected [24]. These data demonstrated that development of rods was prevented totally in *Nrl*^{-/-}

mouse. Accordingly, analogous to *Nrl*'s role in the murine retina, zebrafish *Nr2e3* acts like a “molecular switch” in modulating rod-specific genes. The critical role *Nrl* and *Nr2e3* each play in rod development in, respectively, mouse and zebrafish, may be reflected in the developmental order of expression of these two genes. In mouse, *Nrl* transcripts appear at ~E12.5 [62] and before *Nr2e3* at ~E16.5 [59], while in zebrafish the expression of *Nr2e3* is earlier than *Nrl* in rod lineage. *Nr2e3* appears first in rod progenitors in the INL and in all rod precursors in the ONL. *Nrl* transcripts appear only in the ONL [53]. Unlike the situation in mouse where *Nrl* can activate expression of *Nr2e3* [24], the earlier appearance of *Nr2e3* transcripts in zebrafish means that its expression is not activated by *Nrl*. ESCS patients present with severe deficit of rod function at an early stage and histological analysis of one postmortem retina revealed an absence of rods [26,27,57]. Furthermore, retinal structural analysis led to the conclusion that rods failed to differentiate in ESCS retinas [27]. During fetal development, *Nr2e3* expression follows *Nrl* very closely, and mutations in either *Nrl* or *Nr2e3* can lead to ESCS [63], suggesting these two genes may play parallel roles in rod development.

The phenotypes of ESCS, *Nr2e3*^{-/-} zebrafish, *Nrl*^{-/-} mouse and *rd7* mouse demonstrate that *Nr2e3* has a critical role for rod development in zebrafish and human. By contrast, it is *Nrl* that modulates rod development in mouse. However, *Nrl* alone is not sufficient to maintain rod survival in *rd7* mouse. Therefore, *Nr2e3* may play a role in rod maintenance.

There are two hypotheses that may account for the enhanced S-cones in ESCS and *rd7* mouse. One is that lack of *Nr2e3* function leads the plastic rod precursors to transdifferentiate into S-cones. Two lines of evidence support this hypothesis: 1) *Nr2e3* is predominantly expressed




Fig. 7. L-/M-cones is degenerated progressively in *Nr2e3*^{-/-} retinas. Confocal images stained for red opsin (A) and green opsin (B) at 1 month, 6 months and 12 months. The length of OSs of these two cones was shortened gradually from 1 month. Scale bars: 20 μm. (C and D) The length of L-/M-cones OSs was statistically analyzed at 1 month, 6 months and 12 months. WT, wildtype. Data are presented as mean ± SD. ***P* < 0.01, ****P* < 0.001.

in developing and mature rods and plays a dual role in promoting expression of rod genes and repressing cone genes [25]; 2) there are hybrid rod-cone photoreceptors expressing both rod and cone genes in the *rd7* mouse [36]. The second hypothesis is that losing *Nr2e3* causes aberrant proliferation of cone progenitors. Observations that *Nr2e3* is expressed in cone progenitors and mature cones, as well as the fact that there are no hybrid photoreceptors, support this hypothesis [5]. In zebrafish, we observed *Nr2e3* function only in rod lineage; cone development and proliferation of cone progenitors were not changed. Thus, our data contradict the second hypothesis. However, there are remaining questions about the first hypothesis. Rods account for 97% of the number of photoreceptors in the mouse retina, while the S-cones increased only 2-fold in the *rd7* retina [34]. This suggests only a small

fraction of rod precursors were transformed into S-cones. Because a fraction of rods were derived from the default S-cones in mouse [43], perhaps it was these rods that reverted back to S-cones when *Nrl/Nr2e3* was lost. The enhanced S-cones were actually the default S-cones. By contrast, UV- or S-cone is not the default photoreceptor in zebrafish [10,43] (and the current data). Thus, the number of these two cone types is not increased when rods were lost in *Nr2e3*^{-/-} zebrafish.

The mechanism that causes selective L-/M-cone photoreceptor degeneration remains unknown. In ESCS and *rd7*, arrangement of the ONL was disorganized and this disorganization was present before photoreceptor degeneration [35,64]. Disorganized arrangement of the ONL (especially nuclei of L-/M-cones) was also observed in *Nr2e3*^{-/-} zebrafish, which could have led to the degeneration. This disorganization




Fig. 8. Nr2e3 enhances rhodopsin gene expression but does not repress cone opsin genes. Reporter plasmids contain the promoter sequence of zebrafish *rhodopsin* (A), *UV-opsin* (B) and *S-opsin* (C) were transfected into HEK293 cells alone or combined with the indicated expression vectors of Nr2e3, Crx and Nrl. The relative activity of each promoter is shown. '+', '++' and '+++ ' represent 100, 200 and 400 ng of the indicated vector, respectively. To confirm the total amounts of DNA were constant between pores, empty plasmid was added. Three independent experiments were performed. The results are presented as mean \pm SD.

seemed to have little effect on UV- or S-cone survival. Analysis of a 77-year-old postmortem ESCS retina showed that the number of S-cones was maintained [30]. Besides, S-cones can survive normally to 5 months in *rd7* mouse and 31 weeks in *Nrl*^{-/-} mouse [24,35]. UV- and S-cones can survive to 10 months in *Nr2e3*^{-/-} zebrafish. It is possible that the interaction between rods and L-/M-cones is more critical than that with UV- or S-cones.

Our work has extended the knowledge of Nr2e3 and provides

interpretation of the phenotypes in ESCS and *rd7* mouse. In consideration of the complex phenotypes of ESCS and the *rd7* mouse, causality of the underlying mechanisms that involve multiple cell types is difficult to determine. Thus, it is impossible to identify therapeutic targets to save the cone photoreceptors, which are the key essential cells for useful human vision. By contrast, the *Nr2e3*-null zebrafish model provides a clear target for developing therapeutic intervention, i.e., survival of the L-/M-cone photoreceptors. This feature, in addition to the intrinsic advantages of the zebrafish model, may offer new opportunities in finding treatments for ESCS in particular and retinal degeneration in general.

Conflict of interest statement

The authors declare no conflict of interest.

Transparency document

The Transparency document associated with this article can be found, in online version.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (No. 31571303, 81500762, 81670890, 81870691, 31871260 and 81770102), Research Funding for the Doctoral Program of Higher Education of China (20120142110077).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bbadis.2019.01.022>.

References

- [1] T. Branchek, R. Bremiller, The development of photoreceptors in the zebrafish, *Brachydanio rerio*. I. Structure, *J. Comp. Neurol.* 224 (1984) 107–115.
- [2] K.D. Larison, R. Bremiller, Early onset of phenotype and cell patterning in the embryonic zebrafish retina, *Development* 109 (1990) 567–576.
- [3] P.A. Raymond, L.K. Barthel, M.E. Rounsifer, S.A. Sullivan, J.K. Knight, Expression of rod and cone visual pigments in goldfish and zebrafish: a rhodopsin-like gene is expressed in cones, *Neuron* 10 (1993) 1161–1174.
- [4] J. Robinson, E.A. Schmitt, F.I. Harosi, R.J. Reece, J.E. Dowling, Zebrafish ultraviolet visual pigment: absorption spectrum, sequence, and localization, *Proc. Natl. Acad. Sci. U. S. A.* 90 (1993) 6009–6012.
- [5] N.B. Haider, P. Demarco, A.M. Nystuen, X. Huang, R.S. Smith, M.A. Mccall, J.K. Naggert, P.M. Nishina, The transcription factor Nr2e3 functions in retinal progenitors to suppress cone cell generation, *Vis. Neurosci.* 23 (2006) 917–929.
- [6] Y. Yanagi, S. Takezawa, S. Kato, Distinct functions of photoreceptor cell-specific nuclear receptor, thyroid hormone receptor beta2 and CRX in one photoreceptor development, *Invest. Ophthalmol. Vis. Sci.* 43 (2002) 3489–3494.
- [7] M. Cayouette, B.A. Barres, M. Raff, Importance of intrinsic mechanisms in cell fate decisions in the developing rat retina, *Neuron* 40 (2003) 897–904.
- [8] F.J. Livesey, C.L. Cepko, Vertebrate neural cell-fate determination: lessons from the retina, *Nat. Rev. Neurosci.* 2 (2001) 109–118.
- [9] J.M. Fadool, Development of a rod photoreceptor mosaic revealed in transgenic zebrafish, *Dev. Biol.* 258 (2003) 277–290.
- [10] K. Alvarez-Delfin, A.C. Morris, C.D. Snelson, J.T. Gamse, T. Gupta, F.L. Marlow, M.C. Mullins, H.A. Burgess, M. Granato, J.M. Fadool, Tbx2b is required for ultraviolet photoreceptor cell specification during zebrafish retinal development, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 2023–2028.
- [11] C.L. Cepko, C.P. Austin, X. Yang, M. Alexiades, D. Ezzeddine, Cell fate determination in the vertebrate retina, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 589–595.
- [12] D.L. Turner, C.L. Cepko, A common progenitor for neurons and glia persists in rat retina late in development, *Nature* 328 (1987) 131–136.
- [13] C.L. Freund, C.Y. Gregory-Evans, T. Furukawa, M. Papaioannou, J. Looser, L. Ploder, J. Bellingham, D. Ng, J.A. Herbrick, A. Duncan, S.W. Scherer, L.C. Tsui, A. Loutradis-Anagnostou, S.G. Jacobson, C.L. Cepko, S.S. Bhattacharya, R.R. McInnes, Cone-rod dystrophy due to mutations in a novel photoreceptor-specific homeobox gene (CRX) essential for maintenance of the photoreceptor, *Cell* 91 (1997) 543–553.
- [14] T. Furukawa, E.M. Morrow, C.L. Cepko, Crx, a novel otx-like homeobox gene, shows photoreceptor-specific expression and regulates photoreceptor differentiation, *Cell* 91 (1997) 531–541.
- [15] Y.C. Shen, P.A. Raymond, Zebrafish cone-rod (crx) homeobox gene promotes

- retinogenesis, *Dev. Biol.* 269 (2004) 237–251.
- [16] T. Furukawa, E.M. Morrow, T. Li, F.C. Davis, C.L. Cepko, Retinopathy and attenuated circadian entrainment in *Crx*-deficient mice, *Nat. Genet.* 23 (1999) 466–470.
- [17] Y.Q. Zhang, Y.X. Wang, G.Z. Xu, H.Y. Song, W.J. Wang, Influence of *CRX* gene on development of photoreceptors in zebrafish, *Zhonghua Yan Ke Za Zhi* 44 (2008) 448–454.
- [18] P.K. Swain, S. Chen, Q.L. Wang, L.M. Affatigato, C.L. Coats, K.D. Brady, G.A. Fishman, S.G. Jacobson, A. Swaroop, E. Stone, P.A. Sieving, D.J. Zack, Mutations in the cone-rod homeobox gene are associated with the cone-rod dystrophy photoreceptor degeneration, *Neuron* 19 (1997) 1329–1336.
- [19] M.M. Sohocki, L.S. Sullivan, H.A. Mintz-Hittner, D. Birch, J.R. Heckenlively, C.L. Freund, R.R. McInnes, S.P. Daiger, A range of clinical phenotypes associated with mutations in *CRX*, a photoreceptor transcription-factor gene, *Am. J. Hum. Genet.* 63 (1998) 1307–1315.
- [20] C.L. Freund, Q.L. Wang, S. Chen, B.L. Muskat, C.D. Wiles, V.C. Sheffield, S.G. Jacobson, R.R. McInnes, D.J. Zack, E.M. Stone, De novo mutations in the *CRX* homeobox gene associated with Leber congenital amaurosis, *Nat. Genet.* 18 (1998) 311–312.
- [21] D.A. Bessant, A.M. Payne, K.P. Mitton, Q.L. Wang, P.K. Swain, C. Plant, A.C. Bird, D.J. Zack, A. Swaroop, S.S. Bhattacharya, A mutation in *NRL* is associated with autosomal dominant retinitis pigmentosa, *Nat. Genet.* 21 (1999) 355–356.
- [22] K.M. Nishiguchi, J.S. Friedman, M.A. Sandberg, A. Swaroop, E.L. Berson, T.P. Dryja, Recessive *NRL* mutations in patients with clumped pigmentary retinal degeneration and relative preservation of blue cone function, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 17819–17824.
- [23] K.W. Littink, P. Stappers, F. Riemsdag, H.E. Talsma, M.M. van Genderen, F. Cremers, R. Collin, L.I. van den Born, Autosomal recessive *NRL* mutations in patients with enhanced S-cone syndrome, *Genes (Basel)* 9 (2018).
- [24] A.J. Mears, M. Kondo, P.K. Swain, Y. Takada, R.A. Bush, T.L. Saunders, P.A. Sieving, A. Swaroop, *Nrl* is required for rod photoreceptor development, *Nat. Genet.* 29 (2001) 447–452.
- [25] J. Chen, A. Rattner, J. Nathans, The rod photoreceptor-specific nuclear receptor *Nr2e3* represses transcription of multiple cone-specific genes, *J. Neurosci.* 25 (2005) 118–129.
- [26] G. Peng, O. Ahmad, F. Ahmad, J. Liu, S. Chen, The photoreceptor-specific nuclear receptor *Nr2e3* interacts with *Crx* and exerts opposing effects on the transcription of rod versus cone genes, *Hum. Mol. Genet.* 14 (2005) 747–764.
- [27] N.B. Haider, S.G. Jacobson, A.V. Cideciyan, R. Swiderski, L.M. Streb, C. Seaby, G. Beck, R. Hockey, D.B. Hanna, S. Gorman, D. Duhl, R. Carmi, J. Bennett, R.G. Weleber, G.A. Fishman, A.F. Wright, E.M. Stone, V.C. Sheffield, Mutation of a nuclear receptor gene, *NR2E3*, causes enhanced S cone syndrome, a disorder of retinal cell fate, *Nat. Genet.* 24 (2000) 127–131.
- [28] D.C. Hood, A.V. Cideciyan, A.J. Roman, S.G. Jacobson, Enhanced S cone syndrome: evidence for an abnormally large number of S cones, *Vis. Res.* 35 (1995) 1473–1481.
- [29] A.F. Wright, A.C. Reddick, S.B. Schwartz, J.S. Ferguson, T.S. Aleman, U. Kellner, B. Jurklics, A. Schuster, E. Zrenner, B. Wissinger, A. Lennon, X. Shu, A.V. Cideciyan, E.M. Stone, S.G. Jacobson, A. Swaroop, Mutation analysis of *NR2E3* and *NRL* genes in Enhanced S cone Syndrome, *Hum. Mutat.* 24 (2004) 439.
- [30] A.H. Milam, L. Rose, A.V. Cideciyan, M.R. Barakat, W.X. Tang, N. Gupta, T.S. Aleman, A.F. Wright, E.M. Stone, V.C. Sheffield, S.G. Jacobson, The nuclear receptor *NR2E3* plays a role in human retinal photoreceptor differentiation and degeneration, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 473–478.
- [31] D. Sharon, M.A. Sandberg, R.C. Caruso, E.L. Berson, T.P. Dryja, Shared mutations in *NR2E3* in enhanced S-cone syndrome, Goldmann-Favre syndrome, and many cases of clumped pigmentary retinal degeneration, *Arch. Ophthalmol.* 121 (2003) 1316–1323.
- [32] P. Escher, H.V. Tran, V. Vaclavik, F.X. Borrut, D.F. Schorderet, F.L. Munier, Double concentric autofluorescence ring in *NR2E3*-p.G56R-linked autosomal dominant retinitis pigmentosa, *Invest. Ophthalmol. Vis. Sci.* 53 (2012) 4754–4764.
- [33] S. Gerber, J.M. Rozet, S.I. Takezawa, S.L. Doss, L. Lopes, O. Gribouval, C. Penet, I. Perrault, D. Ducrocq, E. Souied, M. Jeanpierre, S. Romana, J. Frezal, F. Ferraz, R. Yu-Umesono, A. Munnich, J. Kaplan, The photoreceptor cell-specific nuclear receptor gene (*PNR*) accounts for retinitis pigmentosa in the Crypto-Jews from Portugal (Marranos), survivors from the Spanish Inquisition, *Hum. Genet.* 107 (2000) 276–284.
- [34] N.B. Haider, J.K. Naggert, P.M. Nishina, Excess cone cell proliferation due to lack of a functional *NR2E3* causes retinal dysplasia and degeneration in *rd7/rd7* mice, *Hum. Mol. Genet.* 10 (2001) 1619–1626.
- [35] N.B. Akhmedov, N.I. Piriev, B. Chang, A.L. Rapoport, N.L. Hawes, P.M. Nishina, S. Nusinowitz, J.R. Heckenlively, T.H. Roderick, C.A. Kozak, M. Danciger, M.T. Davissou, D.B. Farber, A deletion in a photoreceptor-specific nuclear receptor mRNA causes retinal degeneration in the *rd7* mouse, *Proc. Natl. Acad. Sci. U. S. A.* 97 (2000) 5551–5556.
- [36] J.C. Corbo, C.L. Cepko, A hybrid photoreceptor expressing both rod and cone genes in a mouse model of enhanced S-cone syndrome, *PLoS Genet.* 1 (2005) e11.
- [37] J. Zhu, C. Ming, X. Fu, Y. Duan, D.A. Hoang, J. Rutgard, R. Zhang, W. Wang, R. Hou, D. Zhang, E. Zhang, C. Zhang, X. Hao, W. Xiong, K. Zhang, Gene and mutation independent therapy via CRISPR-Cas9 mediated cellular reprogramming in rod photoreceptors, *Cell Res.* 27 (2017) 830–833.
- [38] W. Yu, S. Mookherjee, V. Chaitankar, S. Hiriyanna, J.W. Kim, M. Brooks, Y. Ataeijannati, X. Sun, L. Dong, T. Li, A. Swaroop, Z. Wu, *Nrl* knockdown by AAV-delivered CRISPR/Cas9 prevents retinal degeneration in mice, *Nat. Commun.* 8 (2017) 14716.
- [39] P.A. Nakamura, S. Tang, A.A. Shimchuk, S. Ding, T.A. Reh, Potential of small molecule-mediated reprogramming of rod photoreceptors to treat retinitis pigmentosa, *Invest. Ophthalmol. Vis. Sci.* 57 (2016) 6407–6415.
- [40] P.A. Nakamura, A.A. Shimchuk, S. Tang, Z. Wang, K. DeGolier, S. Ding, T.A. Reh, Small molecule Photoregulin3 prevents retinal degeneration in the *Rho*(P23H) mouse model of retinitis pigmentosa, *Elife* 6 (2017).
- [41] G.J. Lieschke, P.D. Currie, Animal models of human disease: zebrafish swim into view, *Nat. Rev. Genet.* 8 (2007) 353–367.
- [42] D.L. Stenkamp, The rod photoreceptor lineage of teleost fish, *Prog. Retin. Eye Res.* 30 (2011) 395–404.
- [43] J.W. Kim, H.J. Yang, A.P. Oel, M.J. Brooks, L. Jia, D.C. Plachetzki, W. Li, W.T. Allison, A. Swaroop, Recruitment of rod photoreceptors from short-wavelength-sensitive cones during the evolution of nocturnal vision in mammals, *Dev. Cell* 37 (2016) 520–532.
- [44] S.S. Kitambi, G. Hauptmann, The zebrafish orphan nuclear receptor genes *nr2e1* and *nr2e3* are expressed in developing eye and forebrain, *Gene Expr. Patterns* 7 (2007) 521–528.
- [45] L.E. Jao, S.R. Wente, W. Chen, Efficient multiplex biallelic zebrafish genome editing using a CRISPR nuclease system, *Proc. Natl. Acad. Sci. U. S. A.* 110 (2013) 13904–13909.
- [46] S. Yu, C. Li, L. Biswas, X. Hu, F. Liu, J. Reilly, X. Liu, Y. Liu, Y. Huang, Z. Lu, S. Han, L. Wang, L.J. Yu, T. Jiang, X. Shu, F. Wong, Z. Tang, M. Liu, *CERKL* gene knockout disturbs photoreceptor outer segment phagocytosis and causes rod-cone dystrophy in zebrafish, *Hum. Mol. Genet.* 26 (2017) 2335–2345.
- [47] Z. Lu, X. Hu, F. Liu, D.C. Soares, X. Liu, S. Yu, M. Gao, S. Han, Y. Qin, C. Li, T. Jiang, D. Luo, A.Y. Guo, Z. Tang, M. Liu, Ablation of *EYS* in zebrafish causes mislocalisation of outer segment proteins, F-actin disruption and cone-rod dystrophy, *Sci. Rep.* 7 (2017) 46098.
- [48] M. Gao, Y. Huang, L. Wang, M. Huang, F. Liu, S. Liao, S. Yu, Z. Lu, S. Han, X. Hu, Z. Qu, X. Liu, Y.T. Assefa, L. Yang, Z. Tang, D.W. Li, M. Liu, *HSF4* regulates lens fiber cell differentiation by activating *p53* and its downstream regulators, *Cell Death Dis.* 8 (2017) e3082.
- [49] F. Liu, J. Chen, S. Yu, R.K. Raghupathy, X. Liu, Y. Qin, C. Li, M. Huang, S. Liao, J. Wang, J. Zou, X. Shu, Z. Tang, M. Liu, Knockout of *RP2* decreases *GRK1* and rod transducin subunits and leads to photoreceptor degeneration in zebrafish, *Hum. Mol. Genet.* 24 (2015) 4648–4659.
- [50] X. Cui, L. Wang, J. Zhang, R. Du, S. Liao, D. Li, C. Li, T. Ke, D.W. Li, H. Huang, Z. Yin, Z. Tang, M. Liu, *HSF4* regulates *DLAD* expression and promotes lens degeneration, *Biochim. Biophys. Acta* 1832 (2013) 1167–1172.
- [51] C. Li, L. Wang, J. Zhang, M. Huang, F. Wong, X. Liu, F. Liu, X. Cui, G. Yang, J. Chen, Y. Liu, J. Wang, S. Liao, M. Gao, X. Hu, X. Shu, Q. Wang, Z. Yin, Z. Tang, M. Liu, *CERKL* interacts with mitochondrial *TRX2* and protects retinal cells from oxidative stress-induced apoptosis, *Biochim. Biophys. Acta* 1842 (2014) 1121–1129.
- [52] F. Liu, Y. Qin, S. Yu, D.C. Soares, L. Yang, J. Weng, C. Li, M. Gao, Z. Lu, X. Hu, X. Liu, T. Jiang, J.Y. Liu, X. Shu, Z. Tang, M. Liu, Pathogenic mutations in retinitis pigmentosa 2 predominantly result in loss of *RP2* protein stability in humans and zebrafish, *J. Biol. Chem.* 292 (2017) 6225–6239.
- [53] S.M. Nelson, R.A. Frey, S.L. Wardwell, D.L. Stenkamp, The developmental sequence of gene expression within the rod photoreceptor lineage in embryonic zebrafish, *Dev. Dyn.* 237 (2008) 2903–2917.
- [54] R.L. Bernardos, L.K. Barthel, J.R. Meyers, P.A. Raymond, Late-stage neuronal progenitors in the retina are radial Muller glia that function as retinal stem cells, *J. Neurosci.* 27 (2007) 7028–7040.
- [55] A.C. Morris, T.L. Scholz, S.E. Brockerhoff, J.M. Fadool, Genetic dissection reveals two separate pathways for rod and cone regeneration in the teleost retina, *Dev. Neurobiol.* 68 (2008) 605–619.
- [56] M. Sotolongo-Lopez, K. Alvarez-Delfin, C.J. Saade, D.L. Vera, J.M. Fadool, Genetic dissection of dual roles for the transcription factor *six7* in photoreceptor development and patterning in zebrafish, *PLoS Genet.* 12 (2016) e1005968.
- [57] D.C. Otteson, P.F. Hitchcock, Stem cells in the teleost retina: persistent neurogenesis and injury-induced regeneration, *Vis. Res.* 43 (2003) 927–936.
- [58] G.H. Peng, S. Chen, Chromatin immunoprecipitation identifies photoreceptor transcription factor targets in mouse models of retinal degeneration: new findings and challenges, *Vis. Neurosci.* 22 (2005) 575–586.
- [59] H. Cheng, H. Khanna, E.C. Oh, D. Hicks, K.P. Mitton, A. Swaroop, Photoreceptor-specific nuclear receptor *NR2E3* functions as a transcriptional activator in rod photoreceptors, *Hum. Mol. Genet.* 13 (2004) 1563–1575.
- [60] N.B. Haider, N. Mollema, M. Gaule, Y. Yuan, A.J. Sachs, A.M. Nystuen, J.K. Naggert, P.M. Nishina, *Nr2e3*-directed transcriptional regulation of genes involved in photoreceptor development and cell-type specific phototransduction, *Exp. Eye Res.* 89 (2009) 365–372.
- [61] S. Ueno, M. Kondo, K. Miyata, T. Hirai, T. Miyata, J. Usukura, Y. Nishizawa, Y. Miyake, Physiological function of S-cone system is not enhanced in *rd7* mice, *Exp. Eye Res.* 81 (2005) 751–758.
- [62] M. Akimoto, H. Cheng, D. Zhu, J.A. Brzezinski, R. Khanna, E. Filippova, E.C. Oh, Y. Jing, J.L. Linares, M. Brooks, S. Zarepari, A.J. Mears, A. Hero, T. Glaser, A. Swaroop, Targeting of GFP to newborn rods by *Nrl* promoter and temporal expression profiling of flow-sorted photoreceptors, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 3890–3895.
- [63] O.K. Bumsted, H. Cheng, Y. Jiang, D. Schulte, A. Swaroop, A.E. Hendrickson, Expression of photoreceptor-specific nuclear receptor *NR2E3* in rod photoreceptors of fetal human retina, *Invest. Ophthalmol. Vis. Sci.* 45 (2004) 2807–2812.
- [64] S.G. Jacobson, A. Sumaroka, T.S. Aleman, A.V. Cideciyan, S.B. Schwartz, A.J. Roman, R.R. McInnes, V.C. Sheffield, E.M. Stone, A. Swaroop, A.F. Wright, Nuclear receptor *NR2E3* gene mutations distort human retinal laminar architecture and cause an unusual degeneration, *Hum. Mol. Genet.* 13 (2004) 1893–1902.