



Change in study randomization allocation needs to be included in statistical analysis: comment on ‘Randomized controlled trial of weight loss versus usual care on telomere length in women with breast cancer: the lifestyle, exercise, and nutrition (LEAN) study’

Stephanie L. Dickinson¹ · Lilian Golzarri-Arroyo¹ · Andrew W. Brown² · Bryan McComb³ · Chanaka N. Kahathuduwa⁴ · David B. Allison¹

Received: 23 January 2019 / Accepted: 29 January 2019 / Published online: 5 February 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Data are often combined across multiple studies, sites, strata or phases of data collection, for a variety of reasons. In a randomized controlled trial (RCT), employing proper methods when combining data collected in separate contexts ensures unbiased estimates of the combined treatment effect. Collapsing (or ‘lumping’) data across studies or strata without statistical adjustment can provide misleading results [1], such as occurs in Simpson’s paradox where treatment effects that are consistent across each strata separately are reversed when data are collapsed [2–4]. This paradox occurs specifically when there are differences between the two or more strata (or studies) in the ratio of people in each treatment group [3]. Altman wrote recently of dangers of bias in combining data across studies with varied randomization allocation ratios [5].

Sanft et al. report results from data analysis combining data from an earlier RCT with additional data collected at a later time point, under a different study design, without accounting for these two phases of study in the description of their statistical analysis [6, 7]. The primary study was an RCT with three arms comparing in-person counseling,

telephone counseling, and usual care on weight loss, with an equal 1:1:1 randomization allocation [7]. The second phase of the study randomized participants into only two arms: counseling intervention or usual care, with an ostensibly similar 1:1 equal allocation [6]. However, because the two interventions in the primary study were grouped together for analysis [6], the allocation ratio was effectively 2:1 for intervention in the primary study but changed to 1:1 in the later study. This change in allocation ratio can lead to bias [3, 5].

Because biased estimation of treatment effects can result from differences between the two periods, data should be compared and reported between participants in the two periods, by treatment group. For example, baseline BMI may differ between the two recruitment periods in Sanft et al., where significant differences between treatment groups are reported in the combined data [6] but not the primary data [7]. More critically, it is essential that statistical analyses should be adjusted by including study period as a stratification or blocking variable and testing for interactions [1–5, 8]. Bangdiwala et al. discuss multiple options for pooling data across heterogeneous studies [8].

While the overall conclusions in the Sanft et al. paper may remain unchanged, this is an important methodological issue for researchers to avoid bias in analyses combining data across multiple strata, sites or phases of data collection in RCTs. We encourage Sanft and colleagues to consider re-analyzing their data taking these factors into account and publish corrected results, and we offer to assist in the re-analysis if needed.

Funding This work was funded in part by the National Institute of Health (NIH): R25HL124208, R25DK099080, P30AG050886 and

✉ Stephanie L. Dickinson
sd3@indiana.edu

¹ Department of Epidemiology and Biostatistics, Indiana University School of Public Health-Bloomington, 1025 E. 7th St, Bloomington, IN 47408, USA

² Department of Applied Health Science, Indiana University School of Public Health-Bloomington, Bloomington, IN, USA

³ Division of Biostatistics, Department of Population Health, New York University Langone Health, New York, NY, USA

⁴ Department of Human Development and Family Studies, Texas Tech University, Lubbock, TX, USA

U24AG056053. The opinions expressed are those of the authors and not necessarily of the NIH or any other organization.

Compliance with ethical standards

Conflict of interest In the last 12 months, Dr. Allison has received personal payments or promises for same from: American Society for Nutrition, American Statistical Association, Biofortis, Columbia University, Fish and Richardson, P.C., Frontiers Publishing, Henry Stewart Talks, IKEA, Indiana University, Laura and John Arnold Foundation, Johns Hopkins University, Law Offices of Ronald Marron, MD Anderson Cancer Center, Medical College of Wisconsin, National Institutes of Health (NIH), Sage Publishing, The Obesity Society, Tomasik, Kotin and Kasserman LLC, University of Alabama at Birmingham, University of Miami, Nestle, WW (formerly Weight Watchers International, LLC). Donations to a foundation have been made on his behalf by the Northarvest Bean Growers Association. Dr. Allison is an Unpaid Member of the International Life Sciences Institute North America Board of Trustees. Dr. Allison's institution, Indiana University, has received funds to support his research or educational activities from: NIH, Alliance for Potato Research and Education, American Federation for Aging Research, Dairy Management, Inc., Herbalife, Laura and John Arnold Foundation, Oxford University Press. In the last 12 months, Dr. Brown has received personal payments or paid travel from: American Society for Nutrition, Indiana University, Kentuckiana Health Collaborative, Rippe Lifestyle Institute, Inc. Dr. Brown's institution, Indiana University, has received funds to support his research or educational activities from: American Federation for Aging Research, Dairy Management, Inc., NIH, Oxford University Press, University of Alabama at Birmingham. The other authors declare that they have no disclosures.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

1. DeMets DL (1987) Methods for combining randomized clinical trials: strengths and limitations. *Stat Med* 6(3):341–348
2. Simpson EH (1951) The interpretation of interaction in contingency tables. *J R Stat Soc B* 1:238–241
3. Blyth CR (1972) On Simpson's paradox and the sure-thing principle. *J Am Stat Assoc* 67(338):364–366
4. Norton HJ, Divine G. (2015) Simpson's paradox... and how to avoid it. *Significance* 12(4):40–43
5. Altman DG (2018) Avoiding bias in trials in which allocation ratio is varied. *J R Soc Med* 111(4):143–144
6. Sanft T, Usiskin I, Harrigan M, Cartmel B, Lu L, Li FY, Zhou Y, Chagpar A, Ferrucci LM, Puztai L, Irwin ML (2018) Randomized controlled trial of weight loss versus usual care on telomere length in women with breast cancer: the lifestyle, exercise, and nutrition (LEAN) study. *Breast Cancer Res Treat* 172(1):105–112
7. Harrigan M, Cartmel B, Loftfield E, Sanft T, Chagpar AB, Zhou Y, Playdon M, Li F, Irwin ML (2016) Randomized trial comparing telephone versus in-person weight loss counseling on body composition and circulating biomarkers in women treated for breast cancer: the lifestyle, exercise, and nutrition (LEAN) study. *J Clin Oncol* 34(7):669
8. Bangdiwala SI, Bhargava A, O'Connor DP, Robinson TN, Michie S, Murray DM, Stevens J, Belle SH, Templin TN, Pratt CA (2016) Statistical methodologies to pool across multiple intervention studies. *Transl Behav Med* 6(2):228–235

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.