



## Letter to the Editor

## Raynaud's phenomenon and inflammatory bowel disease: The possible role of microcirculation



## ARTICLE INFO

## Keywords:

Raynaud's phenomenon  
Inflammatory bowel disease  
Ulcerative colitis  
Crohn's disease  
Nailfold capillaroscopy

Vascular pathology is increasingly more recognized as an extra-intestinal manifestation of Inflammatory Bowel Disease (IBD) [1,2]. Raynaud's phenomenon (RP), a syndrome characterized by recurrent episodes of digital artery vasospasm, may be secondary to systemic diseases and is rarely associated with IBD [3,4]. RP occurs in the presence of abnormalities of dermal capillaries, the morphology of which can be examined by nailfold capillaroscopy (NFC), a standardized non-invasive technique [5].

The present study aimed to determine the impact of IBD on the microcirculation. After informed consent, consecutively followed IBD patients were prospectively screened for the presence of RP. NFC was performed with VideoCap® video-capillaroscope, with high magnification lens (x200). According to Cutolo *et al*, we considered significant abnormalities enlarged and giant capillary (loops larger than 20 μm and 50 μm, respectively), microhaemorrhages (defined as dark mass due to haemosiderin deposit), capillary loss (reduction of the number of capillaries to below the normal range), major dystrophies (ramified branched capillary, neoangiogenesis) and minor dystrophies (tortuosities, corkscrew, meandering or bushy loops), over 33% of all capillary bed [6]. Chi Square Test was used for statistical analysis ( $p < 0,05$ ).

We screened 5 patients with ulcerative colitis and 11 with Crohn's disease; overall 12 were female (75%) with a mean age of  $50 \pm 13$  years. IBD patients with RP (IBDRP+,  $n = 5$ ) presented a larger proportion of minor dystrophies and giant capillaries when compared to those without RP (IBDRP-,  $n = 11$ ). In the IBDRP+ group, 20% ( $n = 1$ ) of patients fulfilled criteria for an early scleroderma pattern and 40% ( $n = 2$ ) revealed a non-scleroderma pattern. Unspecific NFC changes were present in 45% ( $n = 5$ ) patients belonging to the IBDRP- group. The most relevant difference between both groups was the presence of persistent inflammatory activity (determined by colonoscopy in the previous 6 months) accompanied by erythrocyte sedimentation rate (ESR) elevation in 40% ( $n = 2$ ) of IBDRP+ patients in contrast to 100% remission in the IBDRP- group. No significant correlation could be established between capillaroscopic findings between IBDRP+ and IBDRP- groups ( $p = 0,197$ ). The groups did not differ as regards immunosuppressive treatment, beta-blocking agents, smoking or contraceptive use. Extra-intestinal known manifestations included

uveitis ( $n = 1$ ), psoriasis ( $n = 4$ ) and spondyloarthritis ( $n = 4$ ).

In our exploratory study, we found nailfold capillaroscopic changes in 25% ( $n = 4$ ) IBD patients. In addition, 38% ( $n = 6$ ) of patients presented unspecific changes, justifying follow up NFC evaluation.

The association between IBD activity, RP and microcirculatory changes has not yet been described and merits further study, as a common immunological mechanism may be responsible for both vascular and enterocytic damage.

## Declarations of interest: none.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## References

- [1] Koutroubaki IE, Sfiridaki A, Tsiolakidou G, et al. Genetic risk factors in patients with inflammatory bowel disease and vascular complications: case-control study. *Inflamm Bowel Dis* 2007;13:410–5. <https://doi.org/10.1002/ibd.20076>.
- [2] Tan V, Chung A, Yan BP, et al. Venous and arterial disease in inflammatory bowel disease. *J Gastroenterol Hepatol* 2013;28:1095–113. <https://doi.org/10.1111/jgh.12260>.
- [3] Prete M, Fatone MC, Favoino E, et al. Raynaud's phenomenon: from molecular pathogenesis to therapy. *Autoimmun Rev* 2014;13(6):655–67. <https://doi.org/10.1016/j.autrev.2013.12.001>.
- [4] Karmiris K, Avgerinos A, Tavernakari A, et al. Prevalence and characteristics of extra-intestinal manifestations in a large cohort of greek patients with inflammatory bowel disease. *J Crohns Colitis* 2016;10(4):429–36. <https://doi.org/10.1093/ecco-jcc/jjv232>.
- [5] Cutolo M, Sulli A, Smith V. How to perform and interpret capillaroscopy. *Best Pract Res Clin Rheumatol* 2013;27:237–48. <https://doi.org/10.1016/j.berh.2013.03.001>.
- [6] Cutolo M, et al. *Atlas of Capillaroscopy in Rheumatic Diseases*. Elsevier; 2010. (eBook ISBN: 9788821433917).

Verara R. Bernardino\*, Ana Catarina Rodrigues, António Panarra  
Unidade de Doenças Auto-Imunes, Internal Medicine Department 7.2,  
Hospital Curry Cabral, Centro Hospitalar Universitário Lisboa Central,  
Lisboa, Portugal  
E-mail address: [verarodriguesb@gmail.com](mailto:verarodriguesb@gmail.com) (V.R. Bernardino).

<https://doi.org/10.1016/j.ejim.2019.02.001>

Received 9 December 2018; Accepted 1 February 2019

Available online 06 February 2019

0953-6205/ © 2019 European Federation of Internal Medicine. Published by Elsevier B.V. All rights reserved.