

References

- [1] Ascani S, Zinzani PL, Gherlinzoni F, et al. Peripheral T-cell lymphomas. Clinicopathologic study of 168 cases diagnosed according to the R.E.A.L. classification. *Ann Oncol* 1997;8(6):583–92.
- [2] Armitage JO, Greer JP, Levine AM, et al. Peripheral T-cell lymphoma. *Cancer* 1989;63(1):158–63.
- [3] Vose J, Armitage J, Weisenburger D, Project IT-CL. International peripheral T-cell and natural killer/T-cell lymphoma study: pathology findings and clinical outcomes. *J Clin Oncol* 2008;26(25):4124–30.
- [4] Weisenburger DD, Savage KJ, Harris NL, et al. Peripheral T-cell lymphoma, not otherwise specified: a report of 340 cases from the International Peripheral T-cell Lymphoma Project. *Blood* 2011;117(12):3402–8.
- [5] Swerdlow SH, Campo E, Harris NL, et al. World Health Organization classification of tumours of haematopoietic and lymphoid tissues. Lyon: IARC Press; 2008.
- [6] Weiss LM, Crabtree GS, Rouse RV, Warnke RA. Morphologic and immunologic characterization of 50 peripheral T-cell lymphomas. *Am J Pathol* 1985;118(2):316–24.
- [7] Schmitz N, Trümper L, Ziepert M, et al. Treatment and prognosis of mature T-cell and NK-cell lymphoma: an analysis of patients with T-cell lymphoma treated in studies of the German High-Grade Non-Hodgkin Lymphoma Study Group. *Blood* 2010;116(18):3418–25.
- [8] Escalón MP, Liu NS, Yang Y, et al. Prognostic factors and treatment of patients with T-cell non-Hodgkin lymphoma: the M.D. Anderson Cancer Center experience. *Cancer* 2005;103(10):2091–8.
- [9] Numata A, Miyamoto T, Ohno Y, et al. Long-term outcomes of autologous PBSCT for peripheral T-cell lymphoma: retrospective analysis of the experience of the Fukuoka BMT group. *Bone Marrow Transplant* 2010;45(2):311–6.
- [10] Sonnen R, Schmidt WP, Müller-Hermelink HK, Schmitz N. The International Prognostic Index determines the outcome of patients with nodal mature T-cell lymphomas. *Br J Haematol* 2005;129(3):366–72.

Timothy S. Krill*

Jenine N. Zaibaq

The University of Texas Medical Branch, Department of Gastroenterology and Hepatology, United States

Harshwardhan Thaker

The University of Texas Medical Branch, Department of Pathology, United States

Shehzad Merwat

The University of Texas Medical Branch, Department of Gastroenterology and Hepatology, United States

*Corresponding author at: The University of Texas Medical Branch, Department of Gastroenterology and Hepatology, 301 University Blvd., Galveston, TX 77555-0570, United States.

E-mail address: tskrill@utmb.edu

(T.S. Krill)

24 February 2019

27 February 2019

<https://doi.org/10.1016/j.dld.2019.02.016>