



## Correction to: Compare two surgical interventions for otitis media with effusion in young children

Jinsheng Hao<sup>1,2,3</sup> · Min Chen<sup>1,2,3</sup> · Bing Liu<sup>1,2,3</sup> · Yang Yang<sup>1,2,3</sup> · Wei Liu<sup>1,2,3</sup> · Ning Ma<sup>1,2,3</sup> · Yang Han<sup>1,2,3</sup> · Qiaoyin Liu<sup>1,2,3</sup> · Xin Ni<sup>1,2,3</sup> · Jie Zhang<sup>1,2,3</sup>

Published online: 18 June 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

**Correction to: European Archives of Oto-Rhino-Laryngology**  
<https://doi.org/10.1007/s00405-019-05421-9>

In the original publication, Tables 5 and 6 were interchanged by mistake. The correct Tables 5 and 6 are given below:

The original article has been corrected.

---

The original article can be found online at <https://doi.org/10.1007/s00405-019-05421-9>.

---

✉ Xin Ni  
nixin@bch.com.cn

✉ Jie Zhang  
stzhangj@263.net

<sup>1</sup> National Center for Children's Health, Beijing,  
People's Republic of China

<sup>2</sup> Department of Otolaryngology, Head and Neck Surgery,  
Beijing Children's Hospital, Capital Medical University,  
Beijing 100045, People's Republic of China

<sup>3</sup> Beijing Key Laboratory for Pediatric Diseases  
of Otolaryngology, Head and Neck Surgery, Beijing Pediatric  
Research Institute, Beijing Children's Hospital, Capital  
Medical University, Beijing, People's Republic of China

**Table 5** Univariate analysis of factors affecting the recurrence of OME

| Factors   | Recurrence (23 cases) | Non-recurrence (161 cases) | <i>T</i> or $\chi^2$ | <i>P</i> |
|---|-----------------------|----------------------------|----------------------|----------|
| Gender (male/female)                                  | 16/7                  | 102/59                     | 0.338                | 0.561    |
| Course of disease (months)                            | 15.43 ± 11.24         | 13.49 ± 9.91               | 1.871                | 0.063    |
| Adjuvant adenoidectomy [ <i>n</i> (%)]                | 6 (26.09)             | 92 (57.14)                 | 7.801                | 0.005    |
| Allergic rhinitis [ <i>n</i> (%)]                     | 18 (78.26)            | 90 (55.90)                 | 4.150                | 0.042    |
| Recurrent respiratory tract infection [ <i>n</i> (%)] | 10 (43.48)            | 27 (16.77)                 | 8.936                | 0.003    |
| Tube retention time < 12 months [ <i>n</i> (%)]       | 16 (69.57)            | 68 (42.24)                 | 6.058                | 0.014    |
| With history of tube placed                           | 3 (13.04)             | 20 (12.42)                 | 0.007                | 0.933    |

**Table 6** Multivariate analysis of factors affecting the recurrence of OME

| Factors                               | <i>B</i> | Wald   | <i>P</i> | OR    | 95% CI      |
|---------------------------------------|----------|--------|----------|-------|-------------|
| Adjuvant adenoidectomy                | − 1.145  | 8.709  | 0.003    | 3.142 | 1.469–6.723 |
| Allergic rhinitis                     | 0.172    | 1.185  | 0.277    | 1.188 | 0.871–1.619 |
| Recurrent respiratory tract infection | 1.089    | 10.113 | 0.002    | 2.987 | 1.515–5.334 |
| Retention time of tube < 12 months    | 0.812    | 6.648  | 0.010    | 2.245 | 1.207–4.093 |

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.