



Nicotine protects against manganese and iron-induced toxicity in SH-SY5Y cells: Implication for Parkinson's disease

Bruk Getachew^a, Antonei B. Csoka^b, Michael Aschner^c, Yousef Tizabi^{a,*}

^a Department of Pharmacology, Howard University College of Medicine, Washington, DC, USA

^b Department of Anatomy, Howard University College of Medicine, Washington, DC, USA

^c Department of Molecular Pharmacology, Albert Einstein College of Medicine, Bronx, NY, USA

ARTICLE INFO

Keywords:

Nicotine
Manganese
Iron
Nicotinic receptors
Neurotoxicity
Neuroprotection
Cell culture
Parkinson's disease

ABSTRACT

Manganese (Mn) and iron (Fe) are trace elements that are essential for proper growth and physiological functions as both play critical role in a variety of enzymatic reactions. At high concentrations, however, they can be toxic and cause neurodegenerative disorders, particularly Parkinson-like syndromes. Nicotine, on the other hand, has been shown to have neuroprotective effects against various endogenous or exogenous toxins that selectively damage the dopaminergic cells. These cells include neuroblastoma-derived SH-SY5Y cells which express significant dopaminergic activity. However, practically no information on possible neuroprotective effects of nicotine against toxicity induced by trace elements is available. Therefore, in this study we investigated the effects of nicotine on toxicity induced by manganese or iron in these cells. Exposure of SH-SY5Y cells for 24 h to manganese (20 μ M) or iron (20 μ M) resulted in approximately 30% and 35% toxicity, respectively. Pretreatment with nicotine (1 μ M) completely blocked the toxicities of Mn and Fe. The effects of nicotine, in turn, were blocked by selective nicotinic receptor antagonists. Thus, dihydro-beta erythroidine (DHBE), a selective alpha 4-beta 2 subtype antagonist and methyllycaconitine (MLA), a selective alpha7 antagonist, as well as mecamylamine, a non-selective nicotinic antagonist all dose-dependently blocked the protective effects of nicotine against both Mn and Fe. These findings provide further support for the potential utility of nicotine or nicotinic agonists in Parkinson's disease-like neurodegenerative disorders, including those that might be precipitated by trace elements, such as Fe and Mn. Moreover, both alpha4-beta2 and alpha7 nicotinic receptor subtypes appear to mediate the neuroprotective effects of nicotine against toxicity induced by these two trace metals.

1. Introduction

Parkinson's disease (PD), associated with loss of dopaminergic neurons in the substantia nigra pars compacta (SNpc), is the second most common progressive neurodegenerative disorder following Alzheimer's disease (Cacabelos, 2017). Main symptoms include motor deficits characterized by akinesia, rigidity, resting tremor and postural instability (Sveinbjornsdottir, 2016). Although its exact etiology remains elusive, involvement of various neurotoxicological agents as well as excess accumulation of trace elements such as manganese and iron have been suggested (Tanner et al., 2011; Wang et al., 2016; Andrade et al., 2017; Vaccari et al., 2017). The most common treatment consists of dopamine replacement (e.g. levodopa = L-DOPA), which not only loses its full efficacy in a few years but may also induce severe dyskinesia (Picconi et al., 2017). Hence more studies on etiology as well as

development of more efficacious interventions aiming at neuroprotection are urgently needed.

Manganese (Mn), an activator or cofactor for a variety of metalloenzymes, is a transition metal that is essential for normal cell growth and development (Yousefi Babadi et al., 2014; Caito and Aschner, 2015; O'Neal and Zheng, 2015; Peres et al., 2016; Andrade et al., 2017). The enzyme or co-enzymes utilizing Mn play critical role in important biological functions such as gluconeogenesis, suppression of oxidative stress (Mn-superoxide dismutase, SOD) and conversion of glutamate into glutamine (glutamine synthetase) (Aschner and Gannon, 1994; Burton and Guilarte, 2009). Thus, Mn deficiency may lead to impaired reproductive function, retarded growth including skeletal abnormalities and seizures (Roth and Garrick, 2003; Aschner and Aschner, 2005). Excessive exposure to Mn, on the other hand, may also cause impaired reproductive function (McMillan, 1999) as well as

* Corresponding author. Department of Pharmacology Howard University College of Medicine, 520 W Street, NW Washington, DC, 20059, USA.
E-mail address: ytizabi@howard.edu (Y. Tizabi).

Abbreviations:

ACh	acetylcholine
DHBE	dihydro-beta-erthroidine
Fe	iron
L-DOPA	levodopa
MEC	mecamylamine
MLA	methyllycaconitine

Mn	manganese
MPTP	1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine
MTT	3, (4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
NIC	nicotine
PD	Parkinson's disease
SNpc	substantia nigra pars compacta
6-OHDA	6-hydroxydopamine

severe neurotoxicity leading to Parkinson's disease-like syndrome (Peres et al., 2016; Andrade et al., 2017).

Iron (Fe), like Mn is also a critical component of a variety of enzymes or co-enzymes including catalases and cytochromes that mediate cellular processes and drug metabolism. Fe is best recognized as an essential element in hemoglobin synthesis, where its deficiency is reflected in wide-spread iron-deficiency anemia (Anand and Gupta, 2018). Excess iron (or iron overload), on the other hand, may lead to conditions such as diabetes, increased infection risk as well as severe neurotoxicological symptoms including Parkinson-like syndrome (Wang et al., 2016; Zhang and Rawal, 2017; Ganz, 2018). Indeed, substantia nigra, where the selective loss of dopaminergic neurons occurs, is the primary region in the brain where iron is deposited (Dexter et al. (1989); Riederer et al. (1989)).

Cellular models of substantia nigra dopamine containing neurons include neuroblastoma-derived SH-SY5Y cells. We and others have used this cell line extensively to investigate mechanism of neurotoxicity as well as potential novel neuroprotective agents (Manavalan et al., 2017; Getachew et al., 2018; Omura et al., 2018). In this regard, protective effects of nicotine or curcumin against toxicity induced by selective dopaminergic toxins, including salsolinol and rotenone have been observed (Copeland et al., 2005; Qualls et al., 2014). Whereas rotenone is a synthetic pesticide, salsolinol is an endogenous compound that is generated from the condensation of dopamine and aldehydes and is selectively toxic to dopaminergic cells (Collins, 1988; Naoi, 2004; Segura-Aguilar and Kostrzewa, 2004). It was shown that protective effects of nicotine against salsolinol are mediated via nicotinic receptors because mecamylamine, a non-selective nicotinic antagonist, could totally block the effects of nicotine (Copeland et al., 2005). In addition, many other in-vitro and in-vivo studies have provided evidence of nicotine's protection against toxicants that lead to neurodegenerative/neurological consequences including PD. These studies, summarized in recent reviews, include genetically modified mice and non-human primates where protective effects of nicotine against neuronal damage induced by 6-hydroxydopamine (6-OHDA), 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP), paraquat, methamphetamine, glutamate and β -amyloid were reported (Tizabi, 2016; Tizabi and Getachew, 2017). Indeed, the findings with neuroprotective effects of nicotine has led to the suggestion of initiating pulsatile nicotine trials in patients suffering from PD (Tizabi and Getachew, 2017).

To better characterize the utility of nicotine in PD-like symptoms that might be brought about by excess exposure to trace elements such as Mn and Fe, we hereby evaluated the effects of nicotine against toxicity brought about by these two metals in SH-SY5Y cells. Moreover, we aimed to determine the involvement of specific nicotinic receptors in potential protective effects of nicotine.

2. Materials and methods

Manganese sulfate, iron sulfate, nicotine (bitartrate salt), dihydro-beta-erthroidine (DHBE), methyllycaconitine (MLA), selective antagonists for α 4 β 2 and α 7 nicotinic receptor subtypes, respectively, mecamylamine (MEC), a non-selective nicotinic antagonist, and other analytical reagents were purchased from Sigma Chemical Company (Sigma-Aldrich, St. Louis, MO). The SH-SY5Y human

neuroblastoma cell line were purchased from American Type Culture Collection (ATCC, Manassas, VA). 3, (4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reagent was purchased from Fisher Scientific (Pittsburgh, PA).

As reported in detail previously (Manavaln et al., 2017; Getachew et al., 2018) and briefly here, SH-SY5Y cells were cultured in a 1:1 mixture of Dulbeccos Modified Eagle Medium (DMEM) and Ham's F12 supplemented with 10% fetal bovine serum, penicillin/streptomycin (100 IU/ml), and gentamicin (50 μ g/ml) at 37 °C in 95% O₂/5% CO₂ humidified incubator. The cells were trypsinized when confluent and plated in 96 well plates (1.2 \times 10⁴ cells/well). Cells were allowed to adhere to bottom surface for 24 h. Then, fresh media containing various concentrations of Mn, Fe, nicotine and nicotinic antagonists were added to the carefully aspirated wells.

We first conducted concentration-response experiments to determine the levels of toxicity of Mn and Fe. Once a suitable toxicity level was obtained, that concentration of Mn or Fe was used in subsequent studies. This was followed by a concentration-response experiment involving nicotine and Mn or Fe. Here also, once an optimal concentration of nicotine (one that totally blocked the effects of Mn or Fe) was obtained, that concentration was used in subsequent studies involving nicotinic antagonists. Thus, various concentrations of DHBE, MLA or MEC were added 1 h prior to nicotine, which was added 1 h prior to Mn or Fe. In all cases, the control group consisted of cells that were maintained in media alone and without any drug treatment. All treatments were carried out for 24 h and the effects on cell viability were determined following the 24 h incubation. Each cell viability study was run in sextuplicate (i.e., 6 replicates) and a minimum of 4 assays were conducted for each experimental manipulation.

Determination of cell viability was done by 3, (4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) colorimetric assay according to the manufacturer's protocol as described previously (Manavalan et al., 2017; Getachew et al., 2018). Briefly, the yellow MTT tetrazolium salt (0.5 mg/ml) was dissolved in phosphate-buffered saline (PBS) with 10 mM (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES). 30 μ l of MTT was added to each well and incubated for 3 h at 37 °C. The live cells cause a reduction of the yellow salt to insoluble purple formazan crystals. The wells were then aspirated, and 50 μ l of dimethyl sulfoxide (DMSO) was added to the wells to solubilize the crystals, and the plates were placed in a shaker for an hour. The plates were then read spectrophotometrically at 570 nm with a background of 630 nm in a plate reader. Cell viability was determined by subtracting the test results from the background and is presented as a percentage of the control.

Data is expressed as mean \pm standard error of the mean (SEM). Statistical differences within and between treatment groups were determined by one-way analysis of variance (ANOVA) followed by post-hoc Newman-Keuls Multiple comparison test, where $P < 0.05$ was considered statistically significant. Data were analyzed using Graphpad Prism 3 (Graphpad Software, Inc., San Diego, CA).

3. Results

Fig. 1 depicts the effect of various concentrations of Mn on SH-SY5Y viability. We used the highest concentration of 20 μ M because it is




Fig. 1. Effect of various concentrations of Mn on SH-SY5Y cell viability. Cells were treated with various concentrations of Mn for 24 h and cell viability was determined by MTT. Values are mean \pm SEM. * $p < 0.05$, ** $p < 0.01$ compared to control. $N = 5$ per treatment.

equivalent to what would be a toxicological exposure in the environment (Caito and Aschner, 2015; O'Neal and Zheng, 2015). As seen, this concentration caused the highest toxicity of approximately 30% [$F(4,20) = 16.1$, $p < 0.01$], and hence was used in all subsequent studies.

Fig. 2 depicts the effect of various concentrations of Fe on SH-SY5Y viability. We used the highest concentration of 20 μM here also because it is equivalent to what would be a toxicological exposure in the environment (Dexter et al., 1989; Fine, 2000). As seen, this concentration caused the highest toxicity of approximately 35% [$F(4,20) = 16.8$, $p < 0.01$], and hence was used in all subsequent studies.

Fig. 3 depicts the effects of various concentrations of nicotine against Mn-induced toxicity in SH-SY5Y cells. As seen, there was a concentration-dependent protection of Mn-induced toxicity by nicotine (NIC), where 1 μM NIC completely blocked the effects of Mn [$F(5,25) = 12.6$, $p < 0.01$]. Thus, this concentration of NIC was used in subsequent studies to investigate the effects of nicotinic antagonists.

Fig. 4 depicts the effects of various concentrations of nicotine against Fe-induced toxicity in SH-SY5Y cells. As seen, there was a concentration-dependent protection of Fe-induced toxicity by nicotine (NIC), where 1 μM NIC completely blocked the effects of Fe [$F(5,25) = 12.8$, $p < 0.01$]. Thus, this concentration of NIC was used in subsequent studies to investigate the effects of nicotinic antagonists.

Fig. 5 depicts the effects of various concentrations of MLA (selective $\alpha 7$ antagonist); DHBE (selective $\alpha 4$ - $\beta 2$ antagonist) and MEC (non-selective nicotinic antagonist) against protective effects of NIC on Mn-induced toxicity in SH-SY5Y cells. All three antagonists concentration-dependently blocked the effects of NIC, where 1 μM of each antagonist completely abolished the effects of NIC [$F(11,48) = 9.3$, $p < 0.01$]. None of the antagonists by themselves had any effect on cell viability (data not shown).

Fig. 6 depicts the effects of various concentrations of MLA, DHBE and MEC against protective effects of NIC on Fe-induced toxicity in SH-SY5Y cells. All three antagonists concentration-dependently blocked the effects of NIC, where 1 μM of each antagonist completely abolished the effects of NIC [$F(11,48) = 10.1$, $p < 0.01$]. None of the antagonists by themselves had any effect on cell viability (data not shown).

4. Discussion

The results of this study provide novel evidence on protective effects of nicotine against trace elements-induced toxicity in SH-SY5Y cells, which further support the potential utility of nicotinic intervention in PD. This contention is based on the ability of nicotine to block the damage that is induced by toxicological concentrations of manganese and iron, two essential elements, implicated in pathophysiology of PD

or PD-like symptoms. The protective effects of nicotine against these two elements was observed in SH-SY5Y cells, which express nicotinic receptors (Gould et al., 1992; Lukas et al., 1993; Elnagar et al., 2018) and are commonly used as a model of substantia nigra dopaminergic neurons that are primarily affected in PD (Brown et al., 2014; Manavalan et al., 2017; Getachew et al., 2018).

Indeed, a number of studies have identified the presence of nicotinic cholinergic receptors in the substantia nigra. These include earlier experiments utilizing autoradiography (Clarke and Pert, 1985) and immunohistochemistry (Göldner et al., 1997; Sorenson et al., 1998). More recently potential roles of these receptors in this area (Fature et al., 2014) as well as in nigrostriatal terminals (Inden et al., 2016) have been provided. In addition, it has been suggested that nicotine, by inhibiting astrocytes activation in the substantia nigra pars compacta (SNpc), can protect dopaminergic neurons against degeneration (Jurado-Coronel et al., 2016). Interestingly, both DHBE and MLA, selective antagonists to nicotinic receptor subtypes, $\alpha 4$ - $\beta 2$ and $\alpha 7$, respectively, appear to equally mediate the effects of nicotine. This is because both of these antagonists at equivalent concentrations, totally blocked the effect of nicotine. Further confirmation of both receptor involvement was provided by results of mecamylamine, a non-selective nicotinic receptor antagonist, which also equi-potently blocked nicotine's protective effects.

It is of relevance to note that nicotine's main targets are nicotinic receptors that belong to the ionotropic class of receptors, which directly regulate the opening of a cation channel in the neuronal membrane. The endogenous ligand for these receptor is the important neurotransmitter acetylcholine (ACh), hence these receptors are commonly referred to as nAChRs. Currently, 11 neuronal nAChR subunits: $\alpha 2$ - $\alpha 7$, $\alpha 9$, $\alpha 10$, $\beta 2$ - $\beta 4$, which assemble into pentameric complexes and provide subtype diversity and distinct anatomical, physiological, and pharmacological characteristics have been identified (Changeux et al., 1998; Dani, 2015). The most predominant and extensively studied subtype in the brain is formed from $\alpha 4$ and $\beta 2$ subunits and is commonly referred to as high-affinity binding site because of its high binding affinity to nicotine and ACh. The other major class with a high affinity for α -bungarotoxin but low affinity for nicotine is formed from $\alpha 7$ homomeric subunits and is commonly referred to as low-affinity binding site. These receptors have been directly implicated not only in nicotine addiction (Tizabi et al., 2002, 2007; Meyerhoff et al., 2006; Picciotto and Mineur, 2014; Liu and Li, 2018), but also in a variety of central functions such as cognitive and attentional processes (Tizabi, 2007; Proulx et al., 2014), mood regulation (Hurley and Tizabi, 2013; Lewis and Picciotto, 2013), pain (Campbell et al., 2006; Bufalo et al., 2014), neuronal plasticity (Serres and Carney, 2006; Antonelli et al., 2012; Barreto et al., 2015; Fuenzalida et al., 2016) and neuronal




Fig. 2. Effect of various concentrations of Fe on SH-SY5Y cell viability. Cells were treated with various concentrations of Mn for 24 h and cell viability was determined by MTT. Values are mean \pm SEM. * $p < 0.05$, ** $p < 0.01$ compared to control. $N = 5$ per treatment.




Fig. 3. Effect of various concentrations of NIC on Mn-induced toxicity in SH-SY5Y cells. Cells were treated with various concentrations of NIC 1 h prior to Mn (20 μM) and cell viability was determined by MTT 24 h later. Values are mean ± SEM. **p < 0.01 compared to control. †† p < 0.01 compared to Mn alone. N = 5 per treatment.




Fig. 4. Effect of various concentrations of NIC on Fe-induced toxicity in SH-SY5Y cells. Cells were treated with various concentrations of NIC 1 h prior to Fe (20 μM) and cell viability was determined by MTT 24 h later. Values are mean ± SEM. **p < 0.01 compared to control. †† p < 0.01 compared to Mn alone. N = 5 per treatment.




Fig. 5. Effects of various concentrations of MLA, DHBE and MEC against protective effects of NIC on Mn-induced toxicity in SH-SY5Y cells. Cells were treated with various concentrations of the antagonists 1 h prior to NIC (1 μM), which was applied 1 h prior to Mn (20 μM) and cell viability was determined by MTT 24 h later. Values are mean ± SEM. **p < 0.01 compared to control. †† p < 0.01 compared to Mn alone. \$p < 0.05, \$\$p < 0.01 compared to NIC + Mn. N = 5 per treatment.

protection (Das and Tizabi, 2009; Dineley et al., 2015; Quik et al., 2015; Tizabi and Getachew, 2017). Although in some cases a specific receptor subtype might be mediating the action of nicotine (e.g. beta2 containing subunit for its addictive property) (Picciotto and Mineur, 2014; Miller et al., 2018; Liu and Li., 2018) or alpha7 for its antipsychotic effects (Freedman, 2014; Wallace and Bertrand, 2015; Wadenberg et al., 2017; Jones, 2018), here, the protective effects of nicotine against Mn and Fe toxicity, appear to be mediated by both alpha 4-beta2 and alpha7 subunits. This finding is similar to observation of nicotine protection against glutamate or beta-amyloid-induced toxicity in cell culture, where both of these nicotinic receptor subtypes were implicated (Kihara et al., 1998; Akaïke et al., 2010; Yu et al., 2011).

Interestingly, both Mn and Fe appear to target primarily the basal ganglia, which includes the globus pallidus and substantia nigra, the latter being the primary site of damage leading to PD (Erikson and Aschner, 2003; Rajagopalan et al., 2016). Thus, manganese associated with increased Mn levels, particularly in basal ganglia, may manifest PD-like symptoms (Peres et al., 2016; Andrade et al., 2017). Although the pathological mechanisms associated with Mn neurotoxicity are poorly understood, several reports have implicated oxidative stress as well as inflammatory processes as important contributory factors (Santos et al., 2012; Dusek et al., 2015). Similarly, iron-induced neurodegenerative diseases including PD has been at least partially ascribed to increases in oxidative stress as well as neuroinflammatory factors (Salvador et al., 2011; Dusek et al., 2015). On the other hand, it has been hypothesized that neuroprotective effects of nicotine are also likely to involve suppression of oxidative stress and pro-inflammatory cytokines (Perez et al., 2010; Chatterjee et al., 2012; Hurley and Tizabi, 2013; Barreto et al., 2015; Tizabi, 2016). Hence, it may be suggested that similar mechanisms play a role in protective effects of nicotine against trace metal induced toxicity. Curiously, it has been shown that in regard to damage induced by other metals such as copper, nicotine's protective mechanism may involve regulation of metal homeostasis. Thus, it has been demonstrated that copper-induced beta-amyloidosis can be attenuated by nicotine via reduction of this metal level in both in-vivo and in-vitro studies (Zhang et al., 2006). Therefore, it would be of significant interest to determine whether nicotine may have a similar effect on Mn and/or Fe. Another important undertaking would be to determine whether the combination of trace metals including tin, copper, zinc, etc. with Mn and/or iron would exacerbate the neurodegenerative condition (Lee et al., 2006; Andrade et al., 2017) and whether nicotine would be equally effective.

It should also be noted that death/survival signaling mechanisms




Fig. 6. Effects of various concentrations of MLA, DHBE and MEC against protective effects of NIC on Fe-induced toxicity in SH-SY5Y cells. Cells were treated with various concentrations of the antagonists 1 h prior to NIC (1 μM), which was applied 1 h prior to Fe (20 μM) and cell viability was determined by MTT 24 h later. Values are mean ± SEM. **p < 0.01 compared to control. †† p < 0.01 compared to Fe alone. \$p < 0.05, \$\$p < 0.01 compared to NIC + Fe. N = 5 per treatment.

have been implicated in the action of nicotine (Wei et al., 2011). Thus, a role for ERK-MAPK and JNK (Dineley et al., 2001; Xue et al., 2014; Li et al., 2015), a role for caspase pathways (Ramlochansingh et al., 2011; Lu et al., 2017), and more recently a role for Wnt/β-catenin signaling (Liu et al., 2017) in protective effects of nicotine have been provided.

In summary, the results of this study provide evidence for neuroprotective effects of nicotine against toxicity induced by Mn or Fe in a cellular model of PD. Moreover, both high and low affinity nicotinic receptors (i.e., alpha 4-beta 2 and alpha7 subtypes) appear to mediate the effects of nicotine. Thus, utility of nicotine or nicotinic agonists in trace element-induced Parkinson-like syndrome may be suggested.

Acknowledgement

Supported by: NIH/NIAAA R03AA022479 (YT), NIA/NIH 1R25AG047843-01 (ABC), NIEHS R01ES10653 (MA).

References

O'Neal, S.L., Zheng, W., 2015. Manganese toxicity upon overexposure: a decade in review. *Current environmental health reports* 2 (3), 315–328.

Akaike, A., Takada-Takatori, Y., Kume, T., Izumi, Y., 2010. Mechanisms of neuroprotective effects of nicotine and acetylcholinesterase inhibitors: role of alpha 4 and alpha7 receptors in neuroprotection. *J. Mol. Neurosci.* 40 (1–2), 211–216.

Anand, I.S., Gupta, P., 2018. Anemia and iron deficiency in heart failure: current concepts and emerging therapies. *Circulation* 138 (1), 80–98.

Andrade, V.M., Aschner, M., Marreilha Dos Santos, A.P., 2017. Neurotoxicity of metal mixtures. *Adv Neurobiol* 18, 227–265.

Antonelli, M.C., Guillemin, G.J., Raisman-Vozari, R., Del-Bel, E.A., Aschner, M., Collins, M.A., et al., 2012. New strategies in neuroprotection and neurorepair. *Neurotox. Res.* 21 (1), 49–56.

Aschner, J.L., Aschner, M., 2005. Nutritional aspects of manganese homeostasis. *Mol. Aspect. Med.* 26 (4–5), 353–362.

Aschner, M., Gannon, M., 1994. Manganese (Mn) transport across the rat blood-brain barrier: saturable and transferrin-dependent transport mechanisms. *Brain Res. Bull.* 33 (3), 345–349.

Barreto, G.E., Iarkov, A., Moran, V.E., 2015. Beneficial effects of nicotine, cotinine and its metabolites as potential agents for Parkinson's disease. *Front. Aging Neurosci.* 6, 340.

Brown, D., Tamas, A., Reglodi, D., Tizabi, Y., 2014. PACAP protects against inflammatory-mediated toxicity in dopaminergic SH-SY5Y cells: implication for Parkinson's disease. *Neurotox. Res.* 26 (3), 230–239.

Bufalo, A.D., Cesario, A., Salinaro, G., Fini, M., Russo, P., 2014. Alpha 9 Alpha 10 nicotinic acetylcholine receptors as target for the treatment of chronic pain. *Curr. Pharmaceut. Des.* 20 (38), 6042–6047.

Burton, N.C., Guilarte, T.R., 2009. Manganese neurotoxicity: lessons learned from longitudinal studies in nonhuman primates. *Environ. Health Perspect.* 117 (3), 325–332.

Caito, S., Aschner, M., 2015. Neurotoxicity of metals. *Handb. Clin. Neurol.* 131, 169–189.

Campbell, V.C., Taylor, R.E., Tizabi, Y., 2006. Antinociceptive effects of alcohol and nicotine: involvement of the opioid system. *Brain Res.* 30 (1), 71–77.

Changeux, J.P., Bertrand, D., Corringier, P.J., Dehaene, S., Edelstein, S., Léna, C., Le Novère, N., Marubio, L., Picciotto, M., Zoli, M., 1998. Brain nicotinic receptors:

structure and regulation, role in learning and reinforcement. *Brain Res. Rev.* 26 (2–3), 198–216.

Chatterjee, P.K., Yeboah, M.M., Dowling, O., Xue, X., Powell, S.R., Al-Abed, Y., Metz, C.N., 2012. Nicotinic acetylcholine receptor agonists attenuate septic acute kidney injury in mice by suppressing inflammation and proteasome activity. *PLoS One* 7 (5), e35361.

Clarke, P.B., Pert, A., 1985. Autoradiographic evidence for nicotine receptors on nigrostriatal and mesolimbic dopaminergic neurons. *Brain Res.* 348 (2), 355–358.

Collins, M.A., 1988. Acetaldehyde and its condensation products as markers in alcoholism. *Recent Dev. Alcohol* 6, 387–403.

Copeland Jr., R.L., Leggett, Y.A., Kanaan, Y.M., Taylor, R.E., Tizabi, Y., 2005. Neuroprotective effects of nicotine against salsolinol-induced cytotoxicity: implications for Parkinson's disease. *Neurotox. Res.* 8 (3–4), 289–293.

Dani, J.A., 2015. Neuronal nicotinic acetylcholine receptor structure and function and response to nicotine. *Int. Rev. Neurobiol.* 124, 3–19.

Das, J.R., Tizabi, Y., 2009. Additive protective effects of donepezil and nicotine against salsolinol-induced cytotoxicity in SH-SY5Y cells. *Neurotox. Res.* 16 (3), 194–204.

Dexter, D.T., et al., 1989. Increased nigral iron content and alterations in other metal ions occurring in brain in Parkinson's disease. *J. Neurochem.* 52, 1830–1836.

Dineley, K.T., Westerman, M., Bui, D., Bell, K., Ashe, K.H., Sweatt, J.D., 2001. Beta-amyloid activates the mitogen-activated protein kinase cascade via hippocampal alpha7 nicotinic acetylcholine receptors: in vitro and in vivo mechanisms related to Alzheimer's disease. *J. Neurosci.* 21 (12), 4125–4133.

Dineley, K.T., Pandya, A.A., Yakel, J.L., 2015. Nicotinic ACh receptors as therapeutic targets in CNS disorders. *Trends Pharmacol. Sci.* 36 (2), 96–108.

Dusek, P., Roos, P.M., Litwin, T., Schneider, S.A., Flaten, T.P., Aaseth, J., 2015. The neurotoxicity of iron, copper and manganese in Parkinson's and Wilson's diseases. *J. Trace Elem. Med. Biol.* 31, 193–203.

Elnagar, M.R., Walls, A.B., Helal, G.K., Hamada, F.M., Thomsen, M.S., Jensen, A.A., 2018. Functional characterization of α7 nicotinic acetylcholine and NMDA receptor signaling in SHSY5Y neuroblastoma cells in an ERK phosphorylation assay. *Eur. J. Pharmacol.* 826, 106–113.

Erikson, K.M., Aschner, M., 2003. Manganese neurotoxicity and glutamate-GABA interaction. *Neurochem. Int.* 43 (4–5), 475–480.

Faure, P., Tolu, S., Valverde, S., Naudé, J., 2014. Role of nicotinic acetylcholine receptors in regulating dopamine neuron activity. *Neuroscience* 282, 86–100.

Fine, J.S., 2000. Iron poisoning. *Curr. Probl. Pediatr.* 30 (3), 71–90.

Freedman, R., 2014. α7-nicotinic acetylcholine receptor agonists for cognitive enhancement in schizophrenia. *Annu. Rev. Med.* 65, 245–261.

Fuenzalida, M., Pérez, M.A., Arias, H.R., 2016. Role of nicotinic and muscarinic receptors on synaptic plasticity and neurological diseases. *Curr. Pharmaceut. Des.* 22 (14), 2004–2014.

Ganz, T., 2018. Iron and infection. *Int. J. Hematol.* 107 (1), 7–15.

Getachew, B., Hudson, T., Heinbockel, T., Csoka, A.B., Tizabi, Y., 2018. Protective effects of donepezil against alcohol-induced toxicity in cell culture: role of caspase-3. *Neurotox. Res.* 34 (3), 757–762.

Göldner, F.M., Dineley, K.T., Patrick, J.W., 1997. Immunohistochemical localization of the nicotinic acetylcholine receptor subunit alpha 6 to dopaminergic neurons in the substantia nigra and ventral tegmental area. *Neuroreport* 8 (12), 2739–2742.

Gould, J., Reeve, H.L., Vaughan, P.F., Peers, C., 1992. Nicotinic acetylcholine receptors in human neuroblastoma (SH-SY5Y) cells. *Neurosci. Lett.* 145 (2), 201–204.

Hurley, L.L., Tizabi, Y., 2013. Neuroinflammation, neurodegeneration, and depression. *Neurotox. Res.* 23 (2), 131–144. *J. Trace Elem. Med. Biol.* 2015;31:193-203.

Inden, M., Takata, K., Yanagisawa, D., Ashihara, E., Tooyama, I., Shimohama, S., Kitamura, Y., 2016. α4 nicotinic acetylcholine receptor modulated by galantamine on nigrostriatal terminals regulates dopamine receptor-mediated rotational behavior. *Neurochem. Int.* 94, 74–81.

- Jones, C., 2018. $\alpha 7$ nicotinic acetylcholine receptor: a potential target in treating cognitive decline in schizophrenia. *J. Clin. Psychopharmacol.* 38 (3), 247–249.
- Jurado-Coronel, J.C., Avila-Rodriguez, M., Capani, F., Gonzalez, J., Moran, V.E., Barreto, G.E., 2016. Targeting the nicotinic acetylcholine receptors (nAChRs) in astrocytes as a potential therapeutic target in Parkinson's disease. *Curr. Pharmaceut. Des.* 22 (10), 1305–1311.
- Kihara, T., Shimohama, S., Urushitani, M., Sawada, H., Kimura, J., Kume, T., Maeda, T., Akaike, A., 1998. Stimulation of $\alpha 4\beta 2$ nicotinic acetylcholine receptors inhibits beta-amyloid toxicity. *Brain Res.* 792 (2), 331–334.
- Lee, J.J., Kim, Y.M., Park, S.K., Lee, M.K., 2006. Effects of tributyltin chloride on L-DOPA-induced cytotoxicity in PC12 cells. *Arch. Pharm. Res. (Seoul)* 29 (8), 645–650.
- Lewis, A.S., Picciotto, M.R., 2013. High-affinity nicotinic acetylcholine receptor expression and trafficking abnormalities in psychiatric illness. *Psychopharmacology* 229 (3), 477–485.
- Li, Z.Z., Guo, Z.Z., Zhang, Z., Cao, Q.A., Zhu, Y.J., Yao, H.L., Wu, L.L., Dai, Q.Y., 2015. Nicotine-induced upregulation of VCAM-1, MMP-2, and MMP-9 through the $\alpha 7$ -nAChR-JNK pathway in RAW264.7 and MOVAS cells. *Mol. Cell. Biochem.* 399 (1–2), 49–58.
- Liu, W., Li, M.D., 2018. Insights into nicotinic receptor signaling in nicotine addiction: implications for prevention and treatment. *Curr. Neuropharmacol.* 16 (4), 350–370.
- Liu, Y., Hao, S., Yang, B., Fan, Y., Qin, X., Chen, Y., Hu, J., 2017. Wnt/ β -catenin signaling plays an essential role in $\alpha 7$ nicotinic receptor-mediated neuroprotection of dopaminergic neurons in a mouse Parkinson's disease model. *Biochem. Pharmacol.* 140, 115–123.
- Lu, J.Y.D., Su, P., Barber, J.E.M., Nash, J.E., Le, A.D., Liu, F., Wong, A.H.C., 2017. The neuroprotective effect of nicotine in Parkinson's disease models is associated with inhibiting PARP-1 and caspase-3 cleavage. *PeerJ* 5, e3933.
- Lukas, R.J., Norman, S.A., Lucero, L., 1993. Characterization of nicotinic acetylcholine receptors expressed by cells of the SH-SY5Y human neuroblastoma clonal line. *Mol. Cell. Neurosci.* 4 (1), 1–12.
- Manavalan, S., Getachew, B., Manaye, K.F., Khundmiri, S.J., Csoka, A.B., McKinley, R., Tamas, A., Reglodi, D., Tizabi, Y., 2017. PACAP protects against ethanol and nicotine toxicity in SH-SY5Y cells: implications for drinking-smoking co-morbidity. *Neurotox. Res.* 32 (1), 8–13.
- McMillan, D.E., 1999. A brief history of the neurobehavioral toxicity of manganese: some unanswered questions. *Neurotoxicology* 20 (2–3), 499–508.
- Meyerhoff, D.J., Tizabi, Y., Staley, J.K., Durazzo, T.C., Glass, J.M., Nixon, S.J., 2006. Smoking comorbidity in alcoholism: neurobiological and neurocognitive consequences. *Alcohol Clin. Exp. Res.* 30 (2), 253–264.
- Miller, M.B., Wilson, R.S., Lam, T.T., Nairn, A.C., Picciotto, M.R., 2018 Oct 15. Evaluation of the phosphoproteome of mouse $\alpha 4\beta 2$ -containing nicotinic acetylcholine receptors in vitro and in vivo. *Proteomes* 6 (4).
- Naoi, M., Maruyama, W., Nagy, G.M., 2004. Dopamine-derived salsolinol derivatives as endogenous monoamine oxidase inhibitors: occurrence, metabolism and function in human brains. *Neurotoxicology* 25 (1–2), 193–204.
- Omura, T., Matsuda, H., Nomura, L., Imai, S., Denda, M., Nakagawa, S., Yonezawa, A., Nakagawa, T., Yano, I., Matsubara, K., 2018 Nov 30. Ubiquitin ligase HMG-CoA reductase degradation 1 (HRD1) prevents cell death in a cellular model of Parkinson's disease. *Biochem. Biophys. Res. Commun.* 506 (3), 516–521.
- Cacabelos, R., 2017. Parkinson's disease: from pathogenesis to pharmacogenomics. *Int. J. Mol. Sci.* 18 (3), 551–579.
- Peres, T.V., Schettinger, M.R., Chen, P., Carvalho, F., Avila, D.S., Bowman, A.B., Aschner, M., 2016. Manganese-induced neurotoxicity: a review of its behavioral consequences and neuroprotective strategies. *BMC Pharmacol Toxicol* 17 (1), 57.
- Perez, X.A., Bordia, T., McIntosh, J.M., Quik, M., 2010. $\alpha 6\beta 2^*$ and $\alpha 4\beta 2^*$ nicotinic receptors both regulate dopamine signaling with increased nigrostriatal damage: relevance to Parkinson's disease. *Mol. Pharmacol.* 78 (5), 971–980.
- Picciotto, M.R., Mineur, Y.S., 2014. Molecules and circuits involved in nicotine addiction: the many faces of smoking. *Neuropharmacology* 76 Pt, 545–553.
- Picconi, B., Calabresi, P., 2017. Switching on the lights of dyskinesia: perspectives and limits of the optogenetic approaches. *Mov. Disord.* 32 (4), 485–486.
- Proulx, E., Piva, M., Tian, M.K., Bailey, C.D., Lambe, E.K., 2014. Nicotinic acetylcholine receptors in attention circuitry: the role of layer VI neurons of prefrontal cortex. *Cell. Mol. Life Sci.* 71 (7), 1225–1244.
- Qualls, Z., Brown, D., Ramlochan Singh, C., Hurley, L.L., Tizabi, Y., 2014. Protective effects of curcumin against rotenone and salsolinol-induced toxicity: implications for Parkinson's disease. *Neurotox. Res.* 25 (1), 81–89.
- Quik, M., Zhang, D., McGregor, M., Bordia, T., 2015. $\alpha 7$ nicotinic receptors as therapeutic targets for Parkinson's disease. *Biochem. Pharmacol.* 97 (4), 399–407.
- Rajagopalan, S., Rane, A., Chinta, S.J., Andersen, J.K., 2016. Regulation of ATP13A2 via PHD2-HIF1 α signaling is critical for cellular iron homeostasis: implications for Parkinson's disease. *J. Neurosci.* 36 (4), 1086–1095.
- Ramlochan Singh, C., Taylor, R.E., Tizabi, Y., 2011. Toxic effects of low alcohol and nicotine combinations in SH-SY5Y cells are apoptotically mediated. *Neurotox. Res.* 20 (3), 263–269.
- Riederer, P., et al., 1989. Transition metals, ferritin, glutathione, and ascorbic acid in parkinsonian brains. *J. Neurochem.* 52, 515–520.
- Roth, J.A., Garrick, M.D., 2003. Iron interactions and other biological reactions mediating the physiological and toxic actions of manganese. *Biochem. Pharmacol.* 66 (1), 1–13.
- Salvador, G.A., Uranga, Romina M., Giusto, Norma M., 2011. Iron and mechanisms of neurotoxicity. *Int. J. Alzheimer's Dis.* 9 Article ID 720658.
- Santos, D., Milatovic, D., Andrade, V., Batoreu, M.C., Aschner, M., Marreilha dos Santos, A.P., 2012. The inhibitory effect of manganese on acetylcholinesterase activity enhances oxidative stress and neuroinflammation in the rat brain. *Toxicology* 292 (2–3), 90–98.
- Segura-Aguilar, J., Kostzewa, R.M., 2004. Neurotoxins and neurotoxic species implicated in neurodegeneration. *Neurotox. Res.* 6 (7–8), 615–630.
- Serres, F., Carney, S.L., 2006. Nicotine regulates SH-SY5Y neuroblastoma cell proliferation through the release of brain-derived neurotrophic factor. *Brain Res.* 1101 (1), 36–42.
- Sorenson, E.M., Shiroyama, T., Kitai, S.T., 1998. Postsynaptic nicotinic receptors on dopaminergic neurons in the substantia nigra pars compacta of the rat. *Neuroscience* 87 (3), 659–673.
- Sveinbjornsdottir, S., 2016. The clinical symptoms of Parkinson's disease. *J. Neurochem.* 139 (Suppl. 1), 318–324.
- Tanner, C.M., Kamel, F., Ross, G.W., et al., 2011. Rotenone, paraquat, and Parkinson's disease. *Environ. Health Perspect.* 119 (6), 866–872.
- Tizabi, Y., 2007. Nicotine and nicotinic system in hypoglutamatergic models of schizophrenia. *Neurotox. Res.* 12 (4), 233–246.
- Tizabi, Y., 2016. Duality of antidepressants and neuroprotectants. *Neurotox. Res.* 30 (1), 1–13.
- Tizabi, Y., Getachew, B., 2017. Nicotinic receptor intervention in Parkinson's disease: future directions. *Clin. Pharmacol. Transl. Med.* 1 (1), 14–19.
- Tizabi, Y., Copeland Jr., R.L., Louis, V.A., Taylor, R.E., 2002. Effects of combined systemic alcohol and central nicotine administration into ventral tegmental area on dopamine release in the nucleus accumbens. *Alcohol Clin. Exp. Res.* 26 (3), 394–399.
- Tizabi, Y., Bai, L., Copeland Jr., R.L., Taylor, R.E., 2007. Combined effects of systemic alcohol and nicotine on dopamine release in the nucleus accumbens shell. *Alcohol* 42 (5), 413–416.
- Vaccari, C., El Dib, R., de Camargo, J.L.V., 2017. Paraquat and Parkinson's disease: a systematic review protocol according to the OHAT approach for hazard identification. *Syst. Rev.* 6, 98.
- Wadenberg, M.G., Manetti, D., Romanelli, M.N., Arias, H.R., 2017. Significance of the nicotinic $\alpha 7$ receptor in cognition and antipsychotic-like behavior in the rat. *Behav. Brain Res.* 333, 129–134.
- Wallace, T.L., Bertrand, D., 2015. Neuronal $\alpha 7$ nicotinic receptors as a target for the treatment of schizophrenia. *Int. Rev. Neurobiol.* 124, 79–111.
- Wang, J.Y., Zhuang, Q.Q., Zhu, L.B., et al., 2016. Meta-analysis of brain iron levels of Parkinson's disease patients determined by postmortem and MRI measurements. *Sci. Rep.* 6, 36669.
- Wei, J., Wang, J., Dwyer, J.B., Mangold, J., Cao, J., Leslie, F.M., Li, M.D., 2011. Gestational nicotine treatment modulates cell death/survival-related pathways in the brains of adolescent female rats. *Int. J. Neuropsychopharmacol.* 14 (1), 91–106.
- Xue, M.Q., Liu, X.X., Zhang, Y.L., Gao, F.G., 2014. Nicotine exerts neuroprotective effects against β -amyloid-induced neurotoxicity in SH-SY5Y cells through the Erk1/2-p38-JNK-dependent signaling pathway. *Int. J. Mol. Med.* 33 (4), 925–933.
- Yousefi Babadi, V., Sadeghi, L., Shirani, K., Malekiran, A.A., Rezaei, M., 2014. The toxic effect of manganese on the acetylcholinesterase activity in rat brains. *J. Toxicol.* 946372.
- Yu, W., Mechawar, N., Krantic, S., Quirion, R., 2011. $\alpha 7$ Nicotinic receptor activation reduces β -amyloid-induced apoptosis by inhibiting caspase-independent death through phosphatidylinositol 3-kinase signaling. *J. Neurochem.* 119 (4), 848–858.
- Zhang, J., Liu, Q., Chen, Q., Liu, N.Q., Li, F.L., Lu, Z.B., Qin, C., Zhu, H., Huang, Y.Y., He, W., Zhao, B.L., 2006. Nicotine attenuates beta-amyloid-induced neurotoxicity by regulating metal homeostasis. *Faseb. J.* 20 (8), 1212–1214.
- Zhang, C., Rawal, S., 2017. Dietary iron intake, iron status, and gestational diabetes. *Am. J. Clin. Nutr.* 106 (Suppl 6), 1672S–1680S.