



Position Paper

Use of biosimilars in inflammatory bowel disease: a position update of the Italian Group for the Study of Inflammatory Bowel Disease (IG-IBD)

Gionata Fiorino^{a,b}, Flavio Caprioli^c, Marco Daperno^d, Filippo Mocciaro^e, Mariabeatrice Principi^f, Angelo Viscido^g, Massimo Claudio Fantini^h, Ambrogio Orlandoⁱ, Claudio Papi^j, Vito Annesse^k, Silvio Danese^{a,b}, Maurizio Vecchi^c, Fernando Rizzello^l, Alessandro Armuzzi^{m,n,*}, National patients' association representatives¹

^a IBD Center, Department of Gastroenterology, Humanitas Clinical and Research Center, Rozzano, Italy

^b Department of Biomedical Sciences, Humanitas University, Rozzano, Italy

^c Department of Pathophysiology and Transplantation, Università degli Studi di Milano, Milan, Italy, and Gastroenterology and Endoscopy Unit, Fondazione IRCCS Cà Granda, Ospedale Policlinico di Milano, Milan, Italy

^d Division of Gastroenterology, Ospedale Ordine Mauriziano di Torino, Turin, Italy

^e Gastroenterology and Endoscopy Unit, A.R.N.A.S. Civico-Di Cristina-Benfratelli Hospital, Palermo, Italy

^f Gastroenterology Unit, Department of Organ Transplantation (DETO), Policlinico Universitario, Bari, Italy

^g Department of Life, Health and Environmental Sciences, University of L'Aquila, L'Aquila, Italy

^h Department of Systems Medicine, Tor Vergata University of Rome, Rome, Italy

ⁱ IBD Unit, A.O. Osp. Riuniti "Villa Sofia-Cervello", Palermo, Italy

^j IBD Unit, San Filippo Neri Hospital, Rome, Italy

^k IG-IBD member, Florence, Italy

^l IBD Unit, DIMEC, University of Bologna, Bologna, Italy

^m IBD Unit, Presidio Columbus, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy

ⁿ Università Cattolica del Sacro Cuore, Rome, Italy

ARTICLE INFO

Article history:

Received 16 January 2019

Accepted 4 February 2019

Available online 19 February 2019

Keywords:

Anti-TNFα

Biosimilars

Crohn's disease

Inflammatory bowel disease

Tumor necrosis factor alpha

Ulcerative colitis

ABSTRACT

The first infliximab biosimilar for the treatment of inflammatory bowel disease (IBD) was introduced in 2013, and today eight anti-TNF alpha biosimilars (three for infliximab and five for adalimumab) have been approved and licensed by the European Medicines Agency. Biosimilars present great potential in terms of cost saving and possible consequential reinvestment in the health care system. The increasing knowledge about the process of biosimilar development and use in IBD and the publication of many prospective clinical studies and real-life clinical experiences have progressively changed the point of view of IBD physicians. In the present position paper, the Italian Group for the Study of Inflammatory Bowel Disease present and discuss their updated statements and positions on this topic, with emphasis on the concepts of biosimilarity and extrapolation across indications, safety and immunogenicity, interchangeability and switching, automatic substitution, and, finally, patient education about biosimilars.

© 2019 Editrice Gastroenterologica Italiana S.r.l. Published by Elsevier Ltd. All rights reserved.

1. Introduction

Biosimilars are biological medicinal products that are similar, but not identical, to an approved biological drug, called “origina-

tor” or “reference product” [1]. Like biological drugs, biosimilars are drugs obtained from biological sources (i.e. they are not synthesized), and thus can take on many biochemical forms, including vaccines, gene therapies and recombinant proteins such as growth factors and monoclonal antibodies. Because biological drugs are complex molecules made by living organisms that are naturally variable [2], biosimilars cannot be considered an exact copy of the originator, as instead synthetic chemical drugs are. Thus, the procedures (i.e. in the preclinical phase) for approval and use of biosimilars are more complex than those for the originators.

* Corresponding author at: IBD Unit, Presidio Columbus, Fondazione Policlinico Universitario A. Gemelli IRCCS — Università Cattolica del Sacro Cuore, Via G. Moscatti 31, I-00168, Rome, Italy.

E-mail addresses: alessandro.armuzzi@policlinicogemelli.it, alessandro.armuzzi@unicatt.it (A. Armuzzi).

¹ See Appendix A.

Monoclonal antibodies (mAb) targeting tumor necrosis factor alpha (TNF α), namely infliximab, adalimumab, certolizumab pegol, and golimumab, are biological drugs used in the treatment of inflammatory bowel disease (IBD) and other immune-mediated inflammatory diseases. In particular, infliximab is an intravenously administered chimeric mAb (75% human and 25% murine sequences; IgG1 κ) that was approved by the European Medicines Agency (EMA) in August 1999 [3]. Adalimumab is a subcutaneously administered human mAb (IgG1) that was approved by EMA in September 2003 [4]. The EU patents on the infliximab originator (Remicade[®], J&J, USA [3]) expired in June 2013 or February 2015, depending on the country, and on the adalimumab originator (Humira[®], Abbvie, USA [4]) in October 2018. Thus, three infliximab biosimilars and five adalimumab biosimilars have become available for the same indications as the originators. For infliximab, the biosimilars are: CT-P13 (Inflectra[®], Pfizer, USA [5]; Remsima[®], Celltrion, South Korea [6]), SB2 (Flixabi[®], Samsung Bioepis, South Korea [7], and Biogen, Denmark), and PF-06438179/GP1111 (Zessly[®], Sandoz, Germany [8]). For adalimumab, the biosimilars are: ABP 501 (Amgevita[®], Solymbic[®], Amgen, USA [9]), SB5 (Imraldi[®], Biogen Denmark-Samsung Bioepis, South Korea [10]), FKB327 (Hulio[®], Mylan, USA [11]; Fujifilm Kyowa Kyrin Biologics, Japan), GP2017 (Hyrimoz[®], Sandoz, Germany [12]), and BI 695501 (Cyltezo[®], Boehringer Ingelheim, Germany [13]). Other biosimilars of infliximab and adalimumab are expected to be approved in the near future [14].

Before the introduction of biosimilar mAbs in the EU market, several concerns had been raised about their use in clinical practice. An online survey conducted by the European Crohn's Colitis Organisation (ECCO) in 2013 showed that more than 60% of clinicians in Europe had little or no confidence in prescribing biosimilar mAb in clinical practice [15]. The main concerns were about the validity of extrapolating across indications, the potentially higher immunogenicity of biosimilar mAb than the originator, traceability, automatic substitution, and switching from the originator to a biosimilar. These concerns were common in the IBD community, and they were addressed and discussed in several position papers by national and international societies [16–20].

The introduction of the first infliximab biosimilar, the increased knowledge of biosimilars, and the publication of the first cohort studies [21–27] have progressively changed the point of view of IBD specialists. In fact, a follow-up of the above-mentioned ECCO survey revealed that in 2015 more than 80% of clinicians felt confident prescribing biosimilar mAb [28]. Such confidence was supported by three large prospective studies: The NOR-SWITCH randomized controlled trial [29] found no differences in efficacy, safety or immunogenicity between patients treated with infliximab and those who switched from the originator to CT-P13. The PROSIT-BIO observational study [30] and the PROSIT follow-up study [31] found that effectiveness, safety and immunogenicity in patients taking CT-P13 were similar to previously reported data for infliximab. Finally, a recently published, large comparative equivalence cohort study [32], using a French nationwide health administrative database, found no significant differences in terms of effectiveness and safety between Crohn's disease patients treated with infliximab originator and those treated with CT-P13.

Based on the increasing evidence, knowledge, and experience with biosimilars in IBD, a working group of the Italian Group for the Study of Inflammatory Bowel Disease (IG-IBD) decided to review the literature and to update the Italian position on the use of biosimilars in IBD patients [16]. Before publication, the updated statements were approved by IG-IBD members and by representatives of the national IBD patients' association (AMICI Onlus).

2. Biosimilarity and extrapolation across indications

2.1. Statement 1

A biosimilar mAb targeting the same molecule as the originator can be considered equivalent in terms of efficacy and safety when such equivalence is supported by in vitro assays and clinical studies. Extrapolation across indications is acceptable when a biosimilar mAb has been tested on one or more approved indications and has been approved and licensed by EMA.

The EMA defines a biosimilar of a mAb as a molecule that is highly similar to the reference product in terms of efficacy, safety and immunogenicity [33]. The approval process for biosimilar mAb in the EU is standardized and guarantees the quality of the product evaluated and licensed for market. The process requires robust preclinical evidence of biosimilarity, whereas clinical efficacy can be confirmed in a limited number of clinical trials conducted on the most “sensitive” indications for the reference product. To document any differences in biological activity between the biosimilar and the originator, data from comparative in vitro assays are required. These studies should demonstrate binding to target antigen(s), binding to representative isoforms of the relevant three Fc gamma receptors, Fab-associated functions (e.g. neutralization of a soluble ligand, receptor activation or blockade), and Fc-associated functions. Based on the in vitro data provided, EMA requests in vivo studies in those conditions where in vitro studies cannot investigate possible effects of the biosimilar drug [34]. Finally, the EMA requires that clinical studies be conducted in sensitive clinical models and study conditions, and the model should be relevant in terms of efficacy, safety, and sensitivity to demonstrate comparability in the indication(s) applied for [34].

After this comparability exercise and step-wise approach, EMA authorizes the extrapolation across indications from the most sensitive to all the other approved indications. Data on specific populations, such as children or the elderly, are not required for approval or extrapolation [34]. CT-P13 was approved for all indications based on two clinical trials, one on ankylosing spondylitis [35] and another on rheumatoid arthritis (RA) [36]. The same approach was used for SB2 with clinical trials on RA [37–39], for PF-06438179/GP1111 with a clinical trials on RA [40], for ABP 501 with clinical trials on RA and plaque psoriasis [41,42], for SB5 with clinical trials on RA [43,44], for BI 695501 with a clinical trial on RA [45], for GP2017 with clinical trials on RA and plaque psoriasis [46,47] and for FKB327 with clinical trials on RA [48,49].

Extrapolation has been strongly debated over the past years, and was one of the main concerns in the IBD community to accept biosimilars in their clinical practice. However, extrapolation is not a new concept, and it is based on solid arguments deriving from pre-clinical and clinical evidence [50]. The basic concept supporting the extrapolation of data on CT-P13, SB2, PF-06438179/GP1111, ABP 501, SB5, BI 695501, GP2017 and FKB327 lies in the mechanism of binding to TNF α , which is common for all immune-mediated diseases (e.g. RA, ankylosing spondylitis, IBD, and psoriasis); possible differences between originator and biosimilars related to Fc-region were carefully addressed and included in the comparability exercise [34]. Clinical data from several cohort studies [21–27,30–32] and the NOR-SWITCH randomized controlled trial [29] support the clinical value of extrapolation from other conditions to IBD. In particular, no significant differences in efficacy, effectiveness, safety or immunogenicity were found between IBD patients who were started with infliximab originator and those with CT-P13 [32] or between IBD patients who maintained treatment with infliximab originator and those who switched from infliximab originator to CT-P13 [29]. Therefore, the same approach should be adopted for the extrapolation of adalimumab biosimilars from other conditions

to IBD. The Italian Regulatory Agency (AIFA) approves this approach and considers it sufficient to demonstrate the quality of a licensed biosimilar so to be reimbursed by the National Healthcare System [51].

3. Safety and immunogenicity

3.1. Statement 2

Licensed biosimilars can be considered as safe as the originator. However, large observational studies are needed to monitor the long-term safety of biosimilars, and registries supported by all involved stakeholders should be developed.

3.2. Statement 3

Any event related to the immunogenicity of a mAb cannot be overcome by a biosimilar of the same molecule.

The safety profile of biosimilars has been the main outcome investigated since biosimilars have been launched on the market. Preliminary data from different countries did not show an increased rate of adverse events related to the use of infliximab biosimilars in IBD [21,23–24,26–27]. The NOR-SWITCH trial [29] also revealed no increased risk of adverse events, loss of response or infusion reactions in patients that switched from the originator to CT-P13.

The large “real-life” cohort of 547 IBD patients treated with a biosimilar, namely the PROSIT-BIO cohort [30], had as its primary outcome the safety of CT-P13, defined as the numbers of all adverse events, adverse events leading to discontinuation, and infusion reactions. This study found that the overall safety profile of CT-P13 was no different from what was expected for the reference product. Although 66 adverse events occurred in this study, there was no difference in their frequency or severity between the 313 infliximab-naïve patients treated with CT-P13, the 97 who switched from the originator to CT-P13, and the 139 who had been previously exposed to other anti-TNF α . However, infusion reactions were significantly more frequent in the patients previously exposed to anti-TNF α , with an incidence rate ratio (IRR) of 4.7 vs. naïve ($p < 0.001$) and 3.25 vs. switched ($p < 0.001$). This increased risk was greater in patients previously exposed to infliximab (IRR = 2.82; 95% CI, 1.05–7.9) than in those exposed to other anti-TNF α , indirectly demonstrating cross-reactivity between CT-P13 and the originator. Further expansion of the original PROSIT-BIO cohort confirmed the safety of CT-P13 in 810 IBD patients through a median follow-up of 327 days (IQR, 161–530) [31]. Infusion reactions leading to discontinuation were again significantly more frequent in patients pre-exposed to infliximab (10.7%) than in naïve patients (6.1%, $p = 0.04$ vs. pre-exposed) and in those who switched to CT-P13 (2.6%, $p = 0.008$ vs. pre-exposed). The rates of serious adverse events were 18.5, 74.0 and 48.5 per 100 person-years in the switched group, anti-TNF α naïve and anti-TNF α pre-exposed patients, respectively.

In the French comparative equivalence cohort study, 2551 patients with Crohn's disease were treated with infliximab originator and another 2499 patients were treated with CT-P13 [32]. Multivariate analyses did not reveal significant differences in serious infections (hazard ratio (HR) = 0.82; 95% CI, 0.61–1.11), tuberculosis (HR = 1.10; 95% CI, 0.36–3.34) or cancer (HR = 0.66; 95% CI, 0.33–1.32) between the two treatment groups.

Recently, a systematic review [52] of clinical studies about switching from biological originators to biosimilars (including 48 studies about switching from infliximab) found no differences in terms of safety and immunogenicity.

A recent in vitro study aimed to determine whether antibodies to infliximab cross-react among infliximab originator and two dif-

ferent infliximab biosimilars [53]. This study examined sera from 34 IBD patients who had been treated with the originator or with CT-P13, or had switched from the original to CT-P13, and had subsequently developed anti-infliximab antibodies. The patients' antibodies were found to recognize the originator, CT-P13 and SB2 equally well, irrespective of which molecule a patient had received in treatment. This finding of cross-reactivity suggests that the three molecules are similarly immunogenic. However, safety data for patients exposed to the two biosimilars are missing.

With regard to adalimumab biosimilars, no safety or immunogenicity data are available for IBD patients, but preclinical and clinical data from RA and psoriasis patients show that the safety and immunogenicity profiles do not differ among the five EMA-approved adalimumab biosimilars and the originator [9–13,41–49]. As is the case for infliximab biosimilars, large observational studies are needed to monitor the long-term safety and immunogenicity of these drugs, and the development of registries supported by all involved stakeholders should be encouraged.

4. Interchangeability and switching

4.1. Statement 4

Once biosimilarity has been confirmed, any biosimilar can be considered interchangeable with the reference product.

4.2. Statement 5

Switching from the originator to a biosimilar of the same molecule is acceptable.

Switching from one biosimilar to another of the same originator and multiple switches among different molecules should be avoided in the absence of direct evidence of efficacy and safety.

One of the main concerns since biosimilars have been licensed is the switching of patients from the originator to a biosimilar. In 2013, only 6% of surveyed clinicians considered the originator and the biosimilars interchangeable [15]. There is no agreement on the definition of interchangeability among regulatory agencies. The World Health Organization (WHO) considers interchangeable any “pharmaceutical product . . . which is therapeutically equivalent to a comparator product” [54]. EMA does not have its own definition of interchangeability, but states that the definition should be set at the national level [33]. The Italian Drug Agency (AIFA) adopts the WHO's definition [54]. Thus, in Italy, a biosimilar should be considered interchangeable with its own originator. This position has recently been stated and confirmed by AIFA [51].

According to a systematic review [52], switching from the infliximab originator to an infliximab biosimilar is supported by various studies, especially NOR-SWITCH [29]. This randomized controlled trial enrolled 482 patients already in treatment with infliximab originator for the maintenance of remission of various immune-mediated diseases including Crohn's disease ($n = 155$), ulcerative colitis ($n = 93$), spondyloarthritis, RA, psoriatic arthritis, and chronic plaque psoriasis. Patients were randomized to continue with infliximab originator or to be switched to CT-P13, and followed for 52 weeks. Disease worsening was observed in 53 patients (26.2%) in the infliximab originator group and 61 patients (29.6%) in the CT-P13 group, for an adjusted treatment difference of -4.4% (95% CI, -12.7% to 3.9%). Because this difference was below the pre-established cutoff of 15%, the study concluded that CT-P13 is noninferior to its originator. Among patients with Crohn's disease, worsening occurred in 21.2% and 36.5%, respectively (adjusted risk difference, -14.3%), and for those with ulcerative colitis disease worsening was seen in 9.1% and 11.9%, respectively (adjusted risk

difference, –2.6%). The frequencies of adverse events were similar between the infliximab originator group and the CT-P13 group: all adverse events, 68% vs. 66%; serious adverse events, 10% vs. 9%; and adverse events leading to discontinuation, 4% vs. 3%, respectively.

In the PROSIT-BIO study [30], a subset of 97 patients switched to CT-P13 after a mean of 18 infliximab infusions (SD = 14). No differences in safety or efficacy were observed between these switched patients and the 313 anti-TNF α naïve patients also treated with CT-P13. In the expanded PROSIT-BIO cohort [31], 155 patients switched to CT-P13 after a mean of 17 infliximab infusions (SD = 13). As in the original PROSIT cohort, no significant differences in safety or efficacy were found between switched and naïve patients. Finally, a prospective study of a Swedish cohort of 313 IBD patients switched from the originator to CT-P13 also found no increased risk of adverse events (including infusion reactions) or loss of response [55].

These data have been confirmed by several studies (reviewed in Ref. [52]), leading to the conclusion that switching between the infliximab originator and one of its biosimilars is acceptable and safe, without concerns about efficacy, safety or immunogenicity. With regard to adalimumab biosimilars currently approved by EMA, data on switching from the originator to a biosimilar in IBD patients are currently lacking. However, the different molecules have negligible differences in structure, and preclinical data show essentially the same biological effects as the originator and very similar pharmacological characteristics [9–13]. Moreover, clinical studies of switching in RA patients [43–46,48,49] and psoriasis patients [42,47] also found that the efficacy, safety and immunogenicity profiles do not differ among the five EMA-approved adalimumab biosimilars and the originator. The totality of evidence from preclinical and clinical studies thus supports the concept of switching from adalimumab originator to an adalimumab biosimilar also in IBD patients. In the absence of data from real-life experiences or large clinical studies, the switching of an IBD patient from adalimumab originator to an adalimumab biosimilar should only be done after a clinical evaluation. Other issues that should be considered, when choosing among available adalimumab formulations, are differences in terms of their excipients and administration devices (e.g. citrate versus citrate-free buffer, low-volume 80 mg versus 40 mg pen/syringe) [4,9–13], because they can potentially impact on patients' treatment adherence/persistence.

Clinical studies on the safety or efficacy of switching from one biosimilar to another of the same originator and multiple switch have not yet been reported. Thus the IG-IBD recommends against switching patients from one biosimilar to another biosimilar and multiple switch until clinical data supporting this practice become available. This position is consistent with that of the ECCO [56].

5. Automatic substitution

5.1. Statement 6

Automatic substitution must be avoided. The clinician alone is responsible for the prescription of a biological drug, and no pharmacist or other health care operator may assume this responsibility.

Automatic substitution is a practice that allows a pharmacist or other health care operator to replace a branded drug prescribed by a physician with a generic drug without consulting the prescribing specialist. This practice is common for synthesized drugs but is not applicable to biological drugs that are subject to different regulations, as stated by AIFA [35]. AIFA also states that the decision to prescribe the originator or a biosimilar must be left to the prescribing clinician, who should take into account the affordability of the prescription itself. Therefore, no other stakeholder is

allowed to change the prescription of a biological drug made by a physician, and hospital policies should not limit the free prescription of these drugs. However, we recommend that the availability of certain biosimilars in a hospital be agreed upon between prescribing physicians and hospital administrators, with the purpose of maintaining costs incurred in the use of biological therapies.

6. Patient education about biosimilars

6.1. Statement 7

Patients' awareness about biosimilars should be fostered through education and the provision of up-to-date information, to let them make informed choices. Switching from an originator to a biosimilar should be done only after the patient has received appropriate information and has agreed.

As for any new medicinal product, biosimilars too have raised questions and concerns among patients. Their skepticism may be related to doubts about the effectiveness or safety of the products or about the way they are approved [57]. Therefore, a physician's proposal for therapy with a biosimilar, whether as a first drug or as replacement of an originator, may be viewed negatively by a patient, in particular if their knowledge about biosimilars is low. This reluctance could translate into low acceptance of biosimilar therapy and may even impair treatment outcomes (i.e. nocebo effect) [58]. A recent survey of patients with IBD, RA, psoriasis and certain cancers in the United States and European Union found low levels of awareness about biosimilars [59]. Moreover, the European Federation of Crohn's and Ulcerative Colitis Associations found that only 38% of 1181 surveyed IBD patients had heard of biosimilars, and many of them expressed concerns about these drugs' safety (46.5%) and efficacy (38.6%) [60]. These results suggest that patients should be provided reliable, up-to-date information to help them understand biosimilars and enable them to make informed choices about their treatment options.

7. Conclusions

This paper states the current position of the IG-IBD, a group of Italian physicians specialized in the management of IBD, on the use of biosimilars in Italy. Biosimilars represent a great opportunity for wisely saving money for the National Healthcare System and also create an opportunity for improving patient care, with reinvestment of the saved money (e.g. "gain-share"). When a biosimilar is approved by EMA according to the strict regulations applied to this drug class, we consider it equivalent to its originator. Switching from the originator to a biosimilar is acceptable, because this approach is safe, efficacious and leads to a significant cost reduction for the health care system and subsequently to the possibility of treating more patients. Careful pharmacovigilance should be done for biosimilars as for all biologics. Clinicians are fully responsible for the prescription of biosimilars, and these drugs cannot be automatically substituted by a pharmacist or other health care operator. Until supportive clinical evidence becomes available, cross-switching or multiple switching among biosimilars is not recommended. Finally, patients should be educated about biosimilars to improve their awareness of this drug class to enable them to make informed choices.

Conflicts of interest

Gionata Fiorino: Humanitas Research Hospital and University, Gastroenterology and IBD Center, Rozzano, Italy. Speaker/Consultant/Advisory Board Abbvie, Ferring, Hospira, Janssen, MSD, Takeda, Mundipharma, Pfizer, Tigenix, Biogen, Celltrion, Sandoz, Roche and Amgen.

Flavio Caprioli: Department of Pathophysiology and Transplantation, Università degli Studi di Milano, Italy and Gastroenterology and Endoscopy Unit, Fondazione IRCCS Cà Granda, Ospedale Policlinico di Milano, Milan, Italy. Consultant to: Mundipharma, Abbvie, MSD, Takeda, Janssen, Roche, Celgene, Biogen; lecture fees from Abbvie, Ferring, Takeda, Allergy Therapeutics, Janssen; unrestricted research grants from Giuliani, Sofar, MS&D, Takeda, Abbvie.

Marco Daperno: Division of Gastroenterology, Ospedale Ordine Mauriziano di Torino, Turin, Italy. Advisory Board/Speakers Bureau: Roche, Abbvie, MSD, Takeda, Celltrion, Hospira, Janssen, Pfizer, Mundipharma, Chiesi, SOFAR; Funding (Grants/Honoraria): Quintiles/Bioclinica; Research/Clinical Trials: Takeda, Janssen, Abbvie, Roche, Gilead; Speaker/Consulting Fees: AbbVie, MSD, Chiesi, Takeda, Hospira, Janssen, Pfizer, Mundipharma, Roche.

Filippo Mocciaro: Gastroenterology and Endoscopy Unit, A.R.N.A.S. Civico-Di Cristina-Benfratelli Hospital, Palermo, Italy. Advisory Board member for Abbvie, MSD, Takeda.

Mariabeatrice Principi: Gastroenterology Unit, Department of Organ Transplantation (DETO), Policlinico Universitario, Bari, Italy. Advisory Board member for: Abbvie, MSD, Takeda; lecture fees from Abbvie, MSD, Takeda, Chiesi.

Angelo Viscido: Department of Life, Health and Environmental Sciences. University of L'Aquila, L'Aquila, Italy. Consultant and/or advisory board member for Abbvie, Janssen, Pfizer.

Massimo Claudio Fantini: Department of Systems Medicine, University of Rome "Tor Vergata", Rome, Italy. Advisory boards Abbvie, MSD, Takeda, Janssen; lecture fees Abbvie, Janssen, Takeda, Pfizer, Zambon; research grants Janssen, Pfizer.

Ambrogio Orlando: IBD Unit, A.O. Osp. Riuniti "Villa Sofia-Cervello" Palermo, Italy. Advisory board member for AbbVie, Janssen-Cilag, MSD, Pfizer, Takeda Pharmaceuticals; lecture fees from AbbVie, Chiesi, Janssen-Cilag, MSD, Sofar and Takeda Pharmaceuticals.

Claudio Papi: IBD Unit, S Filippo Neri Hospital Rome, Italy. Consultant and/or educational grants from Abbvie, MSD, Takeda, Chiesi, Sofar, Ferring, Janssen-Cilag and Pfizer.

Vito Annesse: IG-IBD member, Via P. Metastasio 22, 50124 Florence, Italy. Consultant to: Abbvie, Hospira, Mundipharma, MS&D, Takeda, Janssen, Ferring; lecture fees from Abbvie, Ferring, Hospira, MS&D, Takeda, Medtronic, Menarini; unrestricted research grants from Giuliani, Ferring, Sofar, Roche, Gilead, MS&D, Abbvie, Otsuka.

Maurizio Vecchi: Department of Pathophysiology and Transplantation, Università degli Studi di Milano, Italy and Gastroenterology and Endoscopy Unit, Fondazione IRCCS Cà Granda, Ospedale Policlinico di Milano, Milan, Italy. Consultant and/or lecture fees and/or unrestricted research grants from MSD, Hospira, Mundipharma, Takeda, Janssen, Amgen, Biogen, Abbvie, Chiesi, Zambon, Sofar, Giuliani, Alfa-Wasserman, Ferring.

Silvio Danese: Humanitas Research Hospital and University, Gastroenterology and IBD Center, Rozzano, Italy. Consultant and/or advisory board member and/or lecture fees for Abbvie, Ferring, Hospira, Johnson and Johnson, Janssen, Merck, MSD, Takeda, Mundipharma, Pfizer, Tigenix, UCB Pharma, Vifor, Biogen, Celgene, Allergan, Celltrion, Sandoz, and Boehringer Ingelheim.

Fernando Rizzello: IBD unit, DIMEC, University of Bologna, Bologna, Italy. Consulting and/or advisory board and/or lecture fees from Abbvie, Amgen, Takeda, Ferring, Chiesi, Pfizer, MSD, Sofar.

Alessandro Armuzzi: IBD Unit, Presidio Columbus Fondazione Policlinico Universitario A. Gemelli IRCCS – Università Cattolica del Sacro Cuore, Rome, Italy. Consulting and/or advisory board member for AbbVie, Allergan, Amgen, Biogen, Bristol-Myers Squibb, Celgene, Celltrion, Ferring, Hospira, Janssen, Lilly, MSD, Mundipharma, Mylan, Pfizer, Samsung Bioepis, Sofar and Takeda; lecture and/or speaker bureau fees from AbbVie, Amgen, AstraZeneca, Chiesi, Ferring, Hospira, Janssen, Medtronic, MSD, Mitsubishi-Tanabe, Mundipharma, Nikkiso, Otsuka, Pfizer, Samsung Bioepis, Takeda,

Tigenix, Zambon; and unrestricted research grants from MSD, Pfizer and Takeda.

Sandro Ardizzone: Department of Biochemical and Clinical Sciences "L. Sacco" – University of Milan, ASST Fatebenefratelli Sacco, Milan, Italy. Advisory Board member and/or research support from Abbvie, Ferring, Janssen, MSD, Mundipharma, Sofar, Takeda, Zambon.

Livia Biancone: Gastroenterology Unit Department of Systems Medicine, University Tor Vergata, Rome, Italy. Lecture fees and/or Advisory Board member: Abbvie, Alfa-Wassermann, Ferring, MSD, Sofar, Takeda, Zambon.

Giorgia Bodini: IBD Section, Gastroenterology Unit, Department of Internal Medicine, University of Genoa, Genoa, Italy; Lecture fees from Abbvie, Janssen, MSD.

Fabrizio Bossa: UOC di Gastroenterologia ed Endoscopia Digestiva Ospedale Casa Sollievo della sofferenza IRCCS – San Giovanni Rotondo, Italy. Advisory board member for Janssen and MSD; Lecture fees from MSD, Mundipharma.

Emma Calabrese: Gastroenterology Unit Department of Systems Medicine, University Tor Vergata, Rome, Italy. Lecture fees from Abbvie, Janssen.

Maria Cappello: Ambulatorio IBD, UOC Gastroenterologia ed Epatologia Policlinico di Palermo, Palermo, Italy. Advisory Board member for Abbvie, Janssen, MSD, Shire, Takeda; lecture fees from Abbvie, Chiesi, MSD, Shire, Takeda.

Fabiana Castiglione: Gastroenterology, Department of Clinical Medicine and Surgery, School of Medicine "Federico II" of Naples, Naples, Italy. Advisory Board member and/or Lecture fees from AbbVie, Chiesi, Ferring, MSD, Sofar, Takeda, Janssen, and Zambon.

Francesco Costa: U.O.C. di Gastroenterologia Universitaria – AOUP, Pisa, Italy. Advisory Board member for Abbvie, Amgen, Janssen, MSD, Otsuka, Takeda; Lecture fees from Abbvie, Diasorin, Ferring, MSD, Takeda, ThermoFisher, Zambon.

Salvatore Cucchiara: Pediatric Gastroenterology, Liver and Digestive Endoscopy Unit, Department of Pediatrics and Childhood Neuropsychiatry, Umberto I Hospital – Sapienza University, Rome, Italy; Advisory Board member for Johnson and Johnson, International Registry of Biological Therapy for Pediatric IBD. Vedolizumab Pediatric Program Advisory Board Takeda; unrestricted research grants from DEVELOP Pediatric IBD Registry – Member – Janssen Scientific Affairs, LLC, personal fees from Advisory Boards of Alfa Wasserman, Sanofi.

Renata D'Incà: US Malattie Infiammatorie Croniche dell'Intestino, UOC Gastroenterologia, Azienda Ospedaliera di Padova, Padua, Italy. Lectures fees from Abbvie, Hospira, MSD, Mundipharma, Takeda.

Walter Fries: UOSD Malattie Intestinali Croniche, Dipartimento di Medicina Clinica e Sperimentale, Università di Messina, Messina, Italy. Advisory Board member for Abbvie, MSD, Mundipharma; lecture fees from Takeda, Zambon; educational grant from MSD.

Paolo Gionchetti: IBD unit, DIMEC, University of Bologna, Bologna, Italy; Advisory Board member for Abbvie, Alfa Wassermann, Amgen, Biogen, Celgene, Hospira, Janssen, MSD, Pfizer, Takeda. Lecture fees from Abbvie, Alfa Wassermann, Chiesi, Ferring, MSD, Sofar, Takeda.

Laurino Grossi: Scienze Digestive Ospedale Spirito Santo Pescara, Università G. d'Annunzio Chieti-Pescara, Italy. Advisory Board member for Abbvie, Takeda.

Luisa Guidi: UOS Day Hospital, UOC Medicina Interna e Malattie dell'Apparato Digerente, Presidio Columbus Fondazione Policlinico Universitario A. Gemelli IRCCS – Università Cattolica del Sacro Cuore, Rome, Italy. Consultant and/or lecture fees for AbbVie, Janssen, MSD, Mundipharma, Shire, Takeda, Vifor Pharma.

Giovanni Maconi: Gastroenterology Unit, Department of Biochemical and Clinical Science "L. Sacco" – University of Milan ASST Fatebenefratelli Sacco – Milan, Italy; Consulting and advisory board

fees and/or research support from AbbVie, Janssen, Takeda, Allergan, Alfa Sigma, THD, Novartis.

Silvia Mazzuoli: UOC Gastroenterologia e Nutrizione, Ospedale San Nicola Pellegrino, Trani, Italy; Advisory boards for AbbVie, MSD, Mundipharma, Takeda.

Sara Renna: IBD Unit, “Villa Sofia-Cervello” Hospital, Palermo, Italy. Advisory Board member for AbbVie, Janssen, MSD; Lecture fees from AbbVie, MSD, Takeda.

Simone Saibeni: Gastroenterology Unit, Rho Hospital, ASST Rhodense, Rho, Italy; Lecture fees from Takeda; Advisory Board member for AbbVie and Janssen.

Edoardo Savarino: Department of Surgery, Oncology and Gastroenterology – DiSCOG, University of Padua, Padua, Italy; Lecture/Consulting fees from AbbVie, Janssen, Malesci, Medtronic, MSD, Sofar, Takeda.

Maria Lia Scribano: UOC Gastroenterologia ed Endoscopia Digestiva Diagnostica e Operativa, A.O. San Camillo-Forlanini, Rome, Italy; Advisory board member and/or speaker fees for AbbVie, Janssen, Mundipharma, Pfizer, Takeda.

Anna Testa: Gastroenterology, Department of Clinical Medicine and Surgery, School of Medicine “Federico II” of Naples, Naples, Italy; Advisory Boards for AbbVie, Takeda.

Angela Variola, UOC Gastroenterologia- IBD Unit-IRCCS Ospedale Sacro Cuore Don Calabria, Negrar, Italy; Advisory Boards for MSD, Pfizer, Takeda.

Acknowledgement

The authors are indebted to Valerie Matarese for editing the manuscript.

Appendix A.

Collaborators – National patients’ association representatives

Salvatore Leone, Director of “Associazione nazionale per le Malattie Infiammatorie Croniche dell’Intestino” – A.M.I.C.I. ONLUS.

Enrica Previtali, President of “Associazione nazionale per le Malattie Infiammatorie Croniche dell’Intestino” – A.M.I.C.I. ONLUS.

Collaborators – IG-IBD members

Marina Aloï, Pediatric Gastroenterology Unit, Department of Pediatrics and Childhood Neuropsychiatry, Umberto I Hospital – Sapienza University, Rome, Italy.

Patrizia Alvisi, Pediatric Gastroenterology Unit, Maggiore Hospital, Bologna, Italy.

Elisabetta Antonelli, Gastroenterology Unit, Perugia General Hospital, Perugia, Italy.

Sandro Ardizzone, Department of Biochemical and Clinical Sciences “L. Sacco” – University of Milan, ASST Fatebenefratelli Sacco, Milan, Italy.

Marco Astegiano, Gastroenterology Unit, Department of General and Specialist Medicine, Città della Salute e della Scienza di Torino, Turin, Italy.

Monia Baldoni, Dipartimento di Medicina, Sez. Gastroenterologia Ospedale di Perugia – Università di Perugia, Perugia, Italy.

Marina Beltrami, SOC Medicina Gastroenterologica/SOSD Degenza breve Internistica e Centro MICI Arcispedale S. Maria Nuova – IRCCS, Reggio Emilia, Italy.

Livia Biancone, Gastroenterology Unit, Department of Systems Medicine, Tor Vergata University, Rome, Italy.

Giorgia Bodini, IBD Section, Gastroenterology Unit, Department of Internal Medicine, University of Genoa, Genoa, Italy.

Andrea Buda, Gastroenterology Unit, S. Maria del Prato Hospital, Feltre, Italy.

Fabrizio Bossa, UOC di Gastroenterologia ed Endoscopia Digestiva Ospedale Casa Sollievo della sofferenza IRCCS – San Giovanni Rotondo, Italy.

Fiammetta Bracci, UO Epatogastroenterologia e Nutrizione, Ospedale Pediatrico Bambino Gesù, Rome, Italy.

Emma Calabrese, Gastroenterology Unit, Department of Systems Medicine, Tor Vergata University, Rome, Italy.

Maria Cappello, Ambulatorio IBD, UOC Gastroenterologia ed Epatologia Policlinico di Palermo, Palermo, Italy.

Fabiana Castiglione, Gastroenterology, Department of Clinical Medicine and Surgery, School of Medicine “Federico II” of Naples, Naples, Italy.

Carolina Ciacci, Gastroenterologia AOU San Giovanni di Dio e Ruggi, Dipartimento di Medicina, Chirurgia e Odontoiatria, Scuola Medica Salernitana, Università di Salerno, Salerno, Italy.

Michele Cicala, UOC di Gastroenterology and Digestive Endoscopy, University “Campus Bio Medico di Roma”, Rome, Italy.

Rachele Ciccocioppo, Gastroenterology Unit, Department of Medicine, AOUI Policlinico G.B. Rossi & University of Verona, Verona, Italy.

Michele Comberlato, Department of Gastroenterology and Digestive Endoscopy Central Hospital, Bolzano, Italy.

Claudio Camillo Cortelezzi, SC di Gastroenterologia ed Endoscopia Digestiva ASST Settelaghi – Varese, Italy.

Rocco Cosentino, UOS M.I.C.I., UOC Gastroenterologia ed Endoscopia Digestiva diagnostica ed operativa, A.O. San Camillo-Forlanini, Rome, Italy.

Francesco Costa, U.O.C. di Gastroenterologia Universitaria – AOUP, Pisa, Italy.

Giuseppe Costantino, UOSD Malattie Intestinali Croniche, Dipartimento di Medicina Clinica e Sperimentale, Università di Messina, Messina, Italy.

Salvatore Cucchiara, Pediatric Gastroenterology, Liver and Digestive Endoscopy Unit, Department of Pediatrics and Childhood Neuropsychiatry, Umberto I Hospital – Sapienza University, Rome, Italy.

Antonio Cuomo, UOD Gastroenterologia ed Endoscopia Digestiva, DEA I Livello Umberto I, Nocera Inferiore, Italy.

Renata D’Inca, US Malattie Infiammatorie Croniche dell’Intestino, UOC Gastroenterologia, Azienda Ospedaliera di Padova, Padua, Italy.

Maria Carla Di Paolo, UOC Gastroenterologia ed Endoscopia Digestiva, Azienda Ospedaliera San Giovanni Addolorata, Rome, Italy.

Antonio Di Sabatino, First Department of Medicine, San Matteo Hospital Foundation, University of Pavia, Pavia, Italy.

Antonio Di Sario, Clinica di Gastroenterologia, Università Politecnica delle Marche, Ancona, Italy.

Giuseppe Frieri, UOSD. Gastroenterologia D.U. – PO L’Aquila, Università degli Studi dell’Aquila, L’Aquila, Italy.

Walter Fries, UOSD Malattie Intestinali Croniche, Dipartimento di Medicina Clinica e Sperimentale, Università di Messina, Messina, Italy.

Antonio Gasbarrini, Direttore Area Gastroenterologia e Oncologia, Fondazione Policlinico Universitario A. Gemelli IRCCS – Università Cattolica del Sacro Cuore, Rome, Italy.

Andrea Geccherle, UOC Gastroenterologia- IBD Unit-IRCCS Ospedale Sacro Cuore Don Calabria, Negrar, Italy.

Paolo Gionchetti, IBD Unit, DIMEC, University of Bologna, Bologna, Italy.

Maria Giovanna Graziani, UOC Gastroenterologia ed Endoscopia Digestiva, Azienda Ospedaliera San Giovanni Addolorata, Rome, Italy.

Laurino Grossi, Scienze Digestive Ospedale Spirito Santo Pescara, Università G. d'Annunzio Chieti-Pescara, Italy.

Luisa Guidi, UOS Day Hospital, UOC Medicina Interna e Malattie dell'Apparato Digerente, Presidio Columbus Fondazione Policlinico Universitario A. Gemelli IRCCS – Università Cattolica del Sacro Cuore, Rome, Italy.

Gianni Imperiali, DUOS Malattie Infiammatorie Croniche Intestinali, Gastroenterologia, Ospedale Valduce, Como, Italy.

Giovanni Latella, Gastroenterology Unit, Department of Life, Health and Environmental Sciences, University of L'Aquila, L'Aquila, Italy.

Paolo Lionetti, Department NEUROFARBA, University of Florence, Gastroenterology and Nutrition Unit, Meyer Children's Hospital, Florence, Italy.

Gaetano Inserra, Dipartimento di Medicina Clinica e Sperimentale, Università di Catania, Catania, Italy.

Giovanni Maconi, Gastroenterology Unit, Department of Biochemical and Clinical Science "L. Sacco" – University of Milan ASST Fatebenefratelli Sacco – Milan, Italy.

Francesco Manguso, UOC di Gastroenterologia, A.O.A. Cardarelli di Napoli, Naples, Italy.

Marco Marino, Department for Gastroenterology and Digestive Endoscopy, Academic Center of Udine, Udine, Italy.

Mauro Mastronardi, IBD Unit, UOC Gastroenterologia 2, IRCCS "S. De Bellis", Castellana Grotte, Italy.

Silvia Mazzuoli, UOC Gastroenterologia e Nutrizione, Ospedale San Nicola Pellegrino, Trani, Italy.

Gianmichele Meucci, UO di Gastroenterologia ed Endoscopia Digestiva, Ospedale San Giuseppe, Milan, Italy.

Marco Mendolaro, Division of Gastroenterology, Ospedale Ordine Mauriziano di Torino, Turin, Italy.

Monica Milla, Centro di Riferimento Regionale Malattie Infiammatorie Croniche Intestinali, SOD Gastroenterologia Clinica, AOU Careggi, Florence, Italy.

Giammarco Mocci, SC Gastroenterologia, Ospedale Brotzu, Cagliari, Italy.

Giovanni Monteleone, Gastroenterology Unit, Department of Systems Medicine, Tor Vergata University, Rome, Italy.

Francesco Neri Bortoluzzi, Gastroenterology Unit, Ulss 12 Seregnissima, Venice, Italy.

Cristiano Pagnini, UOC Gastroenterologia ed Endoscopia Digestiva, Azienda Ospedaliera San Giovanni Addolorata, Rome, Italy.

Luca Pastorelli, Gastroenterology Unit, IRCCS Policlinico San Donato, San Donato Milanese, Italy.

Roberta Pica, Centro MICI, UOS Gastroenterologia Territoriale, Ospedale Sandro Pertini, Rome, Italy.

Simona Piergallini, Gastroenterology and Digestive Endoscopy Unit, "Murri" Hospital, Polytechnic University of Marche, Fermo, Italy.

Antonello Privitera, IBD Unit, Azienda Ospedaliera per l'Emergenza "Cannizzaro", Catania, Italy.

Sara Renna, IBD Unit, "Villa Sofia-Cervello" Hospital, Palermo, Italy.

Davide Giuseppe Ribaldone, Division of Gastroenterology, Department of Medical Sciences, University of Turin, Turin, Italy.

Chiara Ricci, Gastroenterology Section, 1st Medical Clinic, Spedali Civili and University of Brescia, Brescia, Italy.

Antonio Rispo, Gastroenterology, Department of Clinical Medicine and Surgery, School of Medicine "Federico II" of Naples, Naples, Italy.

Rodolfo Rocca, Division of Gastroenterology, Ospedale Ordine Mauriziano di Torino, Turin, Italy.

Claudio Romano, Pediatric Gastroenterology Unit, Department, "G. Martino's Hospital", University of Messina, Messina, Italy.

Marco Romano, Dipartimento di Medicina di Precisione ed UP di Endoscopia e Patologie Infiammatorie Croniche Gastrointestinali,

Università della Campania 'Luigi Vanvitelli' ed Azienda Ospedaliera Universitaria, Naples, Italy.

Giovanni Russo, Gastroenterologia ed Endoscopia Digestiva, ASL di Spezzino, La Spezia, Italy.

Renato Sablich, IBD Unit, SC Gastroenterologia, AAS5-Friuli Occidentale – Ospedale di Pordenone, Pordenone, Italy.

Simone Saibeni, Gastroenterology Unit, Rho Hospital, ASST Rhodense, Rho, Italy.

Edoardo Savarino, Department of Surgery, Oncology and Gastroenterology – DISCOG, University of Padua, Padua, Italy.

Maria Lia Scribano, UOC Gastroenterologia ed Endoscopia Digestiva Diagnostica e Operativa, A.O. San Camillo-Forlanini, Rome, Italy.

Rocco Spagnuolo, IBD Unit, Dipartimento di "Scienze Mediche e Chirurgiche", Magna Graecia University, Catanzaro, Italy.

Elisa Stasi, Gastroenterology and Digestive Endoscopy, National Institute of Gastroenterology, "S. De Bellis" Research Hospital, Castellana Grotte, Italy.

Maria Maddalena Terpin, UOC Gastroenterologia ed Endoscopia Digestiva – ASST Ovest Milanese, Legnano, Italy.

Anna Testa, Gastroenterology, Department of Clinical Medicine and Surgery, Federico II School of Medicine, Naples, Italy.

Daniela Valpiani, U.O. Gastroenterologia ed Endoscopia digestiva, Azienda Romagna Ospedale di Forlì, Forlì, Italy.

Angela Variola, UOC Gastroenterologia- IBD Unit-IRCCS Ospedale Sacro Cuore Don Calabria, Negrar, Italy.

Piero Vernia, Gastroenterology Unit, Sapienza University of Rome, Rome, Italy.

Giovanna Vitale, UOC Gastroenterologia e Endoscopia Digestiva, Azienda Ospedaliera Regionale 'San Carlo', Potenza, Italy.

Giorgio Zoli, Direttore Dipartimento di Medicina e Medicina Interna, Centro per la Diagnosi e Cura Malattie dell'Intestino, Ospedale SS. Annunziata, Via Vicini 2, Cento, Italy.

References

- [1] World Health Organization. Guidelines on evaluation of similar biotechnological products (SBPs) Expert Committee on Biological Standardization; 2009 https://www.who.int/biologicals/areas/biological_therapeutics/BIOETHERAPEUTICS.FOR.WEB.22APRIL2010.pdf.
- [2] European Medicine Agency, European Commission. Biosimilars in the EU: information guide for healthcare professionals; 2017 <https://www.ema.europa.eu/documents/leaflet/biosimilars-eu-information-guide-healthcare-professionals.en.pdf>.
- [3] <https://www.ema.europa.eu/medicines/human/EPAR/remicade>.
- [4] <https://www.ema.europa.eu/medicines/human/EPAR/humira>.
- [5] <https://www.ema.europa.eu/medicines/human/EPAR/inflectra>.
- [6] <https://www.ema.europa.eu/medicines/human/EPAR/remsimia>.
- [7] <https://www.ema.europa.eu/medicines/human/EPAR/flixabi>.
- [8] <https://www.ema.europa.eu/en/medicines/human/EPAR/zessly>.
- [9] <https://www.ema.europa.eu/en/medicines/human/EPAR/amgevita/solymbic>.
- [10] <https://www.ema.europa.eu/en/medicines/human/EPAR/imraldi>.
- [11] <https://www.ema.europa.eu/medicines/human/EPAR/hulio>.
- [12] <https://www.ema.europa.eu/en/medicines/human/EPAR/hyrimoz>.
- [13] <https://www.ema.europa.eu/medicines/human/EPAR/cyltezo>.
- [14] Danese S, Bonovas S, Peyrin-Biroulet L. Biosimilars in IBD: from theory to practice. *Nat Rev Gastroenterol Hepatol* 2017;14:22–31.
- [15] Danese S, Fiorino G, Michetti P. Viewpoint: knowledge and viewpoints on biosimilar monoclonal antibodies among members of the European Crohn's and Colitis Organization. *J Crohns Colitis* 2014;8:1548–50.
- [16] Annesse V, Vecchi M. Use of biosimilars in inflammatory bowel disease: statements of the Italian Group for Inflammatory Bowel Disease. *Dig Liver Dis* 2014;46:963–8.
- [17] Argüelles-Arias F, Barreiro-de-Acosta M, Carballo F, et al. Joint position statement by Spanish Society of Gastroenterology and Spanish Society of Pharmacology on biosimilar therapy for inflammatory bowel disease. *Rev Esp Enferm Dig* 2013;105:37–43.
- [18] Danese S, Gomollon F. ECCO position statement: the use of biosimilar medicines in the treatment of inflammatory bowel disease (IBD). *J Crohns Colitis* 2013;7:586–9.
- [19] Fiorino G, Girolomoni G, Lapadula G, et al. The use of biosimilars in immune-mediated diseases: a joint Italian Society of Rheumatology (SIR), Italian Society of Dermatology (SIdMaST), and Italian Group of Inflammatory Bowel Disease (IG-IBD) position paper. *Autoimmun Rev* 2014;13:751–5.

- [20] Vermeire S, Louis E, Dewit O, et al. Clinical and scientific aspects related to biosimilars in inflammatory bowel diseases (IBD): position document of the Belgian IBD Research & Development Group (BIRD). *Acta Gastroenterol Belg* 2015;78:26–9.
- [21] Farkas K, Rutka M, Balint A, et al. Efficacy of the new infliximab biosimilar CT-P13 induction therapy in Crohn's disease and ulcerative colitis – experiences from a single center. *Expert Opin Biol Ther* 2015;15:1257–62.
- [22] Farkas K, Rutka M, Golovics PA, et al. Efficacy of infliximab biosimilar CT-P13 induction therapy on mucosal healing in ulcerative colitis. *J Crohns Colitis* 2016;10:1273–8.
- [23] Kang YS, Moon HH, Lee SE, et al. Clinical experience of the use of CT-P13, a biosimilar to infliximab in patients with inflammatory bowel disease: a case series. *Dig Dis Sci* 2015;60:951–6.
- [24] Park SH, Kim YH, Lee JH, et al. Post-marketing study of biosimilar infliximab (CT-P13) to evaluate its safety and efficacy in Korea. *Expert Rev Gastroenterol Hepatol* 2015;9(Suppl. 1):35–44.
- [25] Siczekowska J, Jarzebicka J, Banaszekiewicz A, et al. Switching between infliximab originator and biosimilar in paediatric patients with inflammatory bowel disease. Preliminary observations. *J Crohns Colitis* 2016;10:127–32.
- [26] Smits LJ, Derikx LA, de Jong DJ, et al. Clinical outcomes following a switch from remicade(R) to the biosimilar CT-P13 in inflammatory bowel disease patients: a prospective observational cohort study. *J Crohns Colitis* 2016;10:1287–93.
- [27] Gecse KB, Lovasz BD, Farkas K, et al. Efficacy and safety of the biosimilar infliximab CT-P13 treatment in inflammatory bowel diseases: a prospective, multicentre, nationwide cohort. *J Crohns Colitis* 2016;10:133–40.
- [28] Danese S, Fiorino G, Michetti P. Changes in biosimilar knowledge among European Crohn's Colitis Organization [ECCO] members: an updated survey. *J Crohns Colitis* 2016;10:1362–5.
- [29] Jorgensen KK, Olsen IC, Goll GL, et al. Switching from originator infliximab to biosimilar CT-P13 compared with maintained treatment with originator infliximab (NOR-SWITCH): a 52-week, randomised, double-blind, non-inferiority trial. *Lancet* 2017;389:2304–16.
- [30] Fiorino G, Manetti N, Armuzzi A, et al. The PROSIT-BIO cohort: a prospective observational study of patients with inflammatory bowel disease treated with infliximab biosimilar. *Inflamm Bowel Dis* 2017;23:233–43.
- [31] Armuzzi A, Fiorino G, Variola A, et al. The PROSIT cohort of infliximab biosimilar in IBD: a prolonged follow-up on the effectiveness and safety across Italy. *Inflamm Bowel Dis* 2018;(August), <http://dx.doi.org/10.1093/ibd/izy264>.
- [32] Meyer A, Rudant J, Drouin J, et al. Effectiveness and safety of reference infliximab and biosimilar in crohn disease: a French equivalence study. *Ann Intern Med* 2019;170:99–107.
- [33] European Medicine Agency. Biosimilar medicines. <https://www.ema.europa.eu/human-regulatory/overview/biosimilar-medicines>.
- [34] European Medicine Agency. Guideline on similar biological medicinal products containing monoclonal antibodies—non-clinical and clinical issues (EMA/CHMP/BMWP/403543/2010 Committee for Medicinal Products for Human Use); 2012.
- [35] Park W, Hrycaj P, Jeka S, et al. A randomised, double-blind, multicentre, parallel-group, prospective study comparing the pharmacokinetics, safety, and efficacy of CT-P13 and innovator infliximab in patients with ankylosing spondylitis: the PLANETAS study. *Ann Rheum Dis* 2013;72:1605–12.
- [36] Yoo DH, Hrycaj P, Miranda P, et al. A randomised, double-blind, parallel-group study to demonstrate equivalence in efficacy and safety of CT-P13 compared with innovator infliximab when coadministered with methotrexate in patients with active rheumatoid arthritis: the PLANETRA study. *Ann Rheum Dis* 2013;72:1613–20.
- [37] Smolen JS, Choe JY, Prodanovic N, et al. Safety, immunogenicity and efficacy after switching from reference infliximab to biosimilar SB2 compared with continuing reference infliximab and SB2 in patients with rheumatoid arthritis: results of a randomised, double-blind, phase III transition study. *Ann Rheum Dis* 2018;77:234–40.
- [38] Smolen JS, Choe JY, Prodanovic N, et al. Comparing biosimilar SB2 with reference infliximab after 54 weeks of a double-blind trial: clinical, structural and safety results. *Rheumatology (Oxford)* 2017;56:1771–9.
- [39] Choe JY, Prodanovic N, Niebrzydowski J, et al. A randomised, double-blind, phase III study comparing SB2, an infliximab biosimilar, to the infliximab reference product Remicade in patients with moderate to severe rheumatoid arthritis despite methotrexate therapy. *Ann Rheum Dis* 2017;76:58–64.
- [40] Cohen SB, Alten R, Kameda H, et al. A randomized controlled trial comparing PF-06438179/GP1111 (an infliximab biosimilar) and infliximab reference product for treatment of moderate to severe active rheumatoid arthritis despite methotrexate therapy. *Arthritis Res Ther* 2018;20:155.
- [41] Cohen S, Genovese MC, Choy E, et al. Efficacy and safety of the biosimilar ABP 501 compared with adalimumab in patients with moderate to severe rheumatoid arthritis: a randomised, double-blind, phase III equivalence study. *Ann Rheum Dis* 2017;76:1679–87.
- [42] Papp K, Bachelez H, Costanzo A, et al. Clinical similarity of biosimilar ABP 501 to adalimumab in the treatment of patients with moderate to severe plaque psoriasis: a randomized, double-blind, multicenter, phase III study. *J Am Acad Dermatol* 2017;76:1093–102.
- [43] Weinblatt ME, Baranaukaite A, Niebrzydowski J, et al. Phase III randomized study of SB5, an adalimumab biosimilar, versus reference adalimumab in patients with moderate-to-severe rheumatoid arthritis. *Arthritis Rheumatol* 2018;70:40–8.
- [44] Weinblatt ME, Baranaukaite A, Dokoupilova E, et al. Switching from reference adalimumab to SB5 (adalimumab biosimilar) in patients with rheumatoid arthritis: fifty-two-week phase III randomized study results. *Arthritis Rheumatol* 2018;70:832–40.
- [45] Cohen SB, Alonso-Ruiz A, Klimiuk PA, et al. Similar efficacy, safety and immunogenicity of adalimumab biosimilar BI 695501 and Humira reference product in patients with moderately to severely active rheumatoid arthritis: results from the phase III randomised VOLTAIRE-RA equivalence study. *Ann Rheum Dis* 2018;77:914–21.
- [46] Wiland P, Jeka S, Dokoupilova E, et al. A randomized, double-blind, parallel-group, multicenter study to compare the efficacy, safety and immunogenicity of a proposed adalimumab biosimilar (GP2017) with reference adalimumab in patients with moderate-to-severe active rheumatoid arthritis. *Arthritis Rheumatol* 2018;70(Suppl. 10):A1936.
- [47] Blauvelt A, Lacour JP, Fowler Jr JF, et al. Phase III randomized study of the proposed adalimumab biosimilar GP2017 in psoriasis: impact of multiple switches. *Br J Dermatol* 2018;179:623–31.
- [48] Genovese MC, Glover J, Matsunaga N, et al. Efficacy, safety and immunogenicity in randomized double-blind (DB) and open-label extension (OLE) studies comparing FKB327, an adalimumab biosimilar, with the adalimumab reference product (Humira®; RP) in patients (pts) with active rheumatoid arthritis (RA). *Arthritis Rheumatol* 2017;69(Suppl. 10):2.
- [49] Alten R, Glover J, Matsunaga N, et al. Efficacy and safety results of a phase III study comparing FKB327, an adalimumab biosimilar, with the adalimumab reference product in patients with active rheumatoid arthritis. *Ann Rheum Dis* 2017;76:59.
- [50] Weise M, Kurki P, Wolff-Holz E, et al. Biosimilars: the science of extrapolation. *Blood* 2014;124:3191–6.
- [51] Agenzia Italiana del Farmaco. Secondo Position Paper AIFA sui Farmaci Biosimilari; 2018 <http://www.agenziafarmaco.gov.it/sites/default/files/2.Position-Paper-AIFA-Farmaci-Biosimilari.pdf>.
- [52] Cohen HP, Blauvelt A, Rifkin RM, Danese S, Gokhale SB, Woollett G. Switching reference medicines to biosimilars: a systematic literature review of clinical outcomes. *Drugs* 2018;78:463–78.
- [53] Fiorino G, Ruiz-Arguello MB, Maguregui A, et al. Full interchangeability in regard to immunogenicity between the infliximab reference biologic and biosimilars CT-P13 and SB2 in inflammatory bowel disease. *Inflamm Bowel Dis* 2018;24:601–6.
- [54] World Health Organisation. Multisource (Generic) pharmaceutical products: guidelines on registration requirements to establish interchangeability World Health Organization WHO technical report series. 937 ed; 2006.
- [55] Bergqvist V, Kadivar M, Molin D, et al. Switching from originator infliximab to the biosimilar CT-P13 in 313 patients with inflammatory bowel disease. *Ther Adv Gastroenterol* 2018;11, 1756284818801244.
- [56] Danese S, Fiorino G, Raine T, et al. ECCO position statement on the use of biosimilars for inflammatory bowel disease—an update. *J Crohns Colitis* 2017;11:26–34.
- [57] Skingle D. Biosimilars: what do patients need to consider? *RMD Open* 2015;1:e000141.
- [58] Pouillon L, Socha M, Demore B, et al. The nocebo effect: a clinical challenge in the era of biosimilars. *Expert Rev Clin Immunol* 2018;14:739–49.
- [59] Jacobs I, Singh E, Sewell KL, Al-Sabbagh A, Shane LG. Patient attitudes and understanding about biosimilars: an international cross-sectional survey. *Patient Prefer Adherence* 2016;10:937–48.
- [60] Peyrin-Biroulet L, Lonnfors S, Roblin X, Danese S, Avedano L. Patient perspectives on biosimilars: a survey by the European federation of Crohn's and ulcerative colitis associations. *J Crohns Colitis* 2017;11:128–33.