




ELSEVIER

Contents lists available at ScienceDirect

Neurotoxicology

journal homepage: www.elsevier.com/locate/neuro

Full Length Article

Adolescent methylmercury exposure: Behavioral mechanisms and effects of sodium butyrate in mice

Steven R. Boomhower^{a,*}, M. Christopher Newland^b^a Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, MA, USA^b Department of Psychology, Auburn University, Auburn, AL, USA

ARTICLE INFO

Keywords:

Adolescence
Fixed ratio
Lysine deacetylase inhibitor
Mathematical principles of reinforcement
Methylmercury
Sodium butyrate

ABSTRACT

Methylmercury (MeHg), an environmental neurotoxicant primarily found in fish, produces neurobehavioral impairment when exposure occurs during gestation. Whether other developmental periods, such as adolescence, display enhanced vulnerability to the behavioral effects of MeHg exposure is only beginning to be explored. Further, little is known about the effects of repeated administration of lysine deacetylase inhibitors, such as sodium butyrate (NaB), on operant behavior. In Experiment 1, male C57BL6/n mice were exposed to 0, 0.3, and 3.0 ppm MeHg ($n = 12$ each) via drinking water from postnatal days 21 to 60 (murine adolescence). As adults, mice were trained to lever press under an ascending series of fixed-ratio schedules of milk reinforcement selected to enable the analysis of three important parameters of operant behavior using the framework provided by Mathematical Principles of Reinforcement. Adolescent MeHg exposure dose-dependently increased saturation rate, a measure of the retroactive reach of a reinforcer, and decreased minimum response time relative to controls. In Experiment 2, the behavioral effects of repeated NaB administration both alone and following adolescent MeHg exposure were examined. Male C57BL6/n mice were given either 0 or 3.0 ppm MeHg during adolescence and, before behavioral testing, two weeks of once daily i.p. injections of saline or 0.6 g/kg NaB ($n = 12$ in each cell). Adolescent MeHg exposure again increased saturation rate but did not significantly alter minimum response time. NaB also increased saturation rate in both MeHg exposure groups. These data suggest that the behavioral mechanisms of adolescent MeHg exposure and NaB may be related to the impact of reinforcement on prior responses. Specifically, MeHg and NaB concentrated the effects of reinforcers onto the most recent responses.

1. Introduction

Adolescence is a time of continued neurobehavioral development in both humans and nonhumans. The density of gray matter decreases in the human frontal cortex (Giedd et al., 1999), and dopamine-receptor pruning occurs in the rodent midbrain (Teicher et al., 1995). In rodents, adolescence ranges between postnatal days 21 and 60 (Spear, 2000) and marks a period of dysregulated choice and sensitivity to reinforcers. Adolescent rodents perform poorly on measures of executive function relative to adult conspecifics in that they prefer small-immediate reinforcers over larger-delayed ones (Doremus-Fitzwater et al., 2012; Pinkston and Lamb, 2011) and demonstrate impaired reversal learning and extradimensional shifting (Newman and McGaughy, 2011). Similar patterns of behavior related to executive-function dysregulation occur in human adolescents (Chambers and Potenza, 2003; Olson et al.,

2007). The neurobehavioral profile of adolescence may escalate vulnerability to neurotoxicants and other chemicals that alter brain function. Thus, understanding how neurotoxic perturbations during adolescence affect behavior in adulthood is important to public health.

Methylmercury (MeHg) is an environmental neurotoxicant found in fish and is a significant public-health concern (National Research Council, 2000). In animal models, gestational exposure to doses of MeHg that encompass human exposures impairs reversal learning and choice in adulthood (Newland et al., 2008, 2004; Reed et al., 2006), suggesting deficiencies in executive functioning (Newland et al., 2015). However, the long-term behavioral effects of MeHg exposure in adolescence remain under explored. This is especially troubling since human adolescents both consume fish known to contain mercury and have higher blood-mercury concentrations compared to younger age groups (Nielsen et al., 2015; Sichert-Hellert et al., 2009). In mice, MeHg

* Corresponding author at: Department of Environmental Health, 665 Huntington Ave, Bldg 1, Room 317, Harvard T.H. Chan School of Public Health, Boston, MA, USA.

E-mail address: sboomhower@hsph.harvard.edu (S.R. Boomhower).

<https://doi.org/10.1016/j.neuro.2018.10.011>

Received 19 June 2018; Received in revised form 25 September 2018; Accepted 27 October 2018

Available online 29 October 2018

0161-813X/ © 2018 Elsevier B.V. All rights reserved.

exposure during adolescence increases the number of trials required to transition through a spatial-discrimination reversal and extradimensional shift in adulthood (Boomhower and Newland, 2017). Adolescent MeHg exposure also dose-dependently reduces choice for a larger reinforcer relative to a smaller reinforcer in adult mice (Boomhower and Newland, 2016). These experiments suggest that MeHg exposure during adolescence can produce long-lasting behavioral effects related to choice in adulthood, but the behavioral mechanisms underlying these effects remain unclear. For example, adolescent MeHg exposure may alter reinforcer efficacy as with gestational exposures (Newland et al., 2015), or it may change some other aspect of reinforcement or behavior.

Mathematical Principles of Reinforcement (MPR) offers one way to characterize the behavioral mechanisms underlying chemical exposure (Killeen, 1994; Killeen and Sitomer, 2003). MPR is a theory-driven model of operant (voluntary) behavior and posits that three fundamental behavioral mechanisms are involved in reinforcement learning. Specifically, response rates under fixed-ratio (FR) schedules of reinforcement are a function of (a) *minimum response time*, which is a measure of the time (in seconds) required to complete a target response (e.g., a lever press), (b) *specific activation*, which is a measure of reinforcer value (in responses), and (c) *saturation rate*, which is a measure of how quickly response rates are maximized as a function of the number of target responses that precede reinforcer delivery. A high saturation rate indicates that relatively few target responses are strengthened by a reinforcer, whereas a low saturation rate indicates that relatively many target responses are strengthened by a reinforcer. Thus, saturation rate provides information about the effects of a reinforcer on target responding, and whether these effects are concentrated (high saturation rate) or dispersed (low saturation rate). MPR predicts response rate (b , responses/sec) as a function of fixed-ratio size (n) using Eq. (1) (Killeen and Sitomer, 2003):

$$b = \frac{1 - e^{-\lambda \delta n}}{\delta} - \frac{n}{\delta a} \quad (1)$$

where a is specific activation, δ is minimum response time, and λ is saturation rate. Fig. 1 illustrates how changes in MPR's parameters affect the shape of the function drawn by Eq. (1). MeHg-induced changes in MPR's parameters would reveal information about the behavioral mechanisms targeted by adolescent MeHg exposure. Whether the lasting behavioral effects following adolescent MeHg exposure can be reversed or mitigated via drug therapy is unclear, however (Fig. 2).

A growing literature suggests lysine deacetylase (LDAC) inhibitors ameliorate cognitive impairment in rodents (Day and Sweatt, 2012) as a result of normal or diseased aging (Fischer et al., 2007; Kilgore et al., 2010; Rane et al., 2012). LDAC inhibitors, such as sodium butyrate (NaB), target both histone deacetylases and DNA methyltransferases in the cell nucleus to increase histone acetylation and neuronal gene expression (Sarkar et al., 2011; Xu et al., 2007). Histone acetylation plays a crucial role in neuronal development, synaptic plasticity, learning,




Fig. 2. Timeline of events for Experiment 1 and Experiment 2. Mice arrived at PND 21 (Experiment 1) or PND 23 (Experiment 2) and were exposed to MeHg during adolescence for 39 days. In Experiment 2, sodium butyrate (NaB; 0.6 g/kg, i.p.) or saline vehicle were injected daily for two weeks prior to behavioral testing. Behavioral testing began in adulthood (PND 90 in Experiment 1, PND 100 in Experiment 2). MeHg = methylmercury, PND = postnatal day, NaB = sodium butyrate.

and memory (Day and Sweatt, 2012; Levenson et al., 2004; Miller et al., 2008), but the effects of repeated LDAC inhibition on operant behavior are underexplored. Some reports show that acute administration of NaB and other LDAC inhibitors do not affect responding under fixed- and progressive-ratio schedules of reinforcement (Romieu et al., 2008; Sun et al., 2008), but it is unclear whether NaB alters other aspects of reinforcement. It is important to characterize the effects of NaB on operant behavior both alone and after neurotoxicant exposure due to the growing use of NaB and other LDAC inhibitors for treating cognitive and behavioral dysfunction (Haggarty and Tsai, 2011). Whether repeated NaB administration mitigates the behavioral effects of adolescent MeHg exposure is unknown.

MeHg exposure has been linked to distorted gene expression and epigenetic dysregulation in some model systems, including cultured cells, nonhuman animals, and humans (Robinson et al., 2011). MeHg reduces *Bdnf* expression, a gene related to synaptic plasticity and learning, and increases *Hdac4* expression in cultured rat cortical neurons (Guida et al., 2017, 2016). Further, the LDAC inhibitor trichostatin A prevents MeHg-induced changes in gene expression and histone deacetylation in vitro (Guida et al., 2016). In behaving animals, gestational MeHg exposure is correlated with reduced *Bdnf* expression in the adult rat hippocampus, with a concurrent decrease in histone H3 acetylation and increase in DNA methylation at the *Bdnf* promoter (Onishchenko et al., 2008). Thus, developmental MeHg exposure may produce its neurobehavioral effects in part by increasing neuroepigenetic marks that suppress gene expression. Treatment with a broad-acting LDAC inhibitor, such as NaB, following MeHg exposure may prevent behavioral impairment.




Fig. 1. The effects of changes in minimum response time (δ), specific activation (a), and saturation rate (λ) on the shape of the MPR function drawn by Eq. (1). The base function (solid line) in each panel is drawn with $\delta = .50$, $a = 300$, and $\lambda = .10$. An increase in minimum response time to $\delta = .80$ (left panel) is evident in a reduction in the maximum response rate ($1/\delta$), which is the extrapolated Y-intercept of the descending limb of the function (dotted lines, left panel). A reduction in specific activation to $a = 200$ (middle panel) is reflected as a decrease in the X-intercept of the descending limb of the function. Finally, an increase in saturation rate to

$\lambda = .75$ shifts the peak of the function to the left. The vertical dotted lines (right panel) indicate that an FR 54 maximizes response rates when $\lambda = .10$ whereas an FR 12 maximizes response rates when $\lambda = .75$.

2. Experiment 1

Experiment 1 was designed to parse the behavioral mechanisms of adolescent MeHg exposure using Mathematical Principles of Reinforcement (Eq. (1); Killeen and Sitomer, 2003). The effects of adolescent MeHg exposure on MPR's parameters—specific activation (a), minimum response time (δ), and saturation rate (λ)—were assessed using response rates derived from a multiple fixed-ratio procedure.

3. Method

3.1. Subjects and MeHg exposure

Thirty-six male C57BL/6n mice were purchased from a commercial vendor (Envigo, Indianapolis, IN). The mice were derived from 12 litters (3 littermates/litter). These mice have previously been reported on (Boomhower and Newland, 2016). Briefly, all mice were pair-housed and maintained on a 12-hr light/dark cycle (lights on at 6:00 AM) in a temperature- and humidity-controlled, AAALAC-accredited animal facility. Upon arrival, 21-day-old littermates were divided among three MeHg-exposure groups (via drinking water): 0 ppm (control), 0.3 ppm, and 3.0 ppm MeHg. Thus, a litter was equally represented across exposure groups. MeHg was delivered as methylmercuric chloride (MeHgCl₂) dissolved in drinking water, and exposure occurred from postnatal day (PND) 21 through 59. All water was replaced with tap water on PND 60. Dosing was calculated by weighing water bottles, and sham water bottles were weighed to account for spillage. The low (0.3 ppm) and high (3.0 ppm) doses of MeHg corresponded to approximately 40 and 400 $\mu\text{g}/\text{kg}/\text{day}$ of MeHg in mice, respectively (Boomhower and Newland, 2017, 2016). The dose range of MeHg used here has been associated with neurobehavioral impairment in past work (Boomhower and Newland, 2017, 2016). Mice were maintained at $25 (\pm 1)$ g body mass by restricting daily food intake to $2.4 (\pm 0.02)$ g chow. The Auburn University Institutional Animal Care and Use Committee approved all procedures.

3.2. Apparatus

Twelve standard operant chambers (Med Associates, St. Albans, VT) modified for mice were used for data collection. Each chamber was equipped with two retractable levers on a front wall panel. Situated between the two levers was an alcove where a dipper system delivered .01-cc presentations of a 3:1 water and sweetened-condensed milk solution (hereafter, milk). Two Sonalert® tone generators (high tone: 4500 Hz, low tone: 2700 Hz) were located at the top of the front wall. A sound-attenuating cubicle enclosed each chamber. All experimental contingencies were controlled within 0.01-sec resolution by a computer in an adjacent room. Each mouse was assigned a particular chamber for the duration of the study. Mice were divided into four squads that ran at approximately the same time every day (± 15 min) Monday through Friday. The number of mice in each exposure group was counter-balanced across chambers and squads.

3.3. Procedure

All mice had prior experience lever pressing as described in Boomhower and Newland (2016). Beginning at 200 days of age, mice were trained under a multiple fixed-ratio (FR) schedule of reinforcement based on Reilly (2003) with some modifications. A session was divided into six components, each associated with a different FR schedule and signaled by a unique, high/low-alternating tone (Table 1). The FR schedule was increased across components in the following order: FR 1, 5, 15, 30, 60, and 120. At the beginning of a session, one lever was inserted into the chamber and the tones sounded. Upon completion of the FR requirement, the lever retracted, the tones were extinguished, and 3 s of access to milk was made available. The lever

Table 1
Tone durations for the FR procedure.

FR	Low/high tone durations (sec)
1	0.15/1.19
5	0.74/0.60
15	0.92/0.42
30	1.04/0.30
60	1.13/0.21
120	1.19/0.15

was then re-inserted into the chamber. A component ended after the delivery of 12 reinforcers or 10 min, whichever came first. Components were separated by 30 s during which the lever was retracted and tones were extinguished. The multiple FR schedule was in effect for 40 sessions to allow responding to stabilize.

3.4. Data analysis

Responding under the multiple FR schedule among exposure groups was assessed by averaging response rates, calculated as the number of responses divided by the duration of the component (in seconds), for each subject from the last 10 sessions. Eq. (1) was fit to individual-subject response-rate functions using nonlinear least-squares regression. Response rates were compared among groups using a repeated-measures ANOVA with exposure and FR as the within-subjects variables. Estimates of a , δ , and λ were compared across exposure groups using a repeated-measures ANOVA. Planned Tukey post-hoc comparisons were conducted between MeHg-exposed animals and controls.

4. Results and discussion

Fig. 3 shows mean response rates as a function of FR for the MeHg exposure groups along with best fits of Eq. (1) (solid lines). Individual-subject data for two mice from each exposure group also are shown to demonstrate the range of goodness of fit of Eq. (1): one was the best fit and one was the worst fit for each exposure group. One 3.0-ppm MeHg exposed mouse, was a statistical outlier due to his low rate of responding (Fig. 1, lower right panel) so his data were not included in the mean for the 3.0-ppm group. There were main effects of both FR [$F(5, 197) = 37.95, p < .001$] and exposure [$F(2, 197) = 7.53, p < .001$] in that response rates were a bitonic function of FR and MeHg-exposed mice responded more quickly on average. Eq. (1) provided a good fit of all individual-subject data with the mean (SEM) R^2 values being 0.93 (0.02), 0.96 (0.01), and 0.94 (0.03) for the 0-, 0.3-, and 3.0-ppm MeHg groups, respectively. The quality of the fits was not significantly different among exposure groups.

Fig. 4 shows mean parameter estimates of λ , δ , and a for the MeHg exposure groups. The outlier is denoted as an open circle. Estimates of λ were significantly different among the exposure groups [$F(2, 33) = 4.16, p = .02$], and post-hoc comparisons revealed that mice in the 3.0-ppm group had higher saturation rates compared to Controls ($p < .05$). Thus, fewer target responses maximized response rates in MeHg-exposed mice than Controls. Estimates of δ were significantly different among exposure groups [$F(2, 33) = 9.15, p < .001$]. Post-hoc comparisons revealed that the 0.3- and 3.0-ppm groups had faster minimum response times than Controls ($p < .02$). This was evident in higher maximum response rates ($1/\delta$) in exposed mice. It should be noted that the inclusion of the statistical outlier in the 3-ppm group rendered δ estimates for this group statistically similar to Controls' δ estimates. Estimates of a were not significantly different among exposure groups [$F(2, 33) = 0.89, p = .42$], indicating that the value of a reinforcer was similar following adolescent MeHg exposure.

As revealed by MPR (Killeen and Sitomer, 2003), adolescent exposure to MeHg manifested as alterations in saturation rate and minimum response time in adulthood. This suggests for the MeHg-




Fig. 3. Left panel: Mean response rate as a function of fixed ratio for mice exposed to MeHg in adolescence. Lines represent mean predictions of Eq. (1). Right panel: Response rate (symbols) and predictions of Eq. (1) (lines) as a function of fixed ratio for individual mice. The left column shows data from mice with the best-fitting curves, and the right column shows data from mice with the worst-fitting curves. Note the Y-axis scaling in the lower-right panel. MeHg = methylmercury.

exposed mice, the reinforcer's effects are more concentrated on the responses that immediately preceded reinforcement than on the earlier responses. In other words, adolescent MeHg exposure reduced the number of responses being strengthened by reinforcement. Further, an enhancement in psychomotor responding following MeHg exposure is consistent with past work showing gestationally-exposed rats respond more quickly on fixed- (Paletz et al., 2006) and progressive-ratio schedules of reinforcement (Reed et al., 2008). These findings suggest that adolescent MeHg exposure may produce long-lasting distortions in both the ability of a reinforcer to strengthen past responses and psychomotor responding. Whether MeHg's behavioral effects can be mitigated by repeated administration of sodium butyrate (NaB), a lysine deacetylase (LDAC) inhibitor, is unknown.

5. Experiment 2

Developmental MeHg exposure may produce its neurobehavioral effects in part by increasing neuroepigenetic marks that suppress gene expression (Guida et al., 2017; Onishchenko et al., 2008). Treatment with a broad-acting LDAC inhibitor, such as NaB, following MeHg exposure may prevent behavioral impairment. Due to the growing use of NaB and other LDAC inhibitors for treating cognitive and behavioral dysfunction (Haggarty and Tsai, 2011), it is also important to characterize the effects of NaB alone on operant behavior. Acute administration of NaB and other LDAC inhibitors does not affect responding under fixed- and progressive-ratio schedules of reinforcement in rodents (Romieu et al., 2008; Sun et al., 2008), but it is unclear whether NaB alters other aspects of reinforcement. The use of Mathematical Principles of Reinforcement (MPR) would parse the effects of NaB on three unique processes of operant behavior: saturation rate, minimum

response time, and reinforcer value. Thus, Experiment 2 was designed to (a) replicate the effects of adolescent MeHg exposure on MPR's parameters in a different cohort of mice, and (b) determine the effects of repeated NaB administration, both alone and after MeHg exposure, on MPR's parameters. We chose to examine the highest dose (3 ppm) of MeHg because it produced significant changes in both saturation rate and minimum response time (Experiment 1) as well as behavioral impairment in other work (Boomhower and Newland, 2017) to determine the degree to which NaB prevented MeHg's effects.

6. Method

6.1. Subjects and MeHg exposure

Forty-eight male C57Bl/6n mice derived from 24 litters (2 littermates/litter) were purchased from a commercial vendor (Envigo, Indianapolis, IN). Mice arrived at our facility on PND 23, and 12 litters (24 mice) were exposed to 3 ppm MeHg beginning on PND 24. MeHg was delivered as methylmercuric chloride dissolved in drinking water. The other 12 litters (24 mice) received control (tap) water. Dosing was calculated by weighing water bottles, and sham water bottles were used to control for spillage. On PND 63, MeHg bottles were removed and replaced with tap water. Thus, mice in Experiment 2 were exposed to MeHg for an identical number of days (39 days) as mice in Experiment 1 except MeHg exposure spanned PND 24–62 in Experiment 2 (a 3-day displacement). All mice were given ad libitum access to food until just prior to behavioral testing, which began on PND 100. Two weeks prior to behavioral testing, mice were maintained at a body mass of 25 (\pm 1) g by food restricting daily chow intake to 2.4 (\pm 0.2) g. All mice were individually-housed and maintained under a 12-hr light/dark cycle




Fig. 4. Mean (\pm SD) parameter estimates from Eq. (1) as a function of dose of MeHg in adolescence. An outlier in the 3.0 ppm group is denoted as an open circle. MeHg = methylmercury * p < 0.05 relative to control.

(lights on at 6:00 AM) in a temperature- and humidity-controlled, AAALAC-approved animal facility. The Auburn University Institutional Animal Care and Use Committee approved all procedures.

6.2. Sodium butyrate (NaB) administration

Two weeks before behavioral testing, half the mice in each MeHg-exposure group received daily acute i.p. injections of 0.9% saline or 0.6 g/kg NaB under a 2 (MeHg) X 2 (NaB) full-factorial design. Litters (2 littermates/litter) within a MeHg exposure group were divided equally between the saline and NaB conditions. Thus, a litter was equally represented across NaB conditions. Drug administration spanned PND 86–99, a regimen similar to past work (Kim et al., 2009; Rane et al., 2012). Sodium butyrate (Sigma) was dissolved in saline. Saline and NaB were injected at a .01 ml/g volume.

6.3. Apparatus

The same apparatuses were used as in Experiment 1.

6.4. Procedure

All mice had previous experience lever pressing under a spatial-discrimination-reversal and visual-discrimination procedure, similar to Boomhower and Newland (2017). Beginning at 200 days of age, mice were trained under the multiple fixed-ratio procedure described in Experiment 1. The number of mice from each group was counter-balanced across chambers and session time. One 0-ppm MeHg mouse (Control) and three 3-ppm MeHg mice (Control, $n = 1$; NaB, $n = 2$) were euthanized before behavioral testing for reasons unrelated to the present experiments.

6.5. Data analysis

Similar to Experiment 1, Eq. (1) was fit to individual-subject response-rate functions using nonlinear least-squares regression. Response rates were analyzed using a linear-mixed effects (LME) model with MeHg, NaB, and FR as fixed effects and litter as a random effect. LME was chosen because it is able to model incomplete repeated-measures data more effectively than does traditional repeated-measures analysis of variance. Estimates of a , δ , and λ were compared across groups using a LME model with MeHg and NaB as fixed effects and litter as a random effect. Planned Tukey post-hoc comparisons were conducted to assess significant differences between groups.

7. Results and discussion

Fig. 5 shows MeHg consumption across adolescence for each MeHg exposure group. The dose of MeHg was highest at the beginning of exposure and gradually decreased to about 400 $\mu\text{g}/\text{kg}/\text{day}$. This pattern of dosing was similar to that seen in Experiment 1 (see Boomhower and Newland, 2016) and Boomhower and Newland (2017) and is due to changes in fluid intake that occurs through early adolescence. Administration of NaB did not significantly alter MeHg consumption.

Fig. 6 shows mean response rates and best fits of Eq. (1) for each treatment group. There was a main effect of FR [$F(5, 239) = 37.37$, $p < .001$], which was reflected in a bitonic relation between response rates and FR size. There were no main effects of MeHg or NaB, and there were no significant interactions. Eq. (1) fit individual response-rate functions well with a mean pseudo $R^2 = 0.94$ ($SD: 0.06$) across all groups.

Fig. 7 shows mean parameter estimates from Eq. (1) for each exposure and treatment group. For saturation rate (λ), there was a main effect of NaB [$F(1, 39) = 4.91$, $p = .03$] with post-hoc comparisons revealing a significant increase in λ following MeHg alone, NaB alone, and the combination relative to the 0-ppm Control group (p 's $< .05$).




Fig. 5. Mean (\pm SD) dose of MeHg as a function of postnatal day for each MeHg-exposure group in Experiment 2. Note the error bars may be obscured by the symbols in some cases. MeHg = methylmercury, NaB = sodium butyrate.




Fig. 6. Mean response rate as a function of fixed ratio for mice exposed to 0 or 3 ppm MeHg in adolescence and treated with control (left panel) and sodium butyrate (right panel) before behavioral testing. Lines represent mean predictions of Eq. (1). MeHg = methylmercury.

Neither minimum response time (δ) nor specific activation (a) estimates were significantly altered relative to 0-ppm Control.

The finding that adolescent MeHg exposure increases saturation rate, or the degree to which prior responses are coupled to reinforcers, was replicated in Experiment 2. Further, MeHg-exposed mice tended to have higher maximum response rates (smaller δ 's) than unexposed mice similar to Experiment 2, but this was more modest than in Experiment 1 and was not reflected in a statistically significant decrease in minimum response time or maximum response rate. Differences in the exposure regimen might explain why the effects on minimum response time were not replicated in Experiment 2. Though mice in Experiment 2 were exposed to MeHg for 39 days, identical to Experiment 1, the exposure duration began 3 days later in development. Some work suggests the effects of MeHg on the postnatal rat brain, particularly dopamine signaling, are mitigated after 7 days of continued striatal development (from PND 14 to 21) (Dreiem et al., 2009). Of all three parameter estimates from MPR, the ability of reinforcers to strengthen past responding (λ) appears to be the one most consistently sensitive to adolescent MeHg exposure. The MeHg-induced increase in λ estimates persisted following treatment with NaB, suggesting no therapeutic benefits of NaB under these levels and durations. NaB treatment alone, however, produced a significant increase in saturation rate relative to 0-ppm Controls, an effect that was similar in magnitude to MeHg's. These findings suggest repeated inhibition of lysine deacetylases prior to behavioral testing changes the extent to which reinforcers strengthen past responding.




Fig. 7. Mean (+SD) parameter estimates from Eq. (1) for each treatment group. MeHg = methylmercury, NaB = sodium butyrate. * $p < 0.05$.

8. General discussion and conclusions

The effects of adolescent methylmercury (MeHg) exposure on reinforcer value, motoric capacity, and the coupling of responses to reinforcement in mice were evaluated using Mathematical Principles of Reinforcement (Killeen and Sitomer, 2003) in two separate experiments. Further, the effects of sodium butyrate (NaB) both alone and after MeHg exposure were examined due to the growing use of NaB and other LDAC inhibitors for treating cognitive and behavioral dysfunction (Haggarty and Tsai, 2011). Overall, MPR described response-rate functions collected from individual mice quite well and captured alterations in responding following the various exposures and treatments.

In both Experiment 1 and 2, adolescent MeHg exposure increased saturation rate (λ), the rate at which response-reinforcer coupling decreased across time. Stated differently, adolescent MeHg exposure decreased the number of responses that were strengthened by reinforcers. Reduced coupling of all responses except the one that immediately preceded reinforcement following adolescent MeHg exposure could be related to past work reporting MeHg-induced impairments in choice. Adolescent MeHg dose-dependently reduced sensitivity to a larger reinforcer compared to a smaller reinforcer in these mice (Boomhower and Newland, 2016), and 3 ppm of MeHg increased the number of trials required to transition through repeated spatial-discrimination reversals and an extradimensional shift (Boomhower and Newland, 2017). These findings suggest a degradation in the ability of a reinforcer to strengthen prior responses and is consistent with past work showing that saturation rate (λ) is correlated with impaired performance on spatial discrimination reversals in mice (Pope et al., 2016).

The finding that adolescent MeHg exposure diminishes the impact of reinforcement on previous responses implicates neural substrates linked to motivation in adolescent MeHg toxicity. The nucleus accumbens, its projections and interaction with the prefrontal cortex, and dopamine signaling are important for reward processing (Everitt and Robbins, 2005; Fiorillo et al., 2003). Lesions of the rat nucleus accumbens core increase a parameter similar to saturation rate (λ) (Bezzina et al., 2008), whereas rat orbital prefrontal-cortex lesions are associated with reductions in specific activation (a) (Kheramin et al., 2005). Changes in the parameters of Mathematical Principles of Reinforcement are linked intimately with dopaminergic function, as both neuroleptics and amphetamines can alter saturation rate, minimum response time, and specific activation (Mobini et al., 2000; Reilly, 2003). Our data suggest adolescent MeHg exposure interferes with the coupling of behavior to reinforcement, which is linked to nucleus accumbens and prefrontal-cortex function. The finding that the effects of reinforcement on behavior are impaired following adolescent MeHg exposure is consistent with Newland et al.'s (2015) hypothesis that developmental MeHg exposure impacts reinforcer processing, primarily through MeHg's effects on dopamine. Though Newland et al.'s (2015) review relied on models of gestational MeHg exposure, the present study suggests that adolescent MeHg exposure also contributes to alterations in brain-behavior function.

Minimum response times (δ) were decreased by adolescent MeHg

exposure in Experiment 1, particularly under the 0.3-ppm dose, but were not significantly altered in Experiment 2. Reduced minimum response time manifests as increased maximum response rate, which is correlated with $1/\delta$ both theoretically (Killeen and Sitomer, 2003) and experimentally (Hutsell and Newland, 2013). Past work in gestationally-exposed rats showed that MeHg increases response rates under fixed-ratio (Paletz et al., 2006) and progressive-ratio schedules of reinforcement (Reed et al., 2008). In Experiment 1, both doses of MeHg reduced minimum response time. The greatest decrease in minimum response time occurred after 0.3 ppm MeHg, and the 3.0-ppm dose of MeHg also produced a significant decrease in δ estimates after a statistical outlier was removed. It is unclear why the same dose of MeHg in Experiment 2 did not produce a significant reduction in minimum response time. We have noted previously dose-specific behavioral effects of adolescent MeHg exposure in the cohort of animals used in Experiment 1. Specifically, delay sensitivity was reduced in the 0.3-ppm group whereas mice in the 3-ppm group displayed similar estimates of delay sensitivity as controls (Boomhower and Newland, 2016). Key differences in the exposure regimen between Experiment 1 and 2 could explain also why our findings related to minimum response time were only partially replicated. We shifted the 39-day exposure window three days later in Experiment 2 because we had shown previously MeHg consumption is dramatically elevated during the first few days (PND 21–23) of exposure (Boomhower and Newland, 2016). We were concerned this could be an artifact of elevated water consumption during shipment from commercial vendors (see Obernier and Baldwin, 2006). Some work suggests the effects of MeHg on the postnatal rat brain, particularly related to striatal dopamine signaling and brain mercury uptake, are mitigated with development (Dreiem et al., 2009; Sakamoto et al., 2018). Specifically, striatal synaptosomes isolated from 7-day-old rats are more susceptible to MeHg's effects on dopamine release and dopamine uptake than striatal synaptosomes from 21-day-old rats (Dreiem et al., 2009). Brain mercury uptake also decreases from the neonatal (PND 1–7) to the mid-adolescent period (PND 35–41) in rats (Sakamoto et al., 2018). The striatum and striatal dopamine have been implicated as targets of MeHg (Faro et al., 2002, 2000; Tiernan et al., 2015) and most likely underlie MeHg's effects on operant behavior in part (Newland et al., 2015). Because both the sensitivity of striatal dopamine to MeHg exposure and MeHg uptake may decrease early in life (Dreiem et al., 2009; Sakamoto et al., 2018), this could explain why mice in Experiment 2 did not show impairment in minimum response time following MeHg exposure and would implicate the early-adolescent period in susceptibility to MeHg. Interestingly, we still observed elevated MeHg consumption early in the exposure regimen in Experiment 2 even after delaying exposure three days.

Repeated administration of sodium butyrate (NaB), a lysine deacetylase inhibitor, increased saturation rate in adulthood and NaB did not interact with adolescent MeHg exposure to change saturation-rate estimates. Past work has shown that MeHg reduces *Bdnf* expression in vitro (Guida et al., 2017) and in vivo (Onishchenko et al., 2008) with concomitant changes in histone acetylation, suggesting MeHg's behavioral impairment may result in part from alterations to

neuroepigenetic mechanisms. Guida et al. (2016) found that trichostatin A, an LDAC inhibitor, devented MeHg-induced changes in gene expression and histone H4 deacetylation in vitro. However, we saw no therapeutic benefit of NaB following MeHg exposure and found that NaB alone increased saturation rate to a comparable level as MeHg exposure alone in behaving mice. Studies using NaB as a treatment for both normal (Peleg et al., 2010; Reolon et al., 2011) and diseased aging (Fischer et al., 2007; Kilgore et al., 2010; Rane et al., 2012) as well as following neurotoxicant exposure (Guida et al., 2016; Sharma and Sharma, 2013; Song et al., 2010) have noted a range of therapeutic effects. One explanation for our findings related to NaB is that LDAC inhibitors often target a broad set of proteins, sometimes related to cell death or oxidative stress (Xu et al., 2007). The broad enzymatic targets of NaB relative to trichostatin A may explain not only why NaB was ineffective at prevent MeHg behavioral toxicity, but also why NaB alone increased saturation rate in our study. NaB induces differentiation in neurons (Balasubramanian et al., 2006; Hsieh et al., 2004) and apoptosis in neuroblastoma cells (Nuydens et al., 1995; Rozental et al., 2004). Further characterization of the neuroprotective and neurotoxic effects of NaB, and other lysine deacetylase inhibitors, on operant behavior are important for future therapeutic treatments of chemical exposure.

Across two studies, exposure to environmentally-relevant levels of MeHg in adolescent mice reduced the degree to which responses were coupled to reinforcement in adulthood. We also found some evidence that adolescent MeHg exposure reduces minimum response time, particularly following exposure to 0.3 ppm MeHg, which is consistent with past work on developmental MeHg exposure's psychomotor effects. Repeated administration of sodium butyrate alone increased saturation rate to a similar extent as MeHg exposure but did not reverse MeHg's effects. The present study suggests adolescence is a time of both behavioral and neurobiological vulnerability to MeHg exposure. The finding that both adolescent MeHg exposure and sodium butyrate administration concentrates the effects of reinforcers on recently-made responses provide insight into the behavioral mechanisms that permit neurotoxicity.

Funding information and disclosures

This work was supported by graduate research grant awards from Psi Chi, Sigma Xi, the American Psychological Association, the American Psychological Foundation, Auburn University Graduate School, the National Science Foundation Graduate Research Fellowship Program [grant number DGE-1414475 to S.R.B.], and the National Institutes of Environmental Health Sciences [grant number ES024850 to M.C.N.]. The authors declare no conflicts of interest.

Acknowledgements

We thank Megan Arnold, Douglas Glenn, Kevin Honeywell, Kate Johnson, Dalisa Kendricks, Sahar Moghadam, Madison Morlan, Alex Sauer, Savannah Simpson, and Kristyn Suelflow for help with data collection and animal care. Portions of this manuscript served as the first author's dissertation and were presented as posters at Society of Toxicology.

References

Balasubramanian, V., Boddeke, E., Bakels, R., Kust, B., Kooistra, S., Veneman, A., Copray, S., 2006. Effects of histone deacetylation inhibition on neuronal differentiation of embryonic mouse neural stem cells. *Neuroscience* 143, 939–951. <https://doi.org/10.1016/j.neuroscience.2006.08.082>.

Bezzina, G., Body, S., Cheung, T.H.C., Hampson, C.L., Deakin, J.F.W., Anderson, I.M., Szabadi, E., Bradshaw, C.M., 2008. Effect of quinolinic acid-induced lesions of the nucleus accumbens core on performance on a progressive ratio schedule of reinforcement: implications for inter-temporal choice. *Psychopharmacology (Berl.)* 197, 339–350. <https://doi.org/10.1007/s00213-007-1036-0>.

Boomhower, S.R., Newland, M.C., 2017. Effects of adolescent exposure to methylmercury

and d-amphetamine on reversal learning and an extradimensional shift in male mice. *Exp. Clin. Psychopharmacol.* 25, 64–73.

Boomhower, S.R., Newland, M.C., 2016. Adolescent methylmercury exposure affects choice and delay discounting in mice. *Neurotoxicology* 57, 136–144.

Chambers, R.A., Potenza, M.N., 2003. Neurodevelopment, impulsivity, and adolescent gambling. *J. Gambl. Stud.* 19, 53–84.

Day, J.J., Sweatt, J.D., 2012. Epigenetic treatments for cognitive impairments. *Neuropsychopharmacology* 37, 247–260. <https://doi.org/10.1038/npp.2011.85>.

Doremus-Fitzwater, T.L., Barreto, M., Spear, L.P., 2012. Age-related differences in impulsivity among adolescent and adult Sprague-Dawley rats. *Behav. Neurosci.* 126, 735–741. <https://doi.org/10.1037/a0029697>.

Dreiem, A., Shan, M., Okoniewski, R.J., Sanchez-Morrissey, S., Seegal, R.F., 2009. Methylmercury inhibits dopaminergic function in rat pup synaptosomes in an age-dependent manner. *Neurotoxicol. Teratol.* 31, 312–317.

Everitt, B.J., Robbins, T.W., 2005. Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat. Neurosci.* 8, 1481–1489. <https://doi.org/10.1038/nn1579>.

Faro, L., do Nascimento, J., Alfonso, M., Durán, R., 2002. Mechanism of action of methylmercury on in vivo striatal dopamine release: possible involvement of dopamine transporter. *Neurochem. Int.* 40, 455–465.

Faro, L., Do Nascimento, J.L.M., San José, J.M., Alfonso, M., Durán, R., 2000. Intra-striatal administration of methylmercury increases in vivo dopamine release. *Neurochem. Res.* 25, 225–229.

Fiorillo, C.D., Tobler, P.N., Schultz, W., 2003. Discrete coding of reward dopamine. *Neurons* Science 80- (299), 1898–1902. <https://doi.org/10.1126/science.1077349>.

Fischer, A., Sananbenesi, F., Wang, X., Dobbin, M., Tsai, L.-H., 2007. Recovery of learning and memory is associated with chromatin remodelling. *Nature* 447, 178–182. <https://doi.org/10.1038/nature05772>.

Giedd, J.N., Blumenthal, J., Jeffries, N.O., Castellanos, F.X., Liu, H., Zijdenbos, A., Paus, T., Evans, A.C., Rapoport, J.L., 1999. Brain development during childhood and adolescence: a longitudinal MRI study. *Nat. Neurosci.* 2, 861–863.

Guida, N., Laudati, G., Anzilotti, S., Sirabella, R., Cuomo, O., Brancaccio, P., Santopalo, M., Galgani, M., Montuori, P., Di Renzo, G., Canzoniero, L.M.T., Formisano, L., 2016. Methylmercury upregulates RE-1 silencing transcription factor (REST) in SH-SY5Y cells and mouse cerebellum. *Neurotoxicology* 52, 89–97. <https://doi.org/10.1016/j.neuro.2015.11.007>.

Guida, N., Laudati, G., Mascolo, L., Valsecchi, V., Sirabella, R., Sella, C., Di Renzo, G., Canzoniero, L.M.T., Formisano, L., 2017. p38/Sp1/Sp4/HDAC4/BDNF Axis is a novel molecular pathway of the neurotoxic effect of the methylmercury. *Front. Neurosci.* 11, 1–10. <https://doi.org/10.3389/fnins.2017.00008>.

Haggarty, S.J., Tsai, L.H., 2011. Probing the role of HDACs and mechanisms of chromatin-mediated neuroplasticity. *Neurobiol. Learn. Mem.* 96, 41–52. <https://doi.org/10.1016/j.nlm.2011.04.009>.

Hsieh, J., Nakashima, K., Kuwabara, T., Mejia, E., Gage, F.H., 2004. Histone deacetylase inhibition-mediated neuronal differentiation of multipotent adult neural progenitor cells. *PNAS* 101, 16659–16664. <https://doi.org/10.1073/pnas.0407643101>.

Hutsell, B.A., Newland, M.C., 2013. A quantitative analysis of the effects of qualitatively different reinforcers on fixed ratio responding in inbred strains of mice. *Neurobiol. Learn. Mem.* 101, 85–93.

Kheramin, S., Body, S., Herrera, F.M., Bradshaw, C.M., Szabadi, E., Deakin, J.F.W., Anderson, I.M., 2005. The effect of orbital prefrontal cortex lesions on performance on a progressive ratio schedule: implications for models of inter-temporal choice. *Behav. Brain Res.* 156, 145–152. <https://doi.org/10.1016/j.bbr.2004.05.017>.

Kilgore, M., Miller, C.A., Fass, D.M., Hennig, K.M., Haggarty, S.J., Sweatt, J.D., Rumbaugh, G., 2010. Inhibitors of class 1 histone deacetylases reverse contextual memory deficits in a mouse model of Alzheimer's disease. *Neuropsychopharmacology* 35, 870–880. <https://doi.org/10.1038/npp.2009.197>.

Killeen, P.R., 1994. Mathematical principles of reinforcement. *Behav. Brain Sci.* 17, 105–172.

Killeen, P.R., Sitomer, M.T., 2003. MPR. *Behav. Processes* 62, 49–64.

Kim, H.J., Leeds, P., Chuang, D.M., 2009. The HDAC inhibitor, sodium butyrate, stimulates neurogenesis in the ischemic brain. *J. Neurochem.* 110, 1226–1240. <https://doi.org/10.1111/j.1471-4159.2009.06212.x>.

Levenson, J.M., O'Riordan, K.J., Brown, K.D., Trinh, M., Molfese, D.L., Sweatt, J.D., 2004. Regulation of histone acetylation during memory formation in the hippocampus. *J. Biol. Chem.* 279, 40545–40559. <https://doi.org/10.1074/jbc.M402229200>.

Miller, C.A., Campbell, S.L., Sweatt, J.D., 2008. DNA methylation and histone acetylation work in concert to regulate memory formation and synaptic plasticity. *Neurobiol. Learn. Mem.* 89, 599–603. <https://doi.org/10.1016/j.nlm.2007.07.016>.

Mobini, S., Chiang, T.-J., Ho, M.-Y., Bradshaw, C.M., Szabadi, E., 2000. Comparison of the effects of clozapine, haloperidol, chlorpromazine and d-amphetamine on performance on a time-constrained progressive ratio schedule and on locomotor behaviour in the rat. *Psychopharmacology (Berl.)* 152, 47–54. <https://doi.org/10.1007/s002130000486>.

National Research Council, 2000. Toxicological Effects of Methylmercury. National Academy Press, Washington, D.C. <https://doi.org/10.17226/9899>.

Newland, M.C., Palez, E.M., Reed, M.N., 2008. Methylmercury and nutrition: adult effects of fetal exposure in experimental models. *Neurotoxicology* 29, 783–801.

Newland, M.C., Reed, M.N., Rasmussen, E., 2015. A hypothesis about how early developmental methylmercury exposure disrupts behavior in adulthood. *Behav. Processes* 114, 41–51.

Newland, M.C., Reile, P.A., Langston, J.L., 2004. Gestational exposure to methylmercury retards choice in transition in aging rats. *Neurotoxicol. Teratol.* 26, 179–194.

Newman, L., McGaughy, J., 2011. Adolescent rats show cognitive rigidity in a test of attentional set shifting. *Dev. Psychobiol.* 53, 391–401.

- Nielsen, S.J., Aoki, Y., Kit, B.K., Ogden, C.L., 2015. More than half of US youth consume seafood and most have blood mercury concentrations below the EPA reference level, 2009–2012. *J. Nutr.* 145, 322–327. <https://doi.org/10.3945/jn.114.203786>.
- Nuydens, R., Heers, C., Chadarevian, A., De Jong, M., Nuyens, R., Cornelissen, F., Geerts, H., 1995. Sodium butyrate induces aberrant tau phosphorylation and programmed cell death in human neuroblastoma cells. *Brain Res.* 688, 86–94. [https://doi.org/10.1016/0006-8993\(95\)00514-Q](https://doi.org/10.1016/0006-8993(95)00514-Q).
- Obnier, J.A., Baldwin, R.L., 2006. Establishing an appropriate period of acclimatization following transportation of laboratory animals. *ILAR J.* 47, 364–369. <https://doi.org/10.1093/ilar.47.4.364>.
- Olson, E.A., Hooper, C.J., Collins, P., Luciana, M., 2007. Adolescents' performance on delay and probability discounting tasks: contributions of age, intelligence, executive functioning, and self-reported externalizing behavior. *Pers. Individ. Dif.* 43, 1886–1897.
- Onishchenko, N., Karpova, N., Sabri, F., Castrén, E., Ceccatelli, S., 2008. Long-lasting depression-like behavior and epigenetic changes of BDNF gene expression induced by perinatal exposure to methylmercury. *J. Neurochem.* 106, 1378–1387. <https://doi.org/10.1111/j.1471-4159.2008.05484.x>.
- Paletz, E.M., Craig-Schmidt, M.C., Newland, M.C., 2006. Gestational exposure to methylmercury and n-3 fatty acids: effects on high- and low-rate operant behavior in adulthood. *Neurotoxicol. Teratol.* 28, 59–73.
- Peleg, S., Sananbenesi, F., Zovoilis, A., Burkhardt, S., Bahari-Javan, S., Agis-Balboa, R.C., Cota, P., Wittmann, J.L., Gogol-Doering, A., Opitz, L., Salinas-Riester, G., Dettenhofer, M., Kang, H., Farinelli, L., Chen, W., Fischer, A., 2010. Altered histone acetylation is associated with age-dependent memory impairment in mice. *Science* 80- (328), 753–756 <https://doi.org/328/5979/753> [pii]r10.1126/science.1186088.
- Pinkston, J.W., Lamb, R.J., 2011. Delay discounting in C57BL/6J and DBA/2J mice: adolescent-limited and life-persistent patterns of impulsivity. *Behav. Neurosci.* 125, 194–201.
- Pope, D.A., Boomhower, S.R., Hutsell, B.A., Teixeira, K., Newland, M.C., 2016. Chronic cocaine exposure in adolescence: effects on spatial discrimination reversal, delay discounting, and performance on fixed-ratio schedules in mice. *Neurobiol. Learn. Mem.* 130, 93–104. <https://doi.org/10.1016/j.nlm.2016.01.017>.
- Rane, P., Shields, J., Heffernan, M., Guo, Y., Akbarian, S., King, J.A., 2012. The histone deacetylase inhibitor, sodium butyrate, alleviates cognitive deficits in pre-motor stage PD. *Neuropharmacology* 62, 2408–2411. <https://doi.org/10.1016/j.neuropharm.2012.01.026>.
- Reed, M.N., Banna, K.M., Donlin, W.D., Newland, M.C., 2008. Effects of gestational exposure to methylmercury and dietary selenium on reinforcement efficacy in adulthood. *Neurotoxicol. Teratol.* 30, 29–37.
- Reed, M.N., Paletz, E.M., Newland, M.C., 2006. Gestational exposure to methylmercury and selenium: effects on a spatial discrimination reversal in adulthood. *Neurotoxicology* 27, 721–732.
- Reilly, M.P., 2003. Extending mathematical principles of reinforcement into the domain of behavioral pharmacology. *Behav. Processes* 62, 75–88.
- Reolon, G.K., Maurmann, N., Werenicz, A., Garcia, V.A., Schröder, N., Wood, M.A., Roesler Rafael, R., 2011. Posttraining systemic administration of the histone deacetylase inhibitor sodium butyrate ameliorates aging-related memory decline in rats. *Behav. Brain Res.* 221, 329–332. <https://doi.org/10.1016/j.bbr.2011.03.033>.
- Robinson, J.F., Theunissen, P.T., van Dartel, D.A.M., Pennings, J.L., Faustman, E.M., Piersma, A.H., 2011. Comparison of MeHg-induced toxicogenomic responses across in vivo and in vitro models used in developmental toxicology. *Reprod. Toxicol.* 32, 180–188. <https://doi.org/10.1016/j.reprotox.2011.05.011>.
- Romieu, P., Host, L., Gobaille, S., Sandner, G., Aunis, D., Zwiller, J., 2008. Histone deacetylase inhibitors decrease cocaine but not sucrose self-administration in rats. *J. Neurosci.* 28, 9342–9348. <https://doi.org/10.1523/JNEUROSCI.0379-08.2008>.
- Rozental, R., Faharani, R., Yu, Y., Johnson, J.M., Chan, S.O., Chiu, F.C., 2004. Sodium butyrate induces apoptosis in MSN neuroblastoma cells in a calcium independent pathway. *Neurochem. Res.* 29, 2125–2134. <https://doi.org/10.1007/s11064-004-6886-9>.
- Sakamoto, M., Tatsuta, N., Chan, H.M., Domingo, J.L., Murata, K., Nakai, K., 2018. Brain methylmercury uptake in fetal, neonate, weanling, and adult rats. *Environ. Res.* 167, 15–20. <https://doi.org/10.1016/j.envres.2018.06.038>.
- Sarkar, S., Abujamra, A.L., Loew, J.E., Forman, L.W., Perrine, S.P., Faller, D.V., 2011. Histone deacetylase inhibitors reverse CpG methylation by regulating DNMT1 through ERK signaling. *Anticancer Res.* 31, 2723–2732.
- Sharma, B., Sharma, P.M., 2013. Arsenic toxicity induced endothelial dysfunction and dementia: pharmacological interdicted by histone deacetylase and inducible nitric oxide synthase inhibitors. *Toxicol. Appl. Pharmacol.* 273, 180–188. <https://doi.org/10.1016/j.taap.2013.07.017>.
- Sichert-Hellert, W., Wicher, M., Kersting, M., 2009. Age and time trends in fish consumption pattern of children and adolescents, and consequences for the intake of long-chain n-3 polyunsaturated fatty acids. *Eur. J. Clin. Nutr.* 63, 1071–1075.
- Song, C., Kanthasamy, A., Anantharam, V., Sun, F., Kanthasamy, a G., 2010. Environmental neurotoxic pesticide increases histone acetylation to promote apoptosis in dopaminergic neuronal cells: relevance to epigenetic mechanisms of neurodegeneration. *Mol. Pharmacol.* 77, 621–632. <https://doi.org/10.1124/mol.109.062174>.
- Spear, L.P., 2000. The adolescent brain and age-related behavioral manifestations. *Neurosci. Biobehav. Rev.* 24, 417–463.
- Sun, J., Wang, L., Jiang, B., Hui, B., Lv, Z., Ma, L., 2008. The effects of sodium butyrate, an inhibitor of histone deacetylase, on the cocaine- and sucrose-maintained self-administration in rats. *Neurosci. Lett.* 441, 72–76. <https://doi.org/10.1016/j.neulet.2008.05.010>.
- Teicher, M.H., Andersen, S.L., Hostetter, J.C., 1995. Evidence for dopamine receptor pruning between adolescence and adulthood in striatum but not nucleus accumbens. *Dev. Brain Res.* 89, 167–172.
- Tiernan, C.T., Edwin, E.A., Hawong, H.Y., Rios-Cabanillas, M., Goudreau, J.L., Atchison, W.D., Lookingland, K.J., 2015. Methylmercury impairs canonical dopamine metabolism in rat undifferentiated pheochromocytoma (pc12) cells by indirect inhibition of aldehyde dehydrogenase. *Toxicol. Sci.* 144, 347–356. <https://doi.org/10.1093/toxsci/kfv001>.
- Xu, W.S., Parmigiani, R.B., Marks, P., 2007. Histone deacetylase inhibitors: molecular mechanisms of action. *Oncogene* 26, 5541–5552. <https://doi.org/10.1038/sj.onc.1210620>.