



## Correction to: In vitro schistosomicidal activity of tamoxifen and its effectiveness in a murine model of schistosomiasis at a single dose

Rosimeire N. Oliveira<sup>1,2</sup> · Sheila A. P. Corrêa<sup>1</sup> · Karen M. Vieira<sup>1</sup> · Tiago Mendes<sup>1</sup> · Silmara M. Allegretti<sup>1</sup> · Danilo C. Miguel<sup>1</sup>

Published online: 22 March 2019  
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

### Correction to: Parasitology Research

<https://doi.org/10.1007/s00436-019-06259-0>

The original published version of this article contains error in Tables 1 and 2. Correct tables are presented here.

**Table 1** Worm recovery from the hepatic portal system and mesenteric intestinal veins of mice experimentally infected with *S. mansoni* and treated with PBS (control group) or one single dose of 100 mg/kg cTAM

Administration route	Treatment	Worm burden (mean ± SD)				Total % reduction
		Couples	Males	Females	Total	
Intraperitoneal	Control	8.5 ± 3	5.4 ± 3.1	3.9 ± 1.4	26.3 ± 5.9	–
	eTAM	2.5 ± 1*	1.3 ± 0.9***	0.7 ± 0.7	7 ± 2.3*	73%
Oral	Control	12.2 ± 3.4	5.8 ± 1.9	1.8 ± 0.7	32 ± 6.4	–
	eTAM	6.7 ± 1.9**	0.9 ± 1****	0.5 ± 0.6	14.7 ± 3.5*	54%

Significant reduction when compared with control group,  $p < 0.05$

\* $p < 0.001$

\*\* $p = 0.0012$

\*\*\* $p = 0.021$

\*\*\*\* $p = 0.0051$

The online version of the original article can be found at <https://doi.org/10.1007/s00436-019-06259-0>

✉ Silmara M. Allegretti  
sallegre@unicamp.br

✉ Danilo C. Miguel  
dcmiguel@unicamp.br

<sup>1</sup> Biology Institute, Department of Animal Biology – Parasitology, University of Campinas – UNICAMP, Campinas, São Paulo, Brazil

<sup>2</sup> Present address: Biological and Health Sciences Section, Department of General Biology, Ponta Grossa State University, Ponta Grossa, Paraná, Brazil

**Table 2** Development singe of eggs retained in the intestinal tissues (oogram pattern); fecal egg count (EPG, number of eggs per gram of feces); and fecal egg count reduction (%) of mice experimentally infected with *S. mansoni* and treated with PBS (control group) or one single dose of 100 mg/kg cTAM

Administration route	Treatment	Oogram pattern			EPG	
		Immature eggs (%)	Mature eggs (%)	Dead ems (%)	Total	% Reduction
Intraperitoneal	Control	68.2 ± 4.6	26.2 ± 4.1	5.8 ± 1.7	1084 ± 322	–
	cTAM	18.1 ± 3.3*	70.5 ± 2*	7.2 ± 2.1	214 ± 53*	80%
Oral	Control	77.6 ± 6	18.4 ± 5.1	4 ± 1.7	1521 ± 339	–
	cTAM	13 ± 6.9*	66 ± 9.6*	13.6 ± 4.5	181 ± 91*	88%

\*Significant reduction when compared with control group,  $p < 0.0001$

The original article has been corrected.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.