



Diagnosis of peripheral bone and prosthetic joint infections: overview on the consensus documents by the EANM, EBJIS, and ESR (with ESCMID endorsement)

Luca Maria Sconfienza^{1,2} · Alberto Signore³ · Victor Cassar-Pullicino⁴ · Maria Adriana Cataldo⁵ · Olivier Gheysens⁶ · Olivier Borens⁷ · Andrej Trampuz⁸ · Klaus Wörtler⁹ · Nicola Petrosillo⁵ · Heinz Winkler¹⁰ · Filip M. H. M. Vanhoenacker^{11,12,13} · Paul C. Jutte¹⁴ · Andor W. J. M. Glaudemans¹⁵

Received: 7 February 2019 / Revised: 27 May 2019 / Accepted: 12 June 2019 / Published online: 27 June 2019
© European Society of Radiology 2019

Abstract

Objectives Peripheral bone infection (PBI) and prosthetic joint infection (PJI) are two different infectious conditions of the musculoskeletal system. They have in common to be quite challenging to be diagnosed and no clear diagnostic flowchart has been established. Thus, a conjoined initiative on these two topics has been initiated by the European Society of Radiology (ESR), the European Association of Nuclear Medicine (EANM), the European Bone and Joint Infection Society (EBJIS), and the European Society of Clinical Microbiology and Infectious Diseases (ESCMID). The purpose of this work is to provide an overview on the two consensus documents on PBI and PJI that originated by the conjoined work of the ESR, EANM, and EBJIS (with ESCMID endorsement).

Methods and results After literature search, a list of 18 statements for PBI and 25 statements for PJI were drafted in consensus on the most debated diagnostic challenges on these two topics, with emphasis on imaging.

Conclusions Overall, white blood cell scintigraphy and magnetic resonance imaging have individually demonstrated the highest diagnostic performance over other imaging modalities for the diagnosis of PBI and PJI. However, the choice of which advanced diagnostic modality to use first depends on several factors, such as the benefit for the patient, local experience of imaging

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00330-019-06326-1>) contains supplementary material, which is available to authorized users.

✉ Luca Maria Sconfienza
io@lucasconfienza.it

¹ IRCCS Istituto Ortopedico Galeazzi, Via Riccardo Galeazzi 4, 20161 Milan, Italy

² Dipartimento di Scienze Biomediche per la Salute, Università degli Studi di Milano, Milan, Italy

³ Nuclear Medicine Unit, Faculty of Medicine and Psychology, Department of Medical-Surgical Sciences and Translational Medicine, “Sapienza” University, Rome, Italy

⁴ Department of Diagnostic Imaging, The Robert Jones and Agnes Hunt Orthopaedic and District Hospital, Oswestry, Shropshire, UK

⁵ Infectious Disease Unit, National Institute for Infectious Diseases “L. Spallanzani”, IRCCS, Rome, Italy

⁶ Department of Nuclear Medicine and Molecular Imaging, University Hospitals Leuven, Leuven, Belgium

⁷ Division of Orthopaedic Surgery and Traumatology, Septic surgical unit, Lausanne University Hospital, University of Lausanne, Lausanne, Switzerland

⁸ Center for Musculoskeletal Surgery (CMSC), Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany

⁹ Department of Diagnostic and Interventional Radiology, Technische Universität München, Munich, Germany

¹⁰ Osteitis-Centre, Privatklinik Döbling, Vienna, Austria

¹¹ Department of Radiology, Antwerp University Hospital and Antwerp University, Antwerp, Belgium

¹² AZ Sint-Maarten, Belgium, Mechelen, Belgium

¹³ University of Ghent, Ghent, Belgium

¹⁴ Department of Orthopaedic Surgery, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

¹⁵ Department of Nuclear Medicine and Molecular Imaging, Medical Imaging Center, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

specialists, costs, and availability. Since robust, comparative studies among most tests do not exist, the proposed flowcharts are based not only on existing literature but also on the opinion of multiple experts involved on these topics.

Key Points

- *For peripheral bone infection and prosthetic joint infection, white blood cell and magnetic resonance imaging have individually demonstrated the highest diagnostic performance over other imaging modalities.*
- *Two evidence- and expert-based diagnostic flowcharts involving variable combination of laboratory tests, biopsy methods, and radiological and nuclear medicine imaging modalities are proposed by a multi-society expert panel.*
- *Clinical application of these flowcharts depends on several factors, such as the benefit for the patient, local experience, costs, and availability.*

Keywords Osteomyelitis · Prosthesis-related infections · Clinical laboratory techniques · Radiology · Nuclear medicine

Abbreviations

AGA	Anti-granulocyte antibody
CRP	C-reactive protein
CT	Computed tomography
EANM	European Society of Nuclear Medicine
EBJIS	European Bone and Joint Infection Society
ESCMID	European Society of Clinical Microbiology and Infectious Diseases
ESR	Erythrocyte sedimentation rate
ESR	European Society of Radiology
FDG-PET	Fluorodeoxyglucose-positron-emission tomography
HMPAO	Hexamethylpropylene amine oxime
MRI	Magnetic resonance imaging
OCEBM	Oxford Center for Evidence-based Medicine
PBI	Peripheral bone infection
PICO	Population/problem, intervention/indicator, comparator, outcome
PJI	Prosthesis joint infection
SPECT	Single-photon emitting computed tomography
WBC	White blood cell

Introduction

Peripheral bone infection (PBI) and prosthetic joint infection (PJI) represent two different infectious conditions of the musculoskeletal system [1, 2].

PBI include osteitis and osteomyelitis. Osteitis is a periosteal bacterial infection, which can develop acutely (< 8 weeks) or chronically (> 8 weeks) after trauma or surgery [3]. Osteomyelitis is classified into acute or chronic, too. Acutely, necrotic bone and bacteria are detected concurrently. Progression to chronic osteomyelitis is characterized by the presence of avascular bony fragments (*sequestra*) [4]. A strong periosteal reaction usually develops around the infection, with or without the presence of a sinus tract. PBI incidence is low, accounting for about 2% per year in developed countries, with slightly increase after surgery (2–4%), or

trauma surgery with potentially contaminated fracture (19%), or in immunocompromised hosts [5–8].

PJI is a complication of joint replacements [9]. Its incidence ranges from 2 to 2.4% for newly implanted prostheses, while it may reach up to 20% for revision procedures. Due to general population aging, the number of replaced joints is increasing. Thus, PJI represents a non-negligible health issue, leading to repeated surgery, prolonged hospitalization, increased morbidity, and increased costs [10, 11]. PJI can be differentiated as early (within 3 months from surgery), delayed (between 3 months and 2 years), and late (over 2 years). At any stage, PJI may present with a quite insidious onset and non-specific symptoms; thus, its diagnosis may be challenging [12–15].

No single routine test alone can diagnose PBI and PJI with sufficient accuracy. In most cases, imaging, clinical, microbiological, and laboratory examinations are performed based on their availability, physicians' experience, and economic considerations. Regarding PBI, current recommendations for its diagnosis are scarce, mainly based on local experiences, or lacking a multidisciplinary approach [5]. Regarding PJI, literature is richer; however, many recommendations still lack multidisciplinary approach, are not updated on the most recent imaging modalities, or fail providing a possible diagnostic flowchart [5].

Thus, an expert panel of radiologists, nuclear medicine physicians, orthopedic surgeons, and infectious disease specialists representing the European Society of Radiology (ESR), the European Association of Nuclear Medicine (EANM), the European Bone and Joint Infection Society (EBJIS), and the European Society of Clinical Microbiology and Infectious Diseases (ESCMID), respectively, performed a thorough systematic literature review and developed two consensus documents on the diagnostic management of PBI and PJI, with emphasis on radiologic and nuclear medicine modalities. These documents were recently published in the *European Journal of Nuclear Medicine and Molecular Imaging* [16, 17].

Our purpose is to provide an overview on the two consensus documents on PBI and PJI produced by the ESR, EANM, and EBJIS (with ESCMID endorsement).

Working group, statements, literature search, and scoring system

Details on these aspects are presented in the original studies [16–19] and provided as Supplementary Material.

Diagnosis of PBI

Pain is the principal acute local symptom of PBI associated with reduced function [20]. Physical examination may show a fistula with pus discharge, although frequently only mild skin redness and swelling can be seen. When a skin breach is present, the probe-to-bone test can be performed, consisting in inserting a metallic probe in the breach trying to reach the bone. The concept behind this simple test is that if the probe can reach the bone, the same can be done by infectious bacteria [21]. In chronic cases, however, symptoms are generally absent.

Statements

Full list of statements regarding PBI and their evidence levels is reported in Table 1.

- Clinical and laboratory parameters (statements no. 1 to no. 4, no. 7, no. 8)

When PBI is suspected, additional diagnostic tests may help confirming the diagnosis. If a skin breach is present, the probe-to-bone test can be performed. This represents a routine practice in the diagnosis of diabetic foot infection, although no clear evidence is published for PBI. Similarly, there is no evidence that a fistula directing to the bone and concurrent purulent discharge represent the proof of an underlying bone infection.

Regarding blood tests, C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), and white blood cell (WBC) count determination should always be performed. Increased ESR and CRP may be present and may suggest the presence of infection. WBC counts are more rarely increased. In patients with acute foot osteomyelitis, positive predictive value for infection of ESR in patients without diabetes was 78%, and in those with diabetes was 81%, with 58% and 31% negative predictive value, respectively [22]. A cutoff of 0.4 ng/ml of serum procalcitonin has been reported to be sensitive and specific in the diagnosis of acute osteomyelitis [23].

Blood cultures become positive mostly in hematogenous osteomyelitis. There is little evidence supporting the use of blood cultures in the diagnosis of PBI outside the spine [24].

To replace bone biopsy, the value of sinus tract and deep tissue cultures has been tested [25–29], finding it to be inadequate to identify the correct pathogen in osteomyelitis. One study showed different result, involving two sinus tract cultures

with bone contact at different times, showing 94% diagnostic accuracy in cases with two concordant cultures with bone contact [30]. Authors however conclude that this approach should not replace biopsy when this can be easily obtained.

Suspending antibiotic therapy prior to microbiological culture is a reasonable and common practice. However, regarding PBI, literature is insufficient to support or discourage this practice [30–32]. Thus, a good clinical practice should recommend that antibiotic administration is suspended or postponed only if feasible and reasonable, being the optimal duration of this suspension still not established.

- Radiological imaging techniques (statements no. 5, no. 6, no. 9 to no. 12)

In PBI, conventional radiography is the first modality to perform. However, conventional radiography becomes positive only after 7–14 days from symptom occurrence and when at least 30–50% of bone mass is lost. Sensitivity and specificity of conventional radiography in PBI range between 43–75% and 75–83%, respectively [33, 34]. Computed tomography (CT) has high resolution in the evaluation of peripheral bone; thus, it can replace conventional radiographs in anatomically complex locations (e.g., shoulder, pelvis). CT is particularly useful to detect bone sequestra and subtle findings such as gas bubbles and tiny area of cortical involvement [35].

In patients with clinical and imaging suspicion of PBI, current clinical practice suggests a percutaneous bone biopsy may be performed to identify the causative microorganism. However, evidence is low and conflicting. Some studies reported high accuracy (85.5–94%) [26, 30, 31] in identifying a single causative microorganism after biopsy. However, another study had low rate of positive specimens obtained by imaging-guided biopsy [32]. Superiority of open surgical biopsy has also been confirmed [26, 27, 36], remaining however a very invasive method of diagnosis.

Non-contrast magnetic resonance imaging (MRI) is pivotal in PBI, having 88–98% sensitivity, 70–96% specificity, and 81–86% accuracy [37–39]. MRI evaluates the involvement of both bone and soft tissues, being also able differentiating infection from other mimickers, such as bone tumors. Intravenous Gadolinium administration does not increase diagnostic performance but may help to better define the presence and extent of soft tissue abscesses and also to avoid overstaging by differentiating infection from edema [40]. Although no specific study addressed the issue of metallic fixation devices in patients with PBI, orthopedic implants in general do not represent a contraindication to MR examinations, with susceptibility artifact mostly limited to the profile of the implant itself [41].

- Nuclear medicine imaging techniques (statements no. 13 to no. 18)

Table 1 Statements on the diagnosis of peripheral bone infection

Item	Statement	Level of evidence
1	Patients presenting with clinical and radiological signs of peripheral bone infection or a positive probe-to-bone test may require further diagnostic procedures	5
2	Fistula direct to the bone and purulent discharge are evidence of bone infection	5
3	C-reactive protein, erythrocyte sedimentation rate, and white blood cell counts should always be performed in patients suspected to have peripheral bone infection for diagnostic purposes	4
4	Blood cultures should be considered in patients with fever suspected to have peripheral bone infection for diagnosing the involved bacteria	4
5	Conventional radiography is the first imaging modality to be performed in patients suspected to have peripheral bone infection for diagnosis and follow-up	3
6	In case of clinical and radiological signs of peripheral bone infection, bone biopsy is the reference standard for confirming infection and identifying the causative microorganism	4
7	In case of clinical and radiological signs of peripheral bone infection, sinus tract cultures and/or superficial swab cultures should be discouraged in the diagnostic workup; bone biopsy is the gold standard	4
8	Antibiotic therapy should be discontinued before biopsy	5
9	CT should be used as an adjunct to conventional radiographs in complex anatomic areas and is useful to detect bone sequestra	4
10	Non-contrast MRI has high diagnostic performance in detecting peripheral bone infection	2
11	Intravenous administration of Gadolinium-based contrast agents does not increase the diagnostic performance of MRI in peripheral bone infection	2
12	The presence of a metallic implant/fixation device is not a contraindication to perform MRI in patients with suspected peripheral bone infection	5
13	Three-phase bone scintigraphy is a sensitive technique in patients suspected for peripheral bone infection although not highly specific	2
14	White blood cell scintigraphy and anti-granulocyte antibody scintigraphy have similar high diagnostic accuracy for diagnosis of peripheral bone infection	2
15	Pre-test probability of infection should be considered for choosing between three-phase bone scan and white blood cell scintigraphy (fractures, recent surgery, osteosynthesis, highly positive serological tests)	5
16a	^{18}F -FDG-PET has high diagnostic accuracy in peripheral bone infection without fracture and osteosynthesis	2
16b	White blood cell scintigraphy is the preferred nuclear medicine imaging technique of choice in patients suspected of peripheral bone infection with recent fracture of hardware in situ	2
17	Hybrid SPECT-CT white blood cell imaging can be performed for exact localization of infection site	2
18	When having a suspicion for hematogenous spread of the infection, ^{18}F -FDG-PET/CT is the first imaging modality of choice	5

Source: reference [16]

CT, computed tomography; MRI, magnetic resonance imaging; ^{18}F -FDG-PET, ^{18}F -fluorodeoxyglucose-positron emitting tomography; SPECT, single-photon emission computed tomography

Three-phase bone scintigraphy is the most classic modality used in the skeletal system, with high sensitivity but low specificity in PBI [42–46]. Notably, the three phases can also be positive because of other reasons, such as post-traumatic edema, fracture healing, or recent surgery, thus explaining the low specificity especially after recent surgery. Single-photon emitting computed tomography (SPECT)/CT can be added in the late phase to improve localization of osteoblastic activity [47–54]. WBC and anti-granulocyte antibody (AGA) scintigraphy have been reported having a high diagnostic performance in PBI [45, 55–67]. However, since acquisition parameters and interpretation criteria differ between studies, there is a wide range in sensitivity and specificity. More recent WBC studies following the correct protocols show high diagnostic accuracies. Based on clinical practice, pre-test probability of infection (i.e., the probability of a patient having PBI before a diagnostic test result is known) should be considered when there is the option to choose between three-phase bone scan and WBC scintigraphy. Since in

the “violated” bone there is evidence of high sensitivity of three-phase bone scan, this should be reserved to patients in whom infection probability is low. Conversely, WBC scintigraphy should be regarded as the preferred nuclear medicine modality after fracture, surgery, or when a metallic implant is present. ^{18}F -fluorodeoxyglucose-positron-emitting tomography (FDG-PET) is a promising and accurate method in PBI without implanted hardware, although there is not enough evidence to propose it as reference standard, particularly in the acute phase [5, 42, 62, 64, 68–73]. Similarly, although evidence is absent and based on clinical practice, ^{18}F -FDG-PET/CT can be used when hematogenous dissemination in patients with PBI is suspected.

Diagnosis of PJI

When early PJI is suspected, local swelling, redness, pain, and pus discharge from the wound are the most common signs

and symptoms, particularly located over the surgical accessions. In later stages and in subacute or chronic phases, inflammatory signs may be absent. Pain and loss of function may be the only symptoms, which are almost impossible to differentiate from those caused by aseptic loosening [12–15, 74].

Statements

Full list of statements regarding PJI and their evidence levels is reported in Table 2.

- Clinical and laboratory parameters (statements no. 1 to no. 4, no. 7 to no. 12)

No specific study addresses the fact that sinus tracts with purulent discharge are signs of PJI. Thus, all patients with clinical suspicion of PJI should be ruled out for infection [63, 75–80]. CRP and ESR blood tests have variable diagnostic performance (sensitivity 21–100% and 58–97%; specificity 20–96% and 33–91%, respectively) in PJI. However, being quick and inexpensive, these tests should always be per-

Table 2 Statements on the diagnosis of prosthetic joint infection

Item	Statement	Level of evidence
1	Prosthetic joint infection should be suspected when one or more of the following symptoms and signs are present: otherwise unexplained pain and/or fever, redness, swelling, scar inflammation, and movement limitations. These symptoms are (especially in the chronic phase) not specific and require other investigations	4
2	Sinus tract and purulent discharge are clear signs of prosthetic joint infection	5
3	C-reactive protein and erythrocyte sedimentation rate should always be performed in patients suspected of prosthetic joint infection. A normal value does not rule out a prosthetic joint infection	2
4	In case of fever, blood cultures should always be performed in patients suspected to have prosthetic joint infection to identify the causative bacteria	5
5	Conventional radiographies are the first imaging modality to perform in patients with suspicion of prosthetic joint infection for diagnosis and follow-up	2
6	Ultrasound can detect complications around the prosthesis, but capability of detecting infection is controversial	2
7	Imaging guidance may be useful to guide joint aspiration or periprosthetic tissue biopsy	2
8	Leukocyte counts and differential of synovial fluid have high diagnostic accuracy to detect prosthetic joint infection	2
9	Bacterial culture from joint aspiration has high diagnostic accuracy to detect prosthetic joint infection	2
10	Measurement of the synovial biomarkers alpha-defensin, leukocyte esterase, interleukin-6, and C-reactive protein is useful in the detection of prosthetic joint infection	2
11	Biopsy of peri-prosthetic tissue for histology and cultures can be performed for pre-operative diagnosis in case erythrocyte sedimentation rate and/or C-reactive protein are positive and aspiration is inconclusive or impossible to test (dry tap)	2
12	Antibiotic therapy should be postponed or discontinued before pre- and intra-operative sampling	4
13	Antibiotic therapy should not be discontinued before white blood cell scintigraphy	4
14	CT can be effectively used to diagnose prosthetic joint infection	2
15	The diagnostic accuracy for 3-phase bone scintigraphy in patients with suspected infection within the first 2 years after hip or knee prosthesis placement is low	2
16	In case of negative 3-phase bone scintigraphy, the diagnosis of prosthetic joint infection can be excluded	2
17	In case of positive 3-phase bone scan, the addition of white blood cell scintigraphy leads to high diagnostic accuracy for prosthetic joint infection	2
18	In case of negative white blood cell scintigraphy, the probability of prosthetic joint infection is low	2
19	¹⁸ F-FDG-PET in patients suspected of prosthetic joint infection has high sensitivity but lower specificity than white blood cell scintigraphy or anti-granulocyte antibodies scintigraphy	2
20	Anti-granulocyte scintigraphy is a good alternative to white blood cell scintigraphy with similar sensitivity and specificity	2
21	Hybrid SPECT/CT imaging can improve localization of infection (and diagnostic accuracy)	2
22	Semi-quantitative analysis of white blood cell accumulation over time in white blood cell scan increases diagnostic accuracy for prosthetic joint infection	3
23	Combining white blood cell scan with bone marrow scan increases diagnostic accuracy for prosthetic joint infection detection	2
24	MRI is fully feasible in patients with suspicion of prosthetic joint infection	2
25	MRI has high diagnostic performance in detecting prosthetic joint infection when clinically suspected with no ionizing radiations	2

Source: reference [17]

CT, computed tomography; MRI, magnetic resonance imaging; ¹⁸F-FDG-PET, ¹⁸F-fluorodeoxyglucose-positron emitting tomography; SPECT, single-photon emission computed tomography

formed when PJI is suspected, using a threshold of 10 mg/l for CRP and 30 mm/h for ESR. The combination of the two markers seems to be even more reliable [81–85], although low CRP and ESR do not exclude PJI. No evidence suggests that blood culture is helpful in patients with fever and suspected PJI. However, when there is suspicion of hematogenous origin of PJI, blood culture might help.

Joint fluid aspiration has been traditionally used to rule out PJI, if there is enough fluid to aspirate. Different biomarkers can be tested. WBC count has 36–100% sensitivity and 80–99% specificity at a suggested threshold > 3000 cells/ μ l. Differential count has 84–100% sensitivity and 80–99% specificity at neutrophil percentage $> 70\%$ [79, 86–98]. Alpha defensin has 95.5–100% sensitivity and 95–100% specificity [89, 99–103] and seems not to be influenced by antibiotic treatment, while interleukin-6 has 62.5–97% sensitivity and 85.7–100% specificity [91, 103–107], although they are both expensive and not widely available tests. Leukocyte esterase has 66–100% sensitivity and 77–100% specificity [108–112]. Synovial CRP has 70–97.3% sensitivity and 78.6–100% specificity [89, 92, 103, 107, 113–116]. Bacterial culture has 43.5–100% sensitivity and 81.2–100% specificity [88, 93, 95, 104, 117–133]. Most authors recommend an incubation period ranging between 7 and 14 days, with at least five culture samples to confirm positivity [134]. Bacterial culture is limited by previous or ongoing antimicrobial therapy, which should be discontinued at least 2 weeks before sampling [96, 135–137].

In case of little or no fluid in the joint space, aspiration may be impossible (“dry tap”). Thus, biopsy of synovial tissue can be performed. This procedure is slightly more invasive than simple aspiration, with 79.1–100% sensitivity and 90–100% [105, 114, 138–141]. Blind procedures may lead to increased complications and lower success rate; thus, ultrasound guidance is highly recommended [141]. Ultrasound-guided aspiration has 67–69% sensitivity and 66–94% specificity [117, 142], while CT-guided aspiration has about 70% sensitivity and 100% specificity [143].

- Radiological imaging techniques (statements no. 5, no. 6, no. 14, no. 24, no. 25)

Conventional radiographs are the first imaging modality to perform in patients with suspected PJI, as they can evaluate bony structure around the implants, potentially showing the presence of abnormal findings, and being able to detect other causes of pain. As conventional radiographs become positive only when at least 30–50% of bone mass has been lost, about half of radiographs performed on infected implants are normal. Serial radiographs have 14% sensitivity and 70% specificity in PJI. The presence of gas bubbles and immature, active

periostitis are highly specific signs for PJI, while implant loosening, soft tissue swelling, and periprosthetic lucency have low specificity [144–146].

Ultrasound can show synovial hypertrophy and fluid around the implant, but its ability in detecting an infection is controversial [147, 148]. In the hip, a capsule-to-bone distance > 4 mm has been reported to be 100% sensitive and 74% specific, being 100% specific when the capsule-to-bone distance is over 3.2 mm and extra-capsular fluid is seen [149]. However, others reported that anterior distension of hip capsule is not predictive of infection [150].

PJI can be also ruled out using CT, which can show soft tissue collections, and distended bursae and joint spaces. When at least one soft tissue finding is used as infection criterion, CT has 100% sensitivity, 87% specificity, and 89% accuracy while it has 83% sensitivity, 96% specificity, and 94% accuracy when joint distention is used as infection criterion [151]. A 100% positive predictive value is assigned to the presence of fluid collections in perimuscular fat and muscle bellies, while 96% negative predictive value is assigned to absence of joint effusion. Periosteal reaction is 100% specific but only 16% sensitive for PJI [152, 153]. However, these figures are valid when these signs are present, but overall accuracy of CT is generally lower than that of MRI.

Joint implants are not a contraindication to MR examinations as signal distortion is generally limited to the area of prosthesis itself [154–156]. Regarding diagnostic performance of MRI in the detection of PJI, 65–92% sensitivity and 85–99% specificity have been reported in the knee, while these figures increase to 94% sensitivity and 97% specificity in the hip [157–160]. Notably, MRI (together with ultrasound) is a modality which does not involve ionizing radiation and may be preferred when possible.

- Nuclear medicine imaging techniques (statements no. 13, no. 15 to no. 23)

Three-phase bone scintigraphy is very sensitive to any bony remodeling. When a joint is replaced, remodeling may proceed for a couple of years. A single study investigated this issue, showing that at 21 months after surgery, three-phase bone scintigraphy had 50% sensitivity and 71% specificity, concluding that this examination should be avoided in the first years after surgery [161]. However, the great advantage of three-phase bone scan is that a negative examination allows excluding the diagnosis of PJI [162–170].

When three-phase bone scan is positive, two studies reported that the addition of WBC scintigraphy may





Fig. 1 **a** Proposed diagnostic flowchart based on published evidence to undertake when peripheral bone infection is suspected. Clearly, not all steps may be required in all cases and some steps may be repeated if necessary. Serological tests can be performed over time since the trend to increase or decrease is more important than a single value. At present, there is not enough clinical evidence to support the use of one advanced diagnostic imaging technique above the other. There is a lack of studies with large numbers of patients and there are hardly no comparative studies. Thus, the choice of which test to use first depends on several

factors, such as the benefit for the patient, local experience, costs, and availability. In many hospitals, magnetic resonance imaging is considered as first advanced imaging modality in daily practice, mainly because of no radiation involved. In patients with metallic hardware, however, there is sufficient literature to support a preferential use of white blood cell scintigraphy. **b** Suggested path to undertake when nuclear medicine techniques are considered in the suspicion of peripheral bone infection, based on literature evidence and expert opinion. Images are reproduced from reference [16] under creative common license

increase specificity [162, 171]. With this association, they found 80% sensitivity and 99.5% specificity for PJI. Within the first years after surgery, three-phase bone scan may be avoided since it will be for sure positive due to bone remodeling and WBC scintigraphy can be used as first nuclear imaging modality. Regarding the issue of suspending antibiotic therapy before WBC scintigraphy, no study addresses specifically this issue. However, studies on antibiotic discontinuation in PBI showed no accuracy difference between the two options [172–174]. If WBC scintigraphy is negative, the diagnosis of PJI is unlikely, with negative predictive values ranging from 92% using ^{99m}Tc -hexamethylpropylene amine oxime (HMPAO)-labeled leukocytes [175] to 100% using sulesumab (AGA) [175]. In addition to conventional qualitative evaluation, semi-quantitative analysis of WBC accumulation at 3–4 h and 20–24 h after leukocyte injection can be performed. Quantification is expressed as a ratio between radioactivity

in the region of interest over background radioactivity. If ratio increases over time, WBC accumulation is reported as active and is interpreted as PJI. Conversely, a ratio decreasing over time suggests a non-infectious inflammatory process [167, 176]. The combination of WBC scintigraphy with bone marrow scintigraphy has been reported to increase the detection of PJI, being able to reduce false-positive cases particularly in doubtful cases at WBC scintigraphy. The correct interpretation criteria to be used have also been recently published by the EANM [177]. The combination of those modalities has been reported having accuracy ranging from 83 to 98% for both ^{99m}Tc -HMPAO-WBC and ^{111}In -oxine-WBC for both knee and hip PJI [168, 178–185]. A valid alternative to WBC scintigraphy is represented by AGA scintigraphy. Two meta-analyses showed that AGA scintigraphy has 83% sensitivity and 79–80% specificity in the diagnosis of PJI [186, 187], which is comparable to those of WBC scintigraphy. The

Fig. 2 **a** Proposed diagnostic flowchart based on published evidence to undertake when prosthetic joint infection is suspected. Some tests can be repeated (i.e., blood cultures, bone biopsies, or soft tissue biopsies) when needed. Serological tests (C-reactive protein, white blood cell count with differential, and erythrocyte sedimentation rate) should be performed over time since the trend to increase or decrease is more important than a single value. The choice of which test to use first depends on several factors, such as the benefit for the patient, local experience, costs, and availability. **b** Suggested path to undertake when nuclear medicine techniques are considered in the suspicion of prosthetic joint infection, based on literature evidence and expert opinion. Initial stratification is based on time after implant (more/less than 2 years). Images are reproduced from reference [17] under creative common license



comparison between ^{99m}Tc -WBC and ^{99m}Tc -besilesomab in peripheral osteomyelitis and PJI revealed no differences [56, 68, 188]. Although current recommendations on WBC scintigraphy include planar image evaluation only, the introduction of SPECT/CT has represented a remarkable advancement. Three studies showed accuracy increased up to 38% in patients with PJI [68, 174, 189]. Thus, in cases of positive WBC scintigraphy, SPECT/CT scan may provide additional information.

Few studies compared ^{18}F -FDG-PET directly to WBC and/or AGA scintigraphy [179, 190–192]. Overall, the comparison showed that ^{18}F -FDG-PET has higher sensitivity but lower specificity when PJI is suspected. However, the wide ranges of reported sensitivities (28–91%) and specificities (34–97%) are justified by different study designs and interpretation criteria. Thus, higher standardization is warranted to increase homogeneity.

Concerns on the use of ionizing radiations

Details on these aspects are presented in the original works [16–19, 193–196] and provided as Supplementary Material.

Conclusions

Overall, WBC scintigraphy and MRI have demonstrated individually the highest diagnostic performance over the other imaging modalities in the diagnosis of both PBI and PJI. However, the choice of which test to use first depends on several factors, such as the benefit for the patient, local experience, costs, and availability.

Figures 1 and 2 summarize the proposed diagnostic flowcharts based on published evidence and the suggested paths to

undertake when nuclear medicine techniques are considered in the suspicion of PBI and PJI, respectively.

Concluding, these flowcharts represent the first evidence, PICO-based proposals to be applied when PBI and PJI are suspected. However, since robust, comparative studies among most tests do not exist, these flowcharts also involve expert opinion based on broad consensus of multiple experts involved on these topics.

Acknowledgements The original guidelines published on the *European Journal of Nuclear Medicine and Molecular Imaging* [16, 17] were drafted by the representatives of the respective Scientific Societies: Luca Maria Sconfienza, Victor Cassar-Pullicino, Klaus Wortler, and Filip MHM Vanhoenacker are delegates of the European Society of Radiology (ESR); Andor W.J.M. Glaudemans, Olivier Gheysens, and Alberto Signore are delegates of the European Association of Nuclear Medicine (EANM); Paul C. Jutte, Olivier Borens, and Heinz Winkler are delegates of the European Bone and Joint Infection Society (EBJIS); Nicola Petrosillo, Maria Adriana Cataldo, and Andrej Trampuz are delegates of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID).

Funding The authors state that this work has not received any funding.

Compliance with ethical standards

Guarantor The scientific guarantor of this publication is Luca Maria Sconfienza.

Conflict of interest The authors of this manuscript declare no relationships with any companies, whose products or services may be related to the subject matter of the article.

Statistics and biometry No complex statistical methods were necessary for this paper.

Informed consent Not applicable.

Ethical approval Institutional Review Board approval was not required because this study does not involve patients directly.

References

1. Tiemann AH, Hofmann GO (2009) Principles of the therapy of bone infections in adult extremities: are there any new developments? *Strategies Trauma Limb Reconstr* 4:57–64
2. Kurtz S, Mowat F, Ong K, Chan N, Lau E, Halpern M (2005) Prevalence of primary and revision total hip and knee arthroplasty in the United States from 1990 through 2002. *J Bone Joint Surg Am* 87:1487–1497
3. Lew DP, Waldvogel FA (2004) Osteomyelitis. *Lancet* 364:369–379
4. Mouzopoulos G, Kanakaris NK, Kontakis G, Obakponwre O, Townsend R, Giannoudis PV (2011) Management of bone infections in adults: the surgeon's and microbiologist's perspectives. *Injury*. 42(Suppl 5):S18–S23
5. Jutte P, Lazzeri E, Sconfienza LM et al (2014) Diagnostic flowcharts in osteomyelitis, spondylodiscitis and prosthetic joint infection. *Q J Nucl Med Mol Imaging* 58:2–19
6. Ochsner P, Bodler P-M, Eich G et al (2014) Infections of the musculoskeletal system. Swiss orthopaedics in-house publisher, Grandvaux
7. Korim MT, Payne R, Bhatia M (2014) A case-control study of surgical site infection following operative fixation of fractures of the ankle in a large U.K. trauma unit. *Bone Joint J* 96-B:636–640
8. Malhotra AK, Goldberg S, Graham J et al (2014) Open extremity fractures: impact of delay in operative debridement and irrigation. *J Trauma Acute Care Surg* 76:1201–1207
9. Uselli FG, Indino C, Maccario C, Manzi L, Liuni FM, Vulcano E (2017) Infections in primary total ankle replacement: anterior approach versus lateral transfibular approach. *Foot Ankle Surg* S1268-7731(17):30790–30797
10. Glaudemans AW, Galli F, Pacilio M, Signore A (2013) Leukocyte and bacteria imaging in prosthetic joint infection. *Eur Cell Mater* 25:61–77
11. Cataldo MA, Petrosillo N, Cipriani M, Cauda R, Tacconelli E (2010) Prosthetic joint infection: recent developments in diagnosis and management. *J Infect* 61:443–448
12. Trampuz A, Zimmerli W (2005) Prosthetic joint infections: update in diagnosis and treatment. *Swiss Med Wkly* 135:243–251
13. Berbari EF, Hanssen AD, Duffy MC et al (1998) Risk factors for prosthetic joint infection: case-control study. *Clin Infect Dis* 27:1247–1254
14. Zimmerli W, Trampuz A, Ochsner PE (2004) Prosthetic-joint infections. *N Engl J Med* 351:1645–1654
15. Drago L, De Vecchi E, Bortolin M, Zagra L, Romanò CL, Cappelletti L (2017) Epidemiology and antibiotic resistance of late prosthetic knee and hip infections. *J Arthroplasty* 32:2496–2500
16. Glaudemans AWJM, Jutte PC, Cataldo MA et al (2019) Consensus document for the diagnosis of peripheral bone infection in adults: a joint paper by the EANM, EBJIS, and ESR (with ESCMID endorsement). *EJNMMI*. <https://doi.org/10.1007/s00259-019-4262-x>
17. Signore A, Sconfienza LM, Borens O et al (2019) Consensus document for the diagnosis of prosthetic joint infections: a joint paper by the EANM, EBJIS, and ESR (with ESCMID endorsement). *EJNMMI*. <https://doi.org/10.1007/s00259-019-4263-9>
18. The 2011 Oxford CEBM levels of evidence (introductory document). <https://www.cebm.net/2016/05/ocebmllevels-of-evidence/> Accessed on 30 Jan 2019
19. OCEBM Levels of Evidence Working Group. The Oxford 2011 Levels of Evidence. <https://www.cebm.net/wp-content/uploads/2014/06/CEBM-Levels-of-Evidence-2.1.pdf> Accessed on 30 Jan 2019
20. Cierny G III, Mader JT, Penninck JJ (2003) The classic: a clinical staging system for adult osteomyelitis. *Clin Orthop* 414:7–24
21. Lavery LA, Armstrong DG (2007) Temperature monitoring to assess, predict, and prevent diabetic foot complications. *Curr Diab Rep* 7:416–419
22. Rabjohn L, Roberts K, Troiano M, Schoenhaus H (2007) Diagnostic and prognostic value of erythrocyte sedimentation rate in contiguous osteomyelitis of the foot and ankle. *J Foot Ankle Surg* 46:230–237
23. Maharajan K, Patro DK, Menon J et al (2013) Serum Procalcitonin is a sensitive and specific marker in the diagnosis of septic arthritis and acute osteomyelitis. *J Orthop Surg Res* 8:19
24. Roger PM, Cua E, Courjon J, Landraud L, Carles M, Bernard E (2014) The impact of bacteremia on the outcome of bone infections. *Med Mal Infect* 44:380–386
25. Zuluaga AF, Galvis W, Saldarriaga JG, Agudelo M, Salazar BE, Vesga O (2006) Etiologic diagnosis of chronic osteomyelitis: a prospective study. *Arch Intern Med* 166:95–100

26. Zuluaga AF, Galvis W, Jaimes F, Vesga O (2002) Lack of microbiological concordance between bone and non-bone specimens in chronic osteomyelitis: an observational study. *BMC Infect Dis* 2:8
27. Akinyoola AL, Adegbehingbe OO, Aboderin AO (2009) Therapeutic decision in chronic osteomyelitis: sinus track culture versus intraoperative bone culture. *Arch Orthop Trauma Surg* 129:449–453
28. Agarwal S, Zahid M, Sherwani MK, Abbas M, Huda N, Khan AQ (2005) Comparison of the results of sinus track culture and sequestrum culture in chronic osteomyelitis. *Acta Orthop Belg* 71:209–212
29. Lindfors N, Geurts J, Drago L et al (2017) Antibacterial bioactive glass, S53P4, for chronic bone infections - a multinational study. *Adv Exp Med Biol* 971:81–92
30. Bernard L, Uckay I, Vuagnat A et al (2010) Two consecutive deep sinus tract cultures predict the pathogen of osteomyelitis. *Int J Infect Dis* 14:e390–e393
31. Khatri G, Wagner DK, Sohnle PG (2001) Effect of bone biopsy in guiding antimicrobial therapy for osteomyelitis complicating open wounds. *Am J Med Sci* 321:367–371
32. Wu JS, Gorbachova T, Morrison WB, Haims AH (2007) Imaging-guided bone biopsy for osteomyelitis: are there factors associated with positive or negative cultures? *AJR Am J Roentgenol* 188: 1529–1534
33. Boutin R (1998) Update on imaging of musculoskeletal infections. *Orthop Clin North Am* 29:41–66
34. David R, Barron BJ, Modewell JE (1987) Osteomyelitis, acute and chronic. *Radiol Clin North Am* 25:1171–1201
35. Gold RH, Hawkins RA, Katz RD (1991) Bacterial osteomyelitis: findings on plain radiography, CT, MR, and scintigraphy. *AJR Am J Roentgenol* 157:365–370
36. Merlet A, Cazanave C, Dauchy FA et al (2014) Prognostic factors of calcaneal osteomyelitis. *Scand J Infect Dis* 46:555–560
37. Erdman WA, Tamburro F, Jayson HT, Weatherall PT, Ferry KB, Peshock RM (1991) Osteomyelitis: characteristics and pitfalls of diagnosis with MR imaging. *Radiology*. 180:533–539
38. Morrison WB, Schweitzer ME, Bock GW et al (1993) Diagnosis of osteomyelitis: utility of fat-suppressed contrast-enhanced MR imaging. *Radiology*. 189:251–257
39. Demirev A, Weijers R, Geurts J, Mottaghy F, Walenkamp G, Brans B (2014) Comparison of 18F-FDG PET/CT and MRI in the diagnosis of acute osteomyelitis. *Skeletal Radiol* 43:665–672
40. Averill LW, Hernandez A, Gonzalez L, Peña AH, Jaramillo D (2009) Diagnosis of osteomyelitis in children: utility of fat-suppressed contrast-enhanced MRI. *AJR Am J Roentgenol* 192: 1232–1238
41. Bellelli A, Silvestri E, Barile A et al (2019) Position paper on magnetic resonance imaging protocols in the musculoskeletal system (excluding the spine) by the Italian College of Musculoskeletal Radiology. *Radiol Med*. <https://doi.org/10.1007/s11547-019-00992-3>
42. Wang GL, Zhao K, Liu ZF, Dong MJ, Yang SY (2011) A meta-analysis of fluorodeoxyglucose-positron emission tomography versus scintigraphy in the evaluation of suspected osteomyelitis. *Nucl Med Commun* 32:1134–1142
43. Hatzenbuehler J, Pulling TJ (2011) Diagnosis and management of osteomyelitis. *Am Fam Physician* 84:1027–1033
44. van der Bruggen W, Bleeker-Rovers CP, Boerman OC, Gotthardt M, Oyen WJ (2010) PET and SPECT in osteomyelitis and prosthetic bone and joint infections: a systematic review. *Semin Nucl Med* 40:3–15
45. Prandini N, Lazzeri E, Rossi B, Erba P, Parisella MG, Signore A (2006) Nuclear medicine imaging of bone infections. *Nucl Med Commun* 27:633–644
46. Termaat MF, Raijmakers PG, Scholten HJ, Bakker FC, Patka P, Haarman HJ (2005) The accuracy of diagnostic imaging for the assessment of chronic osteomyelitis: a systematic review and meta-analysis. *J Bone Joint Surg Am* 87:2464–2471
47. Erba PA, Glaudemans AW, Veltman NC et al (2014) Image acquisition and interpretation criteria for 99mTc-HMPAO-labelled white blood cell scintigraphy: results of a multicentre study. *Eur J Nucl Med Mol Imaging* 41:615–623
48. Thang SP, Tong AK, Lam WW, Ng DC (2014) SPECT/CT in musculoskeletal infections. *Semin Musculoskelet Radiol* 18: 194–202
49. Cyteval C, Bourdon A (2012) Imaging orthopedic implant infections. *Diagn Interv Imaging* 93:547–557
50. Sanlı Y, Ozkan ZG, Unal SN, Türkmen C, Kılıçoğlu O (2011) The additional value of Tc 99m HMPAO white blood cell SPECT in the evaluation of bone and soft tissue infections. *Mol Imaging Radionucl Ther* 20:7–13
51. Vijayanathan S, Butt S, Gnanasegaran G, Groves AM (2009) Advantages and limitations of imaging the musculoskeletal system by conventional radiological, radionuclide, and hybrid modalities. *Semin Nucl Med* 39:357–368
52. Schillaci O (2009) Hybrid imaging systems in the diagnosis of osteomyelitis and prosthetic joint infection. *Q J Nucl Med Mol Imaging* 53:95–104
53. Filippi L, Schillaci O (2006) Usefulness of hybrid SPECT/CT in 99mTc-HMPAO-labeled leukocyte scintigraphy for bone and joint infections. *J Nucl Med* 47:1908–1913
54. Bar-Shalom R, Yefremov N, Guralnik L et al (2006) SPECT/CT using 67Ga and 111In-labeled leukocyte scintigraphy for diagnosis of infection. *J Nucl Med* 47:587–594
55. Glaudemans AW, de Vries EF, Vermeulen LE, Slart RH, Dierckx RA, Signore A (2013) A large retrospective single centre study to define the best image acquisition protocols and interpretation criteria for white blood cell scintigraphy with 99mTc-HMPAO-labelled leukocytes in musculoskeletal infections. *Eur J Nucl Med Mol Imaging* 40:1760–1769
56. Richter WS, Ivancevic V, Meller J et al (2011) 99mTc-besilesomab (Scintimun) in peripheral osteomyelitis: comparison with 99mTc-labelled white blood cells. *Eur J Nucl Med Mol Imaging* 38:899–910
57. Vittorini E, Del Giudice E, Pizzoli A, Caudana R (2005) MRI versus scintigraphy with 99mTc-HMPAO-labelled granulocytes in the diagnosis of bone infection. *Radiol Med* 109:395–403
58. Hughes DK (2003) Nuclear medicine and infection detection: the relative effectiveness of imaging with 111In-oxine, 99mTc-HMPAO-, and 99mTc-stannous fluoride colloid-labeled leukocytes and with 67Ga-citrate. *J Nucl Med Technol* 31:196–201
59. Wolf G, Aigner RM, Schwarz T (2001) Diagnosis of bone infection using 99mTc-HMPAO labeled leukocytes. *Nucl Med Commun* 22:1201–1206
60. Quirce R, Carril JM, Gutiérrez-Mendiguchia C et al (2002) Assessment of the diagnostic capacity of planar scintigraphy and SPECT with 99mTc-HMPAO-labelled leukocytes in superficial and deep sternal infections after median sternotomy. *Nucl Med Commun* 23:453–459
61. Becker W, Meller J (2001) The role of nuclear medicine in infection and inflammation. *Lancet Infect Dis* 1:326–333
62. Sonmezoglu K, Sonmezoglu M, Halac M et al (2001) Usefulness of 99mTc-ciprofloxacin (infecton) scan in diagnosis of chronic orthopedic infections: comparative study with 99mTc-HMPAO leukocyte scintigraphy. *J Nucl Med* 42:567–574
63. Teller RE, Christie MJ, Martin W, Nance EP, Haas DW (2000) Sequential indium-labeled leukocyte and bone scans to diagnose prosthetic joint infection. *Clin Orthop Relat Res* 372:241–247
64. Meller J, Liersch T, Oezerden MM et al (2010) Targeting NCA-95 and other granulocyte antigens and receptors with radiolabeled monoclonal antibodies (Mabs). *Q J Nucl Med Mol Imaging* 54: 582–598

65. Skehan SJ, White JF, Evans JW et al (2003) Mechanism of accumulation of 99mTc-sulesomab in inflammation. *J Nucl Med* 44: 11–18
66. Ryan PJ (2002) Leukoscan for orthopaedic imaging in clinical practice. *Nucl Med Commun* 23:707–714
67. Devillers A, Garin E, Polardt JL et al (2000) Comparison of Tc-99m-labelled antileukocyte fragment Fab' and Tc-99m-HMPAO leukocyte scintigraphy in the diagnosis of bone and joint infections : a prospective study. *Nucl Med Commun* 21:747–753
68. Palestro CJ (2015) Radionuclide imaging of osteomyelitis. *Semin Nucl Med* 45:32–46
69. Demirev A, Weijers R, Geurts J, Mottaghy F, Walenkamp G, Brans B (2014) Comparison of [18 F] FDG PET/CT and MRI in the diagnosis of active osteomyelitis. *Skeletal Radiol* 43:665–672
70. Basu S, Chrysosikis T, Moghadam-Kia S, Zhuang H, Torigian DA, Alavi A (2009) Positron emission tomography as a diagnostic tool in infection: present role and future possibilities. *Semin Nucl Med* 39:36–51
71. Strobel K, Stumpe KD (2007) PET/CT in musculoskeletal infection. *Semin Musculoskelet Radiol* 11:353–364
72. Meller J, Sahlmann CO, Liersch T, Hao Tang P, Alavi A (2007) Nonprosthesis orthopedic applications of (18) F fluoro-2-deoxy-D-glucose PET in the detection of osteomyelitis. *Radiol Clin North Am* 45:719–733
73. Hartmann A, Eid K, Dora C, Trentz O, von Schulthess GK, Stumpe KD (2007) Diagnostic value of 18F-FDG PET/CT in trauma patients with suspected chronic osteomyelitis. *Eur J Nucl Med Mol Imaging* 34:704–714
74. Galbusera F, Volkheimer D, Reitmaier S, Berger-Roscher N, Kienle A, Wilke HJ (2015) Pedicle screw loosening: a clinically relevant complication? *Eur Spine J* 24:1005–1016
75. Gille J, Ince A, González O, Katzer A, Loehr JF (2006) Single-stage revision of peri-prosthetic infection following total elbow replacement. *J Bone Joint Surg Br* 88:1341–1346
76. Achermann Y, Stasch P, Preiss S, Lucke K, Vogt M (2014) Characteristics and treatment outcomes of 69 cases with early prosthetic joint infections of the hip and knee. *Infection*. 42:511–519
77. Garcia RM, Hardy BT, Kraay MJ, Goldberg VM (2010) Revision total knee arthroplasty for aseptic and septic causes in patients with rheumatoid arthritis. *Clin Orthop Relat Res* 468:82–89
78. Magnuson JE, Brown ML, Hauser MF, Berquist TH, Fitzgerald RH Jr, Klee GG (1998) In-111-labeled leukocyte scintigraphy in suspected orthopedic prosthesis infection: comparison with other imaging modalities. *Radiology* 168:235–239
79. Virolainen P, Lahteenmaki H, Hiltunen A, Sipola E, Meurman O, Nelimarkka O (2002) The reliability of diagnosis of infection during revision arthroplasties. *Scand J Surg* 91:178–181
80. Bailey DL, Pichler BJ, Gückel B et al (2016) Combined PET/MRI: from status quo to status go. Summary report of the fifth international workshop on PET/MR imaging; February 15–19, 2016; Tübingen, Germany. *Mol Imaging Biol* 18:637–650
81. Aljanipour P, Bakhshi H, Parvizi J (2013) Diagnosis of periprosthetic joint infection: the threshold for serologic markers. *Clin Orthop Relat Res* 471:3186–3195
82. Berbari E, Mabry T, Tsaras G et al (2010) Inflammatory blood laboratory levels as markers of prosthetic joint infection: a systematic review and meta-analysis. *J Bone Joint Surg Am* 92:2102–2109
83. Baré J, MacDonald SJ, Bourne RB (2006) Preoperative evaluations in revision total knee arthroplasty. *Clin Orthop Relat Res* 446:40–44
84. De Vecchi E, Villa F, Bortolin M et al (2016) Leucocyte esterase, glucose and C-reactive protein in the diagnosis of prosthetic joint infections: a prospective study. *Clin Microbiol Infect* 22:555–560
85. Villa F, Toscano M, De Vecchi E, Bortolin M, Drago L (2017) Reliability of a multiplex PCR system for diagnosis of early and late prosthetic joint infections before and after broth enrichment. *Int J Med Microbiol* 307:363–370
86. Trampuz A, Zimmerli W (2008) Diagnosis and treatment of implant-associated septic arthritis and osteomyelitis. *Curr Infect Dis Rep* 10:394–403
87. Ali S, Christie A, Chapel A (2009) The pattern of procalcitonin in primary total hip and knee arthroplasty and its implication in periprosthetic infection. *J Clin Med Res* 1:90–94
88. Bergin PF, Doppelt JD, Hamilton WG et al (2010) Detection of periprosthetic infections with use of ribosomal RNA-based polymerase chain reaction. *J Bone Joint Surg Am* 92:654–663
89. Cipriano CA, Brown NM, Michael AM, Moric M, Sporer SM, Della Valle CJ (2012) Serum and synovial fluid analysis for diagnosing chronic periprosthetic infection in patients with inflammatory arthritis. *J Bone Joint Surg Am* 94:594–600
90. Della Valle GJ, Sporer SM, Jacobs JJ, Berger RA, Rosenberg AG, Paprosky WG (2007) Preoperative testing for sepsis before revision total knee arthroplasty. *J Arthroplasty* 22(6 suppl 2):90–93
91. Nilsson-Augustinsson A, Briheim G, Herder A, Ljunghusen O, Wahlström O, Ohman L (2007) Inflammatory response in 85 patients with loosened hip prostheses: a prospective study comparing inflammatory markers in patients with aseptic and septic prosthetic loosening. *Acta Orthop* 78:629–639
92. Bottner F, Wegner A, Winkelmann W, Becker K, Erren M, Götze C (2007) Interleukin-6, procalcitonin and TNF- α : markers of periprosthetic infection following total joint replacement. *J Bone Joint Surg Br* 89:94–99
93. Schinsky MF, Della Valle CJ, Sporer SM, Paprosky WG (2008) Perioperative testing for joint infection in patients undergoing revision total hip arthroplasty. *J Bone Joint Surg Am* 90:1869–1875
94. Greidanus NV, Masri BA, Garbus DS et al (2007) Use of erythrocyte sedimentation rate and C-reactive protein level to diagnose infection before revision total knee arthroplasty: a prospective evaluation. *J Bone Joint Surg Am* 89:1409–1416
95. Spanghel MJ, Masri BA, O'Connell JX, Duncan CP (1998) Prospective analysis of preoperative and intraoperative investigations for the diagnosis of infection at the sites of two hundred and two revision total hip arthroplasties. *J Bone Joint Surg Am* 81: 672–683
96. Trampuz A, Piper KE, Hanssen AD et al (2006) Sonication of explanted prosthetic components in bags for diagnosis of prosthetic joint infection is associated with risk of contamination. *J Clin Microbiol* 44:628–631
97. Vasoo S, Cunningham SA, Greenwood-Quaintance KE et al (2015) Evaluation of the film array blood culture ID panel on biofilms dislodged from explanted arthroplasties for prosthetic joint infection diagnosis. *J Clin Microbiol* 53:2790–2792
98. Zmistowsky B, Restrepo C, Huang R, Hozack WJ, Parvizi J (2012) Periprosthetic joint infection diagnosis: a complete understanding of white blood cell count and differential. *J Arthroplasty* 27:1589–1593
99. Drago L, Toscano M, Tacchini L, Banfi G (2017) α -Defensin point-of-care test for diagnosis of prosthetic joint infections: neglected role of laboratory and clinical pathologists. *Clin Chem Lab Med* 27(56):19–24
100. Gallo J, Kolar M, Dendis M et al (2008) Culture and PCR analysis of joint fluid in the diagnosis of prosthetic joint infection. *New Microbiol* 31:97–104
101. Bingham J, Clarke H, Spanghel M, Schwartz A, Beauchamp C, Goldberg B (2014) The alpha defensin-1 biomarker assay can be used to evaluate the potentially infected total joint arthroplasty. *Clin Orthop Relat Res* 472:4006–4009

102. Lenski M, Scherer MA (2014) Synovial IL-6 as inflammatory marker in periprosthetic joint infections. *J Arthroplasty* 29:1105–1109
103. Jacovides CL, Parvizi J, Adeli B, Jung KA (2011) Molecular markers for diagnosis of periprosthetic joint infection. *J Arthroplasty* 26(6 suppl):99–103
104. Ghanem E, Ketonis C, Restrepo C, Joshi A, Barrack R, Parvizi J (2009) Periprosthetic infection: where do we stand with regard to gram stain? *Acta Orthop* 80:37–40
105. Fink B, Makowiak C, Fuerst M, Berger I, Schäfer P, Frommelt L (2008) The value of synovial biopsy, joint aspiration and C-reactive protein in the diagnosis of late peri-prosthetic infection of total knee replacements. *J Bone Joint Surg Br* 90:874–878
106. Mikkelsen DB, Pedersen C, Højbjerg T, Schønheyder HC (2006) Culture of multiple peroperative biopsies and diagnosis of infected knee arthroplasties. *APMIS* 114:449–452
107. Esteban J, Alonso-Rodriguez N, del-Prado G et al (2012) PCR-hybridization after sonication improves diagnosis of implant-related infection. *Acta Orthop* 83:299–304
108. Deirmengian C, Kardos K, Kilmartin P, Cameron A, Schiller K, Parvizi J (2014) Combined measurement of synovial fluid alpha-defensin and C-reactive protein levels: highly accurate for diagnosing periprosthetic joint infection. *J Bone Joint Surg Am* 96:1439–1445
109. Deirmengian C, Kardos K, Kilmartin P, Cameron A, Schiller K, Parvizi J (2014) Diagnosing periprosthetic joint infection: has the era of the biomarker arrived? *Clin Orthop Relat Res* 472:3254–3262
110. Ali F, Wilkinson JM, Cooper JR et al (2006) Accuracy of joint aspiration for the preoperative diagnosis of infection in total hip arthroplasty. *J Arthroplasty* 21:221–226
111. Achermann Y, Vogt M, Leunig M, Wüst J, Trampuz J (2010) Improved diagnosis of periprosthetic joint infection by multiplex PCR of sonication fluid from removed implants. *J Clin Microbiol* 48:1208–1214
112. Wetters NG, Berend KR, Lombardi AV, Morris MJ, Tucker TL, Della Valle CJ (2012) Leukocyte esterase reagent strips for the rapid diagnosis of periprosthetic joint infection. *J Arthroplasty* 27(8 suppl):8–11
113. Parvizi J, Jacovides C, Adeli B, Jung KA, Hozack WJ (2012) Mark B. Coventry award: synovial C-reactive protein: a prospective evaluation of a molecular marker for periprosthetic knee joint infection. *Clin Orthop Relat Res* 470:54–60
114. Tetreault MW, Wetters NG, Moric M, Gross CE, Della Valle CJ (2014) Is synovial C-reactive protein a useful marker for periprosthetic joint infection? *Clin Orthop Relat Res* 472:3997–4003
115. Gomez E, Cazanave C, Cunningham SA et al (2012) Prosthetic joint infection diagnosis using broad-range PCR of biofilms dislodged from knee and hip arthroplasty surfaces using sonication. *J Clin Microbiol* 50:3501–3508
116. Tunney MM, Patrick S, Curran MD et al (1999) Detection of prosthetic hip infection at revision arthroplasty by immunofluorescence microscopy and PCR amplification of the bacterial 16S rRNA gene. *J Clin Microbiol* 37:3281–3290
117. Battaglia M, Vannini F, Guaraldi F, Rossi G, Biondi F, Sudanese A (2011) Validity of preoperative ultrasound-guided aspiration in the revision of hip prosthesis. *Ultrasound Med Biol* 37:1977–1983
118. Hindle P, Davidson E, Biant LC (2012) Septic arthritis of the knee: the use and effect of antibiotics prior to diagnostic aspiration. *Ann R Coll Surg Engl* 94:351–355
119. Atkins BL, Athanasou N, Deeks JJ et al (1998) Prospective evaluation of criteria for microbiological diagnosis of prosthetic-joint infection at revision arthroplasty. *J Clin Microbiol* 36:2932–2939
120. Cazanave C, Greenwood-Quaintance KE, Hanssen AD et al (2013) Rapid molecular microbiologic diagnosis of prosthetic joint infection. *J Clin Microbiol* 51:2280–2287
121. Esteban J, Gomez-Barrena E, Cordero J, Martín-de-Hijas NZ, Kinnari TJ, Fernandez-Roblas R (2008) Evaluation of quantitative analysis of cultures from sonicated retrieved orthopedic implants in diagnosis of orthopedic infection. *J Clin Microbiol* 46:488–492
122. Cheung A, Lachiewicz PF, Renner JB (1997) The role of aspiration and contrast-enhanced arthrography in evaluating the uncemented hip arthroplasty. *AJR Am J Roentgenol* 168:1305–1309
123. Howard CB, Einhorn M, Dagan R, Yagupski P, Porat S (1994) Fine-needle bone biopsy to diagnose osteomyelitis. *J Bone Joint Surg Br* 76:311–314
124. Bonhoeffer J, Haeberle B, Schaad UB, Heininger U (2001) Diagnosis of acute haematogenous osteomyelitis and septic arthritis, 20 years experience at the university children's hospital Basel. *Swiss Med Wkly* 131:575–581
125. Barrack RL, Harris WH (1993) The value of aspiration of the hip joint before revision total hip arthroplasty. *J Bone Joint Surg Am* 75:66–76
126. Portillo ME, Salvadó M, Sorli L et al (2012) Multiplex PCR of sonication fluid accurately differentiates between prosthetic joint infection and aseptic failure. *J Infect* 65:541–548
127. Janz V, Wassilew GI, Kribus M, Trampuz A, Perka C (2015) Improved identification of polymicrobial infection in total knee arthroplasty through sonicate fluid cultures. *Arch Orthop Trauma Surg* 135:1453–1457
128. Minassian AM, Newham R, Kalimeris E, Bejon P, Atkins BL, Bowler ICJW (2014) Use of an automated blood culture system (BD BACTEC) for the diagnosis of prosthetic joint infections: easy and fast. *BMC Infect Dis* 14:233
129. Frangiamore SJ, Gajewski ND, Saleh A, Farias-Kovac M, Barsoum WK, Higuera CA (2016) Alfa-defensin accuracy to diagnose periprosthetic joint infection - best available test? *J Arthroplasty* 31:456–460
130. Ettinger M, Calliess T, Kielstein JT et al (2015) Circulating biomarkers for discrimination between aseptic joint failure, low-grade infection and high grade septic failure. *Clin Infect Dis* 61:332–341
131. Frangiamore SJ, Saleh A, Grosso MJ et al (2015) α -Defensin as a predictor of periprosthetic shoulder infection. *J Shoulder Elbow Surg* 24:1021–1027
132. Cipriano C, Maiti A, Hale G, Jiranek W (2014) The host response: toll-like receptor expression in periprosthetic tissues as a biomarker for deep joint infection. *J Bone Joint Surg Am* 96:1692–1698
133. Liu JZ, Saleh A, Klika AK, Barsoum WK, Higuera CA (2014) Serum inflammatory markers for periprosthetic knee infection in obese versus non-obese patients. *J Arthroplasty* 29:1880–1883
134. Drago L, De Vecchi E, Cappelletti L et al (2015) Prolonging culture to 15 days improves bacterial detection in bone and joint infections. *Eur J Clin Microbiol Infect Dis* 34:1809–1813
135. Trampuz A, Piper KE, Jacobson MJ et al (2007) Sonication of removed hip and knee prostheses for diagnosis of infection. *N Engl J Med* 357:654–663
136. Spangehl MJ, Masterson E, Masri BA, O'Connell JX, Duncan CP (1999) The role of intraoperative gram stain in the diagnosis of infections during revision total hip arthroplasty. *J Arthroplasty* 14:952–956
137. Barrack RL, Jennings RW, Wolfe MW, Bertot AJ (1997) The Coventry Award. The value of preoperative aspiration before total knee revision. *Clin Orthop Relat Res* (345) 8–16
138. Fink B, Gebhard A, Fuerst M, Berger I, Schäfer P (2013) High diagnostic value of synovial biopsy in periprosthetic joint infection of the hip. *Clin Orthop Relat Res* 471:956–964

139. Rak M, Barlic-Maganja D, Kavcic M, Trebse R, Cor A (2013) Comparison of molecular and culture method in diagnosis of prosthetic joint infection. *FEMS Microbiol Lett* 343:42–48
140. Morawietz L, Tiddens O, Mueller M et al (2009) Twenty-three neutrophil granulocytes in 10 high-power fields is the best histopathological threshold to differentiate between aseptic and septic endoprosthesis loosening. *Histopathology* 54:847–853
141. Silvestri E, Barile A, Albano D et al (2018) Interventional therapeutic procedures in the musculoskeletal system: an Italian survey by the Italian College of Musculoskeletal Radiology. *Radiol Med* 123:314–321
142. Eisler T, Svensson O, Engström CF et al (2001) Ultrasound for diagnosis of infection in revision total hip arthroplasty. *J Arthroplasty* 16:1010–1017
143. Tomas X, Bori G, Garcia S et al (2011) Accuracy of CT-guided joint aspiration in patients with suspected infection status post-total hip arthroplasty. *Skeletal Radiol* 40:57–64
144. Lyons CW, Berquist TH, Lyons JC, Rand JA, Brown ML (1985) Evaluation of radiographic findings in painful hip arthroplasties. *Clin Orthop Relat Res* (195) 239–251
145. Tigges S, Stiles RG, Roberson JR (1994) Appearance of septic hip prostheses on plain radiographs. *AJR Am J Roentgenol* 163:377–380
146. Math KR, Zaidi SF, Petchprapa C, Harwin SF (2006) Imaging of total knee arthroplasty. *Semin Musculoskelet Radiol* 10:47–63
147. Tagliafico A, Bignotti B, Rossi F, Sconfienza LM, Messina C, Martinoli C (2017) Ultrasound of the hip joint, soft tissues, and nerves. *Semin Musculoskelet Radiol* 21:582–588
148. Sdao S, Orlandi D, Aliprandi A et al (2014) The role of ultrasonography in the assessment of peri-prosthetic hip complications. *J Ultrasound* 18:245–250
149. Van Holsbeeck MT, Eyler WR, Sherman LS et al (1994) Detection of infection in loosened hip prostheses: efficacy of sonography. *AJR Am J Roentgenol* 163:381–384
150. Weybright PN, Jacobson JA, Murry KH et al (2003) Limited effectiveness of sonography in revealing hip joint effusion: preliminary results in 21 adult patients with native and postoperative hips. *AJR Am J Roentgenol* 181:215–218
151. Cyteval C, Hamm V, Sarabère MP, Lopez FM, Maury P, Taourel P (2002) Painful infection at the site of hip prosthesis: CT imaging. *Radiology*. 224:477–483
152. Cahir JG, Toms AP, Marshall TJ, Wimbust J, Nolan J (2007) CT and MRI of hip arthroplasty. *Clin Radiol* 62:1163–1171
153. Roth TD, Maertz NA, Parr JA, Buckwalter KA, Choplin RH (2012) CT of the hip prosthesis: appearance of components, fixation, and complications. *Radiographics*. 32:1089–1107
154. Talbot BS, Weinberg EP (2016) MR imaging with metal-suppression sequences for evaluation of total joint arthroplasty. *Radiographics*. 36:209–225
155. Sneag DB, Bogner EA, Potter HG (2015) Magnetic resonance imaging evaluation of the painful total knee arthroplasty. *Semin Musculoskelet Radiol* 19:40–48
156. Aliprandi A, Sconfienza LM, Randelli F, Bandirali M, Di Leo G, Sardanelli F (2012) Magnetic resonance imaging of painful total hip replacement: detection and characterisation of periprosthetic fluid collection and interobserver reproducibility. *Radiol Med* 117: 85–95
157. Aliprandi A, Sconfienza LM, Randelli P et al (2011) Magnetic resonance imaging of the knee after medial unicompartmental arthroplasty. *Eur J Radiol* 80:e416–e421
158. White LM, Kim JK, Mehta M et al (2000) Complications of total hip arthroplasty: MR imaging-initial experience. *Radiology*. 215: 254–262
159. Hayter CL, Koff MF, Shah P et al (2011) MRI after arthroplasty: comparison of MAVRIC and conventional fast spin-echo techniques. *AJR Am J Roentgenol* 197:W405–W411
160. He C, Lu Y, Jiang M, Feng J, Wang Y, Liu Z (2014) Clinical value of optimized magnetic resonance imaging for evaluation of patients with painful hip arthroplasty. *Chin Med J (Engl)* 127: 3876–3880
161. Wong MY, Beadsmoore C, Toms A, Smith T, Donell S (2012) Does 99mTc-MDP bone scintigraphy add to the investigation of patients with symptomatic unicompartmental knee replacement? *Knee*. 19:592–596
162. Ikeuchi M, Okanoue Y, Izumi M et al (2013) Diagnostic value of triple-phase bone scintigraphy for the diagnosis of infection around antibiotic-impregnated cement spacers. *Springerplus*. 27(2):401
163. Nagoya S, Kaya M, Sasaki M, Tateda K, Yamashita T (2008) Diagnosis of peri-prosthetic infection at the hip using triple-phase bone scintigraphy. *J Bone Joint Surg Br* 90:140–144
164. Volpe L, Indelli PF, Latella L, Poli P, Yakupoglu J, Marcucci M (2015) Periprosthetic joint infections: a clinical practice algorithm. *Joints*. 13(2):169–174
165. Trevail C, Ravindranath-Reddy P, Sulkin T, Bartlett G (2016) An evaluation of the role of nuclear medicine imaging in the diagnosis of periprosthetic infections of the hip. *Clin Radiol* 71:211–219
166. Ouyang Z, Li H, Liu X, Zhai Z, Li X (2014) Prosthesis infection: diagnosis after total joint arthroplasty with three-phase bone scintigraphy. *Ann Nucl Med* 28:994–1003
167. Glaudemans AW, de Vries EF, Vermeulen LE, Slart RH, Dierckx RA, Signore A (2013) A large retrospective single-centre study to define the best image acquisition protocols and interpretation criteria for white blood cell scintigraphy with ^{99m}Tc-HMPAO-labelled leucocytes in musculoskeletal infections. *Eur J Nucl Med Mol Imaging* 40:1760–1769
168. Lima AL, Oliveira PR, Carvalho VC, Saconi ES, Cabrita HB, Rodrigues MB (2013) Periprosthetic joint infections. *Interdiscip Perspect Infect Dis* 2013:542796
169. Love C, Marwin SE, Palestro CJ (2009) Nuclear medicine and the infected joint replacement. *Semin Nucl Med* 39:66–78
170. Temmerman OP, Raijmakers PG, Berkhof J et al (2006) Diagnostic accuracy and interobserver variability of plain radiography, subtraction arthrography, nuclear arthrography, and bone scintigraphy in the assessment of aseptic femoral component loosening. *Arch Orthop Trauma Surg* 126:316–323
171. Jamar F, Buscombe J, Chiti A et al (2013) EANM/SNMMI guideline for 18F-FDG use in inflammation and infection. *J Nucl Med* 54:647–658
172. Vesco L, Boulahdour H, Hamissa S et al (1999) The value of combined radionuclide and magnetic resonance imaging in the diagnosis and conservative management of minimal or localized osteomyelitis of the foot in diabetic patients. *Metabolism*. 48:922–927
173. Newman LG, Waller J, Palestro CJ et al (1991) Unsuspected osteomyelitis in diabetic foot ulcers. Diagnosis and monitoring by leukocyte scanning with indium 111 oxyquinoline. *JAMA*. 266: 1246–1251
174. Chik KK, Magee MA, Bruce WJ et al (1996) Tc-99m stannous colloid-labeled leukocyte scintigraphy in the evaluation of the painful arthroplasty. *Clin Nucl Med* 21:838–843
175. Kim HO, Na SJ, Oh SJ et al (2014) Usefulness of adding SPECT/CT to 99mTc-hexamethylpropylene amine oxime (HMPAO)-labeled leukocyte imaging for diagnosing prosthetic joint infections. *J Comput Assist Tomogr* 38:313–319
176. Pelosi E, Baiocco C, Pennone M et al (2004) 99mTc-HMPAO-leukocyte scintigraphy in patients with symptomatic total hip or knee arthroplasty: improved diagnostic accuracy by means of semiquantitative evaluation. *J Nucl Med* 45:438–444
177. Signore A, Jamar F, Israel O, Buscombe J, Martin-Comin J, Lazzeri E (2018) Clinical indications, image acquisition and data interpretation for white blood cells and anti-granulocyte

- monoclonal antibody scintigraphy: an EANM procedural guideline. *Eur J Nucl Med Mol Imaging* 45:1816–1831
178. Gemmel F, Van den Wyngaert H, Love C, Welling MM, Gemmel P, Palestro CJ (2012) Prosthetic joint infections: radionuclide state-of-the-art imaging. *Eur J Nucl Med Mol Imaging* 39:892–909
 179. Love C, Marwin SE, Tomas MB et al (2004) Diagnosing infection in the failed joint replacement: a comparison of coincidence detection 18F-FDG and 111In-labeled leukocyte/99mTc-sulfur colloid marrow imaging. *J Nucl Med* 45:1864–1871
 180. Simonsen L, Buhl A, Oersnes T, Duus B (2007) White blood cell scintigraphy for differentiation of infection and aseptic loosening: a retrospective study of 76 painful hip prostheses. *Acta Orthop* 78:640–647
 181. Mulamba L, Ferrant A, Leners N, de Nayer P, Rombouts JJ, Vincent A (1983) Indium-111 leukocyte scanning in the evaluation of painful hip arthroplasty. *Acta Orthop Scand* 54:695–697
 182. Palestro CJ, Kim CK, Swyer AJ, Capozzi JD, Solomon RW, Goldsmith SJ (1990) Total-hip arthroplasty: periprosthetic indium-111-labeled leukocyte activity and complementary technetium-99m-sulfur colloid imaging in suspected infection. *J Nucl Med* 31:1950–1955
 183. Palestro CJ, Love C, Tronco GG, Tomas MB, Rini JN (2006) Combined labeled leukocyte and technetium 99m sulfur colloid bone marrow imaging for diagnosing musculoskeletal infection. *Radiographics*. 26:859–870
 184. Palestro CJ, Swyer AJ, Kim CK, Goldsmith SJ (1991) Infected knee prosthesis: diagnosis with In-111 leukocyte, Tc-99m sulfur colloid, and Tc-99m MDP imaging. *Radiology*. 179:645–648
 185. El Espera I, Blondet C, Moullart V et al (2004) The usefulness of 99mTc sulfur colloid bone marrow scintigraphy combined with 111In leukocyte scintigraphy in prosthetic joint infection. *Nucl Med Commun* 25:171–175
 186. Pakos EE, Trikalinos TA, Fotopoulos AD, Ioannidis JP (2007) Prosthesis infection: diagnosis after total joint arthroplasty with antigranulocyte scintigraphy with 99mTc-labeled monoclonal antibodies—a meta-analysis. *Radiology*. 242:101–108
 187. Xing D, Ma X, Ma J, Wang J, Chen Y, Yang Y (2013) Use of anti-granulocyte scintigraphy with 99mTc-labeled monoclonal antibodies for the diagnosis of periprosthetic infection in patients after total joint arthroplasty: a diagnostic meta-analysis. *PLoS One* 26(8):e69857
 188. Sousa R, Massada M, Pereira A, Fontes F, Amorim I, Oliveira A (2011) Diagnostic accuracy of combined 99mTc-sulesomab and 99mTc-nanocolloid bone marrow imaging in detecting prosthetic joint infection. *Nucl Med Commun* 32:834–839
 189. Filippi L, Schillaci O (2006) Usefulness of hybrid SPECT/CT in 99mTc-HMPAO-labeled leukocyte scintigraphy for bone and joint infections. *J Nucl Med* 47:1908–1913
 190. Vanquickenborne B, Maes A, Nuyts J et al (2003) The value of (18)FDG-PET for the detection of infected hip prosthesis. *Eur J Nucl Med Mol Imaging* 30:705–715
 191. Pill SG, Parvizi J, Tang PH et al (2006) Comparison of fluorodeoxyglucose positron emission tomography and (111)indium-white blood cell imaging in the diagnosis of periprosthetic infection of the hip. *J Arthroplasty* 21(6 Suppl 2): 91–97
 192. Glaudemans AW, Prandini N, DI Girolamo M et al (2018) Hybrid imaging of musculoskeletal infections. *Q J Nucl Med Mol Imaging* 62:3–1
 193. European Union, directive 2013/59/EURATOM. <https://ec.europa.eu/energy/sites/ener/files/documents/CELEX-32013L0059-EN-TXT.pdf>. Accessed on 30 Jan 2019
 194. Messina C, Banfi G, Aliprandi A et al (2016) Ultrasound guidance to perform intra-articular injection of gadolinium-based contrast material for magnetic resonance arthrography as an alternative to fluoroscopy: the time is now. *Eur Radiol* 26:1221–1225
 195. European Society of Radiology, EuroSafe Imaging Campaign. <http://www.eurosafeimaging.org/> Accessed on 30 Jan 2019
 196. European Association of Nuclear Medicine. Radiation protection education program. https://www.eanm.org/content-eanm/uploads/2016/12/EANM_2016-TG_RadiationProtection_lowres.pdf Accessed on 30 Jan 2019

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.