



Heterogeneous tumor features and treatment outcome between males and females with lung cancer (LC): Do gender and sex matter?

Stefano Frega^b, Alessandro Dal Maso^{a,b}, Alessandra Ferro^{a,b}, Laura Bonanno^a, PierFranco Conte^{a,b}, Giulia Pasello^{a,*}

^a Department of Surgical, Oncological and Gastroenterological Sciences, University of Padova, Italy

^b Medical Oncology 2, Istituto Oncologico Veneto IOV-IRCCS, Padova, Italy

ARTICLE INFO

Keywords:

Gender
Sex
Lung cancer

ABSTRACT

Lung cancer (LC) is the leading cause of cancer-related death worldwide, despite a decreasing incidence rate in recent years, especially in men. Most risk factors for LC could be linked to an individual's reproductive system and secondary sex characteristics ('sex-related') and/or to some physical, behavioral and personality traits ('gender-related') peculiar to males rather than females or vice versa. An imbalance of these etiologic factors could explain why some LC features may differ between sexes.

For this review, an extended literature data collection was performed, using keywords to identify 'sex/gender' and 'LC'.

Differences between genders in LC epidemiology, pathological and molecular characteristics, loco-regional and/or systemic treatments outcome and prognosis were systematically analyzed. The possible predictive role of physio-pathological factors in males and females paves the way for a personalized therapeutic approach, emphasizing the need to include gender as a stratification factor in future clinical trials design.

1. Introduction

Lung cancer (LC) is the leading cause of cancer-related death worldwide both in men and women, accounting for 1.8 million deaths in 2018 (Bray et al., 2018). Among all LC cases, about 85% are non-small cell lung cancer (NSCLC), with two main histological types: adenocarcinoma (ADC) and squamous cell carcinoma (SCC) (Bray et al., 2018; Herbst et al., 2018). While tobacco smoking is the most important risk factor for LC, a subset of NSCLC patients (approximately 10–40%) are never-smoker or light-smoker, and in this case the etiology is less clear (Özdemir et al., 2018; Siegel et al., 2019).

Over the last two decades, the therapeutic options for patients affected by LC have greatly expanded, mainly due to the increased use of anti-angiogenic agents, targeted therapies and immune checkpoints inhibitors (ICIs) (Herbst et al., 2018). A patient's sex was not normally used as a preplanned stratification factor in randomized control trials (RCTs) testing different anticancer agents and loco-regional approaches. Current assumptions about a patient's sex as a predictive and/or prognostic factor come from analysis of heterogeneous patient populations or meta-analysis of heterogeneous trials (Jemal et al., 2018). For these reasons, comprehensive knowledge about a possible

predictive/prognostic role of gender in NSCLC is lacking.

In this review, we present data showing differences between men and women who develop LC in terms of epidemiology, risk factors, treatment outcome and prognosis.

2. Review methods

An extended literature data collection through the free search engine PubMed was performed, using keywords related to patient sex ('sex', 'gender', 'male/female', 'men/women') and LC epidemiological, etiological and clinical-pathological features. Data were systematically analyzed to outline putative differences between males and females, highlighting the available level of evidence, when necessary. In the second part of the review, we looked for potential heterogeneous efficacy of loco-regional (radiotherapy, surgery) and systemic treatments (chemotherapy, anti-angiogenic agents, targeted therapies, ICIs) in small cell lung cancer (SCLC) and NSCLC in males *versus* females. Data on limited/early disease stage were separately analyzed from those of extensive/metastatic stage to verify the impact of a patient's sex on outcome in both conditions. The review considered the results of more than 35 randomized controlled trials (RCTs) including patients affected

* Corresponding author at: Division of Medical Oncology 2, Veneto Institute of Oncology (IOV), Via Gattamelata 64, 35128 Padova Italy.

E-mail address: giulia.pasello@iov.veneto.it (G. Pasello).

by LC, across different tumor histologies, disease stages and treatment types; useful contributions coming from non-RCTs were also taken into account, and the meta-analytic and/or retrospective nature of these studies was specified in the text.

3. Epidemiology of LC in males and females

LC has been the most common diagnosed cancer in the world for several decades, and also the most common cause of death due to cancer. Worldwide, LC remains the leading cause of cancer incidence and mortality, with 2.1 million new lung cancer cases and 1.8 million deaths predicted in 2018, accounting for 18.4% of cancer deaths (Bray et al., 2018). LC is the most common cancer in men worldwide (1.3 million new cases/year), with the highest estimated age-standardized incidence rates in Eastern Europe (49.3 per 100,000/year) and Eastern Asia (47.2). Notably, low incidence rates are observed in Middle and Western Africa (3.8 and 2.4 respectively). In women, the incidence rates are generally lower and the geographical pattern is slightly different, mainly reflecting different historical exposure to tobacco smoke. Thus, the highest estimated rates are in Northern America (30.7 per 100,000/year) and Northern Europe (26.9) with a relatively high rate in Eastern Asia (19.2) and the lowest rates again in Western and Middle Africa (1.2 and 2.3 respectively) (Bray et al., 2018).

Over the last 40 years in the U.S., LC incidence rate has constantly declined in men (from 90 per 100,000 in 1975 to 71 per 100,000 in 2010–2015), while it had an increasing trend in women (from 25 per 100,000 in 1975 to 52.3 per 100,000 in 2010–2015), beginning to decline only since the mid-2000's. The decline has been more consistent in the past decade, with rates decreasing from 2011 to 2015 by almost 3% per year in men, but only 1.5% per year in women (Siegel et al., 2019). In the U.S. LC incidence rates are now higher among young women than among young men, with the pattern confined to non-Hispanic whites and Hispanics born since the mid-1960s (Jemal et al., 2018). In contrast to men, most European countries are still observing a rising trend in LC incidence among women (Bray et al., 2018).

In the U.S. the LC death rate has declined by 48% since 1990 in men and by 23% since 2002 in women due to reductions in smoking, and from 2012 to 2016 the rate decreased by about 4% per year in men and 3% per year in women (Siegel et al., 2019). Anyway, among males, LC remains the leading cause of death in most countries in Eastern Europe, Western Asia, Northern Africa, and specific countries in Eastern Asia and South-Eastern Asia. Among females, LC is the leading cause of cancer death in 28 countries, with the highest incidence rates seen in North America, Northern and Western Europe (Bray et al., 2018).

4. Risk factors for LC in males and females

Males and females are possibly differently predisposed to develop LC, due to a series of distinct and/or unbalanced risk factors: those ascribable to male/female biological features, hereafter indicated as 'sex-related', and those indicated as 'gender-related', being more linked to male/female behavior.

4.1. Sex-related risk factors

4.1.1. Sex hormones

Sex hormones could have a key role considering that estrogen receptors (ERs) are often expressed in LC and that estrogens potentially influence tumor growth (Schwartz et al., 2005). In particular, hormone replacement therapy seems to increase incidence of, and mortality from lung cancer in women (Chakraborty et al., 2010).

On the contrary, anti-estrogen use in women was found to be correlated with a reduced risk of lung cancer incidence (Chu et al., 2017). Blood estrogen level is normally higher in females than males, but ERs are not exclusively expressed in lung cancer of women. However, data suggest that estrogens activate lung adenocarcinoma cell lines derived

from women, but not from men (Dougherty et al., 2006; Ivanova et al., 2010). ER alpha (ER α) and ER beta (ER β) are two types of classical ERs, with the latter appearing to be the predominant isoform in lung cancer (Hsu et al., 2017). The location on lung cancer cell (cytoplasm/nucleus) and the prognostic value of each ER have not been univocally reported in literature, and may vary according to a patient's sex (Schwartz et al., 2005). On the contrary, few data exist on an eventual androgen role.

The effects of male sex steroids are mediated by the androgen receptor (AR), expressed in type II pneumocytes and in bronchial epithelium of murine lung model, after androgen stimulation (Mikkonen et al., 2010). In this model, androgens significantly altered lung gene expression profiles, by up-regulating transcripts involved in oxygen transport and down-regulating those responsible for DNA repair and recombination. The carcinogenic effect of androgens might reside in this cytotoxic effect, but it might also reside in the down-regulation of the systemic immune response, as seen in different immunological disorders where the immune-suppressive action of testosterone has been demonstrated (Trigunaite et al., 2015). A population-based cohort study on men aged 70–88 years (n = 3635) showed that a high testosterone level was associated with LC, even after exclusion of current smoker subjects (Hyde et al., 2012). Androgens might play a role in lung cancer pathophysiology, promoting both its origin and progression; androgen deprivation therapy has been found to have a benefic effect on survival after lung cancer diagnosis (HR 0.36; $p = 0.0007$), according to a retrospective analysis (n=3018, APM treated=339) (Harlos et al., 2015).

4.1.2. DNA adducts and DNA repair systems

Tobacco smoke contains a multitude of carcinogens, which exert their biologic effect through the formation of DNA adducts in lung tissue. Carcinogens are metabolized and detoxified within the lung by enzymes that oxidize the hydrocarbons, producing reactive oxygen species or intermediates, later neutralized into water soluble conjugates. Reactive intermediates that are not detoxified bind DNA into DNA adducts, playing a role in lung carcinogenesis (Hemminki, 1993). It has been hypothesized that women are more susceptible to tobacco carcinogens than men. ERs are present in both normal and neoplastic lung tissues and could accelerate the metabolism of tobacco carcinogens in a dose-dependent way, as suggested by higher levels of polycyclic aromatic hydrocarbons (PAH)-related DNA adducts in female smokers compared to males (Berge et al., 2004). Inherited genetic polymorphisms affecting activating and detoxifying enzymes could explain a different susceptibility between sexes to tobacco carcinogens. The cytochrome P-4501A1 (CYP1A1) gene codes for an enzyme involved in DNA adduct formation and induced by cigarette smoke (Kawajiri et al., 1993). Despite substantially fewer pack-years and younger age, female smokers have been found to have significantly higher levels of CYP1A1 expression and of DNA adducts than men. Increased CYP1A1 expression might be a result of hormone induction, mostly estrogens. A cross-talk between ERs and the aryl hydrocarbon receptor, a regulator of CYP1A1, has been demonstrated in breast cancer cell lines (Thomsen et al., 1994). The most common gene involved in neutralizing adducts is glutathione S-transferase M1 (GSTM1). A GSTM1 homozygous deletion (GSTM1 null) genotype, which is present in 40%–60% of the general population, results in a reduced expression and in an increased risk for smoking-related cancers. Polymerase chain reaction analysis of peripheral blood indicated that women had a greater cancer risk than men (odds ratio [OR] 4.98 versus 1.37) if they harbor a mutant CYP1A1 genotype. The absence of a functional GSTM1 enzyme alone was not associated with an increased risk of LC, but the CYP1A1 mutation and the GSTM1 null genotype are significantly more frequent in female cancer patients than female controls. The combined variant genotypes conferred an odds ratio of 6.54 for LC in women compared with 2.36 in men. This risk was not affected by age or by smoking history (Dresler et al., 2000; Seidegård et al., 1990).

DNA repair capacity (DRC) seems to be involved in LC pathogenesis and treatment outcome. Wei et al. looked at the impact of DRC in the pathogenesis of LC by measuring the ability of patients' lung tissue to identify and repair induced DNA adducts. Patients younger than 60 years, those with a family history of LC, and women had lower DRC and a higher risk for LC than the average population (Wei et al., 2000). These findings were confirmed in a similar case-control study by (Spitz et al. (2001).

4.2. Gender-related risk factors

4.2.1. Tobacco use

A smoking habit continues to be the primary risk factor for developing LC.

The World Health Organization (WHO) estimates that 20.2% of the world's population aged ≥ 15 years were current smokers in 2015, indicating that smoking rates have decreased by 6.7% globally since 2000 and by 4.1% since 2005 (Anon., 2015). From 2000 onwards, this decreasing trend in smoking rate was registered for both sexes, being faster in men than in women, and equal to -0.22% for year in women and -0.50% for year in men in the period 2010–2015. Nevertheless, smoking remains globally far less common among adult females (6.4%) than among males (34.1%).

Such differences may relate to a combination of physiological, cultural and behavioral factors. Despite the worldwide predominance of male smokers, female smoking is dominated by higher prevalence in the Americas and European regions. In these countries, the differences in smoking behavior between women and men have decreased over time, and currently 12.2% of American women smoke cigarettes compared with 15.8% of men (Wang et al., 2018). The prevalence of smoking in American women peaked in 1965 at 33% and remained elevated throughout the 70s before beginning to slowly decrease in 1980. By contrast, more than half of American men smoked before 1965, but the prevalence dramatically decreased during the subsequent 20 years (Giovino, 2002). In last decades, the age-standardized prevalence of tobacco smoking has decreased more slowly in women *versus* men in Europe, with this trend expected to be confirmed also in the period 2010–2025 (Anon., 2018).

Second-hand smoke is another established risk factor for LC: approximately 20% of women with LC are non-smokers and women married to men who smoke have been shown to have a 25%–29% increased risk of developing LC (North C and Christiani, 2013).

Besides the difference in tobacco exposure between men and women, several behavioral lifestyles related to smoking habits as well as environmental and occupational exposures are found to be differently expressed, thus potentially representing other gender-related LC risk factors (Kiyohara and Ohno, 2010; Cote et al., 2009; Mollerup et al., 2006; Subramanian and Govindan, 2007).

4.2.2. Environmental exposures

Environmental exposures known to be carcinogenic to the lung include second hand smoke (SHS), asbestos, arsenic, radon, polycyclic aromatic hydrocarbons, cadmium, nickel, metal dusts and vinyl chloride (Kligerman and White, 2011). Moreover, indoor burning of cooking oil and other biomass fuels in poorly ventilated areas produce polycyclic aromatic hydrocarbons (PAHs), which are associated with LC. This effect is seen especially in East and South Asian women, but is significant in all developing countries (Sun et al., 2007). In LC attributed to household coal burning exposure, lung microbiota may play a potential etiopathogenetic role in females (Hosgood et al., 2014).

4.2.3. Other risk factors

Dietary patterns could influence LC risk differently in the two sexes. Among men, inverse relationships with vitamin C, folate, and carotenoids, and positive associations with total fat, monounsaturated and saturated fat were observed after adjusting for age, education,

cigarettes/day, years smoking, and total energy intake (Bandera et al., 1997); diet did not appear to exert a major role on lung cancer risk among women. However, in LC of never-smokers of both gender, a protective effect was suggested for vegetables/carrots and a deleterious effect for cultured milk products (Nyberg et al., 1998); milk resulted a risk factor only among male high-consumers. According to a case-control study based on the results of community mass screening (LC = 363, control subjects = 1089), an inverse association between body mass index (BMI) and LC was observed in men after adjustment for age and smoking, while no association was found in women (Kanashiki et al., 2005). Some preclinical studies have shown that lung microbiome (LM) composition after exposure to external stimuli might differ by sex (Barfod et al., 2015), and recent studies have shown that certain respiratory microbes and microbiota dysbiosis are correlated with development of LC (Mao et al., 2018). However, the impact of LM in LC risk has not yet been described to differ between males and females.

5. Clinical-pathological and molecular features of LC in males and females

SCC, SCLC and large-cell carcinoma (LCC) rates declined since the 1990s for both gender, but less rapidly among females (Lewis et al., 2014).

Since then, the incidence of ADC rose worldwide, and nowadays ADC represents globally the most common type of LC both in men and women (Parkin et al., 2010; Cheng et al., 2016). In last decades in most countries ADC rates remained relatively constant in males, while it increased in females, thus representing an important medical issue (Weiderpass et al., 2014; Meza et al., 2015).

The subset of lung ADC once recognized as bronchoalveolar carcinoma (BAC) disproportionately affects women (Raz et al., 2016) but, since the latest WHO classification of LC discontinued the term BAC in favor of "lepidic" (Travis et al., 2015a), no data is currently available about sex-difference incidence in lepidic ADC subtype; not even any valid information about men-women disparities in other ADC subtypes (acinar, solid, papillary, micropapillary) exists.

Some sex-oriented pathological LC features might depend on different co-etiological factors, such as tobacco exposure (Travis et al., 2015b), carcinogen-metabolizing enzymes expression and/or oncogenic virus prevalence. Moreover, important acquired molecular differences exist between males and females affected by LC.

NSCLC tumor samples of female smoker patients, rather than males, generally present higher frequency of mutations in critical driver genes, such as p53 and KRAS (Mollerup et al., 2006; Foeglé et al., 2007; Barrera and Morales Fuentes, 2012).

Nelson et al. reported a significant association between female sex and KRAS mutation in lung ADC tissue after adjustment for carcinogen exposures (OR 3.3; 95% CI 1.3–7.9), with mutations found only in smokers. They suggested a possible role of estrogen exposure in either the initiation or the selection of KRAS mutant clones in ADC (El Osta et al., 2019). In addition, a large study by Dogan et al. genotyped 3026 lung ADCs showing that KRAS G12C, the most common G > T transversion mutation in smokers, was more frequent in women ($p = 0.007$); these women were younger than men with the same mutation (median 65 years old *versus* 69 years old, $p = 0.0008$) and smoked less than men (Nelson et al., 1999). Regarding p53, LC from smokers shows a distinct TP53 mutation spectrum, such as G to T transversions at codons 157, 158, 179, 248, and 273, which are uncommonly observed in non-smokers (Dogan et al., 2012; Vähäkangas et al., 2001). Indeed, in smokers, 43% of the mutations were G to T transversions, but this number dropped to 13% in never-smokers (Hernandez-Boussard and Hainaut, 1998). Another analysis reported that the difference of p53 mutational spectrum between never-smokers and smokers was detectable only in women (Ding et al., 2008; Kure et al., 1996). p53 mutations in female never-smokers with ADC were predominantly transitions

(83%), while in smokers, they consisted predominantly in transversions (60%) and deletions (20%) (Toyooka et al., 2003).

Generally, the presence of most known driver mutations is more common in non-smokers and female patients. Often, the presence of one of these pathognomonic genetic alterations is recognized as the possible leading cause of cancer itself.

Female tumors more frequently harbor a wide set of targetable alterations, such as mutation of epidermal growth factor (EGFR) (Gealy et al., 1999), human epidermal growth factor receptor 2 (HER2) (Zhang et al., 2016) or serine/threonine-protein kinase B-RAF (Mazières et al., 2013), as well as rearrangement of proto-oncogene 1 receptor tyrosine kinase (ROS1) (Reddy et al., 2017; Rossi et al., 2017) and anaplastic lymphoma kinase (ALK) (Warth et al., 2014; Vidal et al., 2014), even though some discordant data are available on ALK prevalence in females. (Paik et al., 2012; Fallet et al., 2014).

On the other hand, pathogenic mutations of serine/threonine kinase (STK11), RNA-binding protein 10 (RBM10) and SMARCA4 have been reported more frequently in lung ADC samples from males (Shaw et al., 2009); however, no mutation among them is currently targeted by an available agent.

Two additional factors, programmed death-ligand 1 (PD-L1) and tumor mutation burden (TMB) are acquiring increasing relevance in NSCLC, because of their potential predictive value of response to immunotherapy. As known, the expression of PD-L1 on tumor cells promotes down-regulation and self-tolerance of the immune system from rejecting the tumor by suppressing T-cell inflammatory activity through binding to the regulatory T-cell receptor, PD-1 (Collisson et al., 2014).

According to a meta-analysis of nine studies in 1550 patients, gender resulted not correlated with PD-L1 expression (Sharpe et al., 2007). Despite some other experiences reporting no association of PD-L1 with gender (Pan et al., 2015; Lin et al., 2017; Chen et al., 2016), a more recent meta-analysis of fifty-two studies showed male gender as a factor associated with PD-L1 expression (odds ratio [OR] 4.8; 95% CI 3.2–7.2; $p < 0.001$) (Calles et al., 2015a).

Lung cancer has a very high rate of somatic mutations when compared to other tumors; 8.7 mutations per megabase in adenocarcinomas (ADCs) and 9.7 in squamous cell carcinomas (SCC) are reported (Petrelli et al., 2018). Even if TMB did not correlate with PD-L1 expression, they both emerged as key biomarkers of sensitivity to ICIs. Gender difference in TMB has already been reported in patients affected by cutaneous melanoma, and a more recent next-generation sequencing (NGS) analysis in NSCLC tumor samples confirmed a lower TMB in females than males. In particular TMB resulted higher in males and 10-fold higher in smokers than in never-smokers (Lawrence et al., 2014; Govindan et al., 2012). This correlates with the consistently lower TMB observed in NSCLC harboring most oncogenic drivers such as alterations of EGFR, ALK, ROS1, BRAF-V600E and MET exon 14 genes, with the exception of BRAF non-V600E and KRAS mutant tumors (Peters et al., 2017). Moreover, males and females own some differences in NSCLC immune-genes (Spigel et al., 2019) and micro-RNA (miRNA) expression (Araujo et al., 2016), but at the moment it is unknown whether these properties have a predictive/prognostic value.

6. LC treatment outcome by gender

6.1. Small cell-lung cancer (SCLC)

In SCLC, female gender is generally regarded as a positive prognostic factor (Guo et al., 2017; Osterlind et al., 1986; Videtic et al., 2005; Wolf et al., 1991; Albain et al., 1990; Paesmans et al., 2000), whereas it may be related to an increased toxicity from chemotherapy (Johnson et al., 1988; Singh et al., 2005). The role of a patient's sex on SCLC outcome has been fully explored for limited and advanced disease, across clinical trials and real-life evidence.

6.2. Limited-stage disease (LD)

Chemotherapy remains the main option in SCLC management, excluding T1,2 N0,1 M0 tumors where a surgical approach might be proposed (Wheatley-Price et al., 2010; Jett et al., 2013; Rudin et al., 2015). After curative-intent surgical resection, platinum-based adjuvant chemotherapy is recommended (Früh et al., 2013; Shepherd et al., 1988; Tsuchiya et al., 2005).

Two real-life experiences investigated the impact of sex in this setting. A retrospective study, including 1574 patients with pT1-2N0M0 SCLC undergoing surgery, showed that lobectomy followed by adjuvant chemotherapy was associated with improved survival at multivariate analysis, whereas higher age, tumor size, and presence of comorbidities were independently associated with worse survival; in this analysis, sex was not associated with survival (Tsuchiya et al., 2005).

In another retrospective study including stage I SCLCs patients ($n = 2681$) from the National Cancer Database, patient age, female gender, Charlson-Deyo comorbidity score < 2 , tumor size ≤ 3 cm, surgery and chemotherapy were all associated with improved OS (C-FJ et al., 2016).

Recommended treatment in non-surgical limited stage disease is concurrent chemotherapy and thoracic radiotherapy (Paximadis et al., 2017; Saito et al., 2006; Kubota et al., 2014; Skarlos et al., 2001; Pignon et al., 1992; Warde and Payne, 1992). Cox multivariate models were applied to 1363 patients in six limited disease trials conducted by Southwest Oncology Group (SWOG): good performance status (PS), female sex, age less than 70 years, white race, and normal lactate dehydrogenase (LDH) were significant favorable independent predictors (Wolf et al., 1991).

A meta-analysis of 16 RCTs including 2140 patients and evaluating thoracic radiotherapy showed no statistically different treatment effects according to sex (pooled relative risk for women 0.79, 95% CI 0.66–0.94, and for men 0.88, 95% CI 0.78–0.98) (Skarlos et al., 2001).

A RCT comparing twice-daily with once-daily thoracic radiotherapy in 417 patients with limited SCLC treated concurrently with cisplatin and etoposide, showed that male sex and a PS of 2 were significantly associated with shorter failure-free survival at multivariate analysis (no hazard ratio was reported) (Warde and Payne, 1992).

As far as evidence from real-life experiences is concerned, a retrospective study including a cohort of 6271 limited stage SCLCs from the Surveillance, Epidemiology, and End Results (SEER) database reported in multivariate analysis female sex as a positive prognostic factor (HR = 0.91; 95% CI 0.87–0.96; $p = 0.0007$), whereas older age, African American race, and main bronchus location were all associated with a statistically significant increase in mortality hazard (Turrisi et al., 1999).

A retrospective study investigating the role of chemo-radiotherapy on 8637 elderly patients (aged ≥ 70 years) with limited SCLC included female sex among factors associated with improved OS on univariate and on multivariate analysis (HR = 0.83; 95% CI 0.79–0.87; $p = 0.001$) (Lally et al., 2009).

Another retrospective study on 179 limited SCLC treated with definitive chemo-radiotherapy in a real-life scenario showed that female gender was significantly associated with longer brain metastases free survival ($p = 0.023$) and a lower incidence of metachronous brain failure. In this study, males resulted to have an inferior median OS than females (14 months [95% CI, 10–18] versus 20 months [95% CI, 10–22], respectively; $p = 0.021$), and this correlation was also found at multivariate analysis (HR 1.38; 95% CI 1.08–1.92; $p = 0.04$) (Corso et al., 2015).

6.3. Extensive-stage disease (ED)

Palliative systemic treatment of metastatic disease is based on cisplatin-etoposide regimen up to 4–6 cycles (Roengvoraphoj et al., 2018; Pujol et al., 2000); cisplatin can be replaced by carboplatin with no

apparent lack of efficacy and a better tolerance (Mascaux et al., 2000). The combinations irinotecan-cisplatin (Rossi et al., 2012), topotecan-cisplatin (Zatloukal et al., 2010; Fink et al., 2012) and gemcitabine-carboplatin (in poor prognostic patients only) (Eckardt et al., 2006) could also be valid alternatives. Patients having a PFS less than 6 months can benefit in terms of improved symptoms control and survival from use of oral or intravenous topotecan (Lee et al., 2009; O'Brien et al., 2006) or cyclophosphamide plus doxorubicin and vincristine (CAV) (Eckardt et al., 2007). Patients having a late disease relapse instead could benefit from re-challenge with first-line therapy. Independent determinants of improved outcome in patients with ED are traceable in a normal LDH, an intensive multidrug regimen treatment, and a single metastatic lesion (Wolf et al., 1991). Regarding patients' sex, a meta-analysis that included 4 RCTs comparing cisplatin-etoposide versus carboplatin-etoposide showed no evidence of treatment difference between the two arms according to patients' sex (sex interaction for OS $p = 0.42$, for PFS $p = 0.57$) (Mascaux et al., 2000).

In the randomized non-inferiority study comparing gemcitabine-carboplatin with etoposide-cisplatin, no difference was observed in treatment effect on OS or PFS survival in subgroups defined by age, sex, performance status and stage (Eckardt et al., 2006).

Subgroup analyses of the phase III trial comparing oral topotecan to BSC showed a survival benefit in favor of the chemotherapy arm in females, but not so clearly in males (Lee et al., 2009).

A sex-based retrospective analysis of 4 trials including 1006 patients treated with CAV regimen and etoposide-cisplatin conducted by the National Cancer Institute of Canada Clinical Trials Group showed that anemia (55.9% vs 38.6%, $p < 0.0001$) and leukopenia (92.7% vs 84.7% $p = 0.0002$) were significantly higher in female patients (Johnson et al., 1988). Females also experienced stomatitis (16.8% vs 8.3%, $p < 0.0001$) and vomiting (76.8% vs 67.2%, $p = 0.0014$) more often; when logistic regression analysis was performed to adjust the comparisons between the sexes by other factors (age, body surface area, PS, LDH, trial), female sex was still a significant predictor for all these toxicities. Moreover, more females than males had treatment delays of 2 weeks or more (52% vs 43.4%, $p = 0.022$), which was likely due to increased toxicity. Females, compared to males, had a superior overall response rate (RR) and median OS (80.3% vs 55.9% and 1.31 years vs 0.71 years respectively, $p < 0.0001$) (Johnson et al., 1988). Pharmacokinetic differences have been suggested as the cause of this imbalance in toxicity (Von et al., 1999; Kaul et al., 1996).

A pooled analysis of six chemotherapy trials ($n = 1707$) from the Manchester Lung Group and the Medical Research Council Clinical Trials Unit was carried out to validate the results from the previous analysis. This confirmed a significantly longer median survival for women (10.2 vs 9.6 months, $p = 0.006$), and in multivariate analysis female sex remained a significant predictor of improved survival (HR 0.88, 95% CI 0.79–0.99, $p = 0.04$). No differences between the sexes were found for hematological toxicity, but women experienced more frequently grade 3 or 4 emesis (18% vs 9%, $p < 0.0001$) and mucositis (13% vs 8%, $p = 0.005$) (Singh et al., 2005).

Regarding immunotherapy application, nivolumab with or without ipilimumab – an anti Cytotoxic T-Lymphocyte Antigen 4 (CTLA4) inhibitor – has shown antitumor activity in previously chemo-treated patients, leading to Food and Drug Administration (FDA) accelerated approval of nivolumab for third and further line treatment of ED SCLC (Dobbs et al., 1995). TMB was recognized as a predictive factor (Antonia et al., 2016), while it is unknown if males and females benefit equally from ICIs, as no subgroup analyses by gender was performed in Checkmate 032 trial. More recently, IMpower133 phase III randomized trial showed OS and PFS improvement with the addition of atezolizumab to chemotherapy in the first-line treatment of ED SCLC; OS benefit was consistent across patient sex subgroups (males 12.3 mo vs 10.9 mo, HR = 0.74, 95% CI, 0.54–1.00; females 12.5 mo vs 9.5 mo, HR = 0.65, 95% CI, 0.42–1.00) (Hellmann et al., 2018a).

6.4. Non-small cell lung cancer (NSCLC)

6.4.1. Early-locally advanced stage

The subgroup analyses performed during the development of the 7th edition UICC/AJCC/IASLC TNM staging system confirmed male sex as an independent poor prognostic factor in surgically managed patients (unadjusted HR = 1.17, $p < 0.001$; adjusted HR = 1.21, $p < 0.001$) (Horn et al., 2018). At each stage of disease, the relative survival in women was longer than in men, with the largest difference in earlier stages.

According to the national SEER database, which collected data of patients with primary bronchogenic carcinoma (diagnosed from 1975 to 1999), female patients who underwent surgery alone had 2-year and 5-year survival rates of 74.3% and 56.8%, both greater than those of males equal to 66.0% and 48.3%, respectively ($p < 0.0001$) (Chansky et al., 2009). Female gender exerts a significant positive effect on survival of NSCLC patients following lung resection by a prospective collected dataset ($n = 883$). This benefit was more pronounced in earlier stages (stage I, $p = 0.01$; stage II–III, $p = 0.3$) and persisted after adjusting for important differences in the clinical, histo-pathological features and extent of pulmonary resection between male and female patients (Minami et al., 2000). The reason for this advantage in females seems not to be related to a different efficacy of post-surgical treatment. Indeed, subgroup analysis of the main trials of adjuvant chemotherapy showed no different effect of CT on OS according to patient's sex (Alexiou et al., 2002; Group, 2010; Strauss et al., 2008; Artal Cortés et al., 2015; Pignon et al., 2008; Wakelee et al., 2016). Furthermore, a meta-analysis of 35 trials including individuals with histologically confirmed NSCLC who underwent a potentially curative resection and subsequent adjuvant chemotherapy, showed no significant effect of patient's sex on the absolute improvement in 5-year survival (Butts et al., 2010). Sex-based differences in the timing of postoperative recurrence are also reported. A retrospective study conducted on patients who underwent complete pulmonary resection with or without adjuvant chemotherapy found that the risk and the peak time of post-surgical recurrences differed considerably between men and women, with the latter having a longer disease free interval (DFI) at the same disease stage, histological type, and smoking status (Burdett et al., 2015).

The use of neoadjuvant CT appears beneficial in resectable locally advanced disease (Watanabe et al., 2018; Gilligan et al., 2007; Lim et al., 2009). Some trials suggest the effect might be greater in women and others in men, but it is not clear why. A meta-analysis of 15 RCTs ($n = 2385$) did not point out a different outcome by sex, despite a difference in the effect of treatment or treatment by covariate interactions across trials (HR = 1.08, 95% CI, 0.81–1.44; $p = 0.62$; heterogeneity $p = 0.04$) (Lim et al., 2009).

Radiotherapy benefit in localized/locally advanced NSCLC seems not to be so clearly sex-dependent. In patients undergoing radiotherapy within the above cited SEER database, the relative 2-year and 5-year survival rates in men and women were 15.5% vs 12.2% and 5.3% vs 3.4%, respectively ($p \leq 0.0001$) (Chansky et al., 2009). A different pooled prospective data on more than 1300 NSCLC inoperable patients treated within the Radiation Therapy Oncology Group (RTOG) non-operative trials showed that women treated with radiotherapy had longer overall survival (Burdett, 2014). For patients with stage IIIB or stage IIIA NSCLC not resectable, concurrent chemoradiotherapy (CRT) is considered the preferred treatment, when feasible. This option is also valid for patients who, after induction chemotherapy, are not eligible for surgery (Siddiqui et al., 2010). Subgroup analyses for sex reported on two meta-analyses comparing concurrent to sequential CRT did not report any difference of treatment outcome by sex (Stewart, 1995; Aupérin et al., 2006). For elderly or less fit patients, the sequential approach is preferred, regardless of patients' sex (Aupérin et al., 2010).

6.4.2. Advanced-metastatic stage

Platinum-based chemotherapy has been the only treatment available for patients affected by NSCLC for almost 30 years, and currently still plays a major role in advanced and metastatic setting. The realization of the first EGFR tyrosine kinase inhibitor (TKI) and the discovery of its great activity in patients with a sensitizing EGFR gene mutation, allowed the first big revolution in the treatment of this disease (Brunelli et al., 2009). The subsequent identification of other targetable NSCLC genetic alterations led to the development of efficacious drugs for patients harboring them, opening the era of personalized therapy. Nowadays different TKIs targeting a series of genes alterations (EGFR, ALK, ROS-1, BRAF) have already received approval for patients with no radically treatable NSCLC. Meanwhile, the introduction of anti-angiogenic agents led to survival improvements for non-SCC NSCLC when associated with chemotherapy (Bell et al., 2005), although without significant revolution in current clinical practice. Another big innovation for the treatment of NSCLC came with the advent of ICIs, especially those targeting the PD-1/PD-L1 axis. Initially approved only for chemotherapy-pretreated patients (Soria et al., 2013), more recently ICIs alone or combined with chemotherapy have also shown benefit in naïve patients (Ramos-Esquivel et al., 2017), and as consolidation treatment in patients pretreated with definitive chemo-radiotherapy for locally advanced disease (Martinez et al., 2019). The scientific community has tried over the years to detect potential predictive factors of response to these agents. The possible predictive/prognostic value of the variable gender is below that analyzed in NSCLC patients treated with the three main drugs categories described.

6.4.2.1. Chemotherapy with or without anti-angiogenic agents. In patients with advanced NSCLC, CT *plus* best BSC showed survival benefit *over* BSC alone irrespective of age, sex, histology and PS (Siddiqui et al., 2010; Antonia et al., 2017). Subsequent meta-analyses pointed out a greater benefit of two-agents *over* single-agent CT (Group, 2008) and for those patients treated with regimens containing a platinum compound (Delbaldo et al., 2004). Different platinum-doublets with third-generation cytotoxic agents showed comparable efficacy (Pujol et al., 2006), but a slight advantage in OS emerged for cisplatin-based *versus* carboplatin-based chemotherapy (Schiller et al., 2002). Later, a survival benefit was noticed when platinum was combined with the antifolate agent pemetrexed rather than with gemcitabine or docetaxel, in patients with non-SCC histology (Ardizzoni et al., 2007). In all these studies, no subgroup analyses for sex were performed.

Patients are usually treated with four cycles of platinum-based doublet, as adding more cycles does not further improve the outcome in either males or females (Li et al., 2012). In these patients however, RCTs showed a survival benefit for single agent pemetrexed *versus* placebo, both when used as switched maintenance (Rossi et al., 2014) or – limited to non SCC NSCLC – as continuous maintenance after an induction treatment with pemetrexed plus cisplatin (Ciuleanu et al., 2009; Paz-Ares et al., 2012); even if females sometimes resulted to have a much lower PFS HR than males, this did not translate to a different OS HR in subgroup analysis (Ciuleanu et al., 2009; Paz-Ares et al., 2012).

The addition of the anti-vascular growth factor receptor (VGFR) bevacizumab (Bev) to standard platinum-based CT doublets in the first-line setting led to a small but significantly improved OS, PFS and RR for patients with advanced non-SCC NSCLC (Bell et al., 2005; Paz-Ares et al., 2013). The subgroup analyses of the main approval trials of Bev led to conflicting findings about the predictive value of sex. In the Eastern Cooperative Oncology Group (ECOG) 4599 trial, the addition of Bev to paclitaxel-carboplatin was possibly not beneficial on survival among women, while of benefit for men (Lima et al., 2011); further analyses of this trial showed that females younger than 60 years treated with the experimental arm combination had an improved survival compared to females older than 60 years, with menopausal status possibly being a negative predictor of response to Bev (Sandler et al., 2006; Brahmer et al., 2011). In the AVAIL trial, evaluating the addition

of 7.5 or 15 mg/Kg of Bev to cisplatin-gemcitabine, Reck et al. reported subset differences by means of Forest plots of HRs: females had a statistically improved PFS at 15 mg/kg of Bev, but males had a significantly better PFS at the 7.5 mg/kg Bev dose (Wakelee et al., 2012). Retrospective analysis of ECOG 4599 trial showed a possible benefit of maintenance Bev (Reck et al., 2009), while a following RCT failed to demonstrate a superiority of Bev *plus* pemetrexed maintenance *over* Bev alone after induction treatment (Lopez-chavez et al., 2018; Barlesi et al., 2013); no subgroup analysis by gender was reported for either study.

In ECOG PS 2 patients, CT has shown to improve survival and quality of life (QoL) compared with BSC alone (Barlesi et al., 2014). Single-agent CT (gemcitabine, vinorelbine, taxanes) could be the preferred option, although carboplatin-based or low-dose cisplatin-based doublets may represent alternative options in eligible patients (Gridelli et al., 2004; Lilenbaum et al., 2009; Morère et al., 2010; Zukin et al., 2013). No specific gender subgroup analyses were performed in this setting.

For patients 70 years of age or older, therapeutic options include single agent chemotherapy (Bronte et al., 2015; Kudoh et al., 2006) - especially for unfit patients - or combination with carboplatin (Gridelli et al., 2003), while two meta-analyses showed a worse tolerability and contrasting results of OS prolongation with the use of platinum-based regimens (Quoix et al., 2011; Des et al., 2012). The addition of cisplatin to single-agent chemotherapy does not improve OS or QoL in elderly patients, according to a joint analysis of MILES-3 and MILES-4 RCTs (Qi et al., 2012). When reported, subgroup analyses did not show differences in outcome among male and female patients (Kudoh et al., 2006; Gridelli et al., 2003; Qi et al., 2012).

Although prospective data are quite lacking, several retrospective evaluations of gender differences in survival in advanced-stage NSCLC chemo-treated patients have been performed. In an earlier ECOG data analysis of seven phase III RCTs ($n = 893$), the 1-year survival rate was significantly higher in women *versus* men (26% and 16% respectively), even at multivariate analysis ($p = 0.005$) (Gridelli et al., 2018). Female sex resulted a strong independent factor for improved survival ($p < 0.0005$) according to another huge joint analysis from 13 SWOG trials ($n = 2531$) between 1974 and 1988, but this significance was lost when other prognostic variables were considered (Finkelstein et al., 1986). The European Lung Cancer Working Party (ELCWP) evaluated patients receiving platinum-based chemotherapy for either locally advanced or advanced NSCLC from 1980 to 1991 ($n = 1052$) and found female sex to be one of several variables associated with improved survival with a relative risk of death of 0.7 ($p = 0.03$) in multivariate analysis (Albain et al., 1991). A single cancer center study including advanced-staged NSCLC patients chemo-treated from 1978 to 1986 ($n = 378$) further confirmed female sex as an independent predictor of improved survival (median OS of 12.4 months for women *versus* 8.8 months for men, $p = 0.001$) (Paesmans et al., 1995). The ECOG E1594 trial randomized patients with advanced NSCLC ($n = 1157$) to one of four platinum doublets and found that all four regimens had comparable efficacy. The effect of sex on survival was later examined in this population and, despite a similar response rate (19% in both sexes, $p = 0.15$), women had a 1.9-month statistically significant improvement in median OS compared with men and also greater toxicity (O'Connell et al., 1986). The reasons for this apparent major benefit for female sex after a platinum-based chemoregimen among LC patients are currently unknown. Platinum-based agents act through the formation of DNA adducts, which leads to cell cycle arrest; the repair of these adducts is one mechanism of resistance to platinum-based agents. A defective DNA repair and/or recombination function was found to correlate with prolonged survival after cisplatin-based chemotherapy, even in NSCLC patients (Wakelee et al., 2006). Thus, the above seen difference in DRC between sexes (Wei et al., 2000; Spitz et al., 2001) could partially explain why women could benefit more than men from chemotherapy.

Second line CT should be offered to patients progressing after first line CT and with specific contraindications to ICIs. In this setting, combination CT failed to improve OS compared with single-agent CT (Lord et al., 2002), with no evidence of heterogeneity among subgroups of treatment effect around the overall effect. Among single chemotherapeutic agents, docetaxel demonstrated a significant better survival compared to best supportive care (Gridelli et al., 2009). Moreover, treatment with pemetrexed resulted in clinically equivalent efficacy outcomes, but with significantly fewer side effects compared with docetaxel (Shepherd et al., 2000). No subgroup by sex were reported in those studies.

Despite a doubtful clinical relevance, the association of docetaxel with the anti-VEGFR inhibitor Ramucirumab gave a statistically significant outcome benefit over docetaxel alone, similarly between sexes (Hanna et al., 2004). The multi-targeted tyrosine kinase inhibitor Nintedanib in combination with docetaxel is an effective second-line option, especially for patients with adenocarcinoma. No differential efficacy of nintedanib was shown by subgroup analyses by sex (Garon et al., 2014). Prognosis of patients eligible for second-line treatment of advanced NSCLC is significantly conditioned by gender (worse in males), PS, histology, stage, previous use of platinum and response to first-line. A derived prognostic score that also includes a patient's sex could be useful to discriminate subjects with a relatively more favorable prognosis and those with a very short life expectancy (Reck et al., 2014).

6.4.2.2. Targeted therapies

6.4.2.2.1. EGFR inhibitors (EGFRi).

The transmembrane protein with cytoplasmic kinase activity EGFR transduces, upon ligand binding, growth factor signaling to the cancer cell. In around 15%–30% of NSCLC, according to patient ethnicity, EGFR results aberrantly activated due to an acquired mutation (EGFRm). In case of EGFR sensitizing mutations, the best front line therapy consists on an EGFR TKI. EGFR mutation rate is found higher in never-smokers compared with ever-smokers, as well as in women compared to men [185]. Results from pivotal phase III studies, including EGFRm NSCLC patients treated with an EGFRi, showed a reduction in the risk of progression in the female subgroup but not in the male counterpart (Table 1). In the same way, a recent exploratory study showed that women affected by EGFRm NSCLC had a greater benefit from EGFRi than men (HR = 0.34; 95% CI, 0.28–0.40; $p < 0.00001$ vs HR = 0.44; 95% CI, 0.34–0.56; $p < 0.00001$, respectively) (Remon et al., 2017). A possible explanation is that women could benefit more than men from TKI therapy due to their lower tobacco smoking exposure. In fact, a meta-analysis including advanced EGFRm NSCLC patients showed that non-smokers have longer PFS to EGFRi treatment compared to patients who have a smoking history (Pinto et al., 2018). Furthermore, sexual hormones were found to regulate the EGFR signaling pathway, thereby potentially determining different effects on tumor growth and response to TKIs in EGFRm NSCLC patients of different genders. Specifically, growing evidence shows that ERs are expressed in LC cells and are able to interact with the EGFR cascade. In NSCLC cell lines and xenografts, a functional interaction between ERs and the EGFR activity was documented, being the EGFR protein expression down-regulated in response to estrogen and up-regulated in response to estrogen receptor inhibitors (Zhang et al., 2015; Stabile et al., 2005). This suggests that the EGFR pathway is activated when estrogen is depleted. Similarly, ER- β protein expression may be down-regulated after the addition of EGF and up-regulated in response to EGFRi (Zhang et al., 2015). A recent retrospective study investigated the associations between ER α /ER β expression and EGFR mutational status and response to TKI treatment in metastatic lung ADC, and concluded that cytoplasmic ER β expression is a negative predictor of clinical response to EGFR-TKIs in patients with EGFR mutations (Márquez-Garbán et al., 2007a). This awareness suggests that the ER and EGFR pathways might contribute together to the progression of LC, providing the rationale to target both

the ER and EGFR pathways (Ding et al., 2018). Anyway, despite the promising preliminary activity data of antiestrogen plus EGFR-TKI combination (Márquez-Garbán et al., 2007b), two phase II trials failed to show any further survival benefit from adding fulvestrant to EGFR-inhibitors as single agent in EGFR mutant patients (Traynor et al., 2009; Garon et al., 2010). On the other hand, the oncogenic role of androgens in LC development and/or progression was also found in the subset of LC harboring an EGFR mutation. In particular, the androgen receptor resulted over expressed among EGFRm NSCLC and in those acquiring the most common TKI resistance mutation (T790M) (Mazieres et al., 2018).

Thus, despite their different spectrum of interaction with EGFR, both male and female hormonal mediators have a common pro-tumor effect even in EGFRm NSCLC. Further studies are needed to explain the mechanisms involved in a possible different efficacy of TKI therapy between genders, and also if sexual hormones might play a crucial role in this sense or not.

6.4.2.2.2. ALK inhibitors (ALKi).

ALK rearrangements are found in about 5% of patients with NSCLC and are related to younger age, no history of smoking and distinct clinical-pathological features such as adenocarcinoma histology (Fallet et al., 2014). Crizotinib was the first agent approved for front-line therapy of ALK-rearranged NSCLC after its demonstrated superiority over chemotherapy. Next-generation ALKi have been developed to overcome crizotinib resistance. Currently, three second-generation ALKi, ceritinib, alectinib, and brigatinib, and the third-generation inhibitor lorlatinib are FDA approved for use in ALK-rearranged NSCLC following progression or intolerance to crizotinib. Later on, different second generation ALKi demonstrated a survival advantage when used upfront, thus leading FDA to broaden ceritinib and alectinib indication to untreated patients with ALK-rearrangement; more recently brigatinib showed a significantly longer PFS compared to crizotinib in ALKi naïve patients (Blakely et al., 2017).

Analyzing the registration studies of the ALKi, no difference between genders in terms of PFS has been observed (Table 1). This scenario is confirmed by the above mentioned meta-analysis by Pinto et al. (Remon et al., 2017), where ALKi gave a higher benefit over the standard of care independently of patients gender (male, HR = 0.48, 95% CI, 0.39–0.59; $p < 0.00001$; female, HR = 0.51; 95% CI, 0.45–0.61; $p < 0.00001$).

6.4.2.2.3. Immune check point inhibitors (ICIs).

Recently, the advent of ICIs led to substantial and unprecedented changes in the therapeutic algorithms of NSCLC. Thanks to a better outcome and safety profile over previous standard of care chemotherapy regimens, different ICIs - especially those targeting the PD-1/PD-L1 axis - received rapid approval for the treatment of locally advanced/metastatic NSCLC. Subsequent developments led to validation of ICI-ICI and ICI-chemotherapy combinations in metastatic NSCLC (Camidge et al., 2018; Socinski et al., 2018), and to an optimization of chemotherapy in locally advanced NSCLC through anti-PD1 as consolidation approach (Martinez et al., 2019). Moreover, some neoadjuvant and adjuvant clinical ICI trials are currently ongoing in early-stage disease. Beyond a general benefit in median PFS and/or OS, some metastatic NSCLC patients treated with ICIs find, a major benefit, in terms of a more prolonged OS. Predictive criteria of ICI efficacy are not yet fully identified and the efforts of the scientific community are directed to their identification, in order to improve patient selection. Despite their increasing popularity integrated immunograms generally do not take into account a sex. Usually, a patient's sex is considered as a stratification criterion in randomized clinical trials, and considerations about ICI efficacy in males and females mainly derive from post-hoc subgroup analysis (Table 1). A phase III study comparing nivolumab with docetaxel in patients with advanced non-SCCNSCLC showed that nivolumab did not improve overall survival in women (HR = 0.78, 95% CI, 0.58–1.04) (Gandhi et al., 2018). Similarly, lack of survival benefit for females was also found in the same setting for SCC

Table 1
 Outcome by patients' sex according to NSCLC phase III randomized clinical trials. The table reports pivotal trials highlighting the difference between males and females in terms of number-rate of enrolled patients as well as of relative survival benefit. Legend: BEV: bevacizumab; CBDCA: carboplatin; CDDP: cisplatin; (nab)TAX: nab-paclitaxel/paclitaxel; NA: not assessed; ND: not determinate; PEM: pemtreated; Plt: platinum; SCC: squamous cell carcinoma; TAX: paclitaxel; TXT: docetaxel.

| Drug class | Trial | Histology | Stage | Inclusion criteria | Treatment | Comparator | Line | Pts, n | M to F ratio | PFS, male HR (95% CI) | PFS, female HR (95% CI) | PFS, p interact | OS, male HR (95% CI) | OS, female HR (95% CI) | OS, p interact |
|---------------|---------------------|------------|--------------------|-------------------------------------|----------------------------------|------------------|-------|--------|------------------|-----------------------|-------------------------|------------------|----------------------|------------------------|----------------|
| ICI | KEYNOTE 189 | nSCC NSCLC | IIIB, IV | No PD-L1 threshold | PEM-PltCht + Pembrolizumab | PEM-PltCht | 1 | 616 | 1,4 | 0.70 (0.50-0.99) | 0.29 (0.19-0.44) | ND | 0.66 (0.50-0.87) | 0.40 (0.29-0.54) | ND |
| | KEYNOTE 407 | SCC NSCLC | IIIB, IV | No PD-L1 threshold | Pembrolizumab + CBDCA + (nab)TAX | CBDCA + (nab)TAX | 1 | 559 | 4,9 | 0.58 (0.46-0.73) | 0.49 (0.30-0.81) | ND | 0.69 (0.51-0.94) | 0.54 (0.42-0.72) | ND |
| | KEYNOTE-024 | NSCLC | IIIB, IV | PD-L1 TPS ≥ 50% | Pembrolizumab | Plt-based ChT | 1 | 305 | 1,6 | 0.39 (0.26-0.58) | 0.75 (0.46-1.21) | ND | 0.51 (0.36-0.71) | 0.42 (0.29-0.59) | ND |
| | KEYNOTE-042 | NSCLC | IIIB, IV | PD-L1 TPS ≥ 1% | Pembrolizumab | Plt-based ChT | 1 | 1274 | 2,42 | 0.63 (0.46-0.85) | 0.71 (0.40-1.26) | ND | 0.57 (0.44-0.71) | 0.67 (0.52-0.81) | ND |
| CheckMate 057 | nSCC NSCLC | IIIB, IV | No PD-L1 threshold | Nivolumab | TXT | ≥ 2 | 582 | 1,2 | 0.78 (0.56-1.04) | 1.02 (0.78-1.32) | ND | 0.65 (0.52-0.81) | 0.69 (0.52-0.85) | ND | |
| EGFRi | CheckMate 017 | SCC NSCLC | IIIB, IV | No PD-L1 threshold | Nivolumab | TXT | 2 | 272 | 3,3 | 0.56 (0.44-0.71) | 0.54(0.37-0.79) | ND | 0.54 (0.44-0.64) | 0.49 (0.37-0.64) | NA |
| | KEYNOTE-010 | NSCLC | IIIB, IV | PD-L1 TPS ≥ 1% | Pembrolizumab 2 mg/kg | TXT | 2 | 687 | 1,6 | 0.78 (0.64-0.94) | 1.02 (0.78-1.32) | ND | 0.65 (0.52-0.81) | 0.69 (0.52-0.85) | ND |
| | KEYNOTE-010 | NSCLC | IIIB, IV | PD-L1 TPS ≥ 1% | Pembrolizumab 10 mg/kg | TXT | 2 | 689 | 1,6 | 0.78 (0.64-0.94) | 1.02 (0.78-1.32) | ND | 0.65 (0.52-0.81) | 0.69 (0.52-0.85) | ND |
| | OAK | NSCLC | IIIB, IV | No PD-L1 threshold | Atezolizumab | TXT | ≥ 2 | 850 | 1,6 | 0.56 (0.44-0.71) | 0.54(0.37-0.79) | ND | 0.54 (0.44-0.64) | 0.49 (0.37-0.64) | NA |
| EGFRi | PACIFIC | NSCLC | IIIA, IIIB | No PD-L1 threshold | Durvalumab | placebo | cons. | 713 | 2,3 | 0.56 (0.44-0.71) | 0.54(0.37-0.79) | ND | 0.54 (0.44-0.64) | 0.49 (0.37-0.64) | NA |
| | ISEL | NSCLC | IIIB-IV | No selection for EGFR | Gefitinib | placebo | ≥ 1 | 1692 | 2,1 | 0.78 (0.64-0.94) | 1.02 (0.78-1.32) | ND | 0.65 (0.52-0.81) | 0.69 (0.52-0.85) | ND |
| | INTEREST | NSCLC | IIIB-IV | No selection for EGFR | Gefitinib | TXT | ≥ 2 | 1466 | 1,9 | 0.56 (0.44-0.71) | 0.54(0.37-0.79) | ND | 0.54 (0.44-0.64) | 0.49 (0.37-0.64) | 0,27 |
| | IPASS | NSCLC | IIIB-IV | No/former smokers | Gefitinib | CBDCA-TAX | 1 | 1217 | 0,3 | 0.78 (0.64-0.94) | 1.02 (0.78-1.32) | ND | 0.65 (0.52-0.81) | 0.69 (0.52-0.85) | NA |
| | NEJ002 | NSCLC | IIIB-IV | EGFR del19, L858R, others, no T790M | Gefitinib | CBDCA-TAX | 1 | 228 | 0,6 | 0.56 (0.44-0.71) | 0.54(0.37-0.79) | ND | 0.54 (0.44-0.64) | 0.49 (0.37-0.64) | 0,883 |
| | WJTOG3405 | NSCLC | IIIB-IV | EGFR del19, L858R | Gefitinib | CDDP-TXT | 1 | 172 | 0,4 | 0.67 (0.50-0.84) | 0.41 (0.26-0.65) | ND | 0.67 (0.50-0.84) | 0.41 (0.26-0.65) | NA |
| | OPTIMAL, CTONG-0802 | NSCLC | IIIB-IV | EGFR del19, L858R | Erlotinib | CBDCA-GEM | 1 | 154 | 0,7 | 0.26 (0.14-0.50) | 0.13 (0.07-0.24) | ND | 1.31 (0.97-1.76) | 1.2 (0.85-1.71) | NA |
| | EURTAC | NSCLC | IIIB-IV | EGFR del19, L858R | Erlotinib | Plt-based ChT | 1 | 173 | 0,4 | 0.38 (0.17-0.84) | 0.35 (0.22-0.55) | 0.4721 | 0.35 (0.22-0.55) | 0.35 (0.22-0.55) | ND |
| | LUX-Lung 3 | ADC NSCLC | IIIB-IV | EGFR del19, L858R, others | Afatinib | CDDP-PEM | 1 | 345 | 0,5 | 0.61 (0.37-1.01) | 0.54 (0.38-0.76) | 0.85 | 0.61 (0.37-1.01) | 0.54 (0.38-0.76) | ND |
| | LUX-Lung 6 | ADC NSCLC | IIIB-IV | EGFR del19, L858R, others | Afatinib | CDDP-GEM | 1 | 364 | 0,5 | 0.36 (0.21-0.62) | 0.24 (0.16-0.35) | ND | 0.36 (0.21-0.62) | 0.24 (0.16-0.35) | ND |
| AURA3 | NSCLC | IIIB-IV | EGFR T790M | Osimertinib | Plt-PEM | 2 | 419 | 0,6 | 0.43 (0.28-0.65) | 0.34 (0.25-0.47) | ND | 0.43 (0.28-0.65) | 0.34 (0.25-0.47) | NA | |
| FLAURA | NSCLC | IIIB-IV | EGFR del19, L858R | Osimertinib | Gefitinib or Erlotinib | 1 | 556 | 0,6 | 0.58 (0.41-0.82) | 0.4 (0.30-0.52) | ND | 0.58 (0.41-0.82) | 0.4 (0.30-0.52) | NA | |

(continued on next page)

Table 1 (continued)

| Drug class | Trial | Histology | Stage | Inclusion criteria | Treatment | Comparator | Line | Pts, n | M to F ratio | PFS, male HR (95% CI) | PFS, female HR (95% CI) | PFS, p interact | OS, male HR (95% CI) | OS, female HR (95% CI) | OS, p interact | |
|------------|--------------|--------------------|---------|---------------------|------------------------|------------|----------|--------|--------------|-----------------------|-------------------------|-----------------|----------------------|------------------------|----------------|----|
| ALKi | PROFILE 1007 | NSCLC | IIIB-IV | ALK positive | Crizotinib | PEM or TXT | 2 | 347 | 0,8 | 0.52 (0.35-0.77) | 0.48 (0.34-0.68) | ND | ND | ND | ND | |
| | PROFILE 1014 | NSCLC | IIIB-IV | ALK positive | Crizotinib | PEM-PiCht | 1 | 343 | 0,6 | 0.54 (0.36-0.82) | 0.45 (0.32-0.63) | ND | ND | ND | ND | |
| | ASCEND-5 | NSCLC | IIIB-IV | ALK positive | Ceritinib | PEM or TXT | 2 or 3 | 231 | 0,8 | 0.43 (0.26-0.71) | 0.51 (0.33-0.78) | ND | ND | ND | ND | |
| | ASCEND-4 | NSCLC | IIIB-IV | ALK positive | Ceritinib | PEM-PiCht | 1 | 376 | 0,7 | 0.41 (0.27-0.63) | 0.63 (0.43-0.93) | ND | ND | ND | ND | |
| | J-ALEX | NSCLC | IIIB-IV | ALK positive | Alectinib | Crizotinib | 1 or 2 | 207 | 0,7 | 0.35 (0.16-0.77) | 0.31 (0.17-0.57) | ND | ND | ND | ND | |
| | ALEX | NSCLC | IIIB-IV | ALK positive | Alectinib | Crizotinib | 1 | 303 | 0,8 | 0.61 (0.38-0.98) | 0.39 (0.25-0.60) | ND | ND | ND | ND | |
| | ALUR | NSCLC | IIIB-IV | ALK positive | Alectinib | PEM or TXT | 3 | 107 | 1,2 | 0.25 (0.10-0.60) | 0.08 (0.02-0.30) | ND | ND | ND | ND | |
| | ALTA-1 L | NSCLC | IIIB-IV | ALK-positive | Brigatinib | Crizotinib | 1 | 275 | 0,83 | 0.49 (0.28-0.85) | 0.44 (0.24-0.84) | ND | ND | ND | ND | |
| | ChT | Schiller JH et al. | NSCLC | IIIB-IV | See Schiller JH et al. | CDDP-GEM | CDDP-TAX | 1 | 576 | 1,7 | ND | ND | ND | ND | ND | ND |
| | | NEJM 2002 | NSCLC | IIIB-IV | NEJM 2002 | CDDP-TAX | CDDP-TAX | 1 | 577 | 1,7 | ND | ND | ND | ND | ND | ND |
| JMDB | | NSCLC | IIIB-IV | See NCT000887711 | CDDP-PEM | CDDP-TAX | 1 | 578 | 1,7 | ND | ND | ND | ND | ND | ND | |
| PARAMOUNT | | nSCC | NSCLC | Induction CDDP-PEM | PEM maintenance | placebo | 1 | 539 | 1,4 | 0.74(0.55-1.00) | 0.49 (0.34-0.72) | ND | 0.82 | 0.73 | ND | |
| ECOG 4599 | | nSCC | NSCLC | No brain metastases | CBDCA-TAX-BEV | CBDCA-TAX | 1 | 850 | 1,2 | ND | ND | ND | 0.7 | 0.98 | ND | |
| AVAIL | | nSCC | NSCLC | No brain metastases | CDDP-GEM-BEV 7.5 mg/kg | CDDP-GEM | 1 | 692 | 1,8 | 0.76 | 0.8 | ND | 0.94 | 0.91 | ND | |
| AVAIL | | nSCC | NSCLC | No brain metastases | CDDP-GEM-BEV 15 mg/kg | CDDP-GEM | 1 | 698 | 1,7 | 0.98 | 0.57 | ND | 1.12 | 0.84 | ND | |

(HR = 0.67; 95% CI, 0.36–1.25) (Borghaei et al., 2015). Likewise, the results of KEYNOTE-010 trial comparing pembrolizumab *versus* docetaxel in patients previously treated with platinum-doublet chemotherapy showed that pembrolizumab did not increase the progression free survival (PFS) in women (HR = 1.02, 95% CI:0.78–1.32); however, these women benefited in OS from the experimental treatment, just like the men (Brahmer et al., 2015). Pembrolizumab did not confer a PFS benefit in women (HR = 0.75; 95% CI,0.46–1.21) even when administered in naïve patients having a PD-L1 expression $\geq 50\%$ (Herbst et al., 2016). Sex-subgroup analysis for the updated OS results of this latest trial has not been reported (Reck et al., 2016), but another first line study testing pembrolizumab in patients with PD-L1 expression $\geq 1\%$ showed that women did not benefit in OS (HR = 0.89; 95% CI,0.68–1.17) (Brahmer et al., 2017a). Conversely, all these RCTs showed that ICIs improved the outcome in the male subgroup. Differently from the anti-PD-1 agents, the activity of ICIs targeting its ligand seems not to be influenced by a patient's sex (Martinez et al., 2019; Lopes et al., 2018).

Recently some systematic reviews and meta-analyses investigated the correlation between patient sex and survival benefit from ICI. According to a meta-analysis of five RCTs ($N = 3025$) comparing a PD-1/PD-L1 ICI with docetaxel as second line treatment, the immunotherapy benefit resulted similar in men [HR = 0.69] compared to women [HR = 0.70]; interaction, $p = 0.82$) (Rittmeyer et al., 2017). Willis et al. showed no statistically significant association of patient sex with the efficacy of ICIs in terms of OS in the treatment of advanced cancers, and even for NSCLC by subgroup analyses of eleven RCTs involving more than 6000 patients (male HR = 0.79; 95% CI,0.71-0.88; female HR = 0.72; 95% CI,0.56-0.93; $p = 0.79$) (Lee et al., 2018).

However, another meta-analysis showed a greater benefit from ICIs in men than in women across different advanced solid cancer types. This trend was also found for the six RCTs that enrolled NSCLC patients ($N = 3482$), with the pooled hazard ratio of these studies revealing a sex-dependent magnitude of benefit from ICI use (male pooled HR = 0.72; 95% CI,0.61–0.86; female pooled HR = 0.89; 95% CI,0.71–1.11; p heterogeneity = 0.72) (Wallis et al., 2019).

Data about ORR to PD-1/PD-L1-inhibitors as single agents in NSCLC patients included in the main clinical trials are generally unreported distinctly for patients' sex. As known, ICIs confer a variably defined long-term benefit in about a quarter of treated NSCLC patients. Based on data from the ICI trial with the current longest follow-up time, no difference was shown in 5-year survival rate between males and females (Conforti et al., 2018).

More recent, first line setting RCTs showed a benefit from the use of ICIs, when added to standard first line chemotherapy, even in patients with PD-L1 less than 1%. The addition of pembrolizumab to standard chemotherapy doublet of platinum-based drug *plus* pemetrexed resulted in significantly longer OS and PFS than chemotherapy alone in non-squamous NSCLC. Subgroup analysis for OS showed a relative greater benefit, although not significant, for female patients treated with immunotherapy (female HR = 0.44, CI 0.28–0.71; male HR = 0.70, CI 0.50–0.99) (Socinski et al., 2018). The addition of atezolizumab to Bevacicicarboplatin and paclitaxel (BCP) significantly improved PFS compared to BCP alone among patients with metastatic non-squamous NSCLC regardless of PD-L1 expression and gender (female HR = 0.73, 8.2 mo vs 6.8 mo; male HR = 0.55, 8.4 mo vs 6.8 mo), showing also a OS benefit at the interim analysis (Camidge et al., 2018); similar results came from another first line atezolizumab-chemo combo regimen (Brahmer et al., 2017b). Even SCC-NSCLC patients seem to benefit from the use of upfront anti PD-L1 (Papadimitrakopoulou et al., 2019) or anti PD-1 *plus* chemo combinations; the addition of pembrolizumab to carboplatin *plus* paclitaxel or nab-paclitaxel resulted in a significantly longer OS and PFS than chemotherapy alone (Jotte et al., 2019), with no apparent discrepancy between sexes (Table 1). Combining different ICIs together is another emerging strategy: first-line nivolumab *plus* ipilimumab showed a significantly longer PFS compared with

chemotherapy in naïve patients with NSCLC and a high TMB (≥ 10 mutations per megabase), with a greater survival benefit in males than females at the subgroup analysis (male HR = 0.52, 95% CI, 0.36-0.74; female HR = 0.70, 95% CI, 0.41–1.20) (Paz-Ares et al., 2018).

Thus, whether gender could be a factor impacting on ICI benefit remains unclear. Despite this unanswered question, speculations about the role of a patient's sex on immunotherapy efficacy are possible. With various degrees of evidence, an absence of common driver mutations, a positive PD-L1 expression and a high TMB smoking status is known to be associated with smoking status (Hellmann et al., 2018b; Keselman et al., 2017; Davis et al., 2017). Thus, the smoking-related genomic signature, more frequent in males, might be correlated with their major benefit of ICI use.

Furthermore, the physiological differences between male and female immune system responses may also play a role. Across all ages, males have a higher lymphocyteCD8⁺/CD4⁺ ratio than females. The functional expansion and activation of CD8⁺ cytotoxicity by PD-1/PD-L1 ICIs could therefore be more powered in males, thus resulting in their better outcome secondary to immunotherapy use. An additional effect of ICIs seem to be a M1 macrophage polarization (Calles et al., 2015b), known to be correlated to an extended OS in lung cancer patients (Mediavilla-Varela et al., 2019). The classical activated M1-macrophages comprise immune effector cells with an acute inflammatory phenotype, while the alternatively activated M2-macrophages promote cell proliferation and tissue repair. Females, partly due to estrogen signaling (Hellmann et al., 2018b), have a greater M2-macrophage polarization than males, and this could contribute to the lesser benefit from ICIs therapy.

Multiple factors could be responsible for the discrepancies in immune responses observed between males and females (Fig.1), with sexual hormones apparently playing a major role.

For example, the 17 β -estradiol(E2) pathway promotes a protumor microenvironment (TME) through a redundancy of mechanisms (Jackute et al., 2018), among which a PD-L1 increased expression (Rothenberger et al., 2018). Furthermore, under sex hormone stimulation, T cells of males were found to produce higher IFN γ + Th1-type cytokines than those of women (Yang et al., 2017).




Fig. 1. Putative mediators differently expressed in M and F and influencing the immune-system. The differences in the immune system features existing between M and F could be a consequence of various kind of mediators differently expressed in the two sexes, mediators that resulted functionally connected to each other. Legend: BMI: body mass index; miRNA: micro RNA; lncRNAs: long-non-coding RNA; SHS: second hand smoke.


| Sex/gender-based differences in lung cancer |  |  |
|---|--|---|
| Epidemiology  | <ul style="list-style-type: none"> Incidence: 70/100.000 in U.S. Decreasing incidence since mid-80's | <ul style="list-style-type: none"> Incidence: 52/100.000 in U.S. Decreasing incidence since early 2000 Lower decreasing trend than M |
| Etiological concauses  | <ul style="list-style-type: none"> Higher smoking status Androgen carcinogenic effect | <ul style="list-style-type: none"> Higher level of CYP1A1 Estrogen carcinogenic effect Domestic PAH exposure |
| Cancer histology  | <ul style="list-style-type: none"> Higher incidence of SCC-NSCLC | <ul style="list-style-type: none"> Higher incidence of ADK-NSCLC Higher incidence of bronchioalveolar carcinoma |
| Cancer molecular features  | <ul style="list-style-type: none"> Higher incidence of STK11, RBM10, SMARCA4 mutation Higher TMB, PD-L1 expression | <ul style="list-style-type: none"> Higher incidence of more common oncogene driver alterations (KRAS G12C, EGFR, ALK, ROS1, HER2, BRAF) |
| Immune system features  | <ul style="list-style-type: none"> Higher CD8+/CD4+ ratio* M1 oriented macrophages* Th1 oriented CD4+ helper T cells* | <ul style="list-style-type: none"> Higher PD-L1 neg and TILs neg tumors Tumor immune gene sets enrichment (APOBEC3G-F, LAT, CD1D, CCL5) |
| Therapies efficacy  | <ul style="list-style-type: none"> Apparent greater benefit than F from ICIs | <ul style="list-style-type: none"> Apparent greater benefit than M from loco-regional, chemo and targeted therapies |

Fig. 2. Sex/gender-based differences in lung cancer patients. The known differences between males and females in etiological concauses, tumor histology and molecular features, as well as in immune-system characteristics might theoretically be responsible for a sex-based difference in therapy efficacy, especially for NSCLC histology. The asterisk (*) indicates those differences between the sexes evidenced in the general population, but not specifically in lung cancer patients. ADK: adenocarcinoma; APOBEC3 G: apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3 G; CCL5: Chemokine (C-C motif) ligand 5; CD1D: cluster of differentiation 1 D; CYP1A1: cytochrome P450, family 1, subfamily A, polypeptide 1. F: females; ICIs: immune checkpoint inhibitors; LAT: linker for activation T cell; M: males; PAH: polycyclic aromatic hydrocarbons; RCTs: randomized clinical trials; SCC: squamous cell carcinoma; TILs: tumour infiltrating lymphocytes.

The main limitation of this finding is that our knowledge about the male and female immune compartment mostly derives from healthy patients or from patients affected by non-oncological diseases. Noteworthy, the same M2-oriented macrophage differentiation of females has not been found in cancer patients, with tumor associated macrophages (TAM) features not differently represented by sex according to small NSCLC case series (Girón-González et al., 2000).

Finally, on the basis of results from clinical trials investigating ICI-chemo combinations, where similar survival benefit between genders is observed, it is tempting to speculate that the relative greater benefit of chemotherapy described in females (Albain et al., 1991) might balance the lower sensitivity to immunotherapy action, and vice versa in males.

7. Conclusion and perspectives

Available literature data suggest some different features in male and female lung cancer patients, from etiology to anatomo-pathology and treatment outcome (Fig. 2); even some differences between genders at gene-expression and miRNA levels are emerging.

Historically, women with NSCLC exhibit greater survival rates than men regardless of disease stage, histology, treatment modality, or smoking status, even after adjusting for gender-specific life expectancy (Girón-González et al., 2000; Fu et al., 2005; Thomas et al., 2005). With various levels of evidence, a survival advantage in favor of females was found in both NSCLC patients treated with radical aim (surgery, radiotherapy, chemo-radiotherapy) and those treated with a palliative platinum-based chemotherapy regimen or an EGFR TKI (O'Connell et al., 1986). Data on a patient's sex as a predictor of benefit of targeted therapies other than EGFR inhibitors are quite lacking. Also, in SCLC histology, female sex seems to be associated with improved survival.

These suggestions derived mainly from results of a retrospective study including a heterogeneous patient population (Nakamura et al., 2011) and a subgroup analysis of RCTs, in which a patient's sex was not considered as a stratification factor. Females are historically under-represented in RCTs (Viñolas et al., 2007), despite some recent positive data about women enrollment (Murthy et al., 2004; Labots et al., 2018), and trials among uniform populations addressing the role of sex on survival in NSCLC are lacking.

Moreover, most of the available data on treatment outcome refers to

patients submitted to cytotoxic, anti-angiogenic or targeted therapies. Whether female sex remains a positive prognostic factor with the incoming of immunotherapy has to be clarified. First analyses investigating the role of gender in ICIs treated NSCLC patients turned the tide, with males apparently experiencing greater benefit compared to females. Some clinical-pathological features might explain such outcome discrepancies, and these data strongly support the significance of gender as a separate prognostic factor in advanced NSCLC and emphasize its importance as a stratification factor in future phase III NSCLC trial design.

Acknowledgment

We acknowledge Dr. Christina Drace and Dr. Laura Mc. Mahon for linguistic revision.

References

- Albain, K.S., Crowley, J.J., LeBlanc, M., Livingston, R.B., 1990. Determinants of improved outcome in small-cell lung cancer: an analysis of the 2,580-patient Southwest Oncology Group data base. *J. Clin. Oncol.* 8 (9), 1563–1574. <https://doi.org/10.1200/JCO.1990.8.9.1563>.
- Albain, K.S., Crowley, J.J., LeBlanc, M., Livingston, R.B., 1991. Survival determinants in extensive-stage non-small-cell lung cancer: the Southwest Oncology Group experience. *J. Clin. Oncol.* 9 (9), 1618–1626. <https://doi.org/10.1200/JCO.1991.9.9.1618>.
- Alexiou, C., Onyeaka, C.V.P., Beggs, D., Akar, R., Beggs, L., Salama, F.D., et al., 2002. Do women live longer following lung resection for carcinoma? *Eur. J. Cardio-Thorac. Surg.* 21 (February (2)), 319–325. [https://doi.org/10.1016/S1010-7940\(01\)01114-9](https://doi.org/10.1016/S1010-7940(01)01114-9). Oxford University Press.
- Anon, 2015. WHO Global Report on Trends in Prevalence of Tobacco Smoking 2015. Available from: World Health Organization. https://apps.who.int/iris/bitstream/handle/10665/156262/9789241564922_eng.pdf;jsessionid=A4C26C219756E8D2FA3E585A52FA5FB7?sequence=1.
- Anon, 2018. WHO Global Report on Trends in Prevalence of Tobacco Smoking 2000–2025, Second Edition. Available from: World Health Organization, Geneva. <https://apps.who.int/iris/bitstream/handle/10665/272694/9789241514170-eng.pdf>.
- Antonia, S.J., López-Martin, J.A., Bendell, J., Ott, P.A., Taylor, M., Eder, J.P., et al., 2016. Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (CheckMate 032): a multicentre, open-label, phase 1/2 trial. *Lancet Oncol.* 17 (7), 883–895. [https://doi.org/10.1016/S1470-2045\(16\)30098-5](https://doi.org/10.1016/S1470-2045(16)30098-5).
- Antonia, S.J., Villegas, A., Daniel, D., Vicente, D., Murakami, S., Hui, R., et al., 2017. Durvalumab after Chemoradiotherapy in stage III non-Small-Cell Lung Cancer. *N. Engl. J. Med.* 377 (20). <https://doi.org/10.1056/NEJMoa1709937>. NEJMoa1709937.

- Araujo, J.M., Prado, A., Cardenas, N.K., Zaharia, M., Dyer, R., Doimi, F., et al., 2016. Repeated observation of immune gene sets enrichment in women with non-small cell lung cancer. *Oncotarget* 7 (15), 20282. <https://doi.org/10.18632/oncotarget.7943>.
- Ardizzoni, A., Boni, L., Tiseo, M., Fossella, F.V., Schiller, J.H., Paesmans, M., et al., 2007. Cisplatin- versus carboplatin-based chemotherapy in first-line treatment of advanced non-small-cell lung cancer: an individual patient data meta-analysis. *J. Natl. Cancer Inst.* 99 (11), 847–857. <https://doi.org/10.1093/jnci/djk196>.
- Artal Cortés, Á., Calera Urquiza, L., Hernando Cubero, J., 2015. Adjuvant chemotherapy in non-small cell lung cancer: state-of-the-art. *Transl. Lung Cancer Res.* 4 (2), 191–197. <https://doi.org/10.3978/j.issn.2218-6751.2014.06.01>.
- Aupérin, A., Le, P.Échoux C., Pignon, J.P., Koning, C., Jeremic, B., Clamon, G., et al., 2006. Concomitant radio-chemotherapy based on platin compounds in patients with locally advanced non-small cell lung cancer (NSCLC): A meta-analysis of individual data from 1764 patients. *Ann. Oncol.* 17 (3), 473–483. <https://doi.org/10.1093/annonc/mdj117>.
- Aupérin, A., Le, P.Échoux C., Rolland, E., Curran, W.J., Furuse, K., Fournel, P., et al., 2010. Meta-analysis of concomitant versus sequential radiochemotherapy in locally advanced non-small-cell lung cancer. *J. Clin. Oncol.* 28 (13), 2181–2190. <https://doi.org/10.1200/JCO.2009.26.2543>.
- Bandera, E.V., Freudenheim, J.L., Marshall, J.R., Zielezny, M., Priore, R.L., Brasure, J., et al., 1997. Diet and alcohol consumption and lung cancer risk in the New York State Cohort (United States). *Cancer Causes Control* 8 (0957–5243; 0957–5243; 6):828–840.
- Barfod, K.K., Vrankx, K., Mirsepasi-Lauridsen, H.C., Hansen, J.S., Hougaard, K.S., Larsen, S.T., et al., 2015. The murine lung microbiome changes during lung inflammation and intranasal vancomycin treatment. *Open Microbiol. J.* 9 (Lm), 167–179. <https://doi.org/10.2174/1874285801509010167>.
- Barlesi, F., Scherpereel, A., Rittmeyer, A., Pazzola, A., Ferrer Tur, N., Kim, J.H., et al., 2013. Randomized phase III trial of maintenance bevacizumab with or without pemetrexed after first-line induction with bevacizumab, cisplatin, and pemetrexed in advanced nonsquamous non-small-cell lung cancer: AVAPERL (MO22089). *J. Clin. Oncol.* 31 (24), 3004–3011. <https://doi.org/10.1200/JCO.2012.42.3749>.
- Barlesi, F., Scherpereel, A., Gorbunova, V., Gervais, R., Vikström, A., Chouaid, C., et al., 2014. Maintenance bevacizumab-pemetrexed after first-line cisplatin-pemetrexed-bevacizumab for advanced nonsquamous non-small-cell lung cancer: updated survival analysis of the AVAPERL (MO22089) randomized phase III trial. *Ann. Oncol.* 25 (5), 1044–1052. <https://doi.org/10.1093/annonc/mdl098>.
- Barrera, R., Morales Fuentes, J., 2012. Lung cancer in women. *Lung Cancer Targets Ther* 3 (December), 79. <https://doi.org/10.2147/LC.TS37319>.
- Bell, D.W., Lynch, T.J., Haserlat, S.M., Harris, P.L., Okimoto, R.A., Brannigan, B.W., et al., 2005. Epidermal growth factor receptor mutations and gene amplification in non-small-cell lung cancer: molecular analysis of the IDEAL/INTACT gefitinib trials. *J. Clin. Oncol.* 23 (31), 8081–8092. <https://doi.org/10.1200/JCO.2005.02.7078>.
- Berge, G., Mollerup, S., Øvrebo, S., Hewer, A., Phillips, D.H., Eilertsen, E., et al., 2004. Role of estrogen receptor in regulation of polycyclic aromatic hydrocarbon metabolic activation in lung. *Lung Cancer* 45 (3), 289–297. <https://doi.org/10.1016/j.lungcan.2004.02.014>.
- Blakely, C.M., Watkins, T.B.K., Wu, W., Gini, B., Chabon, J.J., McCoach, C.E., et al., 2017. Evolution and clinical impact of co-occurring genetic alterations in advanced-stage EGFR-mutant lung cancers. *Nat. Genet.* NIH Public Access 49 (December (12)), 1693–1704. <https://doi.org/10.1038/ng.3990>.
- Borghaei, H., Paz-Ares, L., Horn, L., Spigel, D.R., Steins, M., Ready, N.E., et al., 2015. Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N. Engl. J. Med.* 373 (17), 1627–1639. <https://doi.org/10.1056/NEJMoa1507643>.
- Brahmer, J.R., Dahlborg, S.E., Gray, R.J., Schiller, J.H., Perry, M.C., Sandler, A., et al., 2011. Sex differences in outcome with bevacizumab therapy: analysis of patients with advanced-stage non-small cell lung cancer treated with or without bevacizumab in combination with paclitaxel and carboplatin in the eastern cooperative oncology group trial 459. *J. Thorac. Oncol. Elsevier* 6 (1), 103–108. <https://doi.org/10.1097/JTO.0b013e3181fa8efd>.
- Brahmer, J., Reckamp, K.L., Baas, P., Crinò, L., Eberhardt, W.E.E., Poddubskaya, E., et al., 2015. Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N. Engl. J. Med.* 373, 123–135. <https://doi.org/10.1056/NEJMoa1504627>.
- Brahmer, J.R., Rodriguez-Abreu, D., Robinson, A.G., Hui, R., Csösz, T., Fülöp, A., et al., 2017a. Progression after the next line of therapy (PFS2) and updated OS among patients with advanced NSCLC and PD-L1 TPS >50% enrolled in KEYNOTE-024. *J. Clin. Oncol.* 35 (suppl). https://doi.org/10.1200/JCO.2017.35.15_suppl.9000. abstr 9000.
- Brahmer, J., Horn, L., Jackman, D., Spigel, D., Antonia, S., Hellmann, M., et al., 2017b. Abstract CT077: five-year follow-up from the CA209-003 study of nivolumab in previously treated advanced non-small cell lung cancer (NSCLC): clinical characteristics of long-term survivors. *Cancer Res. American Association for Cancer Research* 77 (July (13 Supplement)). <https://doi.org/10.1158/1538-7445.AM2017-CT077>. CT077-CT077.
- Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R.L., Torre, L.A., Jemal, A., 2018. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* 68 (6), 394–424. <https://doi.org/10.3322/caac.21492>.
- Bronte, G., Rolfo, C., Passiglia, F., Rizzo, S., Gil-Bazo, I., Fiorentino, E., et al., 2015. What can platinum offer yet in the treatment of PS2 NSCLC patients? A systematic review and meta-analysis. *Crit. Rev. Oncol. Hematol. Elsevier Ireland Ltd.* 95 (3), 306–317. <https://doi.org/10.1016/j.critrevonc.2015.03.010>.
- Brunelli, A., Charloux, A., Bolliger, C.T., Rocco, G., Sculier, J.P., Varela, G., et al., 2009. ERS/ESTS clinical guidelines on fitness for radical therapy in lung cancer patients (surgery and chemo-radiotherapy). *Eur. Respir. J.* 34 (1), 17–41. <https://doi.org/10.1183/09031936.00184308>.
- Burdett, S., 2014. Preoperative chemotherapy for non-small-cell lung cancer: a systematic review and meta-analysis of individual participant data. *Lancet* 383 (9928), 1561–1571. <https://doi.org/10.1097/JTO.0b013e3181b9ecca>. NSCLC Meta-analysis Collaborative Group. Open Access article distributed under the terms of CC BY.
- Burdett, S., Pignon, J.P., Trierney, J., Tribodet, H., Stewart, L., Le Pechoux, C., et al., 2015. Adjuvant chemotherapy for resected early-stage non-small cell lung cancer. *Cochrane Database Syst. Rev.* 2 (March). <https://doi.org/10.1002/14651858.CD011430>.
- Butts, C.A., Ding, K., Seymour, L., Twumasi-Ankrah, P., Graham, B., Gandara, D., et al., 2010. Randomized phase III trial of vinorelbine plus cisplatin compared with observation in completely resected stage IB and II non-small-cell lung cancer: updated survival analysis of JBR-10. *J. Clin. Oncol.* 28 (1), 29–34. <https://doi.org/10.1200/JCO.2009.24.0333>.
- Calles, A., Liao, X., Sholl, L.M., Rodig, S.J., Freeman, G.J., Butaney, M., et al., 2015a. Expression of PD-1 and its ligands, PD-L1 and PD-L2, in smokers and never smokers with KRAS-Mutant lung Cancer. *J. Thorac. Oncol. Elsevier* 10 (December (12)), 1726–1735. <https://doi.org/10.1097/JTO.0000000000000687>.
- Calles, A., Liao, X., Sholl, L.M., Rodig, S.J., Freeman, G.J., Butaney, M., et al., 2015b. Expression of PD-1 and its ligands, PD-L1 and PD-L2, in smokers and never smokers with KRAS-Mutant lung Cancer. *J. Thorac. Oncol. Elsevier* 10 (December (12)), 1726–1735. <https://doi.org/10.1097/JTO.0000000000000687>.
- Camidge, D.R., Kim, H.R., Ahn, M.-J., Yang, J.C.H., Han, J.Y., Lee, J.S., et al., 2018. Brigatinib versus Crizotinib in ALK-Positive non-Small-Cell Lung Cancer. *N. Engl. J. Med.* 379, 2027–2039. <https://doi.org/10.1056/NEJMoa1810171>.
- C-FJ, Yang, Chan, D.Y., Speicher, P.J., Gulack, B.C., Wang, X., Hartwig, M.G., et al., 2016. Role of adjuvant therapy in a population-based cohort of patients with early-stage small-cell lung cancer. *J. Clin. Oncol.* 34 (10), 1057–1064. <https://doi.org/10.1200/JCO.2015.63.8171>.
- Chakraborty, S., Ganti, A.K., Marr, A., Batra, K.S., 2010. Lung cancer in women: role of estrogens. *Expert Rev. Respir. Med.* 4 (4), 509–518. <https://doi.org/10.1586/ers.10.50>.
- Chansky, K., Sculier, J.P., Crowley, J.J., Giroux, D., Van Meerbeeck, J., Goldstraw, P., 2009. The International Association for the Study of Lung Cancer Staging Project: Prognostic Factors and Pathologic TNM Stage in Surgically Managed Non-small Cell Lung Cancer. *J. Thorac. Oncol.* 4 (July (7)), 792–801. <https://doi.org/10.1097/JTO.0b013e3181a7716e>.
- Chen, Z., Mei, J., Liu, L., Wang, G., Li, Z., Hou, J., et al., 2016. PD-L1 expression is associated with advanced non-small cell lung cancer. *Oncol. Lett. Spandidos Publications* 12 (August (2)), 921–927. <https://doi.org/10.3892/ol.2016.4741>.
- Cheng, T.Y.D., Cramb, S.M., Baade, P.D., Youlten, D.R., Nwogu, C., Reid, M.E., 2016. The international epidemiology of lung cancer: latest trends, disparities, and tumor characteristics. *J. Thorac. Oncol. Elsevier Inc* 11 (10), 1653–1671. <https://doi.org/10.1016/j.jtho.2016.05.021>.
- Chu, S.C., Hsieh, C.J., Wang, T.F., Hong, M.K., Chu, T.Y., 2017. Antiestrogen use in breast cancer patients reduces the risk of subsequent lung cancer: a population-based study. *Cancer Epidemiol.* 48 (June), 22–28. <https://doi.org/10.1016/j.canep.2017.02.010>.
- Ciuleanu, T., Brodowicz, T., Zielinski, C., Kim, J.H., Krzakowski, M., Laack, E., et al., 2009. Maintenance pemetrexed plus best supportive care versus placebo plus best supportive care for non-small-cell lung cancer: a randomised, double-blind, phase 3 study. *Lancet. Elsevier Ltd* 374 (9699), 1432–1440. [https://doi.org/10.1016/S0140-6736\(09\)61497-5](https://doi.org/10.1016/S0140-6736(09)61497-5).
- Collisson, E.A., Campbell, J.D., Brooks, A.N., Berger, A.H., Lee, W., Chmielecki, J., et al., 2014. Comprehensive molecular profiling of lung adenocarcinoma. *Nature* 511 (July (7511)), 543–550. <https://doi.org/10.1038/nature13385>.
- Conforti, F., Pala, L., Bagnardi, V., De Pas, T., Martinetti, M., Viale, G., et al., 2018. Metastatic Cancer Immunotherapy efficacy and patients' sex: a systematic review and meta-analysis. *Lancet Oncol. Elsevier Ltd* 4 (18), 1–10. [https://doi.org/10.1016/S1470-2045\(18\)30261-4](https://doi.org/10.1016/S1470-2045(18)30261-4).
- Corso, C.D., Rutter, C.E., Park, H.S., Lester-Coll, N.H., Kim, A.W., Wilson, L.D., et al., 2015. Role of chemoradiotherapy in elderly patients with limited-stage small-cell lung cancer. *J. Clin. Oncol.* 33 (36), 4240–4246. <https://doi.org/10.1200/JCO.2015.62.4270>.
- Cote, M.L., Yoo, W., Wenzlaff, A.S., et al., 2009. Tobacco and estrogen metabolic polymorphisms and risk of non-small cell lung cancer in women. *Carcinogenesis* 30 (January (4)), 626–635. <https://doi.org/10.1093/carcin/bgp033>.
- Davis, A., Kwang Chae, Y., Agte, S., Pan, A., Anjali Mohindra, N., Meucci Villafra, V., et al., 2017. Association of tumor mutational burden with smoking and mutation status in non-small cell lung cancer (NSCLC). *Suppl. 24. J. Clin. Oncol.* 35 (7, suppl (March 1)). https://doi.org/10.1200/JCO.2017.35.7_suppl.24. 24-24.
- Delbaldo, C., Michiels, S., Syz, N., Soria, J.C., Le, Chevalier T., Pignon, J.P., 2004. Benefits of adding drug to a single-agent or a 2-agent chemotherapy regimen in advanced non-small-cell lung cancer: a meta-analysis. *J. Am. Med. Assoc.* 292 (4), 470–484. <https://doi.org/10.1001/jama.292.4.470>.
- Des, Guetz G., Uzzan, B., Nicolas, P., Valeyre, D., Sebbane, G., Morere, J.F., 2012. Comparison of the efficacy and safety of single-agent and doublet chemotherapy in advanced non-small cell lung cancer in the elderly: a meta-analysis. *Crit. Rev. Oncol. Hematol.* 84 (3), 340–349. <https://doi.org/10.1016/j.critrevonc.2012.03.007>.
- Ding, L., Getz, G., Wheeler, D.A., Mardis, E.R., McLellan, M.D., Cibulskis, K., et al., 2008. Somatic mutations affect key pathways in lung adenocarcinoma. *Nature* 455 (October (7216)), 1069–1075. <https://doi.org/10.1038/nature07423>.
- Ding, X., Li, L., Tang, C., Meng, C., Xu, W., Wei, X., et al., 2018. Cytoplasmic expression of estrogen receptor β may predict poor outcome of EGFR-TKI therapy in metastatic lung adenocarcinoma. *Oncol. Lett.* 16 (June (2)), 2382–2390. <https://doi.org/10.3892/ol.2018.8936>.
- Dobbs, N.A., Twelves, C.J., Gillies, H., James, C.A., Harper, P.G., Rubens, R.D., 1995. Gender affects doxorubicin pharmacokinetics in patients with normal liver biochemistry. *Cancer Chemother. Pharmacol.* 36 (6), 473–476. <https://doi.org/10.1007/BF00600000>.

- 1007/BF00685796.
- Dogan, S., Shen, R., Ang, D.C., Johnson, M.L., D'Angelo, S.P., Paik, P.K., et al., 2012. Molecular epidemiology of EGFR and KRAS mutations in 3,026 lung adenocarcinomas: higher susceptibility of women to smoking-related KRAS-mutant cancers. *Clin Cancer Res. American Association for Cancer Research* 18 (November (22)), 6169–6177. <https://doi.org/10.1158/1078-0432.CCR-11-3265>.
- Dougherty, S.M., Mazhawidza, W., Bohn, A.R., Robinson, K.A., Mattingly, K.A., Blankenship, K.A., et al., 2006. Gender difference in the activity but not expression of estrogen receptors alpha and beta in human lung adenocarcinoma cells. *Endocr. Relat. Cancer* 13 (1), 113–134. <https://doi.org/10.1677/erc.1.01118>.
- Dresler, C.M., Fratelli, C., Babb, J., Everley, L., Evans, A.A., Clapper, M.L., 2000. Gender differences in genetic susceptibility for lung cancer. *Lung Cancer* 30 (December (3)), 153–160. [https://doi.org/10.1016/S0169-5002\(00\)00163-X](https://doi.org/10.1016/S0169-5002(00)00163-X).
- Eckardt, J.R., Von, Pawel J., Papai, Z., Tomova, A., Tzekova, V., Crofts, T.E., et al., 2006. Open-label, multicenter, randomized, phase III study comparing oral topotecan/cisplatin versus etoposide/cisplatin as treatment for chemotherapy-naïve patients with extensive-disease small-cell lung cancer. *J. Clin. Oncol.* 24 (13), 2044–2051. <https://doi.org/10.1200/JCO.2005.03.3332>.
- Eckardt, J.R., von, Pawel J., Pujol, J.-L., Papai, Z., Quoix, E., Ardzizoni, A., et al., 2007. Phase III study of oral compared with intravenous topotecan As second-line therapy in small-cell lung Cancer. *J. Clin. Oncol.* 25 (15), 2086–2092. <https://doi.org/10.1200/JCO.2006.08.3998>.
- El Osta, B., Behera, M., Kim, S., Berry, L.D., Sica, G., Pillai, R.N., et al., 2019. Characteristics and outcomes of patients with metastatic KRAS mutant lung adenocarcinomas: the Lung Cancer mutation Consortium experience. *J. Thorac. Oncol.* 5 (February). <https://doi.org/10.1016/j.jtho.2019.01.020>. pii: S1556-0864(19) 30097-8.
- Fallet, V., Cadranel, J., Doubre, H., Toper, C., Monnet, I., Chinnet, T., et al., 2014. Prospective screening for ALK: clinical features and outcome according to ALK status. *Eur J Cancer. Pergamon* 50 (May (7)), 1239–1246. <https://doi.org/10.1016/j.ejca.2014.02.001>.
- Fink, T.H., Huber, R.M., Heigener, D.F., Eschbach, C., Waller, C., Steinhauer, E.U., et al., 2012. Topotecan/cisplatin compared with cisplatin/etoposide as first-line treatment for patients with extensive disease small-cell lung cancer: final results of a randomized phase iii trial. *J. Thorac. Oncol.* 7 (9), 1432–1439. <https://doi.org/10.1097/JTO.0b013e318260de75>.
- Finkelstein, D.M., Ettinger, D.S., Ruckdeschel, J.C., 1986. Long-term survivors in metastatic non-small-cell lung cancer: an Eastern Cooperative Oncology Group study. *J. Clin. Oncol.* 4 (5), 702–709. <https://doi.org/10.1200/JCO.1986.4.5.702>.
- Foeglé, J., Hédelin, G., Lebitasy, M.P., Purohit, A., Velten, M., Quoix, E., 2007. Specific features of non-small cell lung cancer in women: a retrospective study of 1738 cases diagnosed in Bas-Rhin between 1982 and 1997. *J. Thorac. Oncol.* 2 (6), 466–474. <https://doi.org/10.1097/01.JTO.00000275340.39960.25>.
- Früh, M., De, Ruysscher D., Popat, S., Crinò, L., Peters, S., Felip, E., 2013. Small-cell lung cancer (SCLC): ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann. Oncol.* 24 (SUPPL.6), 23813929. <https://doi.org/10.1093/annonc/mdt178>.
- Fu, J.B., Kau, T.Y., Severson, R.K., Kalemkerian, G.P., 2005. Lung Cancer in women. *Chest* 127 (March (3)), 768–777.
- Gandhi, L., Rodríguez-Abreu, D., Gadgeel, S., Esteban, E., Felip, E., de, Angelis F., et al., 2018. Pembrolizumab plus chemotherapy in metastatic non-Small-Cell lung Cancer. *N. Engl. J. Med.* 378 (May (22)), 2078–2092. <https://doi.org/10.1056/NEJMoa1801005>.
- Garon, E.B., Dubinett, S.M., Hosmer, W., Reckamp, K.L., Kabbinavar, F.F., Goodglick, L., et al., 2010. Randomized phase II study of erlotinib (E) alone or combined with fulvestrant (F) in previously treated patients with advanced non-small cell lung cancer (NSCLC). *J Clin Oncol. American Society of Clinical Oncology* 28 (May (15, suppl)). <https://doi.org/10.1158/1538-7445.AM2013-4664>. TPS295-TPS295.
- Garon, E.B., Ciuleanu, T.E., Arrieta, O., Prabhaskar, K., Syrigos, K.N., Goksel, T., et al., 2014. Ramucicrumab plus docetaxel versus placebo plus docetaxel for second-line treatment of stage IV non-small-cell lung cancer after disease progression on platinum-based therapy (REVEL): a multicentre, double-blind, randomised phase 3 trial. *Lancet* 384 (9944), 665–673. [https://doi.org/10.1016/S0140-6736\(14\)60845-X](https://doi.org/10.1016/S0140-6736(14)60845-X).
- Gealy, R., Zhang, L., Siegfried, J.M., Luketich, J.D., Keohavong, P., 1999. Comparison of mutations in the p53 and K-ras genes in lung carcinomas from smoking and non-smoking women. *Cancer Epidemiol. Biomarkers Prev.* 8 (April (4 Pt 1)), 297–302.
- Gilligan, D., Nicolson, M., Smith, I., Groen, H., Dalesio, O., Goldstraw, P., et al., 2007. Preoperative chemotherapy in patients with resectable non-small cell lung cancer: results of the MRC LU22/NVALT 2/EORTC 08012 multicentre randomised trial and update of systematic review. *Lancet* 369 (9577), 1929–1937. [https://doi.org/10.1016/S0140-6736\(07\)60714-4](https://doi.org/10.1016/S0140-6736(07)60714-4).
- Giovino, G.A., 2002. Epidemiology of tobacco use in the United States. *Oncogene* 21 (October (48)), 7326–7340.
- Girón-González, J.A., Moral, F.J., Elvira, J., García-Gil, D., Guerrero, F., Gavilán, I., et al., 2000. Consistent production of a higher TH1:TH2 cytokine ratio by stimulated T cells in men compared with women. *Eur. J. Endocrinol.* 143 (July (1)), 31–36.
- Govindan, R., Ding, L., Griffith, M., Subramanian, J., Dees, N.D., Kanchi, K.L., et al., 2012. Genomic landscape of non-small cell lung cancer in smokers and never-smokers. *Cell. Elsevier* 150 (September (6)), 1121–1134. <https://doi.org/10.1016/j.cell.2012.08.024>.
- Gridelli, C., Perrone, F., Gallo, C., Cigolari, S., Rossi, A., Piantadosi, F., et al., 2003. Chemotherapy for elderly patients with advanced non-small-cell lung cancer: the multicenter Italian lung cancer in the elderly study (MILES) phase III randomized trial. *J. Natl. Cancer Inst.* 95 (5), 362–372. <https://doi.org/10.1093/jnci/95.5.362>.
- Gridelli, C., Ardzizoni, A., Le, Chevalier T., Manegold, C., Perrone, F., Thatcher, N., et al., 2004. Treatment of advanced non-small-cell lung cancer patients with ECOG performance status 2: results of an European Experts Panel. *Ann. Oncol.* 15 (3), 419–426. <https://doi.org/10.1093/annonc/mdh087>.
- Gridelli, C., Di, Maio M., Chiodini, P., Georgoulas, V., Hatzidaki, D., Takeda, K., et al., 2009. Meta-analysis of single-agent chemotherapy compared with combination chemotherapy as second-line treatment of advanced non-small-cell lung cancer. *J. Clin. Oncol.* 27 (11), 1836–1843. <https://doi.org/10.1200/JCO.2008.17.5844>.
- Gridelli, C., Morabito, A., Cavanna, L., Luciani, A., Maione, P., Bonanno, L., et al., 2018. Cisplatin-based first-line treatment of elderly patients with advanced non-small-cell lung Cancer: joint analysis of MILES-3 and MILES-4 phase III trials. *J. Clin. Oncol.* 36 (25), 2585–2592. <https://doi.org/10.1200/JCO.2017.76.8390>.
- Group, N.M.C., 2008. Chemotherapy in addition to supportive care improves survival in advanced non-small-cell lung Cancer: a systematic review and meta-analysis of individual patient data from 16 randomized controlled trials. *J. Clin. Oncol.* 26 (28), 4617–4625. <https://doi.org/10.1200/JCO.2008.17.7162>.
- Group, N.M.C., 2010. Adjuvant chemotherapy, with or without postoperative radiotherapy, in operable non-small-cell lung cancer: two meta-analyses of individual patient data. *Lancet. Elsevier Ltd* 375 (9722), 1267–1277. [https://doi.org/10.1016/S0140-6736\(10\)60059-1](https://doi.org/10.1016/S0140-6736(10)60059-1).
- Guo, L., Zhang, Q., Ma, X., Wang, J., 2017. Liang T. miRNA and mRNA expression analysis reveals potential sex-biased miRNA expression. *Nature* 7 (January (2017)), 39812. <https://doi.org/10.1038/srep39812>.
- Hanna, N., Shepherd, F.A., Fossella, F.V., Pereira, J.R., de, Marinis F., von, Pawel J., et al., 2004. Randomized phase III trial of pemetrexed versus docetaxel in patients with non-small-cell lung cancer previously treated with chemotherapy. *J. Clin. Oncol.* 22 (9), 1589–1597. <https://doi.org/10.1200/JCO.2004.08.163>.
- Harlos, C., Musto, G., Lambert, P., Ahmed, R., Pitz, M.W., 2015. Androgen pathway manipulation and survival in patients with lung Cancer. *Horm Cancer. Springer US* 6 (June (2–3)), 120–127. <https://doi.org/10.1007/s12672-015-0218-1>.
- Hellmann, M.D., Callahan, M.K., Awad, M.M., Calvo, E., Asciero, P.A., Atmaca, A., et al., 2018a. Tumor mutational burden and efficacy of nivolumab monotherapy and in combination with ipilimumab in small-cell lung Cancer. *Cancer Cell. Elsevier Inc.* 33 (5), 853–861. <https://doi.org/10.1016/j.ccell.2018.04.001>. e4.
- Hellmann, M.D., Ciuleanu, T.E., Pluzanski, A., Seok Lee, J., Otterson, G.A., Audigier-Valette, C., et al., 2018b. Nivolumab plus ipilimumab in lung Cancer with a high tumor mutational burden. *N. Engl. J. Med.* <https://doi.org/10.1056/NEJMoa1801946>. NEJMoa1801946.
- Hemminki, K., 1993. DNA adducts, mutations and cancer. *Carcinogenesis* 14 (October (10)), 2007–2012.
- Herbst, R.S., Baas, P., Kim, D.W., Felip, E., Pérez-Gracia, J.L., Han, J.Y., et al., 2016. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet* 387 (10027), 1540–1550. [https://doi.org/10.1016/S0140-6736\(15\)01281-7](https://doi.org/10.1016/S0140-6736(15)01281-7).
- Herbst, R.S., Morgensztern, D., Boshoff, C., 2018. The biology and management of non-small cell lung cancer. *Nature. Nature Publishing Group* 553 (7689), 446–454. <https://doi.org/10.1038/nature25183>.
- Hernandez-Boussard, T.M., Hainaut, P., 1998. A specific spectrum of p53 mutations in lung cancer from smokers: review of mutations compiled in the IARC p53 database. *Environ Health Perspect. National Institute of Environmental Health Science* 106 (July (7)), 385–391. <https://doi.org/10.1289/ehp.98106385>.
- Horn, L., Mansfield, A.S., Szczesna, A., Havel, L., Krzakowski, M., Hochmair, M.J., et al., 2018. First-line atezolizumab plus chemotherapy in extensive-stage small-cell lung Cancer. *N. Engl. J. Med.* <https://doi.org/10.1056/NEJMoa1809064>. NEJMoa1809064.
- Hosgood, H.D., Sapkota, A.R., Rothman, N., Rohan, T., Hu, W., Xu, J., et al., 2014. The potential role of lung microbiota in lung cancer attributed to household coal burning exposures. *Environ. Mol. Mutagen.* 55 (8), 643–651. <https://doi.org/10.1002/em.21878>.
- Hsu, L.H., Chu, N.M., Kao, S.H., 2017. Estrogen, estrogen receptor and lung cancer. *Int. J. Mol. Sci.* 18 (8), 1–17. <https://doi.org/10.3390/ijms18081713>.
- Hyde, Z., Flicker, L., McCaul, K.A., Almeida, O.P., Hankey, G.J., Chubb, S.A., et al., 2012. Associations between testosterone levels and incident prostate, lung, and colorectal Cancer: A Population-Based Study. *Cancer Epidemiol Biomarkers Prev* 21 (August (8)), 1319–1329. <https://doi.org/10.1158/1055-9965.EPI-12-0129>.
- Ivanova, M.M., Mazhawidza, W., Dougherty, S.M., Klinge, C.M., 2010. Sex differences in estrogen receptor subcellular location and activity in lung adenocarcinoma cells. *Am. J. Respir. Cell Mol. Biol.* 42 (3), 320–330. <https://doi.org/10.1165/rcmb.2009-0059OC>.
- Jackute, J., Zemaitis, M., Pranys, D., Sitkauskienė, B., Miliauskas, S., Vaitkiene, S., et al., 2018. Distribution of M1 and M2 macrophages in tumor islets and stroma in relation to prognosis of non-small cell lung cancer. *BMC Immunol. BioMed Central* 19 (January (1)), 3. <https://doi.org/10.1186/s12865-018-0241-4>.
- Jemal, A., Miller, K.D., Ma, J., Siegel, R.L., Fedewa, S.A., Islami, F., et al., 2018. Higher lung Cancer incidence in young women than young men in the United States. *N. Engl. J. Med.* 378 (21), 1999–2009. <https://doi.org/10.1056/NEJMoa1715907>.
- Jett, J.R., Schild, S.E., Kesler, K.A., Kalemkerian, G.P., 2013. Treatment of small cell lung cancer: diagnosis and management of lung cancer, 3rd ed: american college of chest physicians evidence-based clinical practice guidelines. *Chest* 143 (5 SUPPL), 2–3. <https://doi.org/10.1378/chest.12-2363>.
- Johnson, B.E., Steinberg, S.M., Phelps, R., Edison, M., Veach, S.R., Ihde, D.C., 1988. Female patients with small cell lung cancer live longer than male patients. *Am. J. Med.* 85 (2), 194–196. [https://doi.org/10.1016/S0002-9343\(88\)80341-3](https://doi.org/10.1016/S0002-9343(88)80341-3).
- Jotte, R., Cappuzzo, F., Vynnychenko, I., Stroyakovskiy, D., Delvys, R.A.D., Maen, A., et al., 2019. Impower131: Primary PFS and Safety Analysis of a Randomized Phase III Study of Atezolizumab + Carboplatin + Paclitaxel or Nab-Paclitaxel vs Carboplatin + Nab-Paclitaxel as 1L Therapy in Advanced Squamous NSCLC. https://doi.org/10.1200/JCO.2018.36.18_suppl.LBA9000.
- Kanashiki, M., Sairenchi, T., Saito, Y., Ishikawa, H., Satoh, H., Sekizawa, K., 2005. Body

- mass index and lung cancer: a case-control study of subjects participating in a mass-screening program. *Chest* 128 (3), 1490–1496. <https://doi.org/10.1378/chest.128.3.1490>.
- Kaul, S., Srinivas, N.R., Igwemezie, L.N., Barbhaiya, R.H., 1996. A pharmacodynamic evaluation of hematologic toxicity associated with etoposide phosphate in the treatment of cancer patients. *Semin. Oncol.* 23 (6 Suppl 13), 15–22.
- Kawajiri, K., Nakachi, K., Imai, K., Watanabe, J., Hayashi, S., 1993. The CYP1A1 gene and cancer susceptibility. *Crit. Rev. Oncol. Hematol.* 14 (February (1)), 77–87.
- Keselman, A., Fang, X., White, P.B., Heller, N.M., 2017. Estrogen signaling contributes to sex differences in macrophage polarization during asthma. *J. Immunol.* 199 (September (5)), 1573–1583. <https://doi.org/10.4049/jimmunol.1601975>.
- Kiyohara, C., Ohno, Y., 2010. Sex differences in lung cancer susceptibility: a review. *Genet. Med.* 7 (October (5)), 381–401. <https://doi.org/10.1016/j.genn.2010.10.002.30>.
- Kligerman, S., White, C., 2011. Epidemiology of lung Cancer in women: risk factors, survival, and screening. *Am J Roentgenol* 196 (February (2)), 287–295. <https://doi.org/10.2214/AJR.10.5412>.
- Kubota, K., Hida, T., Ishikura, S., Mizusawa, J., Nishio, M., Kawahara, M., et al., 2014. Etoposide and cisplatin versus irinotecan and cisplatin in patients with limited-stage small-cell lung cancer treated with etoposide and cisplatin plus concurrent accelerated hyperfractionated thoracic radiotherapy (JCOG0202): a randomised phase 3 study. *Lancet Oncol.* 15 (1), 106–114. [https://doi.org/10.1016/S1470-2045\(13\)70511-4](https://doi.org/10.1016/S1470-2045(13)70511-4).
- Kudoh, S., Takeda, K., Nakagawa, K., Takada, M., Katakami, N., Matsui, K., et al., 2006. Phase III study of docetaxel compared with vinorelbine in elderly patients with advanced non-small-cell lung cancer: Results of the West Japan Thoracic Oncology Group trial (WJTOG 9904). *J. Clin. Oncol.* 24 (22), 3657–3663. <https://doi.org/10.1200/JCO.2006.06.1044>.
- Kure, E.H., Ryberg, D., Hewer, A., Phillips, D.H., Skaug, V., Baera, R., et al., 1996. p53 mutations in lung tumours: relationship to gender and lung DNA adduct levels. *Carcinogenesis* 17 (October (10)), 2201–2205.
- Labots, G., Jones, A., de Visser S.J., Rissmann, R., Burggraaf, J., 2018. Gender differences in clinical registration trials: is there a real problem? *Br. J. Clin. Pharmacol.* 84 (4), 700–707. <https://doi.org/10.1111/bcp.13497>.
- Lally, B.E., Geiger, A.M., Urbanic, J.J., Butler, J.M., Wentworth, S., Perry, M.C., et al., 2009. Trends in the outcomes for patients with limited stage small cell lung cancer: An analysis of the Surveillance, Epidemiology, and End Results database. *Lung Cancer* 64 (2), 226–231. <https://doi.org/10.1016/j.lungcan.2008.08.010>.
- Lawrence, M.S., Stojanov, P., Mermel, C.H., Robinson, J.T., Garraway, L.A., Golub, T.R., et al., 2014. Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature*. Nature Publishing Group 505 (January (7484)), 495–501. <https://doi.org/10.1038/nature12912>.
- Lee, S.M., James, L.E., Qian, W., Spiro, S., Eisen, T., Gower, N.H., et al., 2009. Comparison of gemcitabine and carboplatin versus cisplatin and etoposide for patients with poorprognosis small cell lung cancer. *Thorax* 64 (1), 75–80. <https://doi.org/10.1136/thx.2007.093872>.
- Lee, C.K., Man, J., Lord, S., Cooper, W., Links, M., GebSKI, V., et al., 2018. Clinical and Molecular Characteristics Associated With Survival Among Patients Treated With Checkpoint Inhibitors for Advanced Non-Small Cell Lung Carcinoma: A Systematic Review and Meta-analysis. *JAMA Oncol.* 4 (February (2)), 210–216. <https://doi.org/10.1001/jamaoncol.2017.4427>.
- Lewis, D.R., Check, D.P., Caporaso, N.E., Travis, W.D., Devesa, S.S., 2014. US lung cancer trends by histologic type. *Cancer.* 120 (18), 2883–2892. <https://doi.org/10.1002/cncr.28749>.
- Li, M., Zhang, Q., Fu, P., Li, P., Peng, A., Zhang, G., et al., 2012. Pemetrexed plus platinum as the first-line treatment option for advanced non-small cell lung cancer: a meta-analysis of randomized controlled trials. *PLoS One* 7 (5).
- Lilenbaum, R., Villafior, V.M., Langer, C., O'Byrne, K., O'Brien, M., Ross, H.J., et al., 2009. Single-agent versus combination chemotherapy in patients with advanced non-small cell lung cancer and a performance status of 2: Prognostic factors and treatment selection based on two large randomized clinical trials. *J Thorac Oncol.* International Association for the Study of Lung Cancer 4 (7), 869–874. <https://doi.org/10.1097/JTO.0b013e3181a9a020>.
- Lim, E., Harris, G., Patel, A., Adachi, I., Edmonds, L., Song, F., 2009. Preoperative versus postoperative chemotherapy in patients with resectable non-small cell lung cancer: systematic review and indirect comparison meta-analysis of randomized trials. *J Thorac Oncol.* International Association for the Study of Lung Cancer 4 (11), 1380–1388.
- Lima, A.B.C., Macedo, L.T., Sasse, A.D., 2011. Addition of bevacizumab to chemotherapy in advanced non-small cell lung cancer: a systematic review and meta-analysis. *PLoS One* 6 (8), 1–8. <https://doi.org/10.1371/journal.pone.0022681>.
- Lin, G., Fan, X., Zhu, W., Huang, C., Zhuang, W., Xu, H., et al., 2017. Prognostic significance of PD-L1 expression and tumor infiltrating lymphocyte in surgically resectable non-small cell lung cancer. *Oncotarget.* Impact Journals 8 (October (48)), 83986–83994. <https://doi.org/10.18632/oncotarget.20233>.
- Lopes, G., Wu, Y., Kudaba, I., Kowalski, D., Cho, B.C., Castro, G., et al., 2018. Pembrolizumab vs platinum-based chemotherapy as first-line therapy for advanced / metastatic NSCLC with a KEYNOTE-042 study pembrolizumab and PD-L1 for metastatic NSCLC. Sliddeck 818, 12–13. https://doi.org/10.1200/JCO.2018.36.18_suppl.LBA4.
- Lopez-chavez, A., Young, T., Fages, S., Leon, L., Schiller, J.H., 2018. Bevacizumab Maintenance in Patients with Advanced Non – Small-Cell Lung Cancer, Clinical Patterns, and Outcomes in the Eastern Cooperative Oncology Group 4599 Study : Results of An Exploratory Analysis. *J. Thorac. Oncol.* 59 (15), 1–7. <https://doi.org/10.1097/JTO.0b013e318265b500>.
- Lord, R.V.N., Brabender, J., Gandara, D., Alberola, V., Camps, C., Domine, M., et al., 2002. Low ERCC1 expression correlates with prolonged survival after cisplatin plus gemcitabine chemotherapy in non-small cell lung cancer. *Clin. Cancer Res.* 8 (July (7)), 2286–2291.
- Mao, Q., Jiang, F., Yin, R., Wang, J., Xia, W., Dong, G., et al., 2018. Interplay between the lung microbiome and lung cancer. *Cancer Lett.* 415, 40–48. <https://doi.org/10.1016/j.canlet.2017.11.036>.
- Márquez-Garbán, D.C., Chen, H.W., Fishbein, M.C., Goodglick, L., Pietras, R.J., 2007a. Estrogen receptor signaling pathways in human non-small cell lung cancer. *Steroids* 72 (2), 135–143. <https://doi.org/10.1016/j.steroids.2006.11.019>.
- Márquez-Garbán, D.C., Chen, H.W., Fishbein, M.C., Goodglick, L., Pietras, R.J., 2007b. ER- β protein expression may be down-regulated after the addition of EGF. *Steroids* 72 (February (2)), 135–143.
- Martinez, P., Solange, P., Stammers, T., Soria, J., 2019. Immunotherapy for the first-line treatment of patients with metastatic non-small cell lung cancer. *Clin. Cancer Res.* <https://doi.org/10.1158/1078-0432.CCR-18-3904>.
- Mascaux, C., Paesmans, M., Berghmans, T., Branle, F., Lafitte, J.J., Lemaître, F., et al., 2000. A systematic review of the role of etoposide and cisplatin in the chemotherapy of small cell lung cancer with methodology assessment and meta-analysis. *Lung Cancer* 30 (1), 23–36. [https://doi.org/10.1016/S0169-5002\(00\)00127-6](https://doi.org/10.1016/S0169-5002(00)00127-6).
- Mazières, J., Peters, S., Lepage, B., Cortot, A.B., Barlesi, F., Beau-Faller, M., et al., 2013. Lung Cancer That harbors an HER2 mutation: epidemiologic characteristics and therapeutic perspectives. *J. Clin. Oncol.* 31 (June (16)), 1997–2003. <https://doi.org/10.1200/JCO.2012.45.6095>.
- Mazieres, J., Barlesi, F., Rouquette, I., Besse, B., 2018. Randomized phase II trial evaluating treatment with EGFR-TKI versus EGFR-TKI associated with anti-estrogen in women with non-squamous advanced stage NSCLC: IFCT-1003 LADIE trial. *J. Clin. Oncol.* 36 (15), 9097_suppl9097. <https://doi.org/10.1200/JCO>.
- Mediavilla-Varela, M., Page, M.M., Kreaahling, J., Antonia, S.J., Altiock, S., 2019. Anti-PD1 treatment to induce M1 polarization of tumor infiltrating macrophages in a 3D ex vivo system of lung cancer patients. *CO20173515_suppl.e23090 J Clin Oncol* 35 (15 suppl). https://doi.org/10.1200/JCO.2017.35.15_suppl.e23090.
- Meza, R., Meernik, C., Jeon, J., Cote, M.L., 2015. Lung cancer incidence trends by gender, race and histology in the United States, 1973–2010. *PLoS One* 10 (3), 1–14. <https://doi.org/10.1371/journal.pone.0121323>.
- Mikkonen, L., Pihlajamaa, P., Sahu, B., Zhang, F.P., Jänne, O.A., 2010. Androgen receptor and androgen-dependent gene expression in lung. *Mol. Cell. Endocrinol.* 317 (April (1–2)), 14–24. <https://doi.org/10.1016/j.mce.2009.12.022>.
- Minami, H., Yoshimura, M., Miyamoto, Y., Matsuoka, H., Tsubota, N., 2000. Lung Cancer in women. *Chest.* Elsevier 118 (December (6)), 1603–1609.
- Mollerup, S., Berge, G., Baera, R., Skaug, V., Hewer, A., Phillips, D.H., et al., 2006. Sex differences in risk of lung cancer: expression of genes in the PAH bioactivation pathway in relation to smoking and bulky DNA adducts. *Int. J. Cancer* 119 (August (4)), 741–744. <https://doi.org/10.1002/ijc.21891>.
- Morère, J.F., Bréchet, J.M., Westeel, V., Goumant, V., Lebeau, B., Vaylet, F., et al., 2010. Randomized phase II trial of gefitinib or gemcitabine or docetaxel chemotherapy in patients with advanced non-small-cell lung cancer and a performance status of 2 or 3 (IFCT-0301 study). *Lung Cancer* 70 (3), 301–307. <https://doi.org/10.1016/j.lungcan.2010.03.003>.
- Murthy, V.H., Krumholz, H.M., Gross, C.P., 2004. Participation in cancer clinical trials: race, sex-, and age-based disparities. *JAMA* 291 (June (22)), 2720–2726. <https://doi.org/10.1001/jama.291.22.2720227>.
- Nakamura, H., Ando, K., Shinmyo, T., Morita, K., Kurimoto, N., Morita, K., et al., 2011. Female gender is an independent prognostic factor in non-small-cell lung cancer: a meta-analysis. *Ann. Thorac. Cardiovasc. Surg.* 17 (5), 469–480. <https://doi.org/10.5761/atcs.0a.101637>.
- Nelson, H.H., Christiani, D.C., Mark, E.J., Wiencke, J.K., Wain, J.C., Kelsey, K.T., 1999. Implications and prognostic value of K-ras mutation for early-stage lung cancer in women. *J. Natl. Cancer Inst.* 91 (23), 2032–2038.
- North C, M., Christiani, D.C., 2013. Women and lung Cancer: what's new? *Semin. Thorac. Cardiovasc. Surg.* 25 (Summer (2)), 87–94. <https://doi.org/10.1053/j>.
- Nyberg, F., Argenius, V., Svartengren, K., Svensson, C., Pershagen, G., 1998. Dietary factors and risk of lung cancer in never-smokers. *Int. J. Cancer* 436 (4), 430–436.
- O'Brien, M.E.R., Ciuleanu, T.-E., Tsekov, H., Shparyk, Y., Cuceviá, B., Juhasz, G., et al., 2006. Phase III trial comparing supportive care alone with supportive care with oral topotecan in patients with relapsed small-cell lung cancer. *J. Clin. Oncol.* 24 (34), 5441–5447. <https://doi.org/10.1200/JCO.2006.06.5821>.
- O'Connell, J.P., Kris, M.G., Gralla, R.J., Groshen, S., Trust, A., Fiore, J.J., et al., 1986. Frequency and prognostic importance of pretreatment clinical characteristics in patients with advanced non-small-cell lung cancer treated with combination chemotherapy. *J. Clin. Oncol.* 4 (11), 1604–1614. <https://doi.org/10.1200/JCO.1986.4.11.1604>.
- Osterlind, K., Hansen, H.H., Hansen, M., Dombernowsky, P., Andersen, P.K., 1986. Long-term disease-free survival in small-cell carcinoma of the lung: a study of clinical determinants. *J. Clin. Oncol.* 4 (9), 1307–1313. <https://doi.org/10.1200/JCO.1986.4.9.1307>.
- Özdemir, B.C., Csajka, C., Dotto, G.P., Wagner, A.D., 2018. Sex differences in efficacy and toxicity of systemic treatments: an undervalued issue in the era of precision oncology. *J. Clin. Oncol.* 36 (26), 2680–2683. <https://doi.org/10.1200/JCO.2018.78.3290>.
- Paesmans, M., Sculier, J.P., Libert, P., Bureau, G., Dabouis, G., Thiriaux, J., et al., 1995. Prognostic factors for survival in advanced non-small-cell lung cancer: univariate and multivariate analyses including recursive partitioning and amalgamation algorithms in 1,052 patients. *J. Clin. Oncol.* 13 (5), 1221–1230. <https://doi.org/10.1200/JCO.1995.13.5.1221>.
- Paesmans, M., Sculier, J.P., Lecomte, J., Thiriaux, J., Libert, P., Sergysels, R., et al., 2000. Prognostic factors for patients with small cell lung carcinoma: analysis of a series of 763 patients included in 4 consecutive prospective trials with a minimum follow-up

- of 5 years. *Cancer* 89 (August (3)), 523–533.
- Paik, J.H., Choi, C.M., Kim, H., Jang, S.J., Choe, G., Kim, D.K., et al., 2012. Clinicopathologic implication of ALK rearrangement in surgically resected lung cancer. A proposal of diagnostic algorithm for ALK-rearranged adenocarcinoma. *Lung Cancer* 76 (3), 403–409. <https://doi.org/10.1016/j.lungcan.2011.11.008>.
- Pan, Z.K., Ye, F., Wu, X., An, H.X., Wu, J.X., 2015. Clinicopathological and prognostic significance of programmed cell death ligand1 (PD-L1) expression in patients with non-small cell lung cancer: a meta-analysis. *J. Thorac. Dis. AME Publications* 7 (March (3)), 462–470. <https://doi.org/10.3978/j.issn.2072-1439.2015.02.13>.
- Papadimitrakopoulou, V.A., Cobo, M., Bordoni, R., Dubray Longeras, P., Szalai, Z., Ursol, G., et al., 2019. IMpower132 : PFS and safety results with 1L atezolizumab + carboplatin / cisplatin + pemetrexed in stage IV Non-squamous NSCLC. Presented at: the IASLC 19th World Conference on Lung Cancer; September 23–26; Toronto, Canada. Abstract OA05.07. <https://doi.org/10.1016/j.jtho.2018.08.262>.
- Parkin, D.M., Tyczyński, J.E., Bunn, P.A., Kato, H., L.J., 2010. Global epidemiology of lung cancer. *Textb Prev. Detect. Early Lung Cancer*. 85 (1), 1–18.
- Paximadis, P.A., Beebe-Dimmer, J.L., George, J., Schwartz, A.G., Wozniak, A.J., Gadgeel, S.M., 2017. Comparing treatment strategies for stage I small cell lung cancer: a National Cancer database study. *J. Clin. Oncol.* 35 (15 suppl (May 20 2017)). https://doi.org/10.1200/JCO.2017.35.15_suppl.8566. 8566–8566.
- Paz-Ares, L., de Marinis F., Dediu, M., Thomas, M., Pujol, J.L., Bidoli, P., et al., 2012. Maintenance therapy with pemetrexed plus best supportive care versus placebo plus best supportive care after induction therapy with pemetrexed plus cisplatin for advanced non-squamous non-small-cell lung cancer (PARAMOUNT): a double-blind, phase 3, random. *Lancet Oncol. Elsevier Ltd.* 13 (3), 247–255. [https://doi.org/10.1016/S1470-2045\(12\)70063-3](https://doi.org/10.1016/S1470-2045(12)70063-3).
- Paz-Ares, L.G., De, Marinis F., Dediu, M., Thomas, M., Pujol, J.L., Bidoli, P., et al., 2013. PARAMOUNT: final overall survival results of the phase III study of maintenance pemetrexed versus placebo immediately after induction treatment with pemetrexed plus cisplatin for advanced nonsquamous non-small-cell lung cancer. *J. Clin. Oncol.* 31 (23). <https://doi.org/10.1200/JCO.2012.47.1102>. 2895–290 2.
- Paz-Ares, L., Luft, A., Vicente, D., Tafreshi, A., Gümiş, M., Mazières, J., et al., 2018. Pembrolizumab plus chemotherapy for squamous non-Small-Cell Lung Cancer. *N. Engl. J. Med.* <https://doi.org/10.1056/NEJMoa1810865>. NEJMoa1810865.
- Peters, S., Creelan, B., Hellmann, M.D., Socinski, M.A., Reck, M., Bhagwatheeswaran, P., et al., 2017. Abstract CT082: Impact of tumor mutation burden on the efficacy of first-line nivolumab in stage iv or recurrent non-small cell lung cancer: An exploratory analysis of CheckMate 026. *Cancer Res. American Association for Cancer Research* 77 (July (13 Supplement)). <https://doi.org/10.1158/1538-7445.AM2017-CT082>. CT082-CT082.
- Petrelli, F., Maltese, M., Tomasello, G., Conti, B., Borgonovo, K., Cabiddu, M., et al., 2018. Clinical and molecular predictors of PD-L1 expression in non-small-Cell lung Cancer: systematic review and meta-analysis. *Clin Lung Cancer. Elsevier Inc.* 19 (4), 315–322. <https://doi.org/10.1016/j.clcc.2018.02.006>.
- Pignon, J.-P., Arriagada, R., Ihde, D.C., Johnson, D.H., Perry, M.C., Souhami, R.L., et al., 1992. A meta-analysis of thoracic radiotherapy for small-cell lung cancer. *N. Engl. J. Med.* 327 (23), 1618–1624.
- Pignon, J.P., Tribodet, H., Scagliotti, G.V., Douillard, J.Y., Shepherd, F.A., Stephens, R.J., et al., 2008. Lung adjuvant cisplatin evaluation: a pooled analysis by the LACE collaborative group. *J. Clin. Oncol.* 26 (21), 3552–3559. <https://doi.org/10.1200/JCO.2007.13.9030>.
- Pinto, J.A., Vallejos, C.S., Raez, L.E., Mas, L.A., Ruiz, R., Torres-Roman, J.S., et al., 2018. Gender and outcomes in non-small cell lung cancer: an old prognostic variable comes back for targeted therapy and immunotherapy? *ESMO Open. BMJ Publishing Group* 3 (3), e000344. <https://doi.org/10.1136/esmoopen-2018-000344>.
- Pujol, J.L., Carestia, L., Daurès, J.P., 2000. Is there a case for cisplatin in the treatment of small-cell lung cancer? A meta-analysis of randomized trials of a cisplatin-containing regimen versus a regimen without this alkylating agent. *Br. J. Cancer* 83 (2), 177–183. <https://doi.org/10.1054/bjoc.2000.1164>.
- Pujol, J.L., Barlesi, F., Daurès, J.P., 2006. Should chemotherapy combinations for advanced non-small cell lung cancer be platinum-based? A meta-analysis of phase III randomized trials. *Lung Cancer* 51 (3), 335–345. <https://doi.org/10.1001/jama.292.4.470>.
- Qi, W.-X., Tang, L., He, A., Shen, Z., Lin, F., Yao, Y., 2012. Doublet versus single cytotoxic agent as first-line treatment for elderly patients with advanced non-small-Cell lung Cancer: a systematic review and meta-analysis. *Lung* 190 (5), 477–485. <https://doi.org/10.1007/s00408-012-9399-3>.
- Quoix, E., Zalcmán, G., Oster, J.P., Westeel, V., Pichon, E., Lavolé, A., et al., 2011. Carboplatin and weekly paclitaxel doublet chemotherapy compared with monotherapy in elderly patients with advanced non-small-cell lung cancer: IFCT-0501 randomised, phase 3 trial. *Lancet* 378 (9796), 1079–1088. [https://doi.org/10.1016/S0140-6736\(11\)60780-0](https://doi.org/10.1016/S0140-6736(11)60780-0).
- Ramos-Esquivel, A., Van Der, Laat A., Rojas-Vigott, R., Juárez, M., Corrales-Rodríguez, L., 2017. Anti-PD-1/anti-PD-L1 immunotherapy versus docetaxel for previously treated advanced non-small cell lung cancer: a systematic review and meta-analysis of randomised clinical trials. *ESMO Open* 2 (3). <https://doi.org/10.1136/esmoopen-2017-000236>.
- Raz, D., He, B., Rosell, R., Jablons, D., 2016. Bronchioloalveolar carcinoma: a review. *Clin. Lung Cancer* 7 (5), 313–322.
- Reck, M., von, Pawel J., Zatlokouk, P., Ramlau, R., Gorbounova, V., Hirsh, V., et al., 2009. Phase III trial of cisplatin plus gemcitabine with either placebo or bevacizumab as first-line therapy for nonsquamous non-small-cell lung cancer: AVAIL. *J. Clin. Oncol.* 27 (8), 1227–1234. <https://doi.org/10.1200/JCO.2007.14.5466>.
- Reck, M., Kaiser, R., Mellemeard, A., Douillard, J.Y., Orlov, S., Krzakowski, M., et al., 2014. Docetaxel plus nintedanib versus docetaxel plus placebo in patients with previously treated non-small-cell lung cancer (LUME-Lung 1): a phase 3, double-blind, randomised controlled trial. *Lancet Oncol.* 15 (2), 143–155. [https://doi.org/10.1016/S1470-2045\(13\)70586-2](https://doi.org/10.1016/S1470-2045(13)70586-2).
- Reck, M., Rodríguez-Abreu, D., Robinson, A.G., Hui, R., Csőszi, T., Fülöp, A., et al., 2016. Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N. Engl. J. Med.* 375 (19), 1823–1833. <https://doi.org/10.1056/NEJMoa1606774>.
- Reddy, V.P., Gay, L.M., Elvin, J.A., Vergilio, J.-A., Suh, J., Ramkissoon, S., et al., 2017. BRAF fusions in clinically advanced non-small cell lung cancer: an emerging target for anti-BRAF therapies. *J. Clin. Oncol. American Society of Clinical Oncology* 35 (May (15 suppl)). https://doi.org/10.1200/JCO.2017.35.15_suppl.9072. 9072–9072.
- Remon, J., Isla, D., Garrido, P., et al., 2017. Efficacy of tyrosine kinase inhibitors in EGFR-mutant lung cancer women in a real-world setting: the WORLD07 database. *Clin. Transl. Oncol.* 19 (December (12)), 1537–1542. <https://doi.org/10.1007/s12094-017-1700-8>.
- Rittmeyer, A., Barlesi, F., Waterkamp, D., Park, K., Ciardiello, F., von, Pawel J., et al., 2017. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. *Lancet* 389 (10066), 255–265. [https://doi.org/10.1016/S0140-6736\(16\)32517-X](https://doi.org/10.1016/S0140-6736(16)32517-X).
- Roengvoraphoj, O., Wijaya, C., Eze, C., Li, M., Dantes, M., Taugner, J., et al., 2018. Analysis of primary tumor metabolic volume during chemoradiotherapy in locally advanced non-small cell lung cancer. *Strahlenther. Onkol.* 194 (2), 107–115. <https://doi.org/10.1007/s00066-017-1229-3>.
- Rossi, A., Maio, M., Di, Chio dini, P., Rudd, R.M., Okamoto, H., Skarlos, D.V., et al., 2012. Carboplatin- or cisplatin-based chemotherapy in first-line treatment of small-cell lung cancer: the COCIS meta-analysis of in individual patient data. *J. Clin. Oncol.* 30 (14), 1692–1698. <https://doi.org/10.1200/JCO.2011.40.4905>.
- Rossi, A., Chiodini, P., Sun, J.M., O'Brien, M.E., von Plessen, C., Barata, F., et al., 2014. Six versus fewer planned cycles of first-line platinum-based chemotherapy for non-small-cell lung cancer: a systematic review and meta-analysis of individual patient data. *Lancet Oncol. Elsevier Ltd* 15 (11), 1254–1262. <https://doi.org/10.1371/journal.pone.0037229>.
- Rossi, G., Jocolli, G., Conti, A., Tiseo, M., Zito, M.F., Donati, G., et al., 2017. Detection of ROS1 rearrangement in non-small cell lung cancer: current and future perspectives. *Lung Cancer (Auckland, NZ)*. Dove Press 8, 45–55. <https://doi.org/10.2147/LCCT.S120172>.
- Rothenberger, N., Somasundaram, A., Stabile, L., 2018. The role of the estrogen pathway in the tumor microenvironment. *Int. J. Mol. Sci.* 19 (February (2)), 611. <https://doi.org/10.3390/ijms19020611>.
- Rudin, C.M., Ismaila, N., Hann, C.L., Malhotra, N., Movsas, B., Norris, K., et al., 2015. Treatment of small-cell lung cancer: american society of clinical oncology endorsement of the American college of chest physicians guideline. *J. Clin. Oncol.* 33 (34), 4106–4111. <https://doi.org/10.1200/JCO.2015.63.7918>.
- Saito, H., Takada, Y., Ichinose, Y., Eguchi, K., Kudoh, S., Matsui, K., et al., 2006. Phase II study of etoposide and cisplatin with concurrent twice-daily thoracic radiotherapy followed by irinotecan and cisplatin in patients with limited-disease small-cell lung cancer: west Japan Thoracic Oncology Group 9902. *J. Clin. Oncol.* 24 (33), 5247–5252. <https://doi.org/10.1200/JCO.2006.07.1605>.
- Sandler, A., Gray, R., Perry, M.C., Brahmer, J., Schiller, J.H., Dowlati, A., et al., 2006. Paclitaxel–Carboplatin alone or with Bevacizumab for non-Small-Cell lung Cancer. *N. Engl. J. Med.* 355 (24), 2542–2550. <https://doi.org/10.1056/NEJMoa061884>.
- Schiller, J.H., Harrington, D., Belani, C.P., Langer, C., Sandler, A., Krook, J., et al., 2002. Comparison of four chemotherapy regimens for advanced non-small-cell lung cancer. *N. Engl. J. Med.* 346 (2), 92–98. <https://doi.org/10.1056/NEJMoa011954>.
- Schwartz, A.G., Prysak, G.M., Murphy, V., Lonardo, F., Pass, H., Schwartz, J., et al., 2005. Nuclear estrogen receptor in lung Cancer: expression and survival differences by sex. *Clin. Cancer Res.* 11 (October (20)), 7280–7287. <https://doi.org/10.1158/1078-0432>.
- Seidegård, J., Pero, R.W., Markowitz, M.M., Roush, G., Miller, D.G., Beattie, E.J., 1990. Isoenzyme(s) of glutathione transferase (class Mu) as a marker for the susceptibility to lung cancer: a follow up study. *Carcinogenesis* 11 (January (1)), 33–36.
- Sharpe, A.H., Wherry, E.J., Ahmed, R., Freeman, G.J., 2007. The function of programmed cell death 1 and its ligands in regulating autoimmunity and infection. *Nat Immunol. Nature Publishing Group* 8 (March (3)), 239–245. <https://doi.org/10.1038/ni1443>.
- Shaw, A.T., Yeap, B.Y., Mino-Kenudson, M., Digumarthy, S.R., Costa, D.B., Heist, R.S., et al., 2009. Clinical features and outcome of patients with non-small-cell lung cancer who harbor EML4-ALK. *J. Clin. Oncol. American Society of Clinical Oncology* 27 (September (26)), 4247–4253. <https://doi.org/10.1200/JCO.2009.22.6993>.
- Shepherd, F.A., Evans, W.K., Feld, R., Young, V., Patterson, A., Ginsberg, R., et al., 1988. Adjuvant chemotherapy following surgical resection for small-cell carcinoma of the lung. *J. Clin. Oncol.* 6 (5), 832–838. <https://doi.org/10.1200/JCO.1988.6.5.832>.
- Shepherd, F.A., Dancy, J., Ramlau, R., Mattson, K., Gralla, R., O'Rourke, M., et al., 2000. Prospective Randomized Trial of Docetaxel Versus Best Supportive Care in Patients With Non-Small-Cell Lung Cancer Previously Treated With Platinum-Based Chemotherapy. *J. Clin. Oncol.* 18 (10), 2095–2103. <https://doi.org/10.1200/JCO.2000.18.10.2095>.
- Siddiqui, F., Bae, K., Langer, C.J., Coyne, J.C., Gamerman, V., Komaki, R., et al., 2010. The influence of gender, race, and marital status on survival in lung Cancer patients: analysis of radiation therapy oncology group trials. *J. Thorac Oncol. Elsevier* 5 (May (5)), 631–639. <https://doi.org/10.1097/JTO.0b013e3181d5e46a>.
- Siegel, R.L., Miller, K.D., Jemal, A., 2019. Cancer statistics, 2019. *CA Cancer J. Clin.* 69 (1), 7–34. <https://doi.org/10.3322/caac.21551>.
- Singh, S., Parulekar, W., Murray, N., Feld, R., Evans, W.K., Tu, D., et al., 2005. Influence of sex on toxicity and treatment outcome in small-cell lung Cancer. *J. Clin. Oncol.* 23 (February (4)), 850–856. <https://doi.org/10.1200/JCO.2005.03.171>.
- Skarlos, D.V., Samantas, E., Briassoulis, E., Panoussaki, E., Pavlidis, N., Kalofonos, H.P., et al., 2001. Randomized comparison of early versus late hyperfractionated thoracic

- irradiation concurrently with chemotherapy in limited disease small-cell lung cancer. *Ann. Oncol.* 12 (9), 1231–1238. <https://doi.org/10.1023/A:1012295131640>.
- Socinski, M.A., Jotte, R.M., Cappuzzo, F., Orlandi, F., Stroyakovskiy, D., Nogami, N., et al., 2018. Atezolizumab for first-line treatment of metastatic nonsquamous NSCLC. *N. Engl. J. Med.* 378 (24), 2288–2301. <https://doi.org/10.1056/NEJMoa1716948>.
- Soria, J.C., Mauguen, A., Reck, M., Sandler, A.B., Saijo, N., Johnson, D.H., et al., 2013. Systematic review and meta-analysis of randomised, phase II/III trials adding bevacizumab to platinum-based chemotherapy as first-line treatment in patients with advanced non-small-cell lung cancer. *Ann. Oncol.* 24 (1), 20–30. <https://doi.org/10.1093/annonc/nds590>.
- Spigel, D., Betzig Schrock, A., Fabrizio, D., Frampton, G., 2019. Total Mutation Burden (TMB) in Lung Cancer (LC) and Relationship With Response to PD-1/PD-L1 Targeted Therapies. https://doi.org/10.1200/JCO.2016.34.15_suppl.9017.
- Spitz, M.R., Wu, X., Wang, Y., Wang, L.E., Shete, S., Amos, C.I., et al., 2001. Modulation of nucleotide excision repair capacity by XPD polymorphisms in lung cancer patients. *Cancer Res.* 61 (4), 1–2.
- Stabile, L.P., Lyker, J.S., Gubish, C.T., Zhang, W., Grandis, J.R., Siegfried, J.M., 2005. Combined targeting of the estrogen receptor and the epidermal growth factor receptor in non-small cell lung Cancer Shows enhanced antiproliferative effects. *Cancer Res.* 65 (February (4)), 1459–1470. <https://doi.org/10.1158/0008-5472.CAN-04-1872>.
- Stewart, L.A., 1995. Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials Introduction Worldwide more than half a million new cases of. *Bmj* 311 (October), 899–909.
- Strauss, G.M., Herndon, J.E., Maddaus, M.A., Johnstone, D.W., Johnson, E.A., Harpole, D.H., et al., 2008. Adjuvant paclitaxel plus carboplatin compared with observation in stage IB non-small-cell lung cancer: CALGB 9633 with the cancer and leukemia group B, radiation therapy oncology group, and North Central cancer treatment group study groups. *J. Clin. Oncol.* 26 (31), 5043–5051. <https://doi.org/10.1200/JCO.2008.16.4855>.
- Subramanian, J., Govindan, R., 2007. Lung cancer in never smokers: a review. *J. Clin. Oncol.* 25 (February (5)), 561–570. <https://doi.org/10.1200/JCO.2006.06.8015>.
- Sun, S., Schiller, J.H., Gazdar, A.F., 2007. Lung cancer in never smokers — a different disease. *Nat. Rev. Cancer* 7 (October (10)), 778–790. <https://doi.org/10.1038/nrc2190>.
- Thomas, L., Doyle, L.A., Edelman, M.J., Martin, J., 2005. Lung Cancer in women. *Chest* 128 (July (1)), 370–381. <https://doi.org/10.1378/chest.128.1.370>.
- Thomsen, J.S., Wang, X., Hines, R.N., Safe, S., 1994. Restoration of aryl hydrocarbon (Ah) responsiveness in MDA-MB-231 human breast cancer cells by transient expression of the estrogen receptor. *Carcinogenesis* 15 (May (5)), 933–937.
- Toyooka, S., Tsuda, T., Gazdar, A.F., 2003. The TP53 gene, tobacco exposure, and lung cancer. *Hum. Mutat.* 21 (March (3)), 229–239. <https://doi.org/10.1002/humu.10177>.
- Travis, W.D., Brambilla, E., Nicholson, A.G., Yatabe, Y., Austin, J.H.M., Beasley, M.B., et al., 2015a. The 2015 world health organization classification of lung tumors: impact of genetic, clinical and radiologic advances since the 2004 classification. *J. Thorac. Oncol. International Association for the Study of Lung Cancer* 10 (9), 1243–1260. <https://doi.org/10.1097/JTO.0000000000000630>.
- Travis, W.D., Brambilla, E., Nicholson, A.G., Yatabe, Y., Austin, J.H.M., Beasley, M.B., et al., 2015b. The 2015 world health organization classification of lung tumors: impact of genetic, clinical and radiologic advances since the 2004 classification. *J. Thorac. Oncol. International Association for the Study of Lung Cancer* 10 (9), 1243–1260. <https://doi.org/10.1097/JTO.0000000000000630>.
- Traynor, A.M., Schiller, J.H., Stabile, L.P., Kolesar, J.M., Eickhoff, J.C., Dacic, S., et al., 2009. Pilot study of gefitinib and fulvestrant in the treatment of post-menopausal women with advanced non-small cell lung cancer. *Lung Cancer* 64 (April (1)), 51–59. <https://doi.org/10.1016/j.lungcan.2008.07.002>.
- Trigunaite, A., Dimo, J., Jørgensen, T.N., 2015. Suppressive effects of androgens on the immune system. *Cell Immunol. Elsevier Inc.* 294 (2), 87–94. <https://doi.org/10.1016/j.cellimm.2015.02.004>.
- Tsuchiya, R., Suzuki, K., Ichinose, Y., Watanabe, Y., Yasumitsu, T., Ishizuka, N., et al., 2005. Phase II trial of postoperative adjuvant cisplatin and etoposide in patients with completely resected stage I-IIIa small cell lung cancer: the Japan Clinical Oncology Lung Cancer Study Group Trial (JCOG9101). *J. Thorac. Cardiovasc. Surg.* 129 (5), 977–983. <https://doi.org/10.1016/j.jtcvs.2004.05.030>.
- Turrisi, A.T., Kim, K., Blum, R., Sause, W.T., Livingston, R.B., Komaki, R., et al., 1999. Twice-daily compared with once-daily thoracic radiotherapy in limited small-cell lung cancer Treated concurrently with cisplatin and etoposide. *N. Engl. J. Med.* 340 (4), 265–271.
- Vähäkangas, K.H., Bennett, W.P., Castrén, K., Welsh, J.A., Khan, M.A., Blömeke, B., et al., 2001. p53 and K-ras mutations in lung cancers from former and never-smoking women. *Cancer Res.* 61 (June (11)), 4350–4356.
- Vidal, J., Clavé, S., De, Muga S., González, I., Pijuan, L., Gimeno, J., et al., 2014. Assessment of ALK status by FISH on 1000 Spanish non-small cell lung cancer patients. *J. Thorac. Oncol.* 9 (12), 1816–1820. <https://doi.org/10.1097/JTO.0000000000000361>.
- Videtic, G.M.M., Truong, P.T., Ash, R.B., Yu EW, Kocha W.I., Vincent, M.D., et al., 2005. Does sex influence the impact that smoking, treatment interruption and impaired pulmonary function have on outcomes in limited stage small cell lung cancer treatment? *Can. Respir. J.* 12 (5), 245–250. <https://doi.org/10.1155/2005/376404>.
- Vinolas, N., Isla, D., Felip, E., Garrido, P., Garcia-Campeloe, R., Lianes, P., et al., 2007. P2-106: Gender differences in non-small cell lung cancer (NSCLC) patients (p): A retrospective study based in Spanish Lung Cancer Group (SLCG) trials. *J. Thorac. Oncol. Elsevier* 2 (August (8)), S533–S534. <https://doi.org/10.1097/01.JTO.0000283570.10829.2d>.
- Von, Pawel J., Schiller, J.H., Shepherd, F.A., Fields, S.Z., Kleisbauer, J.P., Chrysson, N.G., et al., 1999. Topotecan versus cyclophosphamide, doxorubicin, and vincristine for the treatment of recurrent small-cell lung cancer. *J. Clin. Oncol.* 17 (2), 658–667. <https://doi.org/10.1200/JCO.1999.17.2.658>.
- Wakelee, H.A., Wang, W., Schiller, J.H., Langer, C.J., Sandler, A.B., Belani, C.P., et al., 2006. Survival differences by sex for patients with advanced non-small cell lung Cancer on eastern cooperative oncology group trial 1594. *J. Thorac. Oncol. Elsevier* 1 (June (5)), 441–446. [https://doi.org/10.1016/S1556-0864\(15\)31609-9](https://doi.org/10.1016/S1556-0864(15)31609-9).
- Wakelee, H.A., Dahlberg, S.E., Brahmer, J.R., Schiller, J.H., Perry, M.C., Langer, C.J., et al., 2012. Differential effect of age on survival in advanced NSCLC in women versus men: analysis of recent Eastern Cooperative Oncology Group (ECOG) studies, with and without bevacizumab. *Lung Cancer. Elsevier Ireland Ltd* 76 (3), 410–415. <https://doi.org/10.1016/j.lungcan.2011.12.006>.
- Wakelee, H., Dahlberg, S., Keller, S., Tester, W., Gandara, D., 2016. E1505: adjuvant chemotherapy +/- bevacizumab for early stage NSCLC—outcomes based on chemotherapy subsets. *J. Clin. Oncol.* 34 (15). https://doi.org/10.1200/JCO.2016.34.15_suppl.8507.
- Wallis, C.J.D., Butaney, M., Satkunasivam, R., Freedland, S.J., Patel, S.P., Hamid, O., et al., 2019. Association of Patient Sex With Efficacy of Immune Checkpoint Inhibitors and Overall Survival in Advanced Cancers A Systematic Review and Meta-analysis. *JAMA Oncol.* <https://doi.org/10.1001/jamaoncol.2018.5904>.
- Wang, T.W., Asman, K., Gentzke, A.S., Cullen, K.A., Holder-Hayes, E., Reyes-Guzman, C., et al., 2018. Morbidity and mortality weekly report centers for disease control and prevention. *MMWR* 67 (November (44)).
- Warde, P., Payne, D., 1992. Does thoracic irradiation improve survival and local control in limited-stage small-cell carcinoma of the lung? A meta-analysis. *J. Clin. Oncol.* 10 (6), 890–895. <https://doi.org/10.1200/JCO.1992.10.6.890>.
- Warth, A., Muley, T., Dienemann, H., Goepfert, B., Stenzinger, A., Schnabel, P.A., et al., 2014. ROS1 expression and translocations in non-small-cell lung cancer: clinicopathological analysis of 1478 cases. *Histopathology* 65 (August (2)), 187–194. <https://doi.org/10.1111/his.12379>.
- Watanabe, K., Sakamaki, K., Nishii, T., Yamamoto, T., Maehara, T., Nakayama, H., et al., 2018. Gender differences in the recurrence timing of patients undergoing resection for non-small cell lung Cancer. *Asian Pac. J. Cancer Prev. Shahid Beheshti University of Medical Sciences* 19 (Mach (3)), 719–724. <https://doi.org/10.22034/APJCP.2018.19.3.719>.
- Wei, Q., Cheng, L., Amos, C.I., Wang, L.E., Guo, Z., Hong, W.K., et al., 2000. Repair of tobacco carcinogen-induced DNA adducts and lung cancer risk: a molecular epidemiologic study. *J. Natl. Cancer Inst.* 92 (November (21)), 1764–1772.
- Weiderpass, E., Soerjomataram, I., Rutherford, M., Ferlay, J., Bray, F., Lortet-Tieulent, J., 2014. International trends in lung cancer incidence by histological subtype: adenocarcinoma stabilizing in men but still increasing in women. *Lung Cancer. Elsevier Ireland Ltd* 84 (1), 13–22. <https://doi.org/10.1016/j.lungcan.2014.01.009>.
- Wheatley-Pence, P., Blackhall, F., Lee, S.M., Ma, C., Ashcroft, L., Jital, M., et al., 2010. The influence of sex and histology on outcomes in non-small-cell lung cancer: a pooled analysis of five randomized trials. *Ann. Oncol.* 21 (10), 2023–2028. <https://doi.org/10.1093/annonc/mdq067>.
- Wolf, M., Holle, R., Hans, K., Drings, P., Havemann, K., 1991. Analysis of prognostic factors in 766 patients with small cell lung cancer (SCLC): the role of sex as a predictor for survival. *Br. J. Cancer* 63 (6), 986–992.
- Yang, L., Huang, F., Mei, J., Wang, X., Zhang, Q., Wang, H., et al., 2017. Posttranscriptional control of PD-L1 expression by 17β-Estradiol via PI3K/Akt signaling pathway in ERα-Positive Cancer cell lines. *Int. J. Gynecol. Cancer* 27 (February (2)), 196–205. <https://doi.org/10.1097/IGC.0000000000000875>.
- Zatloukal, P., Cardenal, F., Szczesna, A., Gorbunova, V., Moiseyenko, V., Zhang, X., et al., 2010. A multicenter international randomized phase III study comparing cisplatin in combination with irinotecan or etoposide in previously untreated small-cell lung cancer patients with extensive disease. *Ann. Oncol.* 21 (9), 1810–1816. <https://doi.org/10.1093/annonc/mdq036>.
- Zhang, Y., Kang, S., Fang, W., Hong, S., Liang, W., Yan, Y., et al., 2015. Impact of smoking status on EGFR-TKI efficacy for advanced non-small-cell lung cancer in EGFR mutants: a meta-analysis. *Clin. Lung Cancer. Elsevier* 16 (March (2)), 144–151. <https://doi.org/10.1016/j.clc.2014.09.008>.
- Zhang, Y.L., Yuan, J.Q., Wang, K.F., Fu, X.H., Han, X.R., Threapleton, D., et al., 2016. The prevalence of EGFR mutation in patients with non-small cell lung cancer: a systematic review and meta-analysis. *Oncotarget* 7 (48), 9–13. <https://doi.org/10.18632/oncotarget.12587>.
- Zukin, M., Barrios, C.H., Pereira, J.R., Ribeiro Rde, A., Beato, C.A., Nascimento, Y.N., et al., 2013. Randomized phase III trial of single-agent pemetrexed versus carboplatin and pemetrexed in patients with advanced non-small-cell lung cancer and Eastern Cooperative Oncology Group performance status of 2. *J. Clin. Oncol.* 31 (23), 2849–2853. <https://doi.org/10.1200/JCO.2012.48.1911>.

Stefano Frega is a Medical Oncologist at the Division of Medical Oncology 2 at the Veneto Institute of Oncology (IOV) in Padua. He has dedicated his professional activity to treating patients with thoracic tumors, and is also involved as co-investigator in many phase II-III clinical trials principally involving lung cancer and mesothelioma patients. He spent a period abroad at the Oncology Department of Center Hospital University Vaudois (CHUV), Lausanne, Switzerland, directed by Prof. George Coukos, working on a translational project at the Immuno-Oncology Department, and continuing to work as physician at the Service of Medical Oncology directed by Prof. Solange Peters, to gain further knowledge on thoracic malignancies. He published articles on thoracic malignancies topic in peer-reviewed journals.

Alessandro Dal Maso is a resident in Medical Oncology at the Department of Surgery, Oncology and Gastroenterology of the University of Padua, and currently working at

Medical Oncology 2, Veneto Institute of Oncology IOV – IRCCS, Padua. Since 2015, he is committed to medical treatment and clinical research in the field of solid tumors, focusing on thoracic neoplasms. He is involved as sub-investigator in several phase II and III clinical trials on Non-Small Cell Lung Cancer, Small Cell Lung Cancer and Malignant Pleural Mesothelioma.

Alessandra Ferro is a resident in Medical Oncology at the Department of Surgery, Oncology and Gastroenterology of the University of Padua. Since November 2016, she has been committed to the medical treatment and clinical research of solid tumors, focusing on thoracic neoplasms. She has been involved as co-investigator in several phase II-III clinical trials on Non-Small Cell Lung Cancer (NSCLC), Small Cell Lung Cancer and Malignant Pleural Mesothelioma. Currently she is working at the Department of Medical Oncology 2, receiving training in clinical practice.

Laura Bonanno is a Medical Oncologist at the Division of Medical Oncology 2 at the Veneto Institute of Oncology (IOV) in Padua. She graduated with full marks in 2006 and specialized in Medical Oncology with full marks in 2011. Since 2007, she has been actively working at the IOV, involved in clinical and research activity. She has been involved in many clinical trials (phase II-II), both spontaneous and good clinical practice trials. She also has molecular biology expertise, with particular interest in the field of translational research. During her residency, she also worked at the Dexeus University Institute of Barcelona, under the supervision of Dr. Rosell, director of the Catalan Institute of Oncology, where she was head of a translational research project. She has published articles in peer-reviewed journals and she has written e-book chapters in the field of thoracic oncology.

PierFranco Conte is Full Professor of Oncology at the University of Padova and Director

of the Division of Medical Oncology 2 at the Veneto Institute of Oncology (IOV) in Padua. He is specialized in Clinical Oncology, Clinical Immunology and Hematology. Throughout his career, Prof. Conte has been involved in the development of new regimens for the treatment of solid tumors, with special emphasis on breast and ovarian cancer. He was Principal Investigator of numerous research projects supported by the National Research Council (CNR), the Italian Association for Cancer Research (AIRC), the Ministry of Health and the Ministry of Education and Research. He has published more than 300 papers in peer-reviewed journals. The majority of these discuss the biological characterization of breast and ovarian cancer and clinical trials of these disorders. Prof Conte is the recipient of the 2007 Claude Jacquillat Award for Achievement in Clinical Oncology.

Giulia Pasello is a Medical Oncologist at the Division of Medical Oncology 2 at the Veneto Institute of Oncology (IOV) in Padua. She specialized in Medical Oncology in 2010 and received her PhD in Clinical and Experimental Sciences in 2015. Dr. Pasello is committed to medical treatment and clinical research of solid tumours, with particular interest in thoracic neoplasms. She is Coinvestigator and Principal Investigator in more than 40 clinical trials on Non-Small Cell Lung Cancer (NSCLC), Small Cell Lung Cancer (SCLC), Malignant Pleural Mesothelioma and Thymic cancer. She is currently involved as Principal Investigator in basic and translational research projects on rare thoracic cancers, particularly on Malignant Pleural Mesothelioma and neuroendocrine tumors. She is author of about 40 papers in peer-reviewed journals, the majority on clinical and molecular features of thoracic cancers. She was the recipient of the 2010 European Society for Medical Oncology Translational Research Fellowship Award for her work on the Identification of novel molecular targets and biological agents for the chemotherapy of Malignant Pleural Mesothelioma. In 2015, she was recipient of My First AIRC Grant for the translational project MET-targeting in Small Cell Lung Cancer, currently ongoing. She is a reviewer for several peer reviewed journals focused on thoracic oncology.