



Correction to: Diffusion-weighted imaging for assessment of synovial inflammation in juvenile idiopathic arthritis: a promising imaging biomarker as an alternative to gadolinium-based contrast agents

Anouk M. Barendregt^{1,2} · E. Charlotte van Gulik^{1,2} · Cristina Lavini¹ · Charlotte M. Nusman¹ · J. Merlijn van den Berg² · Dieneke Schonenberg-Meinema² · Koert M. Dolman^{3,4,5} · Taco W. Kuijpers² · Robert Hemke¹ · Mario Maas¹

Published online: 30 November 2018
© The Author(s) 2018

Correction to: Eur Radiol (2017) 27:4889–4899

<https://doi.org/10.1007/s00330-017-4876-y>

The original version of this article, published on 12 June 2017, unfortunately contained a wrong y-axis in Figure 4. The following correction has therefore been made in the original: the y-axis of parameter ADC is corrected. The corrected figure is given below.

The online version of the original article can be found at <https://doi.org/10.1007/s00330-017-4876-y>

✉ Anouk M. Barendregt
a.m.barendregt@amc.uva.nl

¹ Department of Radiology, Academic Medical Center/University of Amsterdam, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

² Department of Pediatric Hematology, Immunology, Rheumatology and Infectious Disease, Emma Children's Hospital, Academic Medical Center/University of Amsterdam, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

³ Department of Pediatric Rheumatology, Reade, Dr. Jan van Breemenstraat 2, Amsterdam, The Netherlands

⁴ Department of Pediatric Rheumatology, Onze Lieve Vrouwe Gasthuis West, Jan Tooropstraat 164, Amsterdam, The Netherlands

⁵ Department of Pediatric Rheumatology, Onze Lieve Vrouwe Gasthuis Oost, Oosterpark 9, Amsterdam, The Netherlands

DCE and DWI parameters differentiating inactive from active disease




Fig. 4 Boxplots of dynamic contrast-enhanced (DCE) and diffusion-weighted imaging (DWI) parameters that were significantly different in MRI-inactive JIA when compared to MRI-active JIA. The *y*-axis displays a DCE or DWI parameter (ME, MIS, TTP, TIC5 or ADC). The *x*-axis is

divided in two: first, MRI-inactive patients are displayed (*filled circles*); second patients with MRI-active JIA indicated by a *grey, open square*. * = *p*-value < 0.05