



Longitudinal Effect of Ambient Air Pollution and Pollen Exposure on Asthma Control: The Patient-Reported Outcomes Measurement Information System (PROMIS) Pediatric Asthma Study

Zheng Li, PhD, MPH; Xiaohui Xu, MD, MPH, PhD; Lindsay A. Thompson, MD, MS; Heather E. Gross, MEd; Elizabeth A. Shenkman, PhD; Darren A. DeWalt, MD, MPH; I-Chan Huang, PhD

From the College of Nursing and Health Professions, Valparaiso University, Valparaiso, Ind (Z Li); Department of Epidemiology and Statistics, School of Public Health, Texas A&M Health Science Center, College Station (X Xu); Departments of Pediatrics (LA Thompson); Health Outcomes & Biomedical Informatics (EA Shenkman), College of Medicine, University of Florida, Gainesville; Cecil G. Sheps Center for Health Services Research (HE Gross); Department of Medicine, School of Medicine (DA DeWalt), University of North Carolina at Chapel Hill; and Department of Epidemiology and Cancer Control, St Jude Children's Research Hospital, Memphis, Tenn (I-C Huang)
At the time this study was conducted, Dr Darren DeWalt was an unpaid member of the Board of Directors for the PROMIS Health Organization. Dr DeWalt is also an author of some of the items in the PROMIS instruments and owns the copyright for these items. Dr DeWalt has given an unlimited free license for the use of the materials to the PROMIS Health Organization. All other co-authors declare no conflicts of interest. No honorarium or other form of payment was given to anyone to produce the manuscript.
Address correspondence to I-Chan Huang, PhD, Department of Epidemiology and Cancer Control, St Jude Children's Research Hospital, 262 Danny Thomas Place, Mail Stop # 735, Memphis, TN 38105 (e-mail: i-chan.huang@stjude.org).
Received for publication May 29, 2018; accepted March 31, 2019.

ABSTRACT

OBJECTIVE: Although exposure to air pollution and pollen is associated with asthma exacerbation and increased health care use, longitudinal effects of fine particulate matter 2.5 (PM_{2.5}), ozone (O₃), and pollen exposure on asthma control status in pediatric patients are understudied. This study investigated effects of exposure to PM_{2.5}, O₃, and pollen on asthma control status among pediatric patients with asthma.

METHODS: A total of 229 dyads of pediatric patients with asthma and their parents were followed for 15 months. The Asthma Control and Communication Instrument was used to measure asthma control, which was reported weekly by parents during a 26-week period. PM_{2.5} and O₃ data were collected from the US Environmental Protection Agency Air Quality System. Pollen data were obtained from Intercontinental Marketing Services Health. Mean air pollutant and pollen exposures within 7 days before the reporting of asthma control were used to estimate weekly exposures for each participant. Linear mixed-effects models were performed to test associations of

PM_{2.5}, O₃, and pollen exposure with asthma control status. Sensitivity analyses were performed to evaluate the robustness of findings by different exposure monitoring days per week and distances between monitoring sites and participants' residences.

RESULTS: Elevated PM_{2.5} concentration and pollen severity were associated with poorer asthma control status ($P < .05$), yet elevated O₃ concentration was marginally associated with better asthma control ($P < .1$).

CONCLUSIONS: Poorer asthma control status was associated with elevated PM_{2.5} and pollen severity. Reducing harmful outdoor environmental ambient exposure may improve asthma outcomes in children and adolescents.

KEYWORDS: air pollution; asthma control; children; longitudinal study; patient-reported outcomes; pollen

ACADEMIC PEDIATRICS 2019;19:615–623

WHAT'S NEW

We demonstrate that elevated PM_{2.5} concentration and pollen severity were significantly associated with poorer asthma control in children and adolescents. Effective strategies to prevent individuals from unnecessary exposure to harmful air pollution and pollen are warranted for improving asthma outcomes.

APPROXIMATELY 7 MILLION American children and adolescents younger than 18 years of age lived with asthma in 2012.¹ Respiratory and immune systems of

children are vulnerable to the exposure of air pollutants, including nitrogen dioxide (NO₂), ozone (O₃), particulate matter (PM), carbon dioxide (CO₂), and sulfur dioxide (SO₂).² Increased exposure to NO₂ and PM is linked to early asthma incidence,^{3–5} deteriorated lung functioning,⁶ and substantial economic costs related to asthma health care^{7,8} in pediatric populations.

Although previous studies frequently have examined the relationship between air pollution exposure and health care use, fewer studies have been focused on patient-centered outcomes such as asthma control and daily functioning.⁹ Asthma control status is an important indication of

asthma management because it suggests immediate, direct impacts of environmental exposures before hospitalization and adverse health events. Increased exposure to PM₁₀ has been associated with poorly controlled asthma status, functional status, and quality of life with elevated exposure to O₃ and SO₂ associated with decreased lung function.¹⁰ Others found exposure to elevated PM_{2.5} and O₃ to be associated with poorly controlled asthma.¹¹

In addition to air pollutants, greater pollen levels in grass, trees, and weeds may contribute to increased asthma symptoms (eg, wheezing, coughing, and chest tightness),¹² emergency department visits, and hospitalizations.¹³⁻¹⁵ However, limited studies have examined the effects of both air pollution and pollen on asthmatic outcomes. Some studies show that greater pollen counts rather than air pollution led to more asthma-related emergency department visits.^{14,15} Other studies indicate that ambient air pollution concentrations instead of pollen counts caused pediatric asthma hospitalization¹⁶ or that increased exposure to both air pollution and pollen is associated with asthma-related health care use.¹⁷ In areas with warmer temperatures (eg, Florida), numerous plants release botanic aeroallergens that are triggers of respiratory diseases.¹⁸ Therefore, it is important to examine the effects of both air pollution and pollen exposure on asthma control status by comparing relative impacts in pediatric populations.

This study aimed to test longitudinal associations of air pollution (PM_{2.5} and O₃) and pollen exposure with the change of asthma control status in children/adolescents over a 26-week period by using data collected from the Patient-Reported Outcomes Measurement Information System (PROMIS) Pediatric Asthma Study (PAS). PROMIS PAS evaluated longitudinal validity, including responsiveness to change and minimally important differences, of the PROMIS Pediatric measures for children with asthma ([Supplementary Methods](#)). We hypothesized that increased exposure to PM_{2.5}, O₃, and pollen was associated with poorer asthma control status.

METHODS

POPULATION AND DATA COLLECTION

Study participants were identified from the claims and enrollment files of the Florida Medicaid and the State Children's Health Insurance Program (SCHIP). A total of 1450 pediatric patients with asthma were identified from the database of the Florida Medicaid and SCHIP. The enrollment criteria included being aged 8 to 17.9 years and having parents aged 18 years or older; being continuously enrolled (≥ 6 months) in Florida Medicaid and SCHIP; having a diagnosis of asthma with *International Classification of Diseases, Ninth Revision, Clinical Modification* codes 493.1 (asthma with status asthmaticus), 493.2 (asthma with acute exacerbation), or 493.x listed in claim and enrollment files; having at least 2 asthma-related health care visits during the past 12 months; and having access to Internet and a telephone within the past 6 weeks.

We contacted parents of patients via telephone calls. However, 440 (30%) dyads of patients and parents did not meet the enrollment criteria; 684 (47%) refused to participate, and 326 (23%) verbally agreed to participate in the study. Eventually, 238 dyads consented/assented to study participation, and 26 withdrew by the end of data collection (attrition rate: 11%). According to claims and enrollment files, the child's age and sex were not statistically different between participants and nonparticipants ($P > .05$). In this study, 229 dyads who completed longitudinal data, including asthma control ([Supplementary Methods](#)), were analyzed. The institutional review board at University of Florida approved the study protocol; informed consent for study participation was obtained from parents, and assent was obtained from children/youths.

ASTHMA OUTCOME MEASUREMENT

PROMIS PAS collected data during a 15-month period, and asthma control status information was collected during a 13-week block in the first year and another 13-week block in the second year (26 weeks total). In each week, parents were asked to report their child's asthma control status through our research website. Asthma control status was measured by using the Asthma Control and Communication Instrument (ACCI) with permission from the developer.¹⁹ We slightly modified the ACCI to be content-appropriate for children/adolescents ([Supplemental Survey Instrument](#)). The modified ACCI is in concordance with the pediatric ACCI¹⁹ that was developed after the launch of the PROMIS PAS. We used parent/proxy reports rather than child self-reports to collect asthma control data because our previous studies suggest that the ACCI probes asthma medications and flares, of which parents may be more aware, particularly in cases of younger children.²⁰ ACCI comprises 11 items, including 5 items measuring concepts related to symptoms, use of rescue medicine, asthma attacks, activity limitations, and sleep disturbance due to asthma.¹⁹ The scores of these 5 individual items were summed to represent the overall asthma control status for individuals and ranged from 0 to 19: greater scores indicate worse asthma control status. ACCI has demonstrated satisfactory measurement properties,¹⁹ including adequate concurrent validity associated with other asthma control measures and known-groups validity associated with peak expiratory flow rate.¹⁹

EXPOSURE ASSESSMENT

Air quality data of PM_{2.5} and O₃ were obtained from the U.S. Environmental Protection Agency (EPA) Air Quality System (AQS) between 2010 and 2012 from all counties of Florida.²¹ EPA AQS data contain concentrations of air pollutants, dates of measurements, and locations of specific monitoring sites by latitude and longitude. During the study period, PM_{2.5} and O₃ concentration data were collected from 52 and 59 active monitoring sites in Florida, respectively ([Figure](#)). The 2010 US Census Bureau's 5-digit ZIP code tabulation areas²² were used to estimate the centroid of the ZIP codes in which the study participants

resided. The centroids of ZIP codes were linked to the nearest EPA monitoring sites based on the corresponding latitude and longitude. The distance between each centroid of ZIP codes to air quality monitoring sites was calculated by using ArcGIS software (Version 10.2; Esri, Redlands, Calif). The shortest distance between monitoring sites and the participant's location was used to assign an EPA monitor for generating air pollution data. PM_{2.5} and O₃ exposures were calculated as the mean concentrations within a 7-day window before the reporting of asthma control.

Pollen information was provided by Intercontinental Marketing Services (IMS) Health.²³ Technically, IMS uses the Rotorod Sampler to collect pollen data. First, a rod covered with a sticky substance is attached to the roof of a building. For 24 hours, the rotating rod is tested periodically at different times of the day to determine the amount of adhered pollen. Samples are then analyzed microscopically to determine the count of all types of pollen (eg, grass, tree, etc) in grains per cubic meter in the air over a 24-hour period, and the pollen severity index is subsequently generated. The pollen severity index is a continuous variable ranging from 0 to 12, reflecting the overall severity of pollen exposure. Greater indices indicate greater levels of pollen severity. IMS Health collected the pollen index in nine geographic areas of Florida (Figure). Study participants were linked to IMS Health pollen data on the basis of the ZIP code in which the participants resided. Pollen exposure was calculated as the mean pollen severity index within a 7-day window before the reporting of asthma control.

CONTRIBUTING VARIABLES

Environmental factors and sociodemographic information that may confound the results were collected. Two cohorts were recruited from 2 different seasons to capture the variation of asthma-flare seasons and climatic characteristics in Florida. In the first cohort (N = 135), data were collected in the fall season (from September to December) of 2010 and 2011. In the second cohort (N = 94), data were collected in the spring season (from February to May) of 2011 and 2012. The same weeks and months were applied to the first and second years for the 2 respective cohorts. These months covered the period of school years in Florida. For each cohort, participants' sociodemographic characteristics were collected at baseline of the first year, including children's age, sex, race/ethnicity, and type of chronic conditions and parents' age, race/ethnicity, marital status, and educational background. Household smoking status was self-reported by parents using a binary question (yes/no). Climatic data (daily temperatures and precipitation) of each county in Florida during the study period were obtained from the National Climatic Data Center at the National Oceanic and Atmospheric Administration.²⁴

STATISTICAL ANALYSES

All analyses were performed by using SAS V.9.3 (SAS Institute, Cary, NC). Mean, standard deviation (SD), range,

and interquartile range (IQR) were estimated to characterize the exposure of air pollutants (PM_{2.5} and O₃), the pollen severity index, temperature, and precipitation. PM_{2.5}, O₃, and pollen exposures were measured as the mean concentrations within a 7-day window before the reporting of asthma control.

Linear mixed-effect models were performed to test associations of weekly asthma control scores with ambient air pollution and pollen severity levels. We treated asthma control as a continuous variable and analyzed asthma control in relation to air pollutant and pollen data by examining a change of air pollutant and pollen data from the first quartile to the third quartile corresponding to a change of asthma control scores. The main analyses focused on participants whose exposure data were monitored ≥ 3 days per week and who lived ≤ 30 km between monitoring sites and residences through a balance between the distribution of EPA monitoring days, distances to the monitoring sites, and the number of available participants.

We investigated the effect of individual environmental factors (as a main effect) on asthma control status (as a dependent variable) based on 4 analytic models. The first model included PM_{2.5} as the main effect plus important covariates, including a season/cohort indicator, and participants' sociodemographic characteristics (children/youths' age, sex, race/ethnicity, number of chronic conditions, and their parents' age, marital status, and educational background). The second model included O₃ as the main effect plus the aforementioned covariates. The third model included pollen as the main effect plus the aforementioned covariates. The fourth model investigated the effects of 3 environmental factors (PM_{2.5}, O₃, and pollen) on asthma control status after controlling the influence of covariates. Because the asthma control status of each study participant was observed across multiple time points (at most 26 times), PM_{2.5}, O₃, and pollen were treated as fixed effects in the mixed effect models, with participants treated as random effects to address the clustering effect of outcomes nested within an individual. Technically, we used an unstructured covariance matrix, which is a robust approach to estimating a covariance matrix without an a priori hypothesis.

In EPA AQS, some of the air pollutant concentration data were not collected every day. Therefore, sensitivity analyses based on different monitoring days per week (1, 3, and 5 days) were performed to test the robustness for the effects of PM_{2.5}, O₃, and pollen together on asthma control.

In addition, no gold standard is available to guide the selection of an appropriate distance between residential addresses and monitoring sites. For example, a previous study examining the effects of air pollution and asthma severity focused on participants living within 40 km,²⁵ whereas another study testing the effects of ambient air pollution on asthma exacerbations included participants living within 80 km.²⁶ Based on the distance between the residences of participants and PM_{2.5} monitoring sites (mean: 18 km; SD: 1.2 km; range: 1–82 km) and O₃-monitoring sites (mean: 16 km; SD: 1.2 km; range: 1–63 km), sensitivity analyses based on different distances (20 km, 30 km, any distance) were performed. In total, 9




Figure. Locations of study participants, air pollution–monitoring sites, and pollen-monitoring areas in Florida.

scenarios were considered in sensitivity analyses: 1) monitoring days ≥ 1 days per week and distance ≤ 20 km; 2) monitoring days ≥ 1 days per week and distance ≤ 30 km; 3) monitoring days ≥ 1 days per week and any distance; 4) monitoring days ≥ 3 days per week and distance ≤ 20 km; 5) monitoring days ≥ 3 days per week and distance ≤ 30 km; 6) monitoring days ≥ 3 days per week and any distance; 7) monitoring days ≥ 5 days per week and distance ≤ 20 km; 8) monitoring days ≥ 5 days per week and distance ≤ 30 km; and 9) monitoring days ≥ 5 days per week and any distance.

RESULTS

PARTICIPANT CHARACTERISTICS

Table 1 shows the characteristics of participants ($N = 229$). For children/youths at baseline of the first year,

the mean age was 12.2 years (SD: 2.6); 58.9% were boys; 38.0% were non-Hispanic white. The mean age of parents was 40.6 years (SD: 8.7); most parents were married or living with partners (51.5%) and had completed some college or received an associate's or undergraduate degree (60.2%). The mean distance between the ZIP code centroids and the closest $PM_{2.5}$ monitors was 18.2 km (median: 14.8; IQR: 16.1). The mean distance between the ZIP code centroids and the closest O_3 monitors was 16.4 km (median: 12.6; IQR: 12.3).

SOCIODEMOGRAPHICS, ENVIRONMENTAL EXPOSURES, WEATHER INFORMATION, AND ASTHMA CONTROL BY 2 COHORTS

Supplementary Table 1 reports the distributions of socio-demographic characteristics, $PM_{2.5}$ and O_3 concentrations,

Table 1. Characteristics of Study Participants (N=229)

Characteristics	N (%) or mean (SD)
Children	
Age, y	12.2 (2.6)
Sex	
Boy	135 (58.9%)
Girl	94 (41.1%)
Race/ethnicity	
White, non-Hispanic	87 (38.0%)
Black, non-Hispanic	59 (25.8%)
Hispanic	63 (27.5%)
Other	20 (8.7%)
Top chronic conditions	
Attention-deficit hyperactivity disorder/ attention-deficit disorders	39 (17.0%)
Born premature	26 (11.4%)
Mental health conditions	7 (3.1%)
Epilepsy or other seizure disorders	6 (2.6%)
Inflammatory bowel disease or Crohn's disease	5 (2.2%)
Deaf or hard of hearing	5 (2.2%)
Parents	
Age, y	40.6 (8.7)
Race/ethnicity	
White, non-Hispanic	97 (42.4%)
Black, non-Hispanic	60 (26.2%)
Hispanic	59 (25.8%)
Other	13 (5.7%)
Educational background	
High school or less	74 (32.7%)
Some college/associate's degree/ college degree	136 (60.2%)
Advanced degree	16 (7.1%)
Marital status	
Never married	40 (17.5%)
Married	118 (51.5%)
Living with partner in committed relationship	10 (4.4%)
Separated	9 (3.9%)
Divorced	45 (19.7%)
Widowed	7 (3.1%)
Household smoking status	
Yes	189 (82.5%)
No	40 (17.5%)

SD indicates standard deviation.

pollen severity indices, temperatures, precipitation, and asthma control scores by 2 specific seasonal cohorts. Basically, sociodemographic variables (except for the child's age) were not statistically different between the 2 cohorts ($P > .05$). There were negligible associations of temperature and precipitation with asthma control (correlation coefficients: -0.01 to 0.01). In addition, there were negligible associations of temperature with air pollutants and pollen (correlation coefficients: <0.2) and weak associations of precipitation with air pollutants and pollen (correlation coefficients: -0.26 to -0.19). This finding suggests that weather variables did not confound the associations between air pollutants/pollen and asthma control status. [Supplementary Table 2](#) reports that the mean measurements of PM_{2.5} and pollen were significantly greater in the spring season than in the fall season ($P < .001$). The mean

measurements of temperature and precipitation were greater in the fall season than in the spring season ($P < .001$). For the parsimonious modeling purpose, the seasonal cohort variable was used in the subsequent analyses to control for the variation of climate-related factors.

ASSOCIATIONS BETWEEN THE EXPOSURES AND THE OUTCOME

[Table 2](#) reports the associations of individual environmental exposure with asthma control status after adjusting for the influence of covariates (seasonal cohort and participants' sociodemographic characteristics) across 26 measurement occurrences using linear mixed-effect models. In Model 1, a greater PM_{2.5} concentration was significantly associated with poorer asthma control status (β : 0.32, $P < .001$), indicating that one IQR increase in PM_{2.5} resulted in an increase of 0.32 in asthma control scores. In Model 2, a greater O₃ concentration was associated with better asthma control status after adjusting for the season effect and participants' sociodemographics; however, the effect was not statistically significant (β : -0.02 , $P > .05$). In Model 3, a greater pollen severity index was significantly associated with poorer asthma control after adjusting for the season effect and participants' sociodemographics (β : 0.35, $P < .05$).

Model 4 reports the independent effects of 3 environmental exposures (PM_{2.5}, O₃, and pollen) on asthma control status. Results suggest that exposure to a greater PM_{2.5} concentration (β : 0.37, $P < .001$) and pollen severity (β : 0.44, $P < .05$) was significantly associated with poorer asthma control after adjusting for the season effect and participants' socio-demographics, whereas greater O₃ was marginally associated with better asthma control (β : -0.16 , $P < .1$).

Model 4 also reveals that children and adolescents with more chronic health conditions were likely to have poorer asthma control than those with fewer chronic conditions (β : 0.90, $P < .01$). Having parents who were either unmarried (β : 1.01, $P < .05$) or who had a high school education or less (β : 1.05, $P < .05$) were significantly associated with poorer asthma control than was having parents who were either married/living with a partner or who had at least some college.

SENSITIVITY ANALYSIS

[Table 3](#) reports the results of sensitivity analyses for the effects of PM_{2.5}, O₃, and pollen levels on asthma control status. Results reveal that greater PM_{2.5} concentration was significantly associated with poorer asthma control ($P < .01$) in all scenarios. Exposure to greater O₃ concentration was significantly associated with better asthma control ($P < .05$) in all analytic models, except for the scenario of monitoring days ≥ 1 and distance ≤ 30 km ($P > .05$). Exposure to greater pollen severity was significantly associated with poorer asthma control ($P < .05$) in all analytic models except for the scenarios of monitoring days ≥ 1 , ≥ 3 , and ≥ 5 and distance ≤ 20 km ($P > .05$).

Table 2. Asthma Control Status and Preceding Week's Air Quality and Pollen Severity Index^{*,†}

	Model 1 [‡] Beta (95% CI)	Model 2 [§] Beta (95% CI)	Model 3 Beta (95% CI)	Model 4 [¶] Beta (95% CI)
PM _{2.5}	0.32** (0.16–0.48)	NI	NI	0.37** (0.17–0.56)
O ₃	NI	−0.02 (−0.15 to 0.12)	NI	−0.16 ^{††} (−0.34 to 0.02)
Pollen	NI	NI	0.35 ^{‡‡} (0.05–0.65)	0.44 ^{‡‡} (0.06–0.83)
Season (ref: fall)				
Spring	0.23 (−0.61 to 1.07)	0.15 (−0.62 to 0.93)	−0.19 (−0.93 to 0.54)	−0.03 (−0.96 to 0.90)
Child's age	−0.06 (−0.24 to 0.12)	−0.01 (−0.17 to 0.16)	−0.01 (−0.16 to 0.14)	−0.05 (−0.24 to 0.13)
Child's sex (ref: girl)				
Boy	0.47 (−0.32 to 1.26)	0.43 (−0.33 to 1.19)	0.49 (−0.19 to 1.17)	0.30 (−0.53 to 1.13)
Child's race/ethnicity (ref: white)				
Black	0.78 (−0.23 to 1.79)	0.47 (−0.49 to 1.42)	0.56 (−0.30 to 1.43)	0.89 ^{††} (−0.16 to 1.94)
Hispanic	0.82 ^{††} (−0.15 to 1.78)	0.30 (−0.63 to 1.22)	0.48 (−0.36 to 1.31)	0.82 (−0.18 to 1.83)
Other	−0.06 (−1.56 to 1.44)	−0.16 (−1.65 to 1.33)	−0.19 (−1.55 to 1.18)	−0.06 (−1.60 to 1.48)
Parent's age	−0.03 (−0.08 to 0.03)	−0.03 (−0.08 to 0.02)	−0.03 (−0.07 to 0.02)	−0.03 (−0.08 to 0.02)
Marital status (ref: married)				
Not married	0.96 ^{‡‡} (0.15–1.76)	0.80 ^{‡‡} (0.03–1.57)	0.78 ^{‡‡} (0.08–1.48)	1.01 ^{‡‡} (0.17–1.84)
Education (ref: college or greater)				
High school or less	0.94 ^{‡‡} (0.07–1.81)	1.04 ^{‡‡} (0.20, 1.89)	0.84 ^{‡‡} (0.11–1.57)	1.05 ^{‡‡} (0.13–1.97)
No. of chronic conditions	0.80 ^{§§} (0.32–1.29)	0.70 ^{§§} (0.22–1.19)	0.65 ^{§§} (0.23–1.07)	0.90 ^{**} (0.38–1.43)

CI indicates confidence interval; PM, particular matter; NI, variables were not included in the modeling; and O₃, ozone.

*The results were based on the parameters of ≥ 3 monitoring days/week and a distance ≤ 30 km for measuring air pollutants (see Table 3 for results of sensitivity analyses).

†Greater scores indicate worse asthma control status.

‡Model 1 only includes PM_{2.5}.

§Model 2 only includes O₃.

||Model 3 only includes pollen.

¶Model 4 includes PM_{2.5}, O₃, and pollen.

** $P < .001$.

†† $P < .1$.

‡‡ $P < .05$.

§§ $P < .01$.

DISCUSSION

Asthma control status reflects the direct impacts of air pollutant and pollen exposures before the occurrence of adverse health events such as emergency department visits and hospitalization. Using a longitudinal design, we found that greater PM_{2.5} concentration and pollen severity significantly contributed to poorer asthma control status. In contrast, elevated O₃ concentration was marginally associated with better asthma control. Sensitivity analyses that considered different combinations of monitoring days and the distances between EPA monitors and participants' residences also revealed similar finding with respect to O₃ concentration associated with asthma control.

Previous studies have shown a protective effect of elevated O₃ concentration, including fewer asthma symptoms²⁷ and asthma exacerbations,²⁸ fewer clinical consultations and hospital admissions,²⁹ and lower incidence of other chronic conditions (eg, pulmonary³⁰ and cardiovascular diseases³¹). However, this and the aforementioned studies were conducted during the colder seasons, which is usually accompanied by low O₃ concentrations. In contrast, O₃ concentration is greater in the warmer seasons,^{32,33} and exposure to high levels of O₃ increases asthma incidence among children with frequent outdoor activities.³⁴ Future studies should investigate the association of O₃ concentration with asthma control in different areas with warmer temperatures to determine whether the effects remain protective.

Alternatively, the marginally significant association of elevated O₃ with good asthma control may be because we did not account for the association of O₃ with other air pollutants since greater O₃ concentrations are correlated with lower NO₂, CO, and SO₂ concentrations.^{35,36} NO₂, CO, and SO₂ may confound the association between O₃ and asthma control. Because NO₂, CO, and SO₂ data were not collected in all monitoring sites in Florida, we did not include these air pollutants in analyses. Future studies are encouraged to investigate the influence of exposure to comprehensive ambient air pollutants on asthma control status.

Previous studies have performed sensitivity analyses to understand whether different distances between air pollution monitors and participants' residences led to different associations between air pollutants, pollen levels, and health outcomes.^{37,38} However, sensitivity analyses in previous studies rarely included different monitoring days per week of air pollutants as a parameter. In this study, because children's asthma control status was assessed by using a 7-day recall period and air pollution/pollen data were not collected every day, it is challenging to disentangle the temporal sequence between the dates of air pollution/pollen exposure and the dates of asthma exacerbations. We believe that the use of ≥ 3 monitoring days per week as the main exposure window in this study is the optimal approach.

The harmful effects of PM_{2.5} and pollen on asthma control status are a significant threat to the health of children with asthma. Preventing exposure to these ambient materials

Table 3. Sensitivity Analyses for Associations Between Asthma Control Status and Air Quality and Pollen Severity Index^{*,†}

Scenarios	No. Subjects	No. Observations	PM _{2.5} Beta (95% CI)	O ₃ Beta (95% CI)	Pollen Beta (95% CI)
≥1 monitoring days					
0 ≤ distance ≤20 km	108	1806	0.32 [‡] (0.11–0.53)	–0.20 [§] (–0.39 to –0.02)	0.42 (–0.02 to 0.86)
0 ≤ distance ≤30 km	146	2471	0.31 [¶] (0.13–0.49)	–0.16 (–0.32 to 0.01)	0.50 [‡] (0.13–0.86)
Distance ≥0 km	181	3016	0.22 [‡] (0.07–0.38)	–0.16 [§] (–0.31 to –0.01)	0.37 [§] (0.05–0.68)
≥3 monitoring days					
0 ≤ distance ≤20 km	103	1541	0.37 [‡] (0.13–0.61)	–0.21 [§] (–0.41 to –0.01)	0.38 (–0.09 to 0.84)
0 ≤ distance ≤30 km	141	2153	0.37 [¶] (0.17–0.56)	–0.16 (–0.34 to 0.02)	0.44 [§] (0.06–0.83)
Distance ≥0 km	175	2629	0.26 [‡] (0.08–0.43)	–0.17 [§] (–0.33 to –0.01)	0.35 [§] (0.02–0.68)
≥5 monitoring days					
0 ≤ distance ≤20 km	90	1437	0.42 [‡] (0.16–0.67)	–0.23 [§] (–0.44 to –0.02)	0.35 (–0.13 to 0.84)
0 ≤ distance ≤30 km	124	2013	0.40 [¶] (0.19–0.62)	–0.19 [§] (–0.38 to –0.01)	0.45 [§] (0.06–0.85)
Distance ≥0 km	153	2429	0.31 [‡] (0.12–0.49)	–0.23 [‡] (–0.40 to –0.06)	0.36 [§] (0.01–0.70)

PM indicates particulate matter; O₃, ozone; and CI, confidence interval.

*Sensitivity analyses were adjusted for the season and sociodemographic variables, including the child's age, sex, race/ethnicity, and number of chronic conditions and the parent's age, marital status, and educational background.

†Greater scores indicate worse asthma control status.

‡*P* < .01.

§*P* < .05.

||*P* < .1.

¶*P* < .001.

is an important task in asthma management, both on a practice and policy level. Physicians and health care agencies may develop warning systems to alert children with asthma, parents, and school teachers about the harmful effects of air pollution and pollen on asthma health outcomes. Recent evidence has pinpointed that the implementation of air quality control policies can significantly improve air quality in Southern California, and this policy has shown improvement in pulmonary functioning of children.⁶ In addition, increased emissions of greenhouse gases by motor vehicles and other industries have been linked to global warming, longer pollen seasons, more-severe pollen exposure, and frequent respiratory symptoms.³⁹ It is crucial to develop air quality control policies to address these climate change and air pollution issues.

Several limitations should be considered. First, our participants were recruited from Florida Medicaid and SCHIP and may not be generalizable to other populations. It is critical to evaluate the selection bias related to the characteristics of participants and nonparticipants. Unfortunately, sociodemographic data (other than age and sex) and past asthma events were not available from nonparticipants. Second, the sources and severity of indoor pollutants (eg, radon, carbon monoxide, etc) were not collected in this study. Although this study measured household smoking status using a binary item, we did not include this variable in multivariable analysis, given an unexpected high smoking prevalence (83%) and nonsignificant

associations with asthma control. With the use of sophisticated measures, evidence has shown a harmful effect of smoking on asthma outcomes (eg, impairments of lung development, and worse asthma control).⁴⁰ Third, the pollen index was an aggregate of different origins and types of pollen (eg, tree, grass, and weed pollen). Future studies should distinguish different origins/types of pollen on asthma control status. It is also important to establish a sensitization profile for each child/adolescent by associating different ambient pollutant and pollen data with asthma severity and apply portable/direct devices to collect different sources and types of ambient pollutant and pollen data to establish sensitization profiles for individuals with asthma. Fourth, although we found that exposure to elevated concentrations of PM was associated with worse asthma symptom severity, future studies are warranted to investigate how the sources of PM with varying toxicity contribute to asthma symptoms and quality of life in pediatric population. Fifth, this study relies on the zip codes of the residences to link air pollution data from the nearby EPA monitoring sites rather than the use of portable/direct measurements. However, this ZIP code–based approach provides insightful preliminary evidence for future research to confirm our findings through the portable/direct measurements that could take into consideration the influence of meteorological factors on the movement of pollution plumes. Finally, the resulting poor asthma control associated with PM_{2.5} may not indicate a causal

relationship unless other co-pollutants (SO₂, CO, NO₂, etc.) have been accounted for. Unfortunately, SO₂, CO, and NO₂ data collected from the monitoring sites in Florida are incomplete or insufficient compared with PM_{2.5} and O₃ data. In addition, it is warranted to establish the causality through longitudinal observations spanning multiple year and accounting for the meteorological variations between the seasons and years.

In conclusion, elevated PM_{2.5} concentration and pollen severity were associated with poorer asthma control in pediatric patients with asthma. Effective strategies to prevent children and adolescents from unnecessary exposure to harmful outdoor environmental factors may improve pediatric asthma outcomes.

ACKNOWLEDGMENTS

Financial disclosure: National Institutes of Health U01 AR052181 (to L.A.T., H.E.G., E.A.S., D.A.D., I.-C.H.). The funder has no role or influence in the study design, the collection, analysis, and interpretation of data; the writing of the manuscript; or the decision to submit the manuscript for publication.

SUPPLEMENTARY DATA

Supplementary data related to this article can be found online at <https://doi.org/10.1016/j.acap.2019.03.010>.

REFERENCES

- Bloom B, Jones LI, Freeman G. Summary health statistics for U.S. children: National Health Interview Survey, 2012. *Vital Health Stat 10*. 2013;(258):1–81.
- Bateson TF, Schwartz J. Children's response to air pollutants. *J Toxicol Environ Health A*. 2008;71:238–243.
- Gehring U, Wijga AH, Brauer M, et al. Traffic-related air pollution and the development of asthma and allergies during the first 8 years of life. *Am J Respir Crit Care Med*. 2010;181:596–603.
- Jerrett M, Shankardass K, Berhane K, et al. Traffic-related air pollution and asthma onset in children: a prospective cohort study with individual exposure measurement. *Environ Health Perspect*. 2008;116:1433–1438.
- McConnell R, Islam T, Shankardass K, et al. Childhood incident asthma and traffic-related air pollution at home and school. *Environ Health Perspect*. 2010;118:1021–1026.
- Gauderman WJ, Urman R, Avol E, et al. Association of improved air quality with lung development in children. *N Engl J Med*. 2015;372:905–913.
- Roy A, Sheffield P, Wong K, et al. The effects of outdoor air pollutants on the costs of pediatric asthma hospitalizations in the United States, 1999 to 2007. *Med Care*. 2011;49:810–817.
- Trasande L, Liu Y. Reducing the staggering costs of environmental disease in children, estimated at \$76.6 billion in 2008. *Health Aff*. 2011;30:863–870.
- Cloutier MM, Schatz M, Castro M, et al. Asthma outcomes: composite scores of asthma control. *J Allergy Clin Immunol*. 2012;129(3 suppl):S24–S33.
- Maestrelli P, Canova C, Scapellato ML, et al. Personal exposure to particulate matter is associated with worse health perception in adult asthma. *J Investig Allergol Clin Immunol*. 2011;21:120–128.
- Zora JE, Sarnat SE, Raysoni AU, et al. Associations between urban air pollution and pediatric asthma control in El Paso, Texas. *Sci Total Environ*. 2013;448:56–65.
- DellaValle CT, Triche EW, Leaderer BP, et al. Effects of ambient pollen concentrations on frequency and severity of asthma symptoms among asthmatic children. *Epidemiology*. 2012;23:55–63.
- Gleason JA, BL; Fagliano JA. Associations between ozone, PM_{2.5}, and four pollen types on emergency department pediatric asthma events during the warm season in New Jersey: a case-crossover study. *Environ Res*. 2014;132:421–429.
- Jariwala SP, Kurada S, Moday H, et al. Association between tree pollen counts and asthma ED visits in a high-density urban center. *J Asthma*. 2011;48:442–448.
- Wang HC, Yousef E. Air quality and pediatric asthma-related emergencies. *J Asthma*. 2007;44:839–841.
- Magas OK, Gunter JT, Regens JL. Ambient air pollution and daily pediatric hospitalizations for asthma. *Environ Sci Pollut Res Int*. 2007;14:19–23.
- Tosca MA, Ruffoni S, Canonica GW, et al. Asthma exacerbation in children: relationship among pollens, weather, and air pollution. *Allergol Immunopathol (Madr)*. 2014;42:362–368.
- Phillips JF, Jelks ML, Lockey RF. Important Florida botanical aeroallergens. *Allergy Asthma Proc*. 2010;31:337–340.
- Patino CM, Okelo SO, Rand CS, et al. The Asthma Control and Communication Instrument: a clinical tool developed for ethnically diverse populations. *J Allergy Clin Immunol*. 2008;122:936–943. e936.
- Gandhi PK, Kenzik KM, Thompson LA, et al. Exploring factors influencing asthma control and asthma-specific health-related quality of life among children. *Respir Res*. 2013;14:26.
- United States Environmental Protection Agency. *Download daily data*. Available at: http://www.epa.gov/airdata/ad_data_daily.html. Accessed May 1, 2018.
- United States Census Bureau. *2010 TIGER/LINE® shapefile*. Available at: <http://www.census.gov/cgi-bin/geo/shapefiles2010/main>. Accessed May 1, 2018.
- IMS Health. Available at: <http://www.imshealth.com/portal/site/imshealth>. Accessed May 1, 2018.
- National Climatic Data Center. *Climate data online*. Available at: <http://www.ncdc.noaa.gov/cdo-web>. Accessed May 1, 2018.
- Rage E, Siroux V, Kunzli N, et al. Epidemiological study on the genetics and environment of asthma. Air pollution and asthma severity in adults. *Occup Environ Med*. 2009;66:182–188.
- Schildcrout JS, Sheppard L, Lumley T, et al. Ambient air pollution and asthma exacerbations in children: an eight-city analysis. *Am J Epidemiol*. 2006;164:505–517.
- Studnicka M, Hackl E, Pischinger J, et al. Traffic-related NO₂ and the prevalence of asthma and respiratory symptoms in seven year olds. *Eur Respir J*. 1997;10:2275–2278.
- Evans KA, Halterman JS, Hopke PK, et al. Increased ultrafine particles and carbon monoxide concentrations are associated with asthma exacerbation among urban children. *Environ Res*. 2014;129:11–19.
- Hajat S, Haines A, Goubet SA, et al. Association of air pollution with daily GP consultations for asthma and other lower respiratory conditions in London. *Thorax*. 1999;54:597–605.
- Carey IM, Atkinson RW, Kent AJ, et al. Mortality associations with long-term exposure to outdoor air pollution in a national English cohort. *Am J Respir Crit Care Med*. 2013;187:1226–1233.
- Atkinson RW, Carey IM, Kent AJ, et al. Long-term exposure to outdoor air pollution and incidence of cardiovascular diseases. *Epidemiology*. 2013;24:44–53.
- Kheirbek I, Wheeler K, Walters S, et al. PM_{2.5} and ozone health impacts and disparities in New York City: sensitivity to spatial and temporal resolution. *Air Qual Atmos Health*. 2013;6(2):473–486.
- Raun LH, Ensor KB, Persse D. Using community level strategies to reduce asthma attacks triggered by outdoor air pollution: a case crossover analysis. *Environ Health*. 2014;13:58.
- Guarnieri M, Balmes JR. Outdoor air pollution and asthma. *Lancet*. 2014;383:1581–1592.
- Meng YY, Rull RP, Wilhelm M, et al. Outdoor air pollution and uncontrolled asthma in the San Joaquin Valley, California. *J Epidemiol Community Health*. 2010;64:142–147.
- Moore K, Neugebauer R, Lurmann F, et al. Ambient ozone concentrations cause increased hospitalizations for asthma in children: an 18-year study in Southern California. *Environ Health Perspect*. 2008;116:1063–1070.

37. Chang HH, Reich BJ, Miranda ML. Time-to-event analysis of fine particle air pollution and preterm birth: results from North Carolina, 2001–2005. *Am J Epidemiol.* 2012;175:91–98.
38. Morgenstern V, Zutavern A, Cyrys J, et al. Atopic diseases, allergic sensitization, and exposure to traffic-related air pollution in children. *Am J Respir Crit Care Med.* 2008;177:1331–1337.
39. D'Amato G, Cecchi L. Effects of climate change on environmental factors in respiratory allergic diseases. *Clin Exp Allergy.* 2008;38:1264–1274.
40. Thomson NC. The role of environmental tobacco smoke in the origins and progression of asthma. *Curr Allergy Asthma Rep.* 2007;7:303–309.