



## Correction to: Ultrashort time-to-echo quantitative magnetic resonance imaging of the triangular fibrocartilage: differences in position

Akitaka Fujisaki<sup>1</sup> · Takatoshi Aoki<sup>1</sup> · Hidekuni Narimatsu<sup>1</sup> · Chie Kuwahara<sup>1</sup> · Atsushi Nozaki<sup>2</sup> · Kunitaka Menuki<sup>3</sup> · Akinori Sakai<sup>3</sup> · Yukunori Korogi<sup>1</sup>

Published online: 19 February 2019  
© European Society of Radiology 2019

### European Radiology

<https://doi.org/10.1007/s00330-018-5693-7>

The original version of this article, published on 03 September 2018, unfortunately contained a mistake. The following correction has therefore been made in the original: The presentation of Table 1 was incorrect. The corrected table is given below. The original article has been corrected.

**Table 1** Average T2\* values of TFC of the wrist according to the forearm/wrist position

Position	Neutral	Pronation	Ulnar flexion
mean ± 2SD (ms)	10.08 ± 1.90	7.92 ± 1.37	9.15 ± 1.03

\*                      \*

\* $p < 0.017$  (two-tailed Wilcoxon signed-rank test)

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00330-018-5693-7>

✉ Takatoshi Aoki  
a-taka@med.uoeh-u.ac.jp

<sup>1</sup> Department of Radiology, University of Occupational and Environmental Health, 1-1 Iseigaoka, Yahatanishi-ku, Kitakyushu 807-8555, Japan

<sup>2</sup> MR Applications and Workflow, Asia Pacific, GE Healthcare, Tokyo, Japan

<sup>3</sup> Department of Orthopaedic Surgery, University of Occupational and Environmental Health, Kitakyushu, Japan