



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Revision Roux-en-y hepaticojejunostomy for a post-cholecystectomy complex vasculobiliary injury with complete proper hepatic artery occlusion: A case report and literature review

Gunjan S. Desai^{a,*}, Prasad Pande^a, Rajvilas Narkhede^b, Dattaprasanna R. Kulkarni^a

^a Department of Gastrointestinal Surgery, Lilavati Hospital and Research Center, Mumbai, Maharashtra, 400050, India

^b Department of Gastrointestinal Surgery, Dr. Balabhai Nanavati Super Speciality Hospital, Vile Parle (West), Mumbai, Maharashtra, India

ARTICLE INFO

Article history:

Received 31 January 2019

Received in revised form 13 March 2019

Accepted 21 March 2019

Available online 9 April 2019

Keywords:

Complete hepatic arterial ischemia

Complex biliary injury

Revision hepaticojejunostomy

Case report

ABSTRACT

INTRODUCTION: Complete proper hepatic arterial [PHA] occlusion due to accidental coil migration during embolization of cystic artery stump pseudoaneurysm resulting from a complex vasculobiliary injury [CVBI] post laparoscopic cholecystectomy [LC] is an extremely rare complication with less than 15 cases reported. We present a case depicting our strategy to tackle this obstacle in management of CVBI and review the relevant literature.

PRESENTATION OF CASE: A 35 year old lady presented on sixth postoperative day with an external biliary fistula following Roux-en-y hepaticojejunostomy [RYHJ] for biliary injury during LC. She developed a leaking cystic artery pseudoaneurysm, during angioembolisation of which, one coil accidentally migrated into left hepatic artery resulting in complete PHA occlusion. Fourteen months later, cholangiogram revealed a worsening RYHJ stricture despite repeated percutaneous balloon dilatations. Multiple collaterals had developed. Revision RYHJ was fashioned to the anterior wall of biliary confluence with an extension into left duct. Minimum hilar dissection ensured preservation of collateral supply to the biliary enteric anastomosis. Postoperative recovery was uneventful. The patient is doing well at 1 year follow up.

DISCUSSION: Definitive biliary enteric repair should be delayed till collateral circulation is established within the hilar plate, hepatoduodenal ligament and perihepatic/peribiliary collaterals to provide an adequate arterial blood supply to biliary confluence and extrahepatic portion of the bile duct.

CONCLUSION: Assessment of hepatic arteries should be part of investigation of all complex biliary injuries. Delayed definitive biliary enteric repair ensures a well-perfused anastomosis. Minimum hilar dissection is the key to preserve biliary and hepatic neovascularization.

© 2019 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Complex vasculobiliary injuries [CVBI] constitute 12–61% of biliary injuries, most commonly following laparoscopic cholecystectomy [1]. They present a formidable challenge to the multidisciplinary team treating these patients. During management, complete right and left hepatic arterial occlusion due to accidental coil migration during embolization of a cystic artery stump pseudoaneurysm is an extremely rare complication [2,3]. We present a case depicting our strategy to tackle this obstacle in the management of CVBI. This work has been reported in line with the SCARE criteria [4].

* Corresponding author at: 1301, Building-6, Kesar Harmony Society, Sector 6, Kharghar, Navi Mumbai, Maharashtra, 410210, India.

E-mail addresses: desaigunjan526@gmail.com, dsshish@gmail.com (G.S. Desai), prasadpande2111@gmail.com (P. Pande), rajvilas86@gmail.com (R. Narkhede), surgeonliverpancreas@gmail.com (D.R. Kulkarni).

2. Case

A 35 year old healthy lady presented to our department on sixth postoperative day [POD] with an external biliary fistula and intra-abdominal sepsis. She had undergone laparoscopic converted to open cholecystectomy for acute calculous cholecystitis. She had a biliary injury that was identified intra-operatively, managed by Roux-en-y hepaticojejunostomy [RYHJ]. The anastomosis leaked. An interno-external percutaneous transhepatic biliary drainage [PTBD] extending across the leak was performed at our hospital on POD 7 for both right and left hepatic ducts.

On POD nine, she had an upper gastrointestinal bleed. Esophagogastroduodenoscopy and Contrast enhanced computed tomography [CECT] abdomen did not reveal the source of bleeding. On conventional hepatic arteriogram, a leaking cystic artery pseudoaneurysm was identified (Fig. 1). During angioembolisation, due to a short stump of cystic artery, coils were placed in right hepatic artery [RHA]. However, one of the coils accidentally migrated into the left hepatic artery [LHA] and could not be retrieved. LHA




Fig. 1. Conventional celiac artery angiogram showing the cystic artery pseudoaneurysm[yellow arrow].




Fig. 2. Conventional celiac artery angiography and coiling of cystic artery pseudoaneurysm and right hepatic artery and coil migration into left hepatic artery followed by stenting of the left hepatic artery[yellow arrow].

stenting was performed, with good flow of contrast across the stent (Fig. 2). However, LHA developed spasm in its distal part, resulting in complete block of LHA and RHA.

On first day after coiling, there was significant elevation of liver enzymes with features of ischemic hepatitis. CECT abdomen with arteriography revealed poor enhancement of hepatic arterial tree in the segmental branches with partial revascularization from inferior phrenic and retroperitoneal arteries. The patient's relatives were explained the possibility of a need for an emergency liver transplant. The patient improved over the next 48 h, was transferred out of intensive care unit and oral feeds were started. Abdominal drain was removed after it stopped draining bile. She was discharged on POD 28 with PTBD catheters in situ. On POD 33, her liver function tests were within normal limits, percutaneous transhepatic cholangiogram showed trickle of contrast across the RYHJ, and the PTBD catheters were clamped. PTBD catheters were kept longer than 6 months and intervening periodic cholangiograms revealed anastomotic narrowing. Balloon dilatation of the stenosed anastomosis was performed on multiple occasions. Liver function was normal all along.




Fig. 3. Percutaneous transhepatic cholangiogram showing the Roux-en-y hepaticojejunostomy anastomotic stricture.



Fig. 4. Conventional celiac artery angiogram showing collaterals[red arrows] from the proper hepatic artery, inferior phrenic artery and along the hepatoduodenal ligament.

Fourteen months after surgery, cholangiogram revealed a worsening of RYHJ stricture (Fig. 3). A CECT abdomen and conventional angiogram performed at 18 months showed a blocked RHA and LHA. Multiple collaterals were seen arising from the right inferior phrenic artery, retroperitoneum and along the hepatoduodenal ligament (Fig. 4). In view of the collateral formation and persistent tight stricture with failed multiple dilatations, a surgical revision of the anastomosis was planned. Prior to surgery, right PTBD catheter was maneuvered from right to left duct across the hilum, draining externally.

During surgery, utmost care was taken not to release any adhesions in the hepatoduodenal ligament and not to mobilize liver from the diaphragmatic attachments to preserve all collaterals. Hilar dissection was kept to a minimum and the PTBD catheters guided the identification of biliary hilar confluence. A minor segment IV hepatectomy was done to expose the left hepatic duct and the roof of biliary confluence. Previous anastomosis was resected, and a Roux loop was prepared. A 2 cm single layer, interrupted 5-0 polydioxanone RYHJ was fashioned to the anterior wall of the biliary confluence with an extension onto the left duct.

Her postoperative recovery was uneventful. Cholangiogram on POD 10 showed a good flow across the anastomosis with no sign of a leak, and she was discharged with a clamped PTBD catheter. The PTBD catheter was removed 6 weeks after the surgery. The patient is doing well at 1 year follow up. Liver function is normal. She is being followed up to evaluate for secondary patency.

Table 1

Reported cases of complex vasculobiliary injuries involving common hepatic artery or proper hepatic artery during cholecystectomy, their presentation, management and outcomes.

Sr no.	Year	Arterial injury	Biliary injury	Identification and Presentation	Management	Follow-up	Outcome
1	Frilling et al. [9]	Proper hepatic artery	E2	During first surgery	End-to-end repair of artery and Roux-en-Y hepaticojejunostomy – On POD 16, re-exploration showed ligated right hepatic artery and thrombosis of superior mesenteric vein and portal vein – right hepatectomy and listed for urgent liver transplantation	None	Death on POD 28 – could not undergo liver transplant
2	Frilling et al. [9]	Common hepatic artery	E1	POD 17	Reconstruction of artery using mesenteric venous graft and external drainage of both biliary ducts – Roux-en-Y hepaticojejunostomy after 3 days	None	Death on POD 37
3,4 5	Wudel et al. [10] Gupta et al. [11]	Common hepatic artery Proper hepatic artery	Not available Strasberg E4	Not available Postoperative day 4	Not available Bilateral hepaticojejunostomies without vascular repair/reconstruction	Not available One and half year +	Not available Recovered with intermittent biliary strictures being managed with percutaneous transhepatic balloon dilatations
6	Buell et al. [7]	Common hepatic artery	Transected right hepatic duct	Average 8.5 days	Primary repair of artery and percutaneous external biliary drainage	None	Death
7	Buell et al. [7]	Common hepatic artery	Major biliary injury – Roux-en-Y hepaticojejunostomy in first surgery	Average 8.5 days	Right hepatectomy and posted for liver transplantation	None	Death
8	Buell et al. [7]	Common hepatic artery	Major biliary injury – Roux-en-Y hepaticojejunostomy in first surgery	Average 8.5 days	Orthotopic liver transplantation	Regular	Alive
9	Salman et al. [12]	Proper hepatic artery	E3	First surgery was open cholecystectomy with right hepatic artery ligation	Second surgery on postoperative day 3 – T-tube drainage and hemostasis Percutaneous drainage of cholangitic liver abscess on POD 85 Third surgery on postoperative day 144 – Roux-en-Y hepaticojejunostomy	One year +	Recovered
10	Yan et al. [13]	Proper hepatic artery	E4	Roux-en-Y hepaticojejunostomy during first surgery {laparoscopic cholecystectomy} or first postoperative week of surgery without repair/reconstruction of artery	Secondary biliary cirrhosis and portal hypertension on follow up {Range of 5–148 months} – Underwent liver transplantation	Range of 4–17 months	Recovered
11	Strasberg et al. [14]	Proper hepatic artery	Necrosis of the intrahepatic biliary tree till fourth order branches	Postoperative day 6 with a large biliary fistula	Liver transplantation on Postoperative day 39	None	Death on post transplant day 14
12, 13	Sarno et al. [1]	Common hepatic artery	Major biliary injury	Details not available	Primary or revisional hepaticojejunostomy. Vascular repair/reconstruction not described	Follow up range of 12–245 months	Recovered

Vascular assessment should be part of investigation of all complex biliary injury cases. Delayed definitive repair in cases involving PHA occlusion to allow collateral circulation to be established within the hilar plate, hepatoduodenal ligament and perihepatic/peribiliary collaterals and thereby provide an adequate arterial blood supply to the biliary confluence and the extrahepatic portion

of bile duct is a feasible management option. Minimum dissection should be done during surgery to preserve biliary and hepatic neovasculature.

Conflicts of interest

Authors declare that they have no conflict of interest.

Sources of funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Ethical approval

Institutional ethics committee approval was taken for the publication.

Consent

Informed consent was obtained from the patient for this publication.

Author's contribution

Gunjan Desai: collecting data, analysis of data, preparing the initial draft of the manuscript, critical revision of the manuscript for intellectual content, technical support, material support, study supervision.

Prasad Pande: critical revision of the manuscript for intellectual content, technical support, material support, study supervision.

Rajvilas Narkhede: critical revision of the manuscript for intellectual content, technical support, material support, study supervision.

Dattaprasanna Kulkarni: study concept, critical revision of the manuscript for intellectual content, administrative, technical support, material support, study supervision.

Registration of research studies

This is a case report and does not require such registration. The publication was performed after the approval of research protocols by the Ethics Committee of Lilavati Hospital and Research Centre in accordance with international agreements (World Medical Association Declaration of Helsinki "Ethical Principles for Medical Research Involving Human Subjects," amended in October 2013, www.wma.net).

Guarantor

Gunjan S Desai is the article guarantor.

Provenance and peer review

Not commissioned, externally peer-reviewed.

References

- [1] G. Sarno, A. Al-Sarira, P. Ghaneh, S. Fenwick, H. Malik, G. Poston, Cholecystectomy-related bile duct and vasculobiliary injuries, *Br. J. Surg.* 99 (8) (2012) 1129–1136.
- [2] K. To, E. Lai, D. Chung, O. Chan, C. Tang, Cystic artery pseudoaneurysm with haemobilia after laparoscopic cholecystectomy, *Hong Kong Med. J.* (2018) 203–205.
- [3] W. She, S. Tsang, R. Poon, T. Cheung, Gastrointestinal bleeding of obscured origin due to cystic artery pseudoaneurysm, *Asian J. Surg.* 40 (4) (2017) 320–323.
- [4] R.A. Agha, M.R. Borrelli, R. Farwana, K. Koshy, A. Fowler, D.P. Orgill, For the SCARE Group, The SCARE 2018 statement: updating consensus Surgical Case REport (SCARE) guidelines, *Int. J. Surg.* 60 (2018) 132–136.
- [5] E. De Santibañes, V. Ardiles, J. Pekolj, Complex bile duct injuries: management, *HPB* 10 (1) (2008) 4–12.
- [6] B. Sureka, A. Mukund, Review of imaging in post-laparoscopy cholecystectomy complications, *Indian J. Radiol. Imaging* 27 (4) (2017) 470.
- [7] J. Buell, Devastating and fatal complications associated with combined vascular and bile duct injuries during cholecystectomy, *Arch. Surg.* 137 (6) (2002) 703.
- [8] S. Strasberg, W. Helton, An analytical review of vasculobiliary injury in laparoscopic and open cholecystectomy, *HPB* 13 (1) (2011) 1–14.
- [9] A. Frilling, J. Li, F. Weber, F. Weber, N.R. Fruhauf, J. Engel, et al., Major bile duct injuries after laparoscopic cholecystectomy: a tertiary centre experience, *J. Gastrointest. Surg.* 8 (2004) 679–685.
- [10] L.J. Wudel Jr., J.K. Wright, C.W. Pinson, A. Herline, J. Debelak, S. Seidel, et al., Bile duct injury following laparoscopic cholecystectomy: a cause for continued concern, *Am. Surg.* 67 (2001) 557–563.
- [11] N. Gupta, Management and outcome of patients with combined bile duct and hepatic artery injuries, *Arch. Surg.* 133 (2) (1998) 176.
- [12] B. Salmen, N. Akyurek, B. Onal, M. Cindoruk, Combined proper hepatic artery and common hepatic duct injury in open cholecystectomy: case report and review of the literature, *Adv. Ther.* 24 (2007) 639–647.
- [13] J. Yan, Surgical management in biliary stricture after Roux-en-Y hepaticojejunostomy for bile duct injury, *World J. Gastroenterol.* 13 (48) (2007) 6598.
- [14] S. Strasberg, D. Gouma, 'Extreme' vasculobiliary injuries: association with fundus-down cholecystectomy in severely inflamed gallbladders, *HPB* 14 (1) (2012) 1–8.
- [15] F. Sugihara, S. Murata, F. Uchiyama, J. Watari, Y. Abe, et al., Successful management of a proper hepatic artery aneurysm by embolization without liver dysfunction, *J. Radiol. Radiat. Ther.* 1 (2) (2013) 1007.
- [16] N. Michels, Collateral arterial pathways to the liver after ligation of the hepatic artery and removal of the celiac axis, *Cancer* 6 (4) (1953) 708–724.

Open Access

This article is published Open Access at sciendo.com. It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.