



A grid-cell based fecal sampling scheme reveals: land-use and altitude affect prevalence rates of *Angiostrongylus vasorum* and other parasites of red foxes (*Vulpes vulpes*)

Barbara Koller¹ · Daniel Hegglin¹ · Manuela Schnyder¹

Received: 30 October 2018 / Accepted: 16 April 2019 / Published online: 11 May 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

In view of the role of foxes as a reservoir for *Angiostrongylus vasorum*, a nematode of the heart and lungs of dogs and foxes, its occurrence across Switzerland was investigated in foxes applying a standardized sampling scheme for fox fecal samples. In 72 study areas, which consisted of three 1-km² grid cells, a total of 1481 samples were collected by walking transects following linear features in the terrain, and analyzed by a flotation-sieving method. The overall prevalence rate of *A. vasorum* in fecal samples was 8.8% (95% confidence interval, CI 7.4–10.3%), being significantly higher in the Swiss Plateau (11.4%, CI. 9.4–13.8%) compared with other bioregions, and more prevalent in areas with less than 50% of cultivated land ($p = 0.043$). Prevalence rates increased with decreasing altitudes being significantly higher below 400 m above sea level (20.6%, 95% CI 15.2–26.9%), while all samples collected above 900 m asl were negative. Eggs of *Toxocara* sp. (12.1%), Taeniidae (10.5%), *Capillaria* spp. (8.3%), *Trichuris vulpis* (5.5%), hookworms (5.3%), *Toxascaris leonina* (1.3%) and *Strongyloides* sp. (0.4%) were furthermore identified. Taeniid eggs were positively and *Capillaria* spp. negatively associated with the amount of cultivated land. The prevalence rates based on our fecal analyses were generally lower compared with previous studies from Switzerland which were based on fox necropsies. However, the grid cell-based sampling scheme with replicable transects for fox feces collection proved to be a suitable, non-invasive method for parasitic large-scale surveys providing information on the relative infectious pressure for the local dog population.

Keywords Helminths · Standardized method · Fox feces · Epidemiology · Bioregions · Switzerland

Introduction

Angiostrongylus vasorum is a 13–21 mm long metastrongylid nematode, which mainly affects foxes and dogs but also wolves, badgers, and other wild carnivores. The parasite is transmitted by ingestion of snails of different species (Guilhon and Bressou 1960). In its definitive host, *A. vasorum* resides in the pulmonary arteries and the right side of the heart. The parasite causes cardio-respiratory and neurological signs as well as bleeding disorders especially in dogs, with a potentially fatal outcome (Chapman et al. 2004; Koch and Willeßen 2009; Staebler et al. 2005).

Daniel Hegglin and Manuela Schnyder contributed equally to this work.

Section Editor: David Bruce Conn

✉ Manuela Schnyder
manuela.schnyder@uzh.ch

¹ Institute of Parasitology, Vetsuisse Faculty, University of Zurich, Winterthurerstrasse 266a, 8057 Zürich, Switzerland

Traditionally, *A. vasorum* occurred in a patchy pattern with endemic foci in southern France (Cuillé and Darraspen 1930; Guilhon 1963) and Ireland (Dodd 1973; Roche and Kelliher 1968), followed by reports from Denmark (Bolt et al. 1992). During the last decades, findings from other regions in Great Britain (Helm et al. 2015; Morgan et al. 2008; Schnyder et al. 2013a), on the Iberian peninsula (Alho et al. 2014; Gerrikagoitia et al. 2010; Segovia et al. 2004) and in central (Barutzki and Schaper 2009; Bolt et al. 1992; Jolly et al. 2015; Staebler et al. 2005; Taubert et al. 2008; van Doorn et al. 2009), southern (Guardone et al. 2013; Magi et al. 2009; Papazahariadou et al. 2007; Tieri et al. 2011), northern (Grandi et al. 2017), and eastern Europe (Hurnikova et al. 2013; Majoros et al. 2010; Rajkovic-Janje et al. 2002; Schnyder et al. 2013b) indicate that this parasite becomes more and more important, occurring in areas which were not reported as endemic so far.

Several factors influencing the distribution pattern of *A. vasorum* and the apparently increasing number of reported dogs suffering from the disease are under discussion. Traveling with untreated dogs from endemic to non-endemic

areas could play a role in the expansion of *A. vasorum* into distant, previously parasite-free areas. In contrast, climatic factors, such as mean winter temperatures below $-4\text{ }^{\circ}\text{C}$ may represent a barrier for the establishment of *A. vasorum* (Jeffery et al. 2004; Lurati et al. 2015). Importantly, in areas with a dense fox population, foxes may contribute to the endemic establishment and an increased parasitic infection threat, representing a reservoir and therefore a risk for the dog population. Foxes are well adapted to urban habitats and considered as opportunists concerning their nourishment, and urban areas offer them favorable terms for associations to family groups and to live in high-density populations (Gloor et al. 2001). This fact is of concern because foxes and dogs are sharing most of their parasites: *Toxocara canis*, *Capillaria* spp., *Uncinaria stenocephala*, *Taenia* and *Echinococcus* spp., *Mesocestoides* spp., *Dipylidium caninum* (Fischer et al. 2005; Hofer et al. 2000; Willingham et al. 1996), and *A. vasorum* and *Crenosoma vulpis* as well.

In Switzerland, first cases of *A. vasorum* were reported from a dog breeding station in Zurich in 1968 (Wolff et al. 1969), but only approximately 30 years later Staebler et al. (2005) reported about five dogs infected with *A. vasorum* in the northern part of Switzerland, and about three dogs coming from southern Ticino, all diagnosed between 1999 and 2004. In 2001, two infected foxes originating from the region of Basel were reported (Gottstein 2001). Since 2004 up to now, numerous additional foxes and new cases of canine angiostrongylosis have been identified (Gillis-Germitsch et al. 2017; Sigrist et al. 2017), partly in the same regions as already reported, but also in new areas, as for example in the eastern and French part of Switzerland (Lurati et al. 2015). In dogs, definitive diagnosis is typically attained by isolation of first stage larvae (L1) from feces through larval migration adopting the Baermann-Wetzel technique (Deplazes et al. 2016) or by serological methods (Schnyder et al. 2015; Schnyder et al. 2011). In contrast, prevalence data on foxes mostly rely on culled animals and correlated necropsy findings and/or adopting serological methods adapted to foxes (Gillis-Germitsch et al. 2017; Houpin et al. 2016), therefore implying selection based on killed animals.

In view of the role of foxes as a reservoir for *A. vasorum* and of the increasing number of reports of canine angiostrongylosis, epidemiological investigations on foxes were performed applying a standardized sampling scheme based on the collection of fox fecal samples along transects in 1-km² grid cells dispersed over the whole country, and adopting copromicroscopic analyses allowing the detection of endoparasites. The aim of this study was to investigate the large-scale distribution of *A. vasorum* and other helminths in the Swiss fox population, to identify environmental factors enhancing their presence and therewith to assess the threat of infection for the Swiss dog population.

Materials and methods

Research area and sampling methods

Fox feces were collected in 72 localities distributed across Switzerland (total area 41'285 km²). We selected 54 localities in the low lands of Switzerland (< 700 m asl) and 18 in higher altitudes (range 700–1'800 m, mean 929 m, median 800 m). A total of 58 localities were situated in the north and 14 in the south of the Alpine crest. In all but one locality, where for topographic reasons only one plot could be selected, three sampling plots were determined. Each sample plot of such a triplet corresponded to a grid cell of the Swiss coordinate system which is based on 1-km² grid cells (<http://map.geo.admin.ch>). Each of the three grid cells of a triplet was situated in a distance of 1 km from the next grid cell. This sampling scheme was implemented in order to obtain samples from different fox families at each locality, as typical fox home range sizes vary between 20 and 30 ha in urban areas (Gloor et al. 2001) and are generally smaller than 1 km² for reproductive individuals in landscape with high food resources (Gloor et al. 2001; Henry et al. 2005). The plots were chosen to include agricultural area (cultivated land), forest and human settlements (at least 5% of each category). Fox feces were collected in nine localities between August 2010 and February 2011, in 51 between June 2011 and January 2012 and in 12 localities during both periods (data were pooled for the statistical analyses).

Based on methods in use for monitoring changes in the fox population density (Webbon et al. 2004), a modified technique for a standardized collection of fecal samples was developed. In detail, tracks of 5 to 6 km per each km² were followed using GPS (Garmin, GPSMAP® 60CSx) and the exact positions of fecal samples were logged. The scats were judged concerning their odor, the spot where they were discovered (i.e., in clearly visible and, possibly, elevated spots, characteristic of foxes for territorial marking; or near a cluster of rodents or near carrion indicating successful food recovery), and with details about their content (fruits, pits, bones, hair)-based operating experience and internal documentation to confirm identity and exclude non-fox feces. Samples with no clear characteristics of fox scats were not collected.

Methods

Flotation-sieving method and morphological identification

In order to minimize health risks due to the potential presence of *E. multilocularis* eggs in fox feces, the collected samples were initially stored at $-80\text{ }^{\circ}\text{C}$ for 3 days, afterwards at $-20\text{ }^{\circ}\text{C}$ until examination. Fecal flotation was performed as

previously described for the isolation of eggs of *E. multilocularis* (Deplazes and Eckert 1996), with modifications. First, the samples were centrifuged at 1600g for 10 min, the supernatant was discarded and the sediment mixed with 8 ml of zinc chloride (specific gravity, 1.45). After vortexing, the samples were centrifuged at 1000g for 30 min, followed by two sieving steps: after pouring the supernatant through a 150- μm mesh sieve, the sediment was poured through a 21- μm mesh sieve, in which L1 were supposed to accumulate. Therefore, the material retained in the sieve was washed out with water and poured into a 3.5 ml petri dish. Parasitic stages were identified based on morphological structures (Deplazes et al. 2016) using an inverted light microscopy by a 320-fold magnification (Labovert FS, Leitz Wetzlar, Germany).

Molecular identification

Part (69/115) of the samples in which L1 of *A. vasorum* were morphologically identified were verified by a specific *A. vasorum*-PCR targeting a region of the second internal transcribed spacer (ITS-2) of *A. vasorum* (Jefferies et al. 2009) after DNA-isolation (Stefanic et al. 2004). Also from 110 samples with morphologically unidentified larvae, from 11 out of 317 samples containing morphologically identified L1 of *C. vulpis*, from one sample containing L1 of *C. vulpis* and contemporaneously non-parasitic soil nematode larvae, and from 18 samples containing soil nematode larvae only the same PCR was performed. In detail, the larvae were transferred into tubes and centrifuged for 1 min at 13'000 rpm. The supernatant was discarded and the residue was mixed with 200 μl of TE buffer (1 ml 1 M Tris- HCl pH 8.4, 0.2 ml 0.5 M EDTA pH 8.0, 98.8 ml aqua dest.) and swung twice with a swing mill (model MM 300, Retsch GmbH, 42,781 Haan, Germany) at 30 1/s for 1 min using 3 mm stainless steel balls (Schieritz & Hauenstein AG). Between and after these two swing steps, the samples were promptly centrifuged and cooled down. After incubating at 95 °C for 5 min, the samples were again cooled down, followed by centrifugation at 13'000 rpm for 1 min. Afterwards, 20 μl proteinkinase K and 200 μl of buffer AL (guanidine hydrochloride) were added, using a commercial kit for DNA purification (Qiamp DNA mini kit®, Qiagen, Hilden, Germany). The samples were incubated at 56 °C for 4 h, followed by step 5 of the manufacturer's protocol for purification from blood or body fluid, with a final elution volume of 200 μl . Conventional PCR assays were performed in a final reaction volume of 100 μl . Primers and conditions were used as described by Jefferies et al. (2009) with some adaptations: 35 cycles were performed in a Peltier thermal cycler (PTC-200, Bio-Rad Laboratories AG, Reinach, Switzerland). The final extension step at 72 °C lasted 10 min. A negative control (no DNA) and positive control, consisting of 2 μl PCR confirmed *A. vasorum*-DNA and 8 μl of water, were included in all tests.

Amplicons were detected on 1.5% agarose gels stained with Gel Red (Biotium, Chemie Brunschwig AG, Basel, Switzerland).

Thirteen samples in which the PCR results were not congruent with the microscopic findings were tested for the purpose of discovering an inhibition of the amplification reactions as described by Mathis et al. (1996), and a validation was done by a second PCR using the nematode primers NC1 and NC2, also targeting a part of ITS-2, as previously described (Romstad et al. 1997). The samples ($n = 5$) in which larvae were morphologically determined as L1 of *A. vasorum*, being negative in the specific *A. vasorum*-PCR but positive in the broader PCR for nematodes were sequenced. For this purpose, the amplicons detected on 1.5% agarose gels were excised and extracted following the protocol (MinElute handbook of Qiagen), the concentration measured by spectrophotometry (NanoDrop® ND-1000 spectrometer, Thermo Scientific) and submitted for sequencing (Synergene, Switzerland).

Spatial and statistical analysis

QuantumGIS version 1.8.0 (<http://qgis.org/>) was used to build maps and to characterize the 72 study areas in regard on the land use in these regions. Based on the Swiss Federal area statistics 2004/09, which differentiates 74 land use categories based on a one-ha grid, we calculated for each study area the percentage of urban and cultivated area (urbanized area: categories 20–68, cultivated land: categories 71–89, see Arealstatistik Schweiz on www.bfs.admin.ch). We further attributed to each study area a bioregion (Alps [including Northern, Western Central, and Eastern Central Alps], Central Plateau, Jura, and Southern Alps) corresponding to the biogeographic regions set by Gonseth et al. (2001) and used the mean number of fecal samples found per km^2 (feces frequency) as rough indicator of fox density.

We used an ANCOVA (SPSS 20.0., IBM) to assess environmental factors affecting the presence of the investigated parasites in the fecal samples of the study areas ($n = 72$ areas) with (1) the bioregion as a fixed factor and (2) altitude (mean height above sea level), (3) the percentages of urbanized area and (4) cultivated land, and (5) the feces frequency as the covariates. The number of fecal sample per sampling site was used as weighting factor. Exact binomial 95% confidence intervals (CI) for means of binomial variables were calculated according to the method of Clopper and Pearson (1934). Significance levels were set at $p < 0.05$.

Data availability statement The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Results

A total of 1481 fox fecal samples were collected in the 72 explored areas. Overall, our testing procedures revealed 130 (8.8%, 95% CI 7.4–10.3%) *A. vasorum* positive samples (distribution, see Fig. 1) all from areas below 900 m asl (see Fig. 2), detected by either morphological identification of L1 and/or by genetic analysis. In 7.8% (115/1481) L1 of *A. vasorum* were morphologically identified. Sixty-nine of these samples were tested with the specific *A. vasorum*—PCR, and 60 yielded a positive result (87%, CI 76.7–93.9%). In five out of the remaining nine non-confirmed samples, the broad nematode PCR was positive and sequence analysis confirmed the presence of *A. vasorum* DNA in three samples and the presence of soil nematodes or yeast in another two samples. In none of the samples, inhibition was detected. For the lastly remaining four samples, the material was not sufficient for genetic confirmation. In summary, genetic analyses were able to confirm the morphological identification in 63/69 samples (91.3%, CI 82.0–97.7%). In addition, the specific PCR for detection of *A. vasorum* DNA performed with 110 samples containing morphologically undefined larvae revealed 12 further *A. vasorum* positive samples. Supplementary two *A. vasorum* positive samples were identified by PCR when testing 11 samples containing larvae morphologically identified as L1 of *C. vulpis*, indicating a missed mixed infection with the two lungworms, or a morphological misinterpretation, which could not finally be elucidated. Lastly, 19 samples containing larvae morphologically specified as soil nematodes were all negative in the

specific *A. vasorum*—PCR, with one exception: a sample containing contemporaneously larvae morphologically defined as L1 of *C. vulpis* and soil nematodes was also PCR-positive for *A. vasorum*.

Further identified helminth stages are listed in Table 1. The ANCOVA for assessing the effect of the investigated factors (s. Methods) on the parasite occurrence in fox feces per sampling site ($n = 72$) revealed significant effects of altitude, bioregion, and cultivated land (Table 2). All three factors significantly affected the occurrence of *A. vasorum* (bioregion: $F = 3.6$, $p = 0.018$; altitude: $F = 9.8$, $p = 0.003$; cultivated land: $F = 4.2$, $p = 0.043$) revealing significantly higher prevalence rates of *A. vasorum* in fecal samples from the Central Plateau (Table 3), in regions below 400 m asl (Fig. 2) and in areas with < 50% of cultivated land (Fig. 3). The occurrence of Taeniidae was affected by the percentage of cultivated land with higher occurrence of positive samples in areas with > 50% of cultivated land ($F = 7.0$, $p = 0.010$) (Fig. 3), whereas the bioregion affected the prevalence rates of *C. vulpis* ($F = 11.7$, $p < 0.001$), with higher prevalence rates in the Central Plateau and lower ones south of the Alps, and of *Toxocara* spp. with lower prevalence rates south of the Alps ($F = 2.8$; $p = 0.047$) (Table 3). *Capillaria* spp. were influenced by the bioregion ($F = 4.4$, $p = 0.007$; higher prevalence rates in the Central Plateau, Table 3) and the percentage of cultivated land ($F = 4.0$, $p = 0.049$; higher rates in areas with < than 50% cultivated land, Fig. 3). No significant effect of the investigated factors could be found for *Trichuris vulpis* and hookworm egg positive samples (Table 2).




Fig. 1 Sampling areas and occurrence of *Angiostrongylus vasorum* determined by analysis of fox feces within the four bioregions of Switzerland. Colors and symbol sizes reflect the number of collected samples and parasite frequency (s. legend)




Fig. 2 Percentage and 95% confidence intervals of fox fecal samples positive for *Angiostrongylus vasorum* collected at different altitudes

Discussion

Our study demonstrated how a sampling scheme for collecting fox feces, which was developed for ecological studies of fox populations (Putman 1984; Sadlier et al. 2004), can be applied for epidemiological investigations on fox parasites. In particular, fox scats findings included the registration of the exact route by GPS which allows the repetition of exactly the same pathway and therefore standardization, if repeated over time under the same conditions, without elimination of the animals. Fecal

Table 1 Occurrence of helminths determined in 1481 in fecal samples from fox

Parasite and parasitic stage	N (% and 95% confidence intervals)
<i>Angiostrongylus vasorum</i> (L1)	130 (8.8%; 7.4–10.3%)
<i>Crenosoma vulpis</i> (L1)	317 (21.4%; 19.3–21.6%)
<i>Toxocara canis</i> (eggs)	179 (12.1%; 10.5–13.9%)
Taeniidae (eggs)	156 (10.5%; 9.0–12.2%)
<i>Capillaria</i> spp. (eggs)	123 (8.3%; 6.9–9.8%)
<i>Trichuris vulpis</i> (eggs)	82 (5.5%; 4.4–6.8%)
Hookworms (eggs)	79 (5.3%; 4.2–6.6%)
<i>Toxascaris leonina</i> (eggs)	19 (1.3%; 0.8–2.0%)
<i>Strongyloides</i> sp. (eggs)	6 (0.4%; 0.1–0.9%)

droppings of foxes are frequently placed in visible sites representing territorial marks, but also close to places where foxes were successful in food search; this, together with fecal contents reflecting fox diets, contribute to the correct identification of fox scats (Webbon et al. 2004). To minimize variations, we chose 1-km² areas with alike ecological structure and collected only in autumn and winter, in order to prevent the scats from being exposed to high temperatures. However, a limitation of the study, in contrast to dog data collection, is that the fox scats do not necessarily represent individual animals (unless they are collected from hunted animals, as recently described, Cabanova et al. 2018), as some of the collected samples may have originated from defecations of the same animal(s). This needs to be considered when speaking of “prevalence”. However, with our approach of defining three sampling plots at each site (s. methods), we ensured that our samples originated from different fox families of the investigated study areas.

Our results demonstrate the large-scale occurrence of *A. vasorum* within the Swiss fox population. Serological tests developed for diagnostics in individual dogs and highly suitable for mass-screening lately suggested the broad presence of the parasite in the country, with regional prevalences concerning antibody detection varying from 0 to 7% (Lurati et al. 2015). This latter prevalence was obtained from dogs from the High Rhine area, being part of the Central Plateau where interestingly

Table 2 *P* values for the environmental factors, evaluated by ANCOVA, affecting the occurrence of fox helminths as determined by examinations of 1481 fecal samples of foxes from 72 localities)

Parasite	Bioregion	Altitude	Urban area	Cultivated land	Feces frequency
<i>Angiostrongylus vasorum</i>	0.018* (3.60)	0.003* (9.76)	0.834 (0.04)	0.043* (4.25)	0.135 (2.3)
Taeniidae	0.113 (2.07)	0.282 (1.18)	0.135 (2.29)	0.010* (7.05)	0.942 (0.01)
<i>Crenosoma vulpis</i>	0.000* (11.68)	0.425 (0.65)	0.704 (0.15)	0.312 (1.04)	0.569 (0.33)
<i>Capillaria</i> spp.	0.007* (4.38)	0.803 (0.06)	0.104 (2.71)	0.049* (4.04)	0.399 (0.72)
<i>Toxocara canis</i>	0.047* (2.80)	0.705 (0.15)	0.871 (0.03)	0.458 (0.56)	0.981 (0)
<i>Trichuris vulpis</i>	0.127 (1.97)	0.953 (0)	0.429 (0.64)	0.18 (1.84)	0.62 (0.25)
Hookworms	0.38 (1.04)	0.867 (0.03)	0.406 (0.7)	0.171 (1.92)	0.72 (0.13)

In parentheses, the corresponding *F* values are given; significant values ($p < 0.05$) are marked with an asterisk

the highest percentage of *A. vasorum* positive fox feces were detected in this study (11.4%, CI 9.4–13.8). Towards 90% of the positive samples from dogs originated from areas below 700 m asl (Lurati et al. 2015), while in the here presented study, 97.4% of the positive fox samples were collected below 700 m asl, confirming this trend. The prevalence rate was highest at the lowest altitudes (below 400 m asl), while above 900 m asl, none was positive. Foxes may be present at altitudes up to 2500 m asl (Hausser 1995), while the observed decreasing *A. vasorum* prevalence rates with increasing altitude is suggested to correlate with lower availability of susceptible snail and slug species that act as intermediate hosts (Boschi 2011; Lurati et al. 2015). In fact, previous studies showed that the higher the fox habitat, the less Mollusca are part of fox' diet (Hartova-Nentvichova et al. 2010). The temperature was furthermore considered relevant for the survival of L1 in the environment before transmission to the intermediate hosts (Ferdushy and Hasan 2010). Accordingly, altitude and cold temperatures were identified as limiting factors for *A. vasorum* occurrence (Jeffery et al. 2004; Lurati et al. 2015).

Besides lower altitudes, the occurrence of *A. vasorum* in fox samples was higher in areas with less than 50% of cultivated land. A recent study investigated the occurrence of *A. vasorum* and *C. vulpis* in dogs from Germany by correlating the post-codes of dog locations with a digital landscape model; interestingly, among 12 differently defined land-use classes, “agricultural field” was identified as a protective factor, meaning that dogs living in such areas were at lower risk of infection with *A. vasorum*. Prevalence differences were attributed to

different intermediate host spectra potentially preferring different habitat conditions (Maksimov et al. 2017). In studies performed with 571 frozen fecal samples from red foxes hunted in Slovakia, *A. vasorum* was also detected in areas where temperature reaches up to 25 °C in summer and where the average winter temperature drops below –10 °C, confirming great climatic adaptability of the parasite (Cabanova et al. 2018). Furthermore, in a univariable regression model of environmental variables, the distribution of none of the examined variables influenced the distribution of *C. vulpis*, while *A. vasorum* was identified to occur by trend in dry areas with arable land, avoiding humid areas and forests; however, the corresponding interpretations were considered uncertain because of weak relationships (Cabanova et al. 2018). We hypothesize that in cultivated land, in particular in conventionally used agricultural areas where agrochemical products (fertilizers, herbicides, and pesticides) are employed, food range and shelter options are reduced for snails and slugs, therefore representing a less suitable habitat. This is supported by a study performed in an area of the Central Plateau of Switzerland (canton of Aargau), where the effectiveness of agri-environment schemes (that provide financial incentives for farmers when adopting environmentally friendly agricultural practices (Kleijn et al. 2006)) on biodiversity was evaluated: After 5 years of implementing ecological compensated areas, species richness of less mobile species such as plants and snails significantly increased (Roth et al. 2008).

For successful endemic establishment of *A. vasorum*, overlapping habitats of intermediate and definitive hosts are required. As

Table 3 Occurrence (percentage and 95% confidence intervals, CI) of the four helminths for which significant differences were detected between the four bioregions of Switzerland (s. Table 2)

Parasite	Alps (% , CI)	Central Plateau (% , CI)	Jura (% , CI)	Southern Alps (% , CI)
<i>Angiostrongylus vasorum</i>	1.4 (0.3–4.1)* ^L	11.4 (9.4–13.8)* ^H	1.6 (0.2–5.7)* ^L	6.4 (3.2–11.2)
<i>Capillaria</i> spp.	4.7 (2.3–8.5)	10.1 (8.1–12.3)* ^H	1.6 (0.2–5.7)* ^L	5.2 (2.4–9.7)
<i>Toxocara canis</i>	10.9 (7.0–15.9)* ^H	15.0 (12.7–17.6)* ^H	6.4 (2.8–12.2)	2.3 (0.6–5.8)* ^L
<i>Crenosoma vulpis</i>	10.4 (6.7–15.4)* ^L	28.9 (25.9–32.0)* ^H	4.8 (1.8–10.2) * ^L	0.0 (0.0–1.7)* ^L

Bioregions with the *^H highest and *^L lowest prevalences due to non-overlapping confidence intervals ($p < 0.05$)




Fig. 3 Percentage and 95% confidence intervals of samples positive for *Angiostrongylus vasorum*, Taeniidae and *Capillaria* spp. in sampling sites with less and more than 50% of cultivated land. Percentages for all

three helminths differ significantly between areas with less and more than 50% cultivated land ($p < 0.05$, see ANCOVA results of Table 2)

foxes (in opposition to dogs) are not subjected to anthelmintic treatments and may feed more frequently on snails and slugs, foxes are believed to play the most important role in *A. vasorum* transmission and establishment in suitable areas and therefore represent a highly useful study subject for evaluating the risk for canine angiostrongylosis. This is supported by constantly higher prevalences of *A. vasorum* in fox populations compared with dog data (reviewed in Koch and Willesen 2009). Possibly, the role of dogs is instead substantial for the fast and larger spread of *A. vasorum* by translocations of infected dogs into new areas: the simultaneous presence of dogs and foxes in areas with access to shared intermediate host (Deplazes et al. 2004; Lurati et al. 2015) may facilitate novel establishments of this parasite. The high densities of dogs and foxes in Switzerland can therefore explain the wide-spread establishment and the high prevalence rate of *A. vasorum* within a relatively short time span, especially in the densely populated and urbanized Central Plateau.

The fox lungworm *C. vulpis* occurs in many regions of the world in bronchi, bronchioles and trachea of wild and domestic canids (Deplazes et al. 2016; Nevarez et al. 2005). Prevalence in foxes may reach up to 87% (Newfoundland, Jeffery et al. 2004), but prevalence in Europe mostly ranges between 10 and 25% (Al-Sabi et al. 2014; Rajkovic-Janje et al. 2002; Tolnai et al. 2015), including the here detected prevalence rate of 21.4% (CI 19.3–21.6%). Its life cycle, like *A. vasorum*, involves snails and slugs as intermediate hosts (Stockdale and Hulland 1970); this constellation was suggested to be the reason behind the clumped and correlated occurrence of both parasites (Taylor et al. 2015; Tolnai et al. 2015). Accordingly, the spatial analysis showed a significantly higher occurrence in the Central Plateau, comparable to *A. vasorum*, but contemporaneously a decreased occurrence south of the Alps, as no positive fox sample was identified. First stage larvae of *C. vulpis* were shown to be particularly resistant to harsh cold environmental conditions (Jeffery et al. 2004), supporting findings of higher prevalence in foxes in

colder climates of northern Europe (i.e., Germany, 32.3%, Schug et al. 2018) than in southern Europe (i.e., Italy, 14.7%, Magi et al. 2009).

Also the frequency of fox feces with *Capillaria* spp. eggs was significantly higher in the Central Plateau compared to the other bioregions (s. Table 3), and higher in areas with < than 50% cultivated land (s. Fig. 3). Capillariid infections of carnivores are mainly connected with ingestion of earthworms (although there are still open questions regarding other ways of transmission, Deplazes et al. 2016) which in analogy to snails may be less prevalent in cultivated areas. As no further differentiation was performed, the potential presence of *Capillaria* eggs derived from hunted birds or rodents (intestinal passage) should be also considered.

Toxocara egg prevalence in fox feces (12.1%, CI 10.5–13.9) was lower compared with several other European fox studies, in which the animals were necropsied and adult worms detected: the percentage of positive animals ranged from approximately 25% up to more than 80% (Borgsteede 1984; Bruzinskaite-Schmidhalter et al. 2012; Saeed et al. 2006; Willingham et al. 1996; Wolfe et al. 2001). Accordingly, in previous studies in dissected foxes, i.e., from western Switzerland and Zurich, a prevalence of 44.3% and 47.4%, respectively, was observed (Hofer et al. 2000; Réperant et al. 2007), indicating higher sensitivity of necropsy for detection of roundworm infections.

The same may account for tapeworms: *E. multilocularis* and *Taenia* sp. reached prevalences up to 52.1% and 54.3% in culled foxes from Geneva and up to 66.7% and 16.5% in Zurich, respectively (Hofer et al. 2000; Réperant et al. 2007). Here, taeniid eggs were not differentiated and may therefore include *Taenia* spp. and *Echinococcus multilocularis* as well, but a prevalence of 10.5% (CI 9.0–12.2%) indicate that the adopted coproscopic methods were far less sensitive for tapeworm detection compared with necropsy methods (Deplazes and Eckert 2001). Interestingly, there was a higher occurrence

of positive samples in areas with > 50% of cultivated land. Microtids like *Microtus arvalis* and *Arvicola scherman*, which both are considered as crucial intermediate hosts for *E. multilocularis* (Beerli et al. 2017; Burlet et al. 2011; Guerra et al. 2014), can profit from cultivated land (mainly meadows and pastures), but also other taeniid species (*T. pisiformis*, *T. crassiceps*, *T. polycantha*, and *T. serialis*) have rodent intermediate hosts and canid definitive hosts (Deplazes et al. 2016). Overall, although prevalence rates obtained by fecal examination were lower than the ones obtained by necropsy, the large-scale presence of *Toxocara* and of taeniid eggs in the environment support the important role of wild carnivores for transmission to domestic animals and their zoonotic potential (Otranto et al. 2015).

Additional eggs that were identified in the investigated fox samples were, in descending order, eggs of *T. vulpis* (5.5%, CI 4.4–6.8%), hookworms (5.3%; CI 4.2–6.6%), *T. leonina* (1.3%; CI 0.8–2.0%), and *Strongyloides* sp. (0.4%; CI 0.1–0.9%). They were not significantly associated with any of the investigated factors. For *T. vulpis*, regional studies from Germany or France based on necropsies revealed prevalences up to 10.3% (Petavy et al. 1990; Pilarczyk et al. 2005), while others reported prevalences below 3% (Poglayen et al. 1985; Richards et al. 1995; Vergles Rataj et al. 2013); this places our results as intermediate, taking into account that fecal examination was performed. Ancylostomatids in Swiss foxes are represented by *Uncinaria stenocephala*: adult parasites were identified in a high prevalence (64.8% and 78.2%) in necropsied foxes from Zurich and Geneva, respectively (Hofer et al. 2000; Réperant et al. 2007). In other European countries, the prevalence ranged from 26 to 80% (Al-Sabi et al. 2014; Rajkovic-Janje et al. 2002), also based on necropsies. *T. leonina* may also infect up to 40% of the foxes in other European countries (Al-Sabi et al. 2014; Bruzinskaite-Schmidhalter et al. 2012) including Switzerland (prevalence: 37.3%, Réperant et al. 2007). *Strongyloides* spp. were instead only rarely reported in foxes; generally, in wild canids, they are allocated to *Strongyloides stercoralis*, that is shared not only with dogs but also with humans and other wild canids (Thamsborg et al. 2017). In the Netherlands, fecal examination revealed a prevalence of 0.7% (Borgsteede 1984). In Switzerland, *S. stercoralis* was, to our knowledge, not previously reported in foxes and occurred only rarely in dogs that were mostly imported from Eastern European countries (M. Schnyder, personal communication). In this study, no genetic analyses were performed to confirm the findings.

Fecal examination of fox samples delivers indications on spatial trends for the presence of a given parasite on a large-spatial scale without affecting the population under study by culling animals. For lungworms, larval migration methods are recommended for fresh samples only and therefore not feasible with fox feces collected in the field, while FLOTAC (Schnyder et al. 2010) or the here presented filtration method

with larval accumulation can be alternatively used with older samples (Cabanova et al. 2018). This latter method is based on sieving steps and relies on different sizes of the parasitic stages; it was shown suitable for the detection of small taeniid eggs (Deplazes and Eckert 1996) and was therefore supposed to also separate lungworm larvae and eggs from other material, allowing the contemporaneous detection of helminth and cestode eggs and larvae. However, we observed that due to the flotation solution and possibly the age of the samples, the identification of larvae was often challenging, though 96.2% of selected samples morphologically identified as *A. vasorum* L1 were confirmed by PCR, by which additional positive *A. vasorum* samples were identified (that would have been missed with morphological identification only). Limiting factors for successful PCRs are the quantity of available parasite DNA, DNA denaturation, or PCR inhibition due to fecal material.

In summary, our study confirms that Swiss foxes harbor a wide variety of parasites including zoonotic parasites and parasites representing a risk for the health of the dog population. Among these, *A. vasorum* occurs in all bioregions of Switzerland but is more prevalent in the Central Plateau, concentrated in regions below 700 m asl and less present in areas with a high percentage of cultivated land. The use of the here presented standardized method for fecal collection allows follow-up surveys by collection of fox scats without culling interventions and therefore can be regarded as a useful procedure to assess and monitor over time the infection pressure for dog populations on a large scale.

Acknowledgments We would like to thank Jeannine Hauri, Lucia Kohler and Katharina Huggel for their support with laboratory work; Prof. Alexander Mathis and Dr. Felix Grimm for their advice regarding biomolecular issues; Dr. Iskender Ziadinov for his help with the fox sampling; and Dr. Ramon Eichenberger for support.

Compliance with ethical standards

This article does not contain any studies with human participants or animals performed by any of the authors.

Conflict of interest The authors declare that they have no conflict of interest.

References

- Alho AM, Schnyder M, Meireles J, Belo S, Deplazes P, Madeira de Carvalho L (2014) Preliminary results on the seroprevalence of *Angiostrongylus vasorum* and co-infection with *Dirofilaria immitis* in shelter dogs from Portugal. Parasit Vectors 7(Suppl 1):O26 <http://www.parasitesandvectors.com/content/7/S1/O26>
- Al-Sabi MN, Halasa T, Kapel CM (2014) Infections with cardiopulmonary and intestinal helminths and sarcoptic mange in red foxes from two different localities in Denmark. Acta Parasitol 59:98–107. <https://doi.org/10.2478/s11686-014-0214-6>

- Barutzki D, Schaper R (2009) Natural infections of *Angiostrongylus vasorum* and *Crenosoma vulpis* in dogs in Germany (2007–2009). *Parasitol Res* 105(Suppl 1):S39–S48
- Beerli O, Guerra D, Baltrunaite L, Deplazes P, Hegglin D (2017) *Microtus arvalis* and *Arvicola scherman*: key players in the *Echinococcus multilocularis* life cycle. *Front Vet Sci* 4:216. <https://doi.org/10.3389/fvets.2017.00216>
- Bolt G, Monrad J, Henriksen P, Dietz HH, Koch J, Bindseil E, Jensen AL (1992) The fox (*Vulpes vulpes*) as a reservoir for canine angiostrongylosis in Denmark. Field survey and experimental infections. *Acta Vet Scand* 33:357–362
- Borgsteede FH (1984) Helminth parasites of wild foxes (*Vulpes vulpes* L.) in the Netherlands. *Z Parasitenkd* 70:281–285
- Boschi C (2011) Die Schneckenfauna der Schweiz - Porträts aller Schweizer Schneckenarten. Haupt publisher, Berne
- Bruzinskaite-Schmidhalter R, Sarkunas M, Malakauskas A, Mathis A, Torgerson PR, Deplazes P (2012) Helminths of red foxes (*Vulpes vulpes*) and raccoon dogs (*Nyctereutes procyonoides*) in Lithuania. *Parasitology* 139:120–127. <https://doi.org/10.1017/S0031182011001715>
- Burlet P, Deplazes P, Hegglin D (2011) Age, season and spatio-temporal factors affecting the prevalence of *Echinococcus multilocularis* and *Taenia taeniaeformis* in *Arvicola terrestris*. *Parasit Vectors* 4:6. <https://doi.org/10.1186/1756-3305-4-6>
- Cabanova V, Miterpakova M, Druga M, Humnikova Z, Valentova D (2018) GIS-based environmental analysis of fox and canine lungworm distribution: an epidemiological study of *Angiostrongylus vasorum* and *Crenosoma vulpis* in red foxes from Slovakia. *Parasitol Res* 117:521–530. <https://doi.org/10.1007/s00436-017-5728-z>
- Chapman PS, Boag AK, Guitian J, Boswood A (2004) *Angiostrongylus vasorum* infection in 23 dogs (1999–2002). *J Small Anim Pract* 45:435–440
- Clopper CJ, Pearson ES (1934) The use of confidence or fiducial limits illustrated in the case of the binomial. *Biometrika* 26:404–413
- Cuillé J, Darraspen E (1930) De la Strongylose cardio-pulmonaire du chien. *Rev Gén de Méd Vét* 466:625–639 694–710
- Deplazes P, Eckert J (1996) Diagnosis of the *Echinococcus multilocularis* infection in final hosts. *Appl Parasitol* 37:245–252
- Deplazes P, Eckert J (2001) Veterinary aspects of alveolar echinococcosis—a zoonosis of public health significance. *Vet Parasitol* 98:65–87
- Deplazes P, Eckert J, Mathis A, von Samson-Himmelstjerna G, Zahner H (2016) Parasitology in veterinary medicine. Wageningen Academic Publishers, Wageningen
- Dodd K (1973) *Angiostrongylus vasorum* (Baillet, 1866) infestation in a greyhound kennels. *Vet Rec* 92:195–197
- Ferdushy T, Hasan MT (2010) Survival of first stage larvae (L1) of *Angiostrongylus vasorum* under various conditions of temperature and humidity. *Parasitol Res* 107:1323–1327
- Fischer C, Réperant LA, Weber JM, Hegglin D, Deplazes P (2005) *Echinococcus multilocularis* infections of rural, residential and urban foxes (*Vulpes vulpes*) in the canton of Geneva, Switzerland. *Parasite* 12:339–346. <https://doi.org/10.1051/parasite/2005124339>
- Gerrikagoitia X, Barral M, Juste RA (2010) *Angiostrongylus* species in wild carnivores in the Iberian Peninsula. *Vet Parasitol* 174:175–180
- Gillis-Germitsch N, Kapel CMO, Thamsborg SM, Deplazes P, Schnyder M (2017) Host-specific serological response to *Angiostrongylus vasorum* infection in red foxes (*Vulpes vulpes*): implications for parasite epidemiology. *Parasitology* 144:1–10. <https://doi.org/10.1017/S0031182017000427>
- Gloor S, Bontadina F, Hegglin D, Deplazes P, Breitenmoser U (2001) The rise of urban fox populations in Switzerland. *Mamm Biol* 66:155–164
- Gonseth Y, Wohlgenuth T, Sansonnens B, Buttler A (2001) Die biogeographischen Regionen der Schweiz. Erläuterungen und Einteilungsstandard. Umwelt Materialien 137, Bundesamt für Umwelt, Wald und Landschaft, Bern
- Gottstein B (2001) Lungenwurm *Angiostrongylus vasorum* bei einem Fuchs in der Schweiz. *BVET-Magazin* 1:22
- Grandi G, Osterman Lind E, Schaper R, Schnyder M (2017) Canine angiostrongylosis in Sweden: a summary of five-year diagnostic activity (2011–2015) and a nationwide seroepidemiological survey using combined detection of specific antibodies and circulating antigens by ELISAs. *Acta Vet Scand* 59:85. <https://doi.org/10.1186/s13028-017-0351-7>
- Guardone L, Schnyder M, Macchioni F, Deplazes P, Magi M (2013) Serological detection of circulating *Angiostrongylus vasorum* antigen and specific antibodies in dogs from central and northern Italy. *Vet Parasitol* 192:192–198. <https://doi.org/10.1016/j.vetpar.2012.10.016>
- Guerra D, Hegglin D, Bacciarini L, Schnyder M, Deplazes P (2014) Stability of the southern European border of *Echinococcus multilocularis* in the Alps: evidence that *Microtus arvalis* is a limiting factor. *Parasitology*:1–10. <https://doi.org/10.1017/S0031182014000730>
- Guilhon J (1963) Recherches sur le cycle évolutif du Strongle des vaisseaux du chien. *Bull Acad Vét Fr* 36:431–442
- Guilhon J, Bressou C (1960) Rôle des Limacides dans le cycle évolutif d'*Angiostrongylus vasorum* (Baillet, 1866). *C R Acad Sci* 251:2252–2253
- Hartova-Nentvichova M, Salek M, Cervený J, Koubek P (2010) Variation in the diet of the red fox (*Vulpes vulpes*) in mountain habitats: effects of altitude and season. *Mamm Biol* 75:334–340
- Hausser J (1995) Atlas der Säugetiere der Schweiz. Ed Birkhäuser, Basel
- Helm J, Roberts L, Jefferies R, Shaw SE, Morgan ER (2015) Epidemiological survey of *Angiostrongylus vasorum* in dogs and slugs around a new endemic focus in Scotland. *Vet Rec* 172:46. <https://doi.org/10.1136/vr.103006>
- Henry C, Poulle M-L, Roeder J-J (2005) Effect of sex and female reproductive status on seasonal home range size and stability in rural red foxes (*Vulpes vulpes*). *Écoscience* 12:202–209. <https://doi.org/10.2980/i1195-6860-12-2-202.1>
- Hofer S, Gloor S, Müller U, Mathis A, Hegglin D, Deplazes P (2000) High prevalence of *Echinococcus multilocularis* in urban red foxes (*Vulpes vulpes*) and voles (*Arvicola terrestris*) in the city of Zurich, Switzerland. *Parasitology* 120(Pt 2):135–142
- Houpin E, McCarthy G, Ferrand M, De Waal T, O'Neill EJ, Zintl A (2016) Comparison of three methods for the detection of *Angiostrongylus vasorum* in the final host. *Vet Parasitol* 220:54–58. <https://doi.org/10.1016/j.vetpar.2016.02.023>
- Humnikova Z, Miterpakova M, Mandelik R (2013) First autochthonous case of canine *Angiostrongylus vasorum* in Slovakia. *Parasitol Res* 112:3505–3508. <https://doi.org/10.1007/s00436-013-3532-y>
- Jefferies R, Morgan ER, Shaw SE (2009) A SYBR green real-time PCR assay for the detection of the nematode *Angiostrongylus vasorum* in definitive and intermediate hosts. *Vet Parasitol* 166:112–118
- Jeffery RA, Lankester MW, McGrath MJ, Whitney HG (2004) *Angiostrongylus vasorum* and *Crenosoma vulpis* in red foxes (*Vulpes vulpes*) in Newfoundland, Canada. *Can J Zool* 82:66–74
- Jolly S, Poncelet L, Lempereur L, Caron Y, Bayrou C, Cassart D, Grimm F, Lossou B (2015) First report of a fatal autochthonous canine *Angiostrongylus vasorum* infection in Belgium. *Parasitol Int* 64:97–99. <https://doi.org/10.1016/j.parint.2014.10.008>
- Kleijn D, Baquero RA, Clough Y, Diaz M, De Esteban J, Fernandez F, Gabriel D, Herzog F, Holzschuh A, Johl R, Knop E, Krüess A, Marshall EJ, Steffan-Dewenter I, Tschamtké T, Verhulst J, West TM, Yela JL (2006) Mixed biodiversity benefits of agri-environment schemes in five European countries. *Ecol Lett* 9:243–254; discussion 254–247. <https://doi.org/10.1111/j.1461-0248.2005.00869.x>

- Koch J, Willeßen JL (2009) Canine pulmonary angiostrongylosis: An update. *Vet J* 179:348–359
- Lurati L, Deplazes P, Hegglin D, Schnyder M (2015) Seroepidemiological survey and spatial analysis of the occurrence of *Angiostrongylus vasorum* in Swiss dogs in relation to biogeographic aspects. *Vet Parasitol* 212:219–226. <https://doi.org/10.1016/j.vetpar.2015.08.017>
- Magi M, Macchioni F, Dell'Omodarme M, Prati MC, Calderini P, Gabrielli S, Iori A, Cancrini G (2009) Endoparasites of red fox (*Vulpes vulpes*) in Central Italy. *J Wildl Dis* 45:881–885
- Majoros G, Fukar O, Farkas R (2010) Autochthonous infection of dogs and slugs with *Angiostrongylus vasorum* in Hungary. *Vet Parasitol* 174:351–354
- Maksimov P, Hermosilla C, Taubert A, Staubach C, Sauter-Louis C, Conraths FJ, Vrhovec MG, Pantchev N (2017) GIS-supported epidemiological analysis on canine *Angiostrongylus vasorum* and *Crenosoma vulpis* infections in Germany. *Parasit Vectors* 10:108. <https://doi.org/10.1186/s13071-017-2054-3>
- Mathis A, Deplazes P, Eckert J (1996) An improved test system for PCR-based specific detection of *Echinococcus multilocularis* eggs. *J Helminthol* 70:219–222
- Morgan ER, Tomlinson A, Hunter S, Nichols T, Roberts E, Fox MT, Taylor MA (2008) *Angiostrongylus vasorum* and *Eucoleus aerophilus* in foxes (*Vulpes vulpes*) in Great Britain. *Vet Parasitol* 154:48–57
- Nevarez A, Lopez A, Conboy G, Ireland W, Sims D (2005) Distribution of *Crenosoma vulpis* and *Eucoleus aerophilus* in the lung of free-ranging red foxes (*Vulpes vulpes*). *J Vet Diagn Investig* 17:486–489. <https://doi.org/10.1177/104063870501700516>
- Otranto D, Cantacessi C, Dantas-Torres F, Brianti E, Pfeiffer M, Genchi C, Guberti V, Capelli G, Deplazes P (2015) The role of wild canids and felids in spreading parasites to dogs and cats in Europe. Part II: helminths and arthropods. *Vet Parasitol* 213:24–37. <https://doi.org/10.1016/j.vetpar.2015.04.020>
- Papazahariadou A, Founta A, Papadopoulos E, Chliounakis S, Antoniadou-Sotiriadou K, Theodorides Y (2007) Gastrointestinal parasites of shepherd and hunting dogs in the Serres prefecture, northern Greece. *Vet Parasitol* 148:170–173
- Petavy AF, Deblock S, Prost C (1990) Epidemiology of alveolar echinococcosis in France. 1. Intestinal helminths in the red fox (*Vulpes vulpes* L.) from Haute-Savoie (in French). *Ann Parasitol Hum Comp* 65:22–27. <https://doi.org/10.1051/parasite/1990651022>
- Pilarczyk B, Balicka-Ramisz A, Ramisz A (2005) The occurrence of intestinal nematodes in red foxes in the Western Pomerania (in Polish). *Wiad Parazytol* 51:249–251
- Poglayen G, Guberti V, Leoni B (1985) Parasites present in foxes (*Vulpes vulpes*) of the province of Forlì (in Italian). *Parassitologia* 27:303–311
- Putman RJ (1984) Facts from faeces. *Mammal Rev* 14:79–97
- Rajkovic-Janje R, Marinculic A, Bosnic S, Benic M, Vinkovic B, Mihaljevic Z (2002) Prevalence and seasonal distribution of helminth parasites in red foxes (*Vulpes vulpes*) from the Zagreb County (Croatia). *Z Jagdwiss* 48:151–160
- Réperant LA, Hegglin D, Fischer C, Kohler L, Weber JM, Deplazes P (2007) Influence of urbanization on the epidemiology of intestinal helminths of the red fox (*Vulpes vulpes*) in Geneva, Switzerland. *Parasitol Res* 101:605–611. <https://doi.org/10.1007/s00436-007-0520-0>
- Richards DT, Harris S, Lewis JW (1995) Epidemiological studies on intestinal helminth parasites of rural and urban red foxes (*Vulpes vulpes*) in the United Kingdom. *Vet Parasitol* 59:39–51
- Roche MM, Kelliher DJ (1968) *Angiostrongylus vasorum* infestation in the dog: a case report. *Ir Vet J* 22:108–113
- Romstad A, Gasser RB, Monti JR, Polderman AM, Nansen P, Pit DS, Chilton NB (1997) Differentiation of *Oesophagostomum bifurcum* from *Necator americanus* by PCR using genetic markers in spacer ribosomal DNA. *Mol Cell Probes* 11:169–176
- Roth T, Amrhein V, Peter B, Weber D (2008) A Swiss agri-environment scheme effectively enhances species richness for some taxa over time agriculture. *Ecosyst Environ* 125:167–172. <https://doi.org/10.1016/j.agee.2007.12.012>
- Sadlier LMJ, Webbon CC, Baker PJ, Harris S (2004) Methods of monitoring red foxes *Vulpes vulpes* and badgers *Meles meles*: are field signs the answer? *Mammal Rev* 34:78–98
- Saeed I, Maddox-Hyttel C, Monrad J, Kapel CM (2006) Helminths of red foxes (*Vulpes vulpes*) in Denmark. *Vet Parasitol* 139:168–179
- Schnyder M, Maurelli MP, Morgoglione ME, Kohler L, Deplazes P, Torgerson P, Cringoli G, Rinaldi L (2010) Comparison of faecal techniques including FLOTAC for copromicroscopic detection of first stage larvae of *Angiostrongylus vasorum*. *Parasitol Res* 109:63–69
- Schnyder M, Tanner I, Webster P, Barutzki D, Deplazes P (2011) An ELISA for sensitive and specific detection of circulating antigen of *Angiostrongylus vasorum* in serum samples of naturally and experimentally infected dogs. *Vet Parasitol* 179:152–158
- Schnyder M, Schaper R, Bilbrough G, Morgan ER, Deplazes P (2013a) Seroepidemiological survey for canine angiostrongylosis in dogs from Germany and the UK using combined detection of *Angiostrongylus vasorum* antigen and specific antibodies. *Parasitology* 140:1442–1450. <https://doi.org/10.1017/S0031182013001091>
- Schnyder M, Schaper R, Pantchev N, Kowalska D, Szwedko A, Deplazes P (2013b) Serological detection of circulating *Angiostrongylus vasorum* antigen- and parasite-specific antibodies in dogs from Poland. *Parasitol Res* 112(Suppl 1):109–117. <https://doi.org/10.1007/s00436-013-3285-7>
- Schnyder M, Jefferies R, Schucan A, Morgan ER, Deplazes P (2015) Comparison of coprological, immunological and molecular methods for the detection of dogs infected with *Angiostrongylus vasorum* before and after anthelmintic treatment. *Parasitology* 142:1270–1277. <https://doi.org/10.1017/S0031182015000554>
- Schug K, Kramer F, Schaper R, Hirzmann J, Failing K, Hermosilla C, Taubert A (2018) Prevalence survey on lungworm (*Angiostrongylus vasorum*, *Crenosoma vulpis*, *Eucoleus aerophilus*) infections of wild red foxes (*Vulpes vulpes*) in Central Germany. *Parasit Vectors* 11:85. <https://doi.org/10.1186/s13071-018-2672-4>
- Segovia JM, Torres J, Miquel J (2004) Helminth parasites of the red fox (*Vulpes vulpes* L., 1758) in the Iberian Peninsula: an ecological study. *Acta Parasitol* 49:67–79
- Sigrist NE, Hofer-Inteewom N, Jud Schefer R, Kuemmerle-Fraune C, Schnyder M, Kutter APN (2017) Hyperfibrinolysis and hypofibrinogenemia diagnosed with rotational thromboelastometry in dogs naturally infected with *Angiostrongylus vasorum*. *J Vet Intern Med* 31:1091–1099. <https://doi.org/10.1111/jvim.14723>
- Staebler S, Ochs H, Steffen F, Naegeli F, Borel N, Sieber-Ruckstuhl N, Deplazes P (2005) Autochthonous infections with *Angiostrongylus vasorum* in dogs in Switzerland and Germany (in German). *Schweiz Arch Tierheilkd* 147:121–127
- Stefanic S, Shaikenov BS, Deplazes P, Dinkel A, Torgerson PR, Mathis A (2004) Polymerase chain reaction for detection of patent infections of *Echinococcus granulosus* ("sheep strain") in naturally infected dogs. *Parasitol Res* 92:347–351
- Stockdale PH, Hlland TJ (1970) The pathogenesis route of migration, and development of *Crenosoma vulpis* in the dog. *Pathol Vet* 7:28–42
- Taubert A, Pantchev N, Vrhovec MG, Bauer C, Hermosilla C (2008) Lungworm infections (*Angiostrongylus vasorum*, *Crenosoma vulpis*, *Aelurostrongylus abstrusus*) in dogs and cats in Germany and Denmark in 2003–2007. *Vet Parasitol* 159:175–180
- Taylor CS, Garcia Gato R, Learmount J, Aziz NA, Montgomery C, Rose H, Coulthwaite CL, McGary JW, Forman DW, Allen S, Wall R,

- Morgan ER (2015) Increased prevalence and geographic spread of the cardiopulmonary nematode *Angiostrongylus vasorum* in fox populations in Great Britain. *Parasitology* 142:1190–1195. <https://doi.org/10.1017/S0031182015000463>
- Thamsborg SM, Ketzis J, Horii Y, Matthews JB (2017) *Strongyloides* spp. infections of veterinary importance. *Parasitology* 144:274–284. <https://doi.org/10.1017/S0031182016001116>
- Tieri E, Pomilio F, Di Francesco G, Saletti MA, Totaro P, Troilo M, Menna S, Tampieri MP, Morelli D (2011) *Angiostrongylus vasorum* in 20 dogs in the province of Chieti, Italy. *Vet Ital* 47:77–88 65–76
- Tolnai Z, Szell Z, Sreter T (2015) Environmental determinants of the spatial distribution of *Angiostrongylus vasorum*, *Crenosoma vulpis* and *Eucoleus aerophilus* in Hungary. *Vet Parasitol* 207:355–358. <https://doi.org/10.1016/j.vetpar.2014.12.008>
- van Doorn DC, van de Sande AH, Nijssse ER, Eysker M, Ploeger HW (2009) Autochthonous *Angiostrongylus vasorum* infection in dogs in the Netherlands. *Vet Parasitol* 162:163–166
- Vergles Rataj A, Posedi J, Zele D, Vengust G (2013) Intestinal parasites of the red fox (*Vulpes vulpes*) in Slovenia. *Acta Vet Hung* 61:454–462. <https://doi.org/10.1556/AVet.2013.029>
- Webbon CC, Baker PJ, Harris S (2004) Faecal density counts for monitoring changes in red fox numbers in rural Britain. *J Appl Ecol* 41:768–779
- Willingham AL, Ockens NW, Kapel CM, Monrad J (1996) A helminthological survey of wild red foxes (*Vulpes vulpes*) from the metropolitan area of Copenhagen. *J Helminthol* 70:259–263
- Wolfe A, Hogan S, Maguire D, Fitzpatrick C, Vaughan L, Wall D, Hayden TJ, Mulcahy G (2001) Red foxes (*Vulpes vulpes*) in Ireland as hosts for parasites of potential zoonotic and veterinary significance. *Vet Rec* 149:759–763
- Wolff K, Eckert J, Leemann W (1969) Beitrag zur Angiostrongylose des Hundes. In: Dtsch Vet-med Ges, Zürich, 1969. Vortrag Tagung Fachgr. "Kleintierkrankheiten"

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.