



Contents lists available at ScienceDirect

Critical Reviews in Oncology / Hematology

journal homepage: www.elsevier.com/locate/critrevonc

Efficacy and safety of combination immunotherapy for malignant solid tumors: A systematic review and meta-analysis

Yuhan Wei^{a,1}, Qi Du^{a,1}, Xiaoyue Jiang^a, Li Li^a, Teng Li^a, Mengqi Li^a, Xueke Fan^b, Yingrui Li^c, Seyed Kariminia^d, Qin Li^{a,*}^a Department of Oncology, Capital Medical University Affiliated Beijing Friendship Hospital, Beijing, 100050, China^b Gastroenterology Department, Jincheng People's Hospital, Jincheng, 048000, China^c Biochemistry and Molecular Biology, Basic Medicine College, Shanxi Medical University, Taiyuan, 050001, China^d Molecular and Cellular Oncology, MD Anderson Cancer Center, 1515 Holcombe Blvd, Houston 77030, USA

ARTICLE INFO

Keywords:

Combination immunotherapy
Immune checkpoint blockade
PD-1/PD-L1
CTLA-4
Meta-analysis

ABSTRACT

Background: Combination immunotherapy has become an actively growing field of clinical investigation.**Methods:** We searched for clinical trials of combination immunotherapy and calculated the pooled hazard ratio (HR), odds ratio (OR) of clinical outcomes and safety by subgroups of different combination regimens.**Results:** Totally 28 clinical trials were analyzed. The study showed that the pooled HRs of overall survival and progression-free survival for combination therapy were 0.77 (95% CI: 0.70–0.84, $p < 0.001$) and 0.72 (95% CI: 0.66–0.79, $p < 0.001$) while the pooled OR of high-grade adverse effects was 1.45 ($p = 0.004$). Subgroup analysis showed that the HR of overall survival were 0.74 ($p = 0.005$), 0.79 ($p < 0.001$), 0.70 ($p = 0.003$) and 0.85 ($p = 0.052$) for immunotherapy combined with immunotherapy, chemotherapy, targeted therapy and radiotherapy group, respectively.**Conclusions:** The meta-analysis indicated that combination immunotherapy could bring more clinical benefits with increased high-grade adverse effects.

1. Introduction

In recent years, remarkable progress has been made in tumor treatment with the discovery of immunotherapy, which has been used to a wide range of solid tumors such as melanoma, non-small cell lung cancer (NSCLC), renal cell carcinoma (RCC), small cell lung cancer (SCLC), squamous-cell carcinoma of the head and neck, urothelial carcinoma and prostate cancer (Ferris et al., 2016; Kantoff et al., 2010; Motzer et al., 2015; Reck et al., 2016b; Robert et al., 2015a,b; Sharma et al., 2016). In 2013, *Science* magazine announced immunotherapy a turning point in cancer treatment (Couzin-Frankel, 2013). Immunotherapy stimulates and enhances antitumor responses to inhibit tumor growth or kill tumor cells by active or passive ways such as immune checkpoint blockade (ICB), antitumor vaccine, adoptive immunotherapy, cytokine, etc. Among them, ICB, which restores antitumor effects by blocking cytotoxic T lymphocyte antigen 4 (CTLA-4), programmed death 1 (PD-1), T cell immunoglobulin mucin 3 (TIM-3), lymphocyte activation gene 3 (LAG-3), indoleamine 2,3 dioxygenase (IDO) and other inhibitory receptor-ligand interactions, is most

promising due to its unprecedented durable responses and manageable adverse events (AEs) (Pennock and Chow, 2015). Particularly, anti-CTLA-4 antibodies (ipilimumab), anti-PD-1 antibodies (nivolumab and pembrolizumab) and anti-PD-L1 antibodies (atezolizumab, avelumab and durvalumab) have been approved for clinical use.

Despite the remarkable progress that has been achieved, the efficacy of immunotherapy is not entirely satisfactory. Even for ICB, there are approximately 60%–70% of patients who do not respond to monotherapy (Larkin et al., 2015; Larkins et al., 2017; Robert et al., 2015a,b; Topalian et al., 2012, 2014). Immune tolerance may be the main reason for the phenomenon. Therefore, combination strategies are necessary to overcome the resistance and enlarge the clinical utility of immunotherapy. It is demonstrated that traditional treatments such as chemotherapy, radiotherapy and targeted therapy have immunomodulatory effects (Demaria et al., 2015; Fukumura et al., 2018; Pfirschke et al., 2016) and combination therapies with immunotherapy may achieve synergistic effects and thus maximize clinical efficacy. Some combinatorial regimens are demonstrated statistically significant improvements in survival and have been approved in clinical

* Corresponding author at: Beijing Friendship Hospital, 95 Yong'an Road, Beijing 100050, China.

E-mail address: oncologistinbj@163.com (Q. Li).¹ Yuhan Wei and Qi Du contributed equally as first authors.

application (e.g., ipilimumab + nivolumab for melanoma and NSCLC, pembrolizumab + pemetrexed and platinum-based drug for non-squamous NSCLC) (Gandhi et al., 2018; Hodi et al., 2016; Wolchok et al., 2017).

Nevertheless, it remains controversial whether combination therapy with immunotherapy is superior to monotherapy indeed, because not all studies showed significant improvement (Hodi et al., 2014; Reck et al., 2016a) and some even showed reduction of efficacy (Hodi et al., 2010). Besides, the increased AEs and the added costs of combinatorial regimens make it doubtful. It is of great significance to explore the efficacy and safety of combination therapy from evidence-based medicine but there are few relevant studies. We conducted a systematic review and meta-analysis of clinical data that compared the combination immunotherapy with monotherapy and explored relevant mechanisms to guide the application of combination approaches.

2. Methods

2.1. Search strategy

The study was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statement (Moher et al., 2015). Three researchers (Yuhan Wei, Mengqi Li and Qi Du) searched for relevant studies on PubMed, EMBASE and the Central Registry of Controlled Trials of the Cochrane Library. We also searched abstracts from the meetings of American Society of Clinical Oncology (ASCO). Proposed MeSH terms were: “immunotherapy OR immune therapy OR immune checkpoint inhibitor OR immune checkpoint blockade OR immune vaccine OR nivolumab OR ipilimumab OR pembrolizumab OR atezolizumab OR tremelimumab OR avelumab OR durvalumab” AND “combination OR combined with OR plus” AND “tumor OR cancer OR carcinoma OR neoplasm OR malignancy OR sarcoma”. Qualifier was “clinical trial”. The language was restricted to English. The cut-off date was February 20, 2019.

2.2. Eligibility

The inclusion criteria was as follows: (1) prospective study; (2) phase II or III clinical trials; (3) the subjects of the trials must be patients with histologically or cytologically confirmed malignant solid tumor; (4) the experimental intervention group was immunotherapy + immunotherapy, immunotherapy + chemotherapy, immunotherapy + targeted therapy or immunotherapy + radiotherapy; (5) the study should analyze the efficacy and safety of combination immunotherapy versus monotherapy; (6) the study must present any of the following measurements: overall survival (OS), progression-free survival (PFS), objective response rate (ORR), disease control rate (DCR) and the rate of grade 3–5 AEs. Where multiple articles analyzed the same trial, the most recent data was used.

2.3. Data extraction

Relevant information was extracted independently by two reviewers (Yuhan Wei and Qi Du) according to a predefined form including study characteristics, patient characteristics, treatment regimens and outcome data. Data from relevant appendix was extracted as well. All disagreements were resolved by discussion with a third reviewer (Teng Li). The primary outcomes were OS, PFS and secondary outcomes were ORR, DCR and the rate of grade 3–5 AEs. An exploratory analysis included evaluating the survival benefit of combination immunotherapy according to PD-L1 expression.

2.4. Statistical analysis

Two reviewers (Yuhan Wei and Qi Du) independently assessed the risk of bias using the Cochrane Reviewer Handbook. Any discrepancy

was resolved by consensus. The evaluation of the risk of bias was conducted by Review Manager (RevMan5.3; The Nordic Cochrane Centre, The Cochrane Collaboration, Copenhagen, Denmark).

Statistical analysis, forest plots, detection of publication bias and sensitivity analysis were done with Stata SE 12 (Stata Corporation, College Station, TX, USA). We calculated the pooled hazard ratio (HR) and 95% CI for PFS and OS and the pooled odds ratio (OR) and 95% CI for ORR and DCR and the rate of grade 3–5 AEs. An exploratory analysis evaluating the pooled HR of OS according to PD-L1 expression bounded by 1% was done as well. I-squared (I^2) tests were used to evaluate the heterogeneity between trials. Either $p < 0.1$ or $I^2 > 50\%$ indicated the existence of heterogeneity, then the randomized-effects model was used. Otherwise, the fixed-effects model was used. $p < 0.05$ was considered statistically significant. The publication bias was assessed by both Bag's test and Egger's test.

3. Results

3.1. Search results

A total of 5462 original articles as well as 452 ASCO meeting abstracts were obtained from electronic searches. After screening, duplicates and irrelevant studies were excluded and 24 original articles and 4 ASCO conference abstracts remained for final meta-analysis (Fig. 1).

3.2. Main characteristics and the risk of bias in the included studies

Totally 28 clinical trials published from 2011 to 2019 involving 14,070 patients were identified in the systematic evaluation, including seven immunotherapy + immunotherapy (D'Angelo et al., 2018; Hodi et al., 2016, 2014; Hodi et al., 2010; Motzer et al., 2018a,b; Peters et al., 2016; Wolchok et al., 2017), thirteen immunotherapy + chemotherapy (eleven full-text articles (Borghaei et al., 2019; Gandhi et al., 2018; Govindan et al., 2017; Hersh et al., 2011; Horn et al., 2018; Lynch et al., 2012; Paz-Ares et al., 2018; Reck et al., 2013, 2016a; Robert et al., 2011; Schmid et al., 2018) and two ASCO meeting abstracts (Borghaei et al., 2018; Jotte et al., 2018)), five immunotherapy + targeted therapy (four full-text articles (McDermott et al., 2018; Motzer et al., 2019; Rini et al., 2019; Socinski et al., 2018) and one ASCO meeting abstracts (Motzer et al., 2018a,b) and three immunotherapy + radiotherapy (two full-text articles (Kwon et al., 2014; Slovin et al., 2013) and one ASCO meeting abstracts (Theelen et al., 2018)). There were seven different tumor types such as NSCLC (N = 9), melanoma (N = 6), RCC (N = 5), SCLC (N = 4), prostate cancer (N = 2), breast cancer (N = 1) and sarcoma (N = 1). There were 21 trials done in first-line settings and seven with second or additional lines of therapy (Table 1, Supplementary Table S1). The assessment of risk of bias was presented in Supplementary Fig. S1.

3.3. Efficacy

Among 28 identified studies, 19 trials reported OS with 23 comparisons; 23 trials reported PFS with 27 comparisons; data on ORR were available in 27 trials with 31 comparisons; 21 trials with 25 comparisons provided data of DCR. Randomized-effects model was used in all measurements because of the significant heterogeneity.

Overall analysis indicated that combination immunotherapy could significantly improve OS (HR = 0.77, 95% CI: 0.70–0.84, $p < 0.001$), PFS (HR = 0.72, 95% CI: 0.66–0.79, $p < 0.001$), ORR (OR = 1.79, 95% CI: 1.45–2.21, $p < 0.001$) and DCR (OR = 1.32, 95% CI: 1.08–1.63, $p = 0.008$) in comparison to monotherapy.

Subgroup analysis demonstrated that immunotherapy + immunotherapy was associated with better OS (HR = 0.74, 95% CI: 0.59–0.91, $p = 0.005$), PFS (HR = 0.72, 95% CI: 0.53–0.97, $p = 0.031$), ORR (OR = 2.33, 95% CI: 1.33–4.05, $p = 0.003$) and DCR (OR = 1.54, 95% CI: 1.03–2.29, $p = 0.034$) compared to monotherapy.




Fig. 1. Flow diagram representing the selection process.

Immunotherapy + chemotherapy could significantly prolong OS (HR = 0.79, 95% CI: 0.70–0.89, $p < 0.001$), PFS (HR = 0.72, 95% CI: 0.65–0.80, $p < 0.001$), improve ORR (OR = 1.56, 95% CI: 1.22–2.01, $p < 0.001$) when compared with immunotherapy or chemotherapy alone, but didn't increase DCR (OR = 1.10, 95% CI: 0.78–1.53, $p = 0.591$). Immunotherapy + targeted therapy brought more clinical benefits in OS (HR = 0.70, 95% CI: 0.55–0.88, $p = 0.003$), PFS (HR = 0.73, 95% CI: 0.64–0.84, $p < 0.001$) and ORR (OR = 1.88, 95% CI: 1.28–2.78, $p = 0.001$) and DCR (OR = 1.58, 95% CI: 1.28–1.95, $p < 0.001$) than targeted therapy. In addition, PFS (HR = 0.69, 95% CI: 0.60–0.80, $p < 0.001$) was significantly longer in the immunotherapy + radiotherapy group than that in monotherapy group, while OS improvement was marginally significant (HR = 0.85, 95% CI: 0.72–1.00, $p = 0.052$), ORR (OR = 0.87, 95% CI: 0.05–14.78, $p = 0.921$) and DCR (OR = 1.56, 95% CI: 0.59–4.12, $p = 0.369$) were not statistically improved (Fig. 2 and Fig. 3).

When grouped by tumor types, OS was significantly prolonged in melanoma, RCC and NSCLC groups (pooled HR were 0.75, 0.64 and 0.76, respectively). Combination immunotherapy was associated with longer PFS in RCC, NSCLC, breast cancer, SCLC and prostate cancer groups (pooled HR were 0.77, 0.66, 0.80, 0.81 and 0.70, respectively). For melanoma, RCC, NSCLC and breast cancer groups, the pooled OR of ORR were 2.30, 1.91, 1.91 and 1.50, which were superior to monotherapy (Supplementary Fig. S2, S3).

Four RCTs were available for analysis of OS according to PD-L1

expression bounded by 1%. For patients of PD-L1 < 1%, the pooled HR of survival was 0.66 (95% CI: 0.55–0.80, $p < 0.001$). For patients of PD-L1 $\geq 1\%$, the pooled HR of survival was 0.53 (95% CI: 0.43–0.64, $p < 0.001$) (Supplementary Fig. S5).

3.4. Toxicity

In total 22 trials reported the incidence of grade 3–5 AEs. Randomized-effects model was used because of the substantial heterogeneity. It showed that the rate of grade 3–5 AEs in the combination group was significantly higher than that in the monotherapy group (OR = 1.45, 95% CI: 1.12–1.86, $p = 0.004$).

According to the subgroup analysis, immunotherapy + immunotherapy, immunotherapy + chemotherapy and immunotherapy + targeted therapy were associated with higher risk of the incidence of grade 3–5 AEs with the pooled OR of 1.81, 1.51 and 1.16, respectively. However, the rate of grade 3–5 AEs did not increase in immunotherapy + radiotherapy (OR = 0.93). Moreover, melanoma and NSCLC groups had higher incidence of grade 3–5 AEs. Main adverse events included general adverse events related to immune activation (diarrhea, fatigue and rash) and organ specific adverse events (colitis, hepatitis, pneumonia, hypothyroidism and hypophysitis) (Baxi et al., 2018), among which the rate of grade 3–5 diarrhea, fatigue and colitis drastically increased in all combination regimens (Fig. 4).

Table 1
Main characteristics of included studies.

| Author | Year | Phase | Tumor | Line | Sample Size | Age | Interventions | |
|--------------|------|-------|--------------------|--------|-------------|-----|------------------------|---|
| Hodi FS | 2016 | II | Melanoma | 1 L | Arms | N | Nivolumab + ipilimumab | |
| | | | | | Study | 95 | | 64 |
| Wolchok JD | 2017 | III | Melanoma | 1 L | Control | 47 | Ipilimumab | |
| | | | | | Study | 314 | 61 | Nivolumab + ipilimumab |
| | | | | | Control | 316 | 60 | Nivolumab |
| Antonia SJ | 2016 | I/II | SCLC | 2L | Control | 315 | Ipilimumab | |
| | | | | | Study | 54 | 61 | Nivolumab + ipilimumab |
| | | | | | Control | 98 | 63 | Nivolumab |
| Motzer RJ | 2018 | III | RCC | 1 L | Study | 550 | 62 | Nivolumab + ipilimumab |
| | | | | | Control | 546 | 62 | Sunitinib |
| Hodi FS | 2010 | III | Melanoma | 2L | Study | 403 | 55.6 | Ipilimumab + gp100 |
| | | | | | Control | 137 | 56.8 | Ipilimumab + placebo |
| | | | | | Control | 136 | 57.4 | Gp100 |
| Hodi FS | 2014 | II | Melanoma | 1 L/2L | Study | 123 | 61 | Ipilimumab + sargramostim |
| | | | | | Control | 122 | 64 | Ipilimumab |
| D'Angelo SP | 2018 | II | Sarcoma | 2L | Study | 42 | 57 | Nivolumab + ipilimumab |
| | | | | | Control | 43 | 56 | Nivolumab |
| Borghaei H* | 2018 | III | NSCLC | 1 L | Study | 177 | – | Nivolumab + platinum doublet chemotherapy |
| | | | | | Control | 186 | – | Platinum doublet chemotherapy |
| Schmid P | 2018 | III | Breast cancer | 1 L | Study | 451 | 55 | Atezolizumab + nab-paclitaxel |
| | | | | | Control | 451 | 56 | Nab-paclitaxel |
| Jotte R* | 2018 | III | Squamous NSCLC | 1 L | Study | 343 | – | Atezolizumab + carboplatin + nab-paclitaxel |
| | | | | | Control | 340 | – | Carboplatin + nab-paclitaxel |
| Horn L | 2018 | | SCLC | 1 L | Study | 201 | 64 | Atezolizumab + carboplatin + etoposide |
| | | | | | Control | 202 | 64 | Carboplatin + etoposide |
| Paz-Ares L | 2018 | III | Squamous NSCLC | 1 L | Study | 278 | – | Pembrolizumab + carboplatin-paclitaxel/nab-paclitaxel |
| | | | | | Control | 281 | – | Carboplatin-paclitaxel/nab-paclitaxel |
| Gandhi L | 2018 | III | Non-squamous NSCLC | 1 L | Study | 410 | 65 | Cisplatin/carboplatin + pemetrexed + pembrolizumab |
| | | | | | Control | 206 | 63.5 | Cisplatin/carboplatin + pemetrexed |
| Govindan R | 2017 | III | Squamous NSCLC | 1 L | Study | 388 | 64 | Paclitaxel + carboplatin + ipilimumab |
| | | | | | Control | 361 | 64 | Paclitaxel + carboplatin |
| Reck M | 2016 | III | SCLC | 1 L | Study | 478 | 62 | Etoposide + cisplatin/carboplatin + ipilimumab |
| | | | | | Control | 476 | 63 | Etoposide + cisplatin/carboplatin |
| Borghaei H | 2019 | II | Non-squamous NSCLC | 1 L | Study | 60 | 62.5 | Pembrolizumab + pemetrexed + carboplatin |
| | | | | | Control | 63 | 63.2 | Pemetrexed + carboplatin |
| Reck M | 2013 | II | SCLC | 1 L | Study | 43 | – | Paclitaxel + carboplatin + concurrent-ipilimumab |
| | | | | | Study | 42 | – | Paclitaxel + carboplatin + phased-ipilimumab |
| | | | | | Control | 45 | – | Paclitaxel + carboplatin |
| Lynch TJ | 2012 | II | NSCLC | 1 L | Study | 70 | 59 | Paclitaxel + carboplatin + concurrent-ipilimumab |
| | | | | | Study | 68 | 61 | Paclitaxel + carboplatin + phased-ipilimumab |
| | | | | | Control | 66 | 62 | Paclitaxel + carboplatin |
| Robert C | 2011 | III | Melanoma | 1 L | Study | 250 | 57.5 | Ipilimumab + dacarbazine |
| | | | | | Control | 252 | 56.4 | Dacarbazine |
| Hersh EM | 2011 | II | Melanoma | 1 L | Study | 35 | 60 | Ipilimumab + dacarbazine |
| | | | | | Control | 37 | 66 | Ipilimumab |
| Rini BI | 2019 | III | RCC | 1 L | Study | 432 | 62 | Pembrolizumab + axitinib |
| | | | | | Control | 429 | 61 | Sunitinib |
| Motzer RJ | 2019 | III | RCC | 1 L | Study | 442 | 62 | Avelumab + axitinib |
| | | | | | Control | 444 | 61 | Sunitinib |
| McDermott DF | 2018 | II | RCC | 1 L | Study | 101 | 62 | Atezolizumab + bevacizumab |
| | | | | | Control | 101 | 61 | Sunitinib |
| Motzer RJ* | 2018 | III | RCC | 1 L | Study | 454 | – | Atezolizumab + bevacizumab |
| | | | | | Control | 461 | – | Sunitinib |
| Socinski MA | 2018 | III | Non-squamous NSCLC | 1 L | Study | 356 | 63 | Atezolizumab + bevacizumab + paclitaxel + carboplatin |
| | | | | | Control | 336 | 63 | Bevacizumab + paclitaxel + carboplatin |
| Slovin SF | 2013 | I/II | Prostate cancer | 2L | Study | 34 | 66 | Ipilimumab 10 mg/kg + radiotherapy |
| | | | | | Control | 16 | 65 | Ipilimumab |
| Kwon ED | 2014 | III | Prostate cancer | 2L | Study | 399 | 69 | Radiotherapy + ipilimumab |
| | | | | | Control | 400 | 67.5 | Radiotherapy |
| Theelen W* | 2018 | II | NSCLC | 2L | Study | 32 | – | Pembrolizumab + SBRT |
| | | | | | Control | 32 | – | Pembrolizumab |

Abbreviation: SCLCsmall-cell lung cancer; RCCrenal cell carcinoma; NSCLCnon-small-cell lung cancer; 1 Lfirst line; 2Lsecond line or beyond; SBRTstereotactic body radiotherapy.

* showed on ASCO; - unclear.

3.5. Publication bias and sensitivity analysis

There was no publication bias based on our analysis. The funnel plot was basically symmetrical and the *p*-values of both the Bag's test and the Egger's test were over 0.05 in each outcome measurement. Sensitivity analysis was conducted for all outcomes and revealed that no individual study appeared to change the pooled effect estimate

dramatically (Supplementary Fig. S4).

4. Discussion

Despite the durable responses observed in tumor immunotherapies, most patients do not respond to the initial treatment (primary resistance) and some of the responders relapse after a period of response




Fig. 2. Subgroup analysis of OS and PFS by combination regimens. A, OS; B, PFS. IO, immunotherapy; CT, chemotherapy; RT, radiotherapy; TT, targeted therapy; CRT, chemoradiotherapy.




Fig. 3. Subgroup analysis of ORR and DCR by combination regimens. **A**, ORR; **B**, DCR. IO, immunotherapy; CT, chemotherapy; RT, radiotherapy; TT, targeted therapy; CRT, chemoradiotherapy.





Fig. 4. A, OR of grade 3–5 AEs rates. B, Incidence of grade 3–5 irAEs. IO, immunotherapy; CT, chemotherapy; RT, radiotherapy; TT, targeted therapy; CRT, chemoradiotherapy.

(acquired resistance) (Sharma et al., 2017). Insufficient infiltration of cytotoxic T lymphocytes, lack of tumor-associated antigens (TAAs) and activation of other immunosuppressive pathways are major causes of resistance to immunotherapy. Researchers tried to overcome the resistance to improve clinical benefits of immunotherapy. Chemotherapy, radiotherapy and targeted therapy have immunomodulatory effects. Their clinical utilities in combination with immunotherapy to achieve synergetic effects and improve clinical outcomes have been hot research topics at present. However, it remains controversial as outcomes regarding the efficacy and safety of combination immunotherapy are not always consistent. Our meta-analysis showed that combination immunotherapy could significantly improve the anti-tumor efficacy. Although the incidence of adverse events above grade 3 increased, it was manageable.

The combination of different immunotherapies is an actively growing field of clinical investigation, as many studies showed great clinical efficacy. In checkmate 067 and 069, the combination group was associated with significantly better responses and longer PFS: the ORR of nivolumab + ipilimumab combination group were 58% and 59%, respectively and PFS were 11.5 months and NR (not reached), respectively (Hodi et al., 2016; Wolchok et al., 2017). We further confirmed this in the meta-analysis. Our study showed that combination immunotherapy resulted in a significantly improved ORR and OS. Among all immunotherapy + immunotherapy research, CTLA-4 blockade + PD-1 blockade is most frequent in research: CTLA-4 blockade primarily acts at sites of priming of T cell (e.g., tumor draining lymph nodes). It blocks the competitive inhibition signal of CTLA-4, inhibits the function of regulatory T cell (Treg), thus promoting T cell activation. PD-1/PD-L1 blockade primarily acts in inflamed peripheral tissues (e.g., tumor) by blocking the binding of PD-1 on T cells to PD-L1 on tumor cells to reverse the depletion of T cells and restore the activity of T cells (Fig. 5A) (Hargadon et al., 2018; Wei et al., 2018). Distinct mechanisms that act on different targets and at different stages of T cell differentiation make it possible to realize synergistic effect. In addition, tumor vaccines or immune cellular therapy stimulating T cells, combined with immune checkpoint inhibitors releasing brake of T cell, may improve therapeutic efficacy (Dammeijer et al., 2016). All above explain our results very well. However, NCT00094653 showed that gp100 vaccine + ipilimumab did not improve therapeutic efficacy. On the contrary, additional gp100 appeared to attenuate responses (Hodi et al., 2010). Researchers demonstrated that it was due to a persisting antigen depot produced by gp100 that primes T cells and induces T cell isolation, dysfunction and deletion, thus reducing the efficacy of ICB. Researchers further proved that non-persistent, virus, or water-based vaccine would not cause this inhibitory effect and could play a synergistic role with ICB (Hailemichael et al., 2013, 2018).

Whether the combination of immunotherapy and chemotherapy exert synergetic effects is highly concerned by researchers. Our meta-analysis gave a positive answer: immunotherapy + chemotherapy did show significantly longer survival than chemotherapy alone (HR = 0.79). Among them, Keynote-189, which evaluated the combination of pembrolizumab with pemetrexed and platinum-based drug in the treatment of advanced non-squamous NSCLC, showed that the combination regimen extended the median PFS by 3.9 months (8.8 months vs 4.9 months) and reduced the risk of death by 51% (HR for survival: 0.49), and ORR in the combined group was significantly superior to that in the single group (47.6% vs 18.9%). More importantly, additional pembrolizumab does not increase the incidence of adverse events (Gandhi et al., 2018). In addition to the direct cytotoxic effects on tumor cells, some chemotherapeutic agents can also stimulate immune system. This mainly including: increasing tumor antigenicity by releasing TAAs, inducing immunogenic cell death (ICD) with releasing damage-associated molecular patterns (DAMPs) and eliminate immunosuppressive cells such as Treg and myeloid-derived suppressor cell (MDSC) (Fig. 5B) (Pfirschke et al., 2016; Sistigu et al., 2014). In addition, several studies reported upregulation of PD-L1 on tumor cells

induced by chemotherapy (Galluzzi et al., 2015; Peng et al., 2015), which may be one of the mechanisms of resistance to chemotherapy. Blocking PD-1/PD-L1 signaling pathway may reverse the resistance. Therefore, the combination can play a facilitating role. It is worth noting that different chemotherapeutic drugs have different immune mechanisms and effect duration, so it is particularly important to select appropriate drugs, dosage and timing of administration to improve the efficacy of combination therapy. The optimal combination therapy strategy is still under exploration.

In recent years, studies have reported abscopal effect of radiotherapy, that is, regression of non-irradiated metastatic lesions at a distance from the primary site of irradiation (Fig. 5C). This phenomenon is closely related to immune responses, which gives researchers an idea for immunotherapy + radiotherapy combination to improve clinical efficacy. The analysis showed that immune + radiotherapy only prolonged the PFS yet did not show statistically significant improvements on OS and ORR. Dying tumor cells after radiation can release TAAs so that irradiative tumor can be converted to in situ vaccine. They may also release danger signals and chemokines, promote the infiltration of dendritic cells (DCs) and cytotoxic T lymphocytes and enhance antitumor immune responses (Demaria et al., 2015). On the other hand, the radiation can also attract immunosuppressive cells such as Treg to tumor microenvironment, making negative regulation of the immune system, which may explain the rarity of abscopal effect (Muroyama et al., 2017). Preclinical studies showed that radiotherapy combined with immunotherapy may improve the response rate of distal tumors and enhance the anti-tumor effect (Dovedi et al., 2017; Hao et al., 2016; Twyman-Saint Victor et al., 2015). Current exploration of immunotherapy combined with radiotherapy is still in its early stage.

This study showed that immunotherapy + targeted therapy increased PFS and OS by 27% and 30%, respectively. Anti-angiogenesis therapy and immune system interact with each other as well: Anti-angiogenesis therapy induced normalization of abnormal tumor vascular, thereby reducing tumor microenvironment hypoxia, promoting the infiltration of CD8⁺ T lymphocytes and transforming immunosuppressed microenvironment into immune-activated microenvironment to improve the effect of immunotherapy (Fukumura et al., 2018). Meanwhile, immune cells promote the normalization of tumor blood vessels as well. The mutual regulation between tumor blood vessel normalization and immune reprogramming forms positive circulation and induces long-lasting anti-tumor immunity (Huang et al., 2018). Immunotherapy combined with anti-angiogenic drugs is a promising approach, with a large number of relevant clinical trials ongoing.

In the presence of remarkable efficacy, we also found that combination immunotherapy appeared to increase toxicity. The most common AEs were diarrhea, fatigue and colitis, etc. Therefore, in the implementation of the combination regimens, attention should be paid to the prevention, early detection and proper management of these side effects. In addition, the cost-utility analysis of the combination should be concerned.

What's more, there is no satisfactory biomarker for immunotherapy with high sensitivity and specificity up to now. The exploratory analysis indicated that patients with PD-L1 \geq 1% derived more survival benefits than those with PD-L1 < 1%, which was consistent with previous studies (Fehrenbacher et al., 2016; Motzer et al., 2018a,b). Nevertheless, in some studies, such as Keynote-407, the survival benefit of pembrolizumab in combination with chemotherapy was independent from PD-L1 expression (Paz-Ares et al., 2018). TMB as another promising biomarker has showed some clinical predictive value in some clinical trials (Carbone et al., 2017). Nevertheless, IMpower133 showed that the OS and PFS benefits of first-line atezolizumab plus chemotherapy in extensive-stage small-cell lung cancer were consistent in patients with mutations greater or less than 10 and 16 mutations per megabase (Horn et al., 2018). Because of the complex interactions between tumor immune system, the predictive effects of these biomarkers vary by tumor types and drug regimens and there is not yet an




Fig. 5. Relevant mechanisms. A, CTLA-4 blockade + PD-1/PD-L1 blockade; B, Immunotherapy + chemotherapy; C, Immunotherapy + radiotherapy.

appropriate cut-off for these biomarkers to distinguish the efficacy of ICB (Havel et al., 2019). It was found that PD-L1 expression and TMB were independent predictors of the response and a multivariate model containing PD-L1 expression and TMB could improve the sensitivity and specificity of the prediction (Hellmann et al., 2018). A predictive model that takes into account multiple factors may be necessary for patient selection.

Although some important conclusions have been drawn from this study, we must pay attention to its limitations: 1. Differences in combination regimens, drug dosage, lines of treatment and tumor histotype may lead to heterogeneity between studies; 2. Two I/II studies were included; 3. Some subgroups, such as the immunotherapy + radiotherapy group, had small sample sizes; 4. The follow-up time of some

studies was not long enough, resulting in lack of long-term outcome data. We expect more complete follow-up data in the future to confirm our results.

To maximize the benefits for patients, how to optimize the therapeutic mode, select the appropriate treatment population and weigh the risks and benefits should be taken into account in future study designs and clinical practice to ensure the selection of the best beneficial population, the most synergistic combinatorial regimes and the most appropriate sequencing of therapies.

5. Conclusion

Combination immunotherapy significantly improved survival.

Though the adverse effects above grade 3 increased, they were manageable in clinic. Further research is required to identify sensitive biomarkers, as well as explore more combinatorial regimens, dosages and sequencing of therapies for combination therapy.

Conflict of interest statement

There is no potential conflict of interest to disclose.

Authors' contributions

Qin Li contributed to the design of the study and was responsible for the integrity of the data and accuracy of the data analysis. Yuhan Wei, Qi Du and Mengqi Li collected and extracted the clinical data. Qi Du and Teng Li were responsible for the statistical analysis. Li Li, Xueke Fan, Yingrui Li, Seyed Kariminia prepared the figures and tables. Yuhan Wei, Qi Du and Xiaoyue Jiang wrote the manuscript. All authors contributed to the review and approved the paper. All authors agreed to be accountable for the content of this paper.

Funding

This study was funded by the National Natural Science Foundation of China (Grant NO. 81301912), Beijing Municipal Health System High-level Health Person Foundation Project (Grant NO. 2014-3-005), Beijing Municipal Science and Technology Commission Foundation (Capital Features, Z16110000516083, to Qin Li) and Natural Science Foundation of Capital Medical University (to Qin Li).

Acknowledgements

We thank Shanshan Wu for her guidance on the statistical analysis.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.critrevonc.2019.04.008>.

References

- Baxi, S., Yang, A., Gennarelli, R.L., Khan, N., Wang, Z., Boyce, L., Korenstein, D., 2018. Immune-related adverse events for anti-PD-1 and anti-PD-L1 drugs: systematic review and meta-analysis. *Bmj* 360, k793.
- Borghaei, H., Hellmann, M.D., Paz-Ares, L., Ramalingam, S.S., Reck, M., O'Byrne, K.J., Bhagavatheswaran, P., Nathan, F., Brahmer, J., 2018. Nivolumab (Nivo) + platinum-doublet chemotherapy (Chemo) vs chemo as first-line (1L) treatment (Tx) for advanced non-small cell lung cancer (NSCLC) with < 1% tumor PD-L1 expression: results from CheckMate 227. *J. Clin. Oncol.* 36 (15, suppl) 9001-9001.
- Borghaei, H., Langer, C.J., Gadgeel, S., Papadimitrakopoulou, V.A., Patnaik, A., Powell, S.F., Gentzler, R.D., Martins, R.G., Stevenson, J.P., Jalal, S.I., Panwalkar, A., Yang, J.C., Gubens, M., Sequist, L.V., Awad, M.M., Fiore, J., Saraf, S., Keller, S.M., Gandhi, L., 2019. 24-month overall survival from KEYNOTE-021 cohort G: pemetrexed and carboplatin with or without Pembrolizumab as first-line therapy for advanced non-squamous non-small cell lung cancer. *J. Thorac. Oncol.* 14 (1), 124–129.
- Carbone, D.P., Reck, M., Paz-Ares, L., Creelan, B., Horn, L., Steins, M., Felip, E., van den Heuvel, M.M., Ciuleanu, T.E., Badin, F., Ready, N., Hiltermann, T.J.N., Nair, S., Juergens, R., Peters, S., Minenza, E., Wrangle, J.M., Rodriguez-Abreu, D., Borghaei, H., Blumenschein Jr., G.R., Villaruz, L.C., Havel, L., Krejci, J., Corral Jaime, J., Chang, H., Geese, W.J., Bhagavatheswaran, P., Chen, A.C., Socinski, M.A., CheckMate, I., 2017. First-line nivolumab in stage IV or recurrent non-small-cell lung cancer. *N. Engl. J. Med.* 376 (25), 2415–2426.
- Couzin-Frankel, J., 2013. Cancer immunotherapy. *Science* 342 (6165), 1432–1433.
- D'Angelo, S.P., Mahoney, M.R., Van Tine, B.A., Atkins, J., Milhem, M.M., Jahagirdar, B.N., Antonescu, C.R., Horvath, E., Tap, W.D., Schwartz, G.K., Streicher, H., 2018. Nivolumab with or without Ipilimumab treatment for metastatic sarcoma (Alliance A091401): two open-label, non-comparative, randomised, phase 2 trials. *Lancet Oncol.* 19 (3), 416–426.
- Dammeijer, F., Lievense, L.A., Veerman, G.D., Hoogsteden, H.C., Hegmans, J.P., Arends, L.R., Aerts, J.G., 2016. Efficacy of tumor vaccines and cellular immunotherapies in non-small-cell lung cancer: a systematic review and meta-analysis. *J. Clin. Oncol.* 34 (26), 3204–3212.
- Demaria, S., Golden, E.B., Formenti, S.C., 2015. Role of local radiation therapy in cancer immunotherapy. *JAMA Oncol.* 1 (9), 1325–1332.
- Dovedi, S.J., Cheadle, E.J., Popple, A.L., Poon, E., Morrow, M., Stewart, R., Yusko, E.C., Sanders, C.M., Vignali, M., Emerson, R.O., Robins, H.S., Wilkinson, R.W., Honeychurch, J., Illidge, T.M., 2017. Fractionated radiation therapy stimulates anti-tumor immunity mediated by both resident and infiltrating polyclonal T-cell populations when combined with PD-1 blockade. *Clin. Cancer Res.* 23 (18), 5514–5526.
- Fehrenbacher, L., Spira, A., Ballinger, M., Kowanetz, M., Vansteenkiste, J., Mazieres, J., Park, K., Smith, D., Artal-Cortes, A., Lewanski, C., Braiteh, F., Waterkamp, D., He, P., Zou, W., Chen, D.S., Yi, J., Sandler, A., Rittmeyer, A., 2016. Atezolizumab versus docetaxel for patients with previously treated non-small-cell lung cancer (POPLAR): a multicentre, open-label, phase 2 randomised controlled trial. *Lancet* 387 (10030), 1837–1846.
- Ferris, R.L., Blumenschein, G., Fayette, J., Guigay, J., Colevas, A.D., Licitra, L., Harrington, K., Kasper, S., Vokes, E.E., Even, C., Worden, F., Saba, N.F., Iglesias Docampo, L.C., Haddad, R., Rordorf, T., Kiyota, N., Tahara, M., Monga, M., Lynch, M., Geese, W.J., Kopit, J., Shaw, J.W., Gillison, M.L., 2016. Nivolumab for recurrent squamous-cell carcinoma of the head and neck. *N. Engl. J. Med.* 375 (19), 1856–1867.
- Fukumura, D., Kloepper, J., Amoozgar, Z., Duda, D.G., Jain, R.K., 2018. Enhancing cancer immunotherapy using antiangiogenics: opportunities and challenges. *Nat. Rev. Clin. Oncol.* 15 (5), 325–340.
- Galluzzi, L., Buque, A., Kepp, O., Zitvogel, L., Kroemer, G., 2015. Immunological effects of conventional chemotherapy and targeted anticancer agents. *Cancer Cell* 28 (6), 690–714.
- Gandhi, L., Rodriguez-Abreu, D., Gadgeel, S., Esteban, E., Felip, E., De Angelis, F., Domine, M., Clingan, P., Hochmair, M.J., Powell, S.F., Cheng, S.Y., Bischoff, H.G., Peled, N., Grossi, F., Jennens, R.R., Reck, M., Hui, R., Garon, E.B., Boyer, M., Rubio-Viqueira, B., Novello, S., Kurata, T., Gray, J.E., Vida, J., Wei, Z., Yang, J., Raftopoulos, H., Pietanza, M.C., Garassino, M.C., Investigators, K., 2018. Pembrolizumab plus chemotherapy in metastatic non-small-cell lung cancer. *N. Engl. J. Med.* 378 (22), 2078–2092.
- Govindan, R., Szczesna, A., Ahn, M.J., Schneider, C.P., Gonzalez Mella, P.F., Barlesi, F., H.B. Ganea, D.E., Von Pawel, J., Vladimirov, V., Fadeeva, N., Lee, K.H., Kurata, T., Zhang, L., Tamura, T., Postmus, P.E., Jassem, J., O'Byrne, K., Kopit, J., Li, M., Tschaika, M., Reck, M., 2017. Phase III trial of ipilimumab combined with paclitaxel and carboplatin in advanced squamous non-small-cell. *Lung Cancer J. Clin. Oncol.* 35 (30), 3449–3457.
- Hailemichael, Y., Dai, Z., Jaffarzar, N., Ye, Y., Medina, M.A., Huang, X.F., Dorta-Estremera, S.M., Greeley, N.R., Nitti, G., Peng, W., Liu, C., Lou, Y., Wang, Z., Ma, W., Rabinovich, B., Sowell, R.T., Schluns, K.S., Davis, R.E., Hwu, P., Overwijk, W.W., 2013. Persistent antigen at vaccination sites induces tumor-specific CD8(+) T cell sequestration, dysfunction and deletion. *Nat. Med.* 19 (4), 465–472.
- Hailemichael, Y., Woods, A., Fu, T., He, Q., 2018. Cancer vaccine formulation dictates synergy with CTLA-4 and PD-L1 checkpoint blockade therapy. *J. Clin. Invest.* 128 (4), 1338–1354.
- Hao, Y., Yasmin-Karim, S., Moreau, M., Sinha, N., Sajo, E., Ngwa, W., 2016. Enhancing radiotherapy for lung cancer using immunoadjuvants delivered in situ from new design radiotherapy biomaterials: a preclinical study. *Phys. Med. Biol.* 61 (24), N697–N707.
- Hargadon, K.M., Johnson, C.E., Williams, C.J., 2018. Immune checkpoint blockade therapy for cancer: an overview of FDA-approved immune checkpoint inhibitors. *Int. Immunopharmacol.* 62, 29–39.
- Havel, J.J., Chowell, D., Chan, T.A., 2019. The evolving landscape of biomarkers for checkpoint inhibitor immunotherapy. *Nat. Rev. Cancer* 19 (3), 133–150.
- Hellmann, M.D., Nathanson, T., Rizvi, H., Creelan, B.C., Sanchez-Vega, F., Ahuja, A., Ni, A., Novik, J.B., Mangarin, L.M.B., Abu-Akeel, M., Liu, C., Sauter, J.L., Rehtman, N., Chang, E., Callahan, M.K., Chaft, J.E., Voss, M.H., Tenet, M., Li, X.M., Covello, K., Renninger, A., Vitazka, P., Geese, W.J., Borghaei, H., Rudin, C.M., Antonia, S.J., Swanton, C., Hammerbacher, J., Merghoub, T., McGranahan, N., Snyder, A., Wolchok, J.D., 2018. Genomic features of response to combination immunotherapy in patients with advanced non-small-cell lung cancer. *Cancer Cell* 33 (5), 843–852 e844.
- Hersh, E.M., O'Day, S.J., Powderly, J., Khan, K.D., Pavlick, A.C., Cranmer, L.D., Samlowski, W.E., Nichol, G.M., Yellin, M.J., Weber, J.S., 2011. A phase II multicenter study of ipilimumab with or without dacarbazine in chemotherapy-naïve patients with advanced melanoma. *Invest. New Drugs* 29 (3), 489–498.
- Hodi, F.S., O'Day, S.J., McDermott, D.F., Weber, R.W., Sosman, J.A., Haanen, J.B., Gonzalez, R., Robert, C., Schadendorf, D., Hassel, J.C., Akerley, W., van den Eertwegh, A.J., Lutzky, J., Lorigan, P., Vaubel, J.M., Linette, G.P., Hogg, D., Ottensmeier, C.H., Lebbe, C., Peschel, C., Quirt, I., Clark, J.I., Wolchok, J.D., Weber, J.S., Tian, J., Yellin, M.J., Nichol, G.M., Hoos, A., Urba, W.J., 2010. Improved survival with ipilimumab in patients with metastatic melanoma. *N. Engl. J. Med.* 363 (8), 711–723.
- Hodi, F.S., Lee, S., McDermott, D.F., Rao, U.N., Butterfield, L.H., Tarhini, A.A., Leming, P., Puzanov, I., Shin, D., Kirkwood, J.M., 2014. Ipilimumab plus sargramostim vs ipilimumab alone for treatment of metastatic melanoma: a randomized clinical trial. *Jama* 312 (17), 1744–1753.
- Hodi, F.S., Chesney, J., Pavlick, A.C., Robert, C., Grossmann, K.F., McDermott, D.F., Linette, G.P., Meyer, N., Giguere, J.K., Agarwala, S.S., Shaheen, M., Ernstoff, M.S., Minor, D.R., Salama, A.K., Taylor, M.H., Ott, P.A., Horak, C., Gagnier, P., Jiang, J., Wolchok, J.D., Postow, M.A., 2016. Combined Nivolumab and Ipilimumab versus Ipilimumab alone in patients with advanced melanoma: 2-year overall survival outcomes in a multicentre, randomised, controlled, phase 2 trial. *Lancet Oncol.* 17 (11), 1558–1568.
- Horn, L., Mansfield, A.S., Szczesna, A., Havel, L., Krzakowski, M., Hochmair, M.J., Huemer, F., Losonczy, G., Johnson, M.L., Nishio, M., Reck, M., Mok, T., Lam, S.,

- Shames, D.S., Liu, J., Ding, B., Lopez-Chavez, A., Kabbinavar, F., Lin, W., Sandler, A., Liu, S.V., Group, I.M.S., 2018. First-line Atezolizumab plus chemotherapy in extensive-stage small-cell lung cancer. *N. Engl. J. Med.*
- Huang, Y., Kim, B.Y.S., Chan, C.K., Hahn, S.M., Weissman, I.L., Jiang, W., 2018. Improving immune-vascular crosstalk for cancer immunotherapy. *Nat. Rev. Immunol.* 18 (3), 195–203.
- Jotte, R.M., Cappuzzo, F., Vynnychenko, I., Stroyakovskiy, D., Abreu, D.R., Hussein, M.A., Soo, R.A., Conter, H.J., Kozuki, T., Silva, C., Graupner, V., Sun, S., Lin, R.S., Kelsch, C., Kowanetz, M., Hoang, T., Sandler, A., Socinski, M.A., 2018. IMpower131: primary PFS and safety analysis of a randomized phase III study of Atezolizumab + carboplatin + paclitaxel or nab-paclitaxel vs carboplatin + nab-paclitaxel as 1L therapy in advanced squamous NSCLC. *J. Clin. Oncol.* 36 (18 suppl) LBA9000-LBA9000.
- Kantoff, P.W., Higano, C.S., Shore, N.D., Berger, E.R., Small, E.J., Penson, D.F., 2010. Sipuleucel-t immunotherapy for castration-resistant prostate cancer. *N. Engl. J. Med.* 363 (5), 411–422.
- Kwon, E.D., Drake, C.G., Scher, H.I., Fizazi, K., Bossi, A., van den Eertwegh, A.J.M., Krainer, M., Houede, N., Santos, R., Mohammed, H., Ng, S., Maio, M., Franke, F.A., Sundar, S., Agarwal, N., Bergman, A.M., Ciuleanu, T.E., Korbenfeld, E., Sengelov, L., Hansen, S., Logothetis, C., Beer, T.M., McHenry, M.B., Gagnier, P., Liu, D., Gerritsen, W.R., 2014. Ipilimumab versus placebo after radiotherapy in patients with metastatic castration-resistant prostate cancer that had progressed after docetaxel chemotherapy (CA184-043): a multicentre, randomised, double-blind, phase 3 trial. *Lancet Oncol.* 15 (7), 700–712.
- Larkin, J., Chiarion-Sileni, V., Gonzalez, R., Grob, J.J., Cowey, C.L., Lao, C.D., Schadendorf, D., Dummer, R., Smylie, M., Rutkowski, P., Ferrucci, P.F., Hill, A., Wagstaff, J., Carlino, M.S., Haanen, J.B., Maio, M., Marquez-Rodas, I., McArthur, G.A., Ascierto, P.A., Long, G.V., Callahan, M.K., Postow, M.A., Grossmann, K., Sznol, M., Dreno, B., Bastholt, L., Yang, A., Rollin, L.M., Horak, C., Hodi, F.S., Wolchok, J.D., 2015. Combined nivolumab and Ipilimumab or monotherapy in untreated melanoma. *N. Engl. J. Med.* 373 (1), 23–34.
- Larkins, E., Blumenthal, G.M., Yuan, W., He, K., Sridhara, R., Subramanian, S., Zhao, H., Liu, C., Yu, J., Goldberg, K.B., McKee, A.E., Keegan, P., Pazdur, R., 2017. FDA approval summary: pembrolizumab for the treatment of recurrent or metastatic head and neck squamous cell carcinoma with disease progression on or after platinum-containing chemotherapy. *Oncologist* 22 (7), 873–878.
- Lynch, T.J., Bondarenko, I., Luft, A., Serwatowski, P., Barlesi, F., Chacko, R., Sebastian, M., Neal, J., Lu, H., Cuillerot, J.M., Reck, M., 2012. Ipilimumab in combination with paclitaxel and carboplatin as first-line treatment in stage IIIB/IV non-small-cell lung cancer: results from a randomized, double-blind, multicenter phase II study. *J. Clin. Oncol.* 30 (17), 2046–2054.
- McDermott, D.F., Huseni, M.A., Atkins, M.B., Motzer, R.J., Rini, B.I., Escudier, B., Fong, L., Joseph, R.W., Pal, S.K., Reeves, J.A., Sznol, M., Hainsworth, J., Rathmell, W.K., Stadler, W.M., Hutson, T., Gore, M.E., Ravaud, A., Bracarda, S., Suarez, C., Danielli, R., Gruenewald, V., Choueiri, T.K., Nickles, D., Jhunjunwala, S., Piau-Louis, E., Thobhani, A., Qiu, J., Chen, D.S., Hegde, P.S., Schiff, C., Fine, G.D., Powles, T., 2018. Clinical activity and molecular correlates of response to Atezolizumab alone or in combination with bevacizumab versus Sunitinib in renal cell carcinoma. *Nat. Med.* 24 (6), 749–757.
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A.P.M., 2015. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst. Rev.* 4 (1), 1–9.
- Motzer, R.J., Escudier, B., McDermott, D.F., George, S., Hammers, H.J., Srinivas, S., Tykodi, S.S., Sosman, J.A., Procopio, G., Plimack, E.R., Castellano, D., Choueiri, T.K., Gurney, H., Donskov, F., Bono, P., Wagstaff, J., Gauler, T.C., Ueda, T., Tomita, Y., Schutz, F.A., Kollmannsberger, C., Larkin, J., Ravaud, A., Simon, J.S., Xu, L.A., Waxman, I.M., Sharma, P., CheckMate, I., 2015. Nivolumab versus Everolimus in advanced renal-cell carcinoma. *N. Engl. J. Med.* 373 (19), 1803–1813.
- Motzer, R.J., Powles, T., Atkins, M.B., Escudier, B., McDermott, D.F., Suarez, C., Bracarda, S., Stadler, W.M., Donskov, F., Lee, J.-L., Hawkins, R.E., Ravaud, A., Alekseev, B.Y., Staehler, M.D., Uemura, M., Donaldson, F., Li, S., Huseni, M.A., Schiff, C., Rini, B.I., 2018a. IMmotion151: a randomized phase III study of Atezolizumab plus bevacizumab vs Sunitinib in untreated metastatic renal cell carcinoma (mRCC). *J. Clin. Oncol.* 36 (6 suppl) 578-578.
- Motzer, R.J., Tannir, N.M., McDermott, D.F., Aren Frontera, O., Melichar, B., Choueiri, T.K., Plimack, E.R., Barthelmy, P., Porta, C., George, S., Powles, T., Donskov, F., Neiman, V., Kollmannsberger, C.K., Salman, P., Gurney, H., Hawkins, R., Ravaud, A., Grimm, M.O., Bracarda, S., Barrios, C.H., Tomita, Y., Castellano, D., Rini, B.I., Chen, A.C., Mekan, S., McHenry, M.B., Wind-Rotolo, M., Doan, J., Sharma, P., Hammers, H.J., Escudier, B., CheckMate, I., 2018b. Nivolumab plus Ipilimumab versus Sunitinib in advanced renal-cell carcinoma. *N. Engl. J. Med.* 378 (14), 1277–1290.
- Motzer, R.J., Penkov, K., Haanen, J., Rini, B., Albiges, L., Campbell, M.T., Venugopal, B., Kollmannsberger, C., Negrier, S., Uemura, M., Lee, J.L., Vasiliev, A., Miller Jr., W.H., Gurney, H., Schmidinger, M., Larkin, J., Atkins, M.B., Bedke, J., Alekseev, B., Wang, J., Mariani, M., Robbins, P.B., Chudnovsky, A., Fowst, C., Hariharan, S., Huang, B., di Pietro, A., Choueiri, T.K., 2019. Avelumab plus axitinib versus sunitinib for advanced renal-cell carcinoma. *N. Engl. J. Med.*
- Muroyama, Y., Nirschl, T.R., Koehl, C.M., Lopez-Bujanda, Z., Theodoros, D., Mao, W., Carrera-Haro, M.A., Ghasemzadeh, A., Marciscano, A.E., Velarde, E., Tam, A.J., Thoburn, C.J., Uddin, M., Meeker, A.K., Anders, R.A., Pardoll, D.M., Drake, C.G., 2017. Stereotactic radiotherapy increases functionally suppressive regulatory T cells in the tumor microenvironment. *Cancer Immunol. Res.* 5 (11), 992–1004.
- Paz-Ares, L., Luft, A., Vicente, D., Tafreshi, A., Gumus, M., Mazieres, J., Hermes, B., Cay Senler, F., Csozsi, T., Fulop, A., Rodriguez-Cid, J., Wilson, J., Sugawara, S., Kato, T., Lee, K.H., Cheng, Y., Novello, S., Halmos, B., Li, X., Lubiniecki, G.M., Piperdi, B., Kowalski, D.M., Investigators, K., 2018. Pembrolizumab plus chemotherapy for squamous non-small-cell lung cancer. *N. Engl. J. Med.*
- Peng, J., Hamanishi, J., Matsumura, N., Abiko, K., Murat, K., Baba, T., Yamaguchi, K., Horikawa, N., Hosoe, Y., Murphy, S.K., Konishi, I., Mandai, M., 2015. Chemotherapy induces programmed cell death-ligand 1 overexpression via the nuclear Factor-KappaB to foster an immunosuppressive tumor microenvironment in ovarian cancer. *Cancer Res.* 75 (23), 5034–5045.
- Pennock, G.K., Chow, L.Q., 2015. The evolving role of immune checkpoint inhibitors in cancer treatment. *Oncologist* 20 (7), 812–822.
- Peters, S., Antonia, S., Goldberg, S.B., Heymach, J.V., Kim, E.S., Nakagawa, K., Papadimitrakopoulou, V., Mukhopadhyay, P., McIntosh, S., Rizvi, N.A., 2016. 191TIP: MYSTIC: a global, phase 3 study of durvalumab (MED14736) plus tremelimumab combination therapy or durvalumab monotherapy versus platinum-based chemotherapy (CT) in the first-line treatment of patients (pts) with advanced stage IV NSCLC. *J. Thorac. Oncol.* 11 (4), S139–S140.
- Pfirschke, C., Engblom, C., Rickelt, S., Cortez-Retamozo, V., Garriss, C., Pucci, F., Yamazaki, T., Poirier-Colame, V., Newton, A., Redouane, Y., Lin, Y.J., Wojtkiewicz, G., Iwamoto, Y., Mino-Kenudson, M., Huynh, T.G., Hynes, R.O., Freeman, G.J., Kroemer, G., Zitvogel, L., Weissleder, R., Pittet, M.J., 2016. Immuno-genic chemotherapy sensitizes tumors to checkpoint blockade therapy. *Immunity* 44 (2), 343–354.
- Reck, M., Bondarenko, I., Luft, A., Serwatowski, P., Barlesi, F., Chacko, R., Sebastian, M., Lu, H., Cuillerot, J.M., Lynch, T.J., 2013. Ipilimumab in combination with paclitaxel and carboplatin as first-line therapy in extensive-disease-small-cell lung cancer: results from a randomized, double-blind, multicenter phase 2 trial. *Ann. Oncol.* 24 (1), 75–83.
- Reck, M., Luft, A., Szczesna, A., Havel, L., Kim, S.W., Akerley, W., Pietanza, M.C., Wu, Y.L., Zielinski, C., Thomas, M., Felip, E., Gold, K., Horn, L., Aerts, J., Nakagawa, K., Lorigan, P., Pieters, A., Kong Sanchez, T., Fairchild, J., Spigel, D., 2016a. Phase III randomized trial of ipilimumab plus etoposide and platinum versus placebo plus etoposide and platinum in extensive-stage small-cell lung cancer. *J. Clin. Oncol.* 34 (31), 3740–3748.
- Reck, M., Rodriguez-Abreu, D., Robinson, A.G., Hui, R., Czoszi, T., Fulop, A., Gottfried, M., Peled, N., Tafreshi, A., Cuffe, S., O'Brien, M., Rao, S., Hotta, K., Leiby, M.A., Lubiniecki, G.M., Shentu, Y., Rangwala, R., Brahmer, J.R., Investigators, K., 2016b. Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N. Engl. J. Med.* 375 (19), 1823–1833.
- Rini, B.I., Plimack, E.R., Stus, V., Gafanov, R., Hawkins, R., Nosov, D., Pouliot, F., Alekseev, B., Soulieres, D., Melichar, B., Vynnychenko, I., Kryzhanivska, A., Bondarenko, I., Azevedo, S.J., Borchiellini, D., Szczylik, C., Mark, C., McDermott, R.S., Bedke, J., Tartas, S., Chang, Y.H., Tamada, S., Shou, Q., Perini, R.F., Chen, M., Atkins, M.B., Powles, T., Investigators, K., 2019. Pembrolizumab plus Axitinib versus Sunitinib for advanced renal-cell carcinoma. *N. Engl. J. Med.*
- Robert, C., Thomas, L., Bondarenko, I., O'Day, S., Weber, J., Garbe, C., Wolchok, J.D., 2011. Ipilimumab plus Dacarbazine for previously untreated metastatic melanoma. *N. Engl. J. Med.* 364 (26), 2517–2526.
- Robert, C., Long, G.V., Brady, B., Dutriaux, C., Maio, M., Mortier, L., Hassel, J.C., Rutkowski, P., McNeil, C., Kalinka-Warchoła, E., Savage, K.J., HERNBERG, M.M., Lebbe, C., Charles, J., Mihalciou, C., Chiarion-Sileni, V., Mauch, C., Cognetti, F., Arance, A., Schmidt, H., Schadendorf, D., Gogas, H., Lundgren-Eriksson, L., Horak, C., Sharkey, B., Waxman, I.M., Atkinson, V., Ascierto, P.A., 2015a. Nivolumab in previously untreated melanoma without BRAF mutation. *N. Engl. J. Med.* 372 (4), 320–330.
- Robert, C., Schachter, J., Long, G.V., Arance, A., Grob, J.J., Mortier, L., Daud, A., Carlino, M.S., McNeil, C., Lotem, M., Larkin, J., Lorigan, P., Neyns, B., Blank, C.U., Hamid, O., Mateus, C., Shapira-Frommer, R., Kosh, M., Zhou, H., Ibrahim, N., Ebbinghaus, S., Ribas, A., 2015b. Pembrolizumab versus Ipilimumab in advanced melanoma. *N. Engl. J. Med.* 372 (26), 2521–2532.
- Schmid, P., Adams, S., Rugo, H.S., Schneeweiss, A., Barrios, C.H., Iwata, H., Dieras, V., Hegg, R., Im, S.A., Shaw Wright, G., Henschel, V., Molinero, L., Chui, S.Y., Funke, R., Husain, A., Winer, E.P., Loi, S., Emens, L.A., Investigators, I.M.T., 2018. Atezolizumab and nab-paclitaxel in advanced triple-negative breast cancer. *N. Engl. J. Med.*
- Sharma, P., Callahan, M.K., Bono, P., Kim, J., Spiliopoulou, P., Calvo, E., Pillai, R.N., Ott, P.A., de Braud, F., Morse, M., Le, D.T., Jaeger, D., Chan, E., Harbison, C., Lin, C.-S., Tschaike, M., Azrilevich, A., Rosenberg, J.E., 2016. Nivolumab monotherapy in recurrent metastatic urothelial carcinoma (CheckMate 032): a multicentre, open-label, two-stage, multi-arm, phase 1/2 trial. *Lancet Oncol.* 17 (11), 1590–1598.
- Sharma, P., Hu-Lieskovan, S., Wargo, J.A., Ribas, A., 2017. Primary, adaptive, and acquired resistance to cancer immunotherapy. *Cell* 168 (4), 707–723.
- Sistigu, A., Yamazaki, T., Vacchelli, E., Chaba, K., Enot, D.P., Adam, J., Vitale, I., Goubar, A., Baracco, E.E., Remedios, C., Fend, L., Hannani, D., Aymeric, L., Ma, Y., Niso-Santano, M., Kepp, O., Schultze, J.L., Tuting, T., Belardelli, F., Bracci, L., La Sorsa, V., Zicheddu, G., Sestili, P., Urbani, F., Delorenzi, M., Lacroix-Triki, M., Quidville, V., Conforti, R., Spano, J.P., Pusztai, L., Poirier-Colame, V., Delalogue, S., Penault-Llorca, F., Ladoire, S., Arnould, L., Cyrta, J., Dessoliers, M.C., Eggermont, A., Bianchi, M.E., Pittet, M., Engblom, C., Pfirschke, C., Preville, X., Uze, G., Schreiber, R.D., Chow, M.T., Smyth, M.J., Proietti, E., Andre, F., Kroemer, G., Zitvogel, L., 2014. Cancer cell-autonomous contribution of type I interferon signaling to the efficacy of chemotherapy. *Nat. Med.* 20 (11), 1301–1309.
- Slovin, S.F., Higano, C.S., Hamid, O., Tejwani, S., Harzstark, A., Alumkal, J.J., Scher, H.I., Chin, K., Gagnier, P., McHenry, M.B., Beer, T.M., 2013. Ipilimumab alone or in combination with radiotherapy in metastatic castration-resistant prostate cancer: results from an open-label, multicenter phase I/II study. *Ann. Oncol.* 24 (7), 1813–1821.
- Socinski, M.A., Jotte, R.M., Cappuzzo, F., Orlandi, F., Stroyakovskiy, D., Nogami, N., Rodriguez-Abreu, D., Moro-Sibilot, D., Thomas, C.A., Barlesi, F., Finley, G., Kelsch, C., Lee, A., Coleman, S., Deng, Y., Shen, Y., Kowanetz, M., Lopez-Chavez, A., Sandler,

- A., Reck, M., Group, I.M.S., 2018. Atezolizumab for first-line treatment of metastatic nonsquamous NSCLC. *N. Engl. J. Med.* 378 (24), 2288–2301.
- Theelen, W., Lalezari, F., Vries, J., Langen, J.D., Aerts, J., Monkhorst, K., Baas, P., 2018. Randomized phase II study of pembrolizumab after stereotactic body radiotherapy (SBRT) versus pembrolizumab alone in patients with advanced non-small cell lung cancer: the PEMBRO-RT study. *J. Clin. Oncol.* 36 (15_suppl) 9023-9023.
- Topalian, S.L., Hodi, F.S., Brahmer, J.R., Gettinger, S.N., Smith, D.C., McDermott, D.F., Powderly, J.D., Carvajal, R.D., Sosman, J.A., Atkins, M.B., Leming, P.D., Spigel, D.R., Antonia, S.J., Horn, L., Drake, C.G., Pardoll, D.M., Chen, L., Sharfman, W.H., Anders, R.A., Taube, J.M., McMiller, T.L., Xu, H., Korman, A.J., Jure-Kunkel, M., Agrawal, S., McDonald, D., Kollia, G.D., Gupta, A., Wigginton, J.M., Sznol, M., 2012. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N. Engl. J. Med.* 366 (26), 2443–2454.
- Topalian, S.L., Sznol, M., McDermott, D.F., Kluger, H.M., Carvajal, R.D., Sharfman, W.H., Brahmer, J.R., Lawrence, D.P., Atkins, M.B., Powderly, J.D., Leming, P.D., Lipson, E.J., Puzanov, I., Smith, D.C., Taube, J.M., Wigginton, J.M., Kollia, G.D., Gupta, A., Pardoll, D.M., Sosman, J.A., Hodi, F.S., 2014. Survival, durable tumor remission, and long-term safety in patients with advanced melanoma receiving nivolumab. *J. Clin. Oncol.* 32 (10), 1020–1030.
- Twyman-Saint Victor, C., Rech, A.J., Maity, A., Rengan, R., Pauken, K.E., Stelekati, E., Benci, J.L., Xu, B., Dada, H., Odorizzi, P.M., Herati, R.S., Mansfield, K.D., Patsch, D., Amaravadi, R.K., Schuchter, L.M., Ishwaran, H., Mick, R., Pryma, D.A., Xu, X., Feldman, M.D., Gangadhar, T.C., Hahn, S.M., Wherry, E.J., Vonderheide, R.H., Minn, A.J., 2015. Radiation and dual checkpoint blockade activate non-redundant immune mechanisms in cancer. *Nature* 520 (7547), 373–377.
- Wei, S.C., Duffy, C.R., Allison, J.P., 2018. Fundamental mechanisms of immune checkpoint blockade therapy. *Cancer Discov.* 8 (9), 1069–1086.
- Wolchok, J.D., Chiarion-Sileni, V., Gonzalez, R., Rutkowski, P., Grob, J.J., Cowey, C.L., Lao, C.D., Wagstaff, J., Schadendorf, D., Ferrucci, P.F., Smylie, M., Dummer, R., Hill, A., Hogg, D., Haanen, J., Carlino, M.S., Bechter, O., Maio, M., Marquez-Rodas, I., Guidoboni, M., McArthur, G., Lebbe, C., Ascierto, P.A., Long, G.V., Cebon, J., Sosman, J., Postow, M.A., Callahan, M.K., Walker, D., Rollin, L., Bhole, R., Hodi, F.S., Larkin, J., 2017. Overall survival with combined Nivolumab and Ipilimumab in advanced melanoma. *N. Engl. J. Med.* 377 (14), 1345–1356.