

drugs (AEDs) in elderly individuals with NCSE diagnosis, in accordance with ILAE, at a significance level of $p < 0.05$.

Procedures: Thirty-six elderly patients (70.8 ± 7.8 years) with clinical manifestation and critical record of NCSE, treated at the PUC Hospital, Campinas, SP, Brazil, were included in the study.

Results: Eight patients with NCSE with coma and 28 patients with NCSE were not comatose. Change in basal activity occurred in 30 cases, rhythmic delta activity (RDA) occurred in 13 cases and periodic patterns (PD) occurred in 19 cases (lateralized in 18 cases and generalized in 1 case), and electrographic SE was observed in 13 (53.1%) cases. Initial clinical improvement after BZD and/or antiepileptic drug (AEDs) therapy was observed in 20 (52.6%) cases; improvement in EEG background activity occurred in 10 (27.8%) cases and in EEG patterns in 20 (55.6%) cases after 3.9 days. A total of 14 deaths occurred. No significant association was observed between initial clinical improvement and background activity improvement (Fisher's exact test, $p = 0.456$) and EEG patterns (RDA, PD, unequivocally clear focal electrographic SE) ($p = 0.091$), death ($n = 9$ vs $n = 5$, $p = 0.501$) and presence of acute or remote brain injury. The predictive factor for the occurrence of death was the diagnosis of NCSE in comatose patients ($p = 0.016$).

Conclusion: No specific electrographic discharge patterns were observed according to the type of NCSE. No relationship between EEG patterns and initial clinical improvement after BZD/AED therapy was observed in elderly patients with NCSE. Death rate was high and related to NCSE with coma.

Keywords: non-convulsive status epilepticus, EEG, elderly.

doi:10.1016/j.yebeh.2019.08.043

Epilepsy & Behavior 101 (2019) 106769

Electroencephalographic status epilepticus is common in critically ill children undergoing continuous EEG monitoring

F. Fung, M. Jacobowitz, L. Vala, D. Parikh, M. Donnelly, M. Fitzgerald, S. Massey, A. Topjian, N. Abend
Children's Hospital of Philadelphia and the University of Pennsylvania, Philadelphia, United States

Objectives: Electroencephalographic seizures are common in critically ill children undergoing continuous EEG monitoring. However, few data are available regarding seizure characteristics which would inform management decisions. We aimed to describe electroencephalographic seizure characteristics with management implications.

Methods: We performed a single-center prospective observational study of consecutive critically ill children undergoing continuous EEG monitoring.

Results: This interim analysis included 498 consecutive critically ill children who underwent EEG monitoring from April – November 2017. Subjects were 56% male, the median (IQR) age 5.9 years (1.5, 13.1), and categorical diagnoses included acute structural (40%), acute non-structural (30%), and epilepsy-related (30%). Electrographic seizures occurred in 137 (28%) patients. Seizure duration was less than 1 minute in 69 subjects (51%), 1-5 minutes in 39 subjects (29%), 6-30 minutes in 17 subjects (13%), and >30 minutes in 10 subjects (7%). Electrographic status epilepticus occurred in 29 subjects (21%) and consisted of continuous seizures in 9 subjects (31%) and frequent recurrent seizures in 18 subjects (62%). Seizure onset was focal in 69 subjects (51%), generalized in 63 subjects (46%) and multifocal in 4 subjects (3%).

Seizure spread was focal/unilateral in 51 subjects (37%) and bilateral in 85 subjects (62%).

Conclusions: Electroencephalographic status epilepticus is common and has been associated with unfavorable neurodevelopmental outcomes. However, brief seizures (< 1 minute) are also very common. These seizures may not induce secondary brain injury may not provide benefit while exposing patients to unnecessary anti-seizure medication adverse effects. Further studies are warranted to determine the optimal management of seizures of varying durations.

doi:10.1016/j.yebeh.2019.08.044

Epilepsy & Behavior 101 (2019) 106770

The EEG diagnosis of NCSE: concordance between Salzburg Criteria and clinical practice

Giada Giovannini^a, Alshimaa Othman^b, Niccolo' Orlandi^{a,c}, Stefano Meletti^{a,c}

^aUnit of Neurology, OCSAE Hospital, AOU, Modena, Italy

^bKasr-Alainy School of Medicine, Cairo University, Cairo, Egypt

^cDepartment of Biomedical, Metabolic and Neural Sciences, Center for Neuroscience and Neurotechnology, University of Modena and Reggio Emilia, Modena, Italy

Background: The diagnosis of Non Convulsive Status Epilepticus (NCSE) in everyday clinical practice can be challenging. To help identify NCSE, Salzburg criteria (SC) have been recently validated. Here we evaluate the concordance in NCSE diagnosis between neurologists not trained in SC use, who take care of the patient in the clinical setting, and an expert panel who retrospectively evaluated the EEG according to SC.

Methods: All consecutive urgent EEG done from January 1st to March 30th 2018 were considered.

A pool of three epileptologists trained in the use of SC, and not involved in the clinical evaluation of the incident case (and of the corresponding EEG), retrospectively classified the EEG pattern according to SC in three categories: definite NCSE, no NCSE, possible NCSE. Finally, we defined the degree of concordance between the diagnosis made by the neurologist who took care of the patient and the classification made by the expert.

Results: Among the 574 consecutive EEG done in emergency conditions in the examined period, 187 (33%) have been made to rule out a NCSE (105 male; median age of 73 yrs) and were evaluated by 15 physicians. The neurologist made a diagnosis of NCSE in 19 (10%) out of 187 cases. The expert panel classified 9 EEG (5%) as "definite NCSE", 96 (51%) as "no NCSE" and 82 (44%) as "possible NCSE". Among the 82 cases defined as "Possible NCSE" by the expert, 10 (12%) were diagnosed as NCSE, while 72 (88%) were considered without status.

Concordance was 100% evaluating the "no NCSE" and the "definite NCSE" categories.

Conclusions: our data show that there is a considerable discrepancy between diagnosis made by neurologists not trained in the use of SC and the expert panel. The "Possible NCSE" category is a grey zone and further studies are needed.

doi:10.1016/j.yebeh.2019.08.045