



Comparative Effectiveness of Brief Alcohol Interventions for College Students: Results from a Network Meta-Analysis

Emily Alden Hennessy^{1,2} · Emily E. Tanner-Smith^{1,3} · Dimitris Mavridis⁴ · Sean P. Grant⁵

Published online: 2 January 2019
© Society for Prevention Research 2019

Abstract

Late adolescence is a time of increased drinking, and alcohol plays a predominant role in college social experiences. Colleges seeking to prevent students' hazardous drinking may elect to implement brief alcohol interventions (BAIs). However, numerous manualized BAIs exist, so an important question remains regarding the comparative effectiveness of these different types of BAIs for college students. This study uses network meta-analyses (NMA) to compare seven manualized BAIs for reducing problematic alcohol use among college students. We systematically searched multiple sources for literature, and we screened studies and extracted data in duplicate. For the quantitative synthesis, we employed a random-effects frequentist NMA to determine the effectiveness of different BAIs compared to controls and estimated the relative effectiveness ranking of each BAI. A systematic literature search resulted in 52 included studies: On average, 58% of participants were male, 75% were binge drinkers, and 20% were fraternity/sorority-affiliated students. Consistency models demonstrated that BASICS was consistently effective in reducing students' problematic alcohol use (ES range: $g = -0.23$, 95%CI $[-0.36, -0.16]$ to $g = -0.36$, 95% CI $[-0.55, -0.18]$), but AlcoholEDU ($g = -0.13$, 95%CI $[-0.22, -0.04]$), e-CHUG ($g = -0.35$, 95%CI $[-0.45, -0.05]$), and THRIVE ($g = -0.47$, 95%CI $[-0.60, -0.33]$) were also effective for some outcomes. Intervention rankings indicated that BASICS, THRIVE, and AlcoholEDU hold the most promise for future trials. Several BAIs appear effective for college students. BASICS was the most effective but is resource intensive and may be better suited for higher risk students; THRIVE and e-CHUG are less resource intensive and show promise for universal prevention efforts.

Keywords Brief alcohol intervention · College students · Network meta-analysis

Consumption of alcohol peaks for most people during late adolescence and early adulthood (SAMHSA 2018), the

developmental period in which many US youth are enrolled in college. In 2016, approximately 11% of full-time college students reported heavy drinking and 38% of full-time college students reported binge drinking at least once in the past 30 days (SAMHSA 2016a, b). Drinking rates on college campuses over the past 30 years have remained fairly stable (Hingson and White 2014), and evidence suggests that college students engage in heavier drinking than their non-student peers (Schulenberg et al. 2017). Given that college attendance is often a period of experimentation with alcohol and that alcohol plays a predominant role in many college social experiences, college administrators often focus on harm reduction by implementing programs aimed at preventing hazardous or heavy levels of drinking. Hazardous drinking patterns such as binge drinking (i.e., drinking that brings blood alcohol concentration levels to 0.08 g/dL) can result in a variety of negative alcohol-related consequences including aggression and assaultive behaviors, unplanned sex, injuries, sexual victimization, and future alcohol use disorder (Abbey et al. 2014;

✉ Emily Alden Hennessy
Emily.hennessy@uconn.edu

¹ Department of Human and Organizational Development, Vanderbilt University, PMB 90, 230 Appleton Place, Nashville, TN 37203-5721, USA

² Present address: Institute for Collaboration on Health, Intervention, and Policy, Department of Psychological Sciences, University of Connecticut, Storrs, CT 06269-1248, USA

³ Present address: Department of Counseling Psychology and Human Services, University of Oregon, 5251 University of Oregon, Eugene, OR 97403-5251, USA

⁴ Department of Primary Education, University of Ioannina, University Campus, 45110 Ioannina, Greece

⁵ Department of Social & Behavioral Sciences, Indiana University Richard M. Fairbanks School of Public Health, 1050 Wishard Blvd, RG 6046, Indianapolis, IN 46202, USA

Foster et al. 2014; Jennison 2004; Kingree and Thompson 2015; Valenstein-Mah et al. 2015; Voloshyna et al. 2016; Wechsler et al. 1994).

Prevention and Intervention

Given the public health impact of heavy and hazardous drinking among students, colleges and universities are increasingly implementing brief alcohol interventions (BAIs), broadly defined here as interventions delivered in a circumscribed time frame that aim to promote changes in alcohol use behaviors or their determinants. Evidence suggests that these briefer modalities may be preferable among this age group (Buscemi et al. 2010). A number of evidence-based BAIs exist, including manualized programs available for free or purchase.

One example of a highly studied BAI is the Brief Alcohol Screening Intervention for College Students (BASICS; Dimeff et al. 1999), which aims to reduce alcohol consumption and related negative consequences for students who already drink heavily or are at-risk of experiencing such problems. The intervention is typically delivered in two 50-min counseling sessions, wherein the provider uses a motivational interviewing therapeutic style to enhance students' motivations and skills to change drinking behaviors. Other manualized BAIs that are often implemented on college campuses include Alcohol 101/Alcohol 101 Plus (Century Council 1997, 2003), AlcoholEdu (www.alcohol.edu), Check Your Drinking (CYD; www.checkyourdrinking.net), Electronic CHECKUP TO GO (e-CHUG; www.e-chug.com), College Drinker's Check-up (CDCU; www.collegedrinkerscheckup.com), and Tertiary Health Research Intervention Via Email (THRIVE; <http://ceriph.curtin.edu.au/thrive>). These BAIs vary in length between 5 min and 2 h, and can vary in delivery mode and personnel; thus, effectiveness of these BAIs may differ for programs with different implementation features. For example, six of these seven BAIs involve computer activities, while BASICS is provided through an in-person encounter. THRIVE is the shortest, a 5-min survey, that gives brief personalized feedback, while the other computer programs are longer surveys with more comprehensive feedback or involve interactive decision making (e.g., such as at a bar). See Table 1 for other key components of these interventions. Additionally, other study and participant characteristics, such as study quality (e.g., allocation concealment to reduce selection bias: Pildal et al. 2007), length of time studied (Tanner-Smith and Lipsey 2015), membership in a fraternity or sorority organization (Barry 2007; Meisel and Barnett 2017; Voloshyna et al. 2016), or gender (Ham and

Hope 2003; Pedersen 2013), may also moderate the effectiveness of BAIs for college students.

Previous Syntheses and Extensions to Address Gaps

Meta-analytic reviews quantitatively synthesize findings across multiple primary studies and thus provide a useful method for evaluating the current best evidence regarding intervention effectiveness and potential moderators of effects. Previous reviews demonstrate that alcohol interventions, including BAIs, can reduce drinking among college students (Carey et al. 2016; Fachini et al. 2012; Huh et al. 2015; Tanner-Smith and Lipsey 2015; Steinka-Fry et al. 2015). However, these reviews focus on comparing manualized interventions to a control group or a single suboptimal intervention(s); that is, they do not assess the relative effectiveness among manualized interventions and rank them accordingly. Thus, an assessment of which manualized BAI, if any, outperforms others will be useful for colleges wishing to maximize use of limited funds and most effectively change problematic drinking behaviors among students. Whereas traditional pairwise meta-analysis synthesizes direct evidence about a single intervention-comparison contrast (e.g., BASICS vs. e-CHUG reported in one or more trials), network meta-analysis (Petropoulou et al. 2017) can assess the comparative effectiveness of multiple interventions simultaneously by pooling direct and indirect evidence, as long as the included studies form a connected network of both direct and indirect information (Mavridis et al. 2015). That is, a network meta-analysis incorporates (a) direct evidence from studies where two groups were compared with (b) indirect evidence, which is generated across evidence loops in a network where different studies compared different groups, but have at least one group in common with at least one other study in the network. For example, if the two BAIs, BASICS and e-CHUG, have both separately been compared with a common other treatment (AlcoholEdu), in two different sets of trials, then the relative effectiveness between BASICS and e-CHUG can be estimated indirectly through their common comparator, AlcoholEdu. Network meta-analyses can therefore provide more precise estimates of effects and permit comparison of interventions that were not directly compared in any one trial (Cipriani et al. 2013; Mavridis et al. 2015). One key assumption of network meta-analysis is transitivity: i.e., the distributions of potential effect modifiers are balanced and randomly distributed across all pairwise comparisons within the network of interventions being compared (Salanti 2012). Transitivity is empirically explored by comparing the

Table 1 Intervention characteristics

Brand	Aim	Description	Delivery	Dose
Alcohol 101/Alcohol 101 Plus	Educate students about effects of alcohol misuse and “normal” peer drinking	1. Make decisions for virtual characters at a party 2. Feedback given based on participant behavior and peer norms	Interactive CD rom program set on a virtual campus	45–60 min
AlcoholEdu	Alcohol misuse prevention course	1. Pretest of alcohol knowledge 2. Precourse survey on drinking behavior, attitudes, demographics 3. Interactive alcohol education: students must receive a grade of $\geq 65\%$ on postcourse knowledge test to pass	Online	2 h
Brief Alcohol Screening and Intervention for College Students (BASICS)	Alcohol harm reduction	1. Assessment of drinking patterns, attitudes, motivations 2. Feedback on personal risk factors and advice to moderate drinking	Interviews; brief assessment survey	2 interviews (50–60 min each)
Check Your Drinking (CYD)	Reduce high risk drinking	1. Complete survey: demographics; drinking consumption, behavior, consequence 2. Personalized feedback and sensible drinking guidelines	Online: http://www.checkyourdrinking.net	15 min
Electronic CHECKUP TO GO (e-CHUG)	Personalized prevention intervention to motivate individuals to reduce alcohol or marijuana consumption	1. Complete online assessment 2. Personalized feedback report: quantity/frequency drinking; comparison to U.S. drinking norms; estimated risk level (AUDIT score, genetic risk of alcoholism, tolerance); money per year on alcohol; cigarettes smoked per month; explanation, advice, local referral	Online: http://www.echeckuptogo.com/usa/	20–30 min (optional 15–20 min. Personal reflection)
College Drinker’s Check-up (CDCU)	Intervention for heavy drinking college students collegedrinkerscheckup.com/	Video “interviewer” gives instructions, offers encouragement and interpretive information about the feedback, and asks open-ended questions 40 min	CD (Windows) or Internet: http://www.collegedrinkerscheckup.com/	
Tertiary Health Research Intervention Via Email (THRIVE)	Target unhealthy alcohol use in college students	1. Complete survey: demographics; drinking last year; AUDIT; largest number of drinks consumed once last month, duration of the drinking episode, height weight; secondhand effects; opinions on alcohol beverage labeling; smoking history 2. Personalized feedback	Online: http://ceriph.curtin.edu.au/thrive/	5 min

distribution of pre-specified effect modifiers across treatment comparisons (Chaimani et al. 2017; Salanti 2012). Network meta-analysis also assumes consistency in the network, which is the statistical manifestation of transitivity (Cipriani et al. 2013) whereby the direct and indirect evidence agree (Mavridis et al. 2015). In the “Methods” section, we further discuss specific methods we used to assess whether these assumptions were met for this analysis.

Objective This network meta-analysis examines the comparative effectiveness of seven manualized BAIs for college students and aims to identify which of these BAIs are more or less effective in reducing students’ problematic alcohol consumption. The following BAIs were all

initially identified by reviewing manualized programs developed specifically for use with college students from a comprehensive meta-analysis examining the overall effectiveness of BAIs for adolescents and young adults (see Tanner-Smith and Lipsey 2015): Alcohol 101/Alcohol 101 Plus, AlcoholEdu, BASICS, CYD, CDCU, e-CHUG, and THRIVE. A new systematic literature search was then conducted to identify primary trials that evaluated the effects of those interventions (see methods). A secondary aim of the study is to assess whether heterogeneity in intervention effectiveness could be explained with any of the following factors: fraternity/sorority membership, sex (male/female), sample mandated to the intervention, and allocation concealment risk of bias.

Methods

The review protocol (including full strategy for our PubMed search) was registered in the online database for systematic reviews (see PROSPERO 2016 CRD42016035952 for the protocol and a link to the data and syntax needed to reproduce the analysis). The PRISMA-NMA statement document guided review steps and reporting (Hutton et al. 2015).

Eligibility Criteria

There were no geographic or language limitations on eligibility. Studies were eligible if they were conducted in 1980 or later to be relevant to current intervention practices. Eligible studies were those that assessed intervention effects for undergraduate college students from any country, who were no older than 30 years of age. Given the analytical assumptions required for a network meta-analysis, only randomized controlled trials (RCTs) were eligible for inclusion; that is, we only included RCTs due to their ability to reduce threats of selection bias and to prevent concerns that including multiple types of study design would further increase the chances of inconsistency. Eligible trials were those that evaluated one of the seven BAIs in Table 1 along with a comparison condition consisting of either a control group or an eligible comparator BAI. Finally, studies had to report at least one of the following alcohol consumption outcomes to be eligible for inclusion: (1) frequency of heavy alcohol consumption (e.g., how many times per week a large number of drinks were consumed); (2) quantity of alcohol consumption (e.g., number of drinks consumed in a week); and (3) quantity of alcohol consumption during a peak drinking period (e.g., alcohol consumption at social events). Studies were ineligible if they only provided outcomes for a single occasion (e.g., 21st birthday celebration).

Information Sources and Search Strategy

Using a comprehensive search strategy, published and unpublished studies that met the above criteria were identified for inclusion in the review. The following electronic bibliographic databases were searched through April 18 2016: ERIC, PsycARTICLES, PsycINFO, Sociological Abstracts, PubMed, Dissertation Abstracts International, Clinical Trials Register, and NIH RePORTER. Search terms were adapted to be appropriate for each database but generally used three blocks of terms that described (1) intervention name, (2) alcohol outcomes, and (3) research design. The following gray literature sources were also searched: Australasian Medical Index, Campbell Collaboration Library, Cochrane Collaboration CENTRAL, College on Problems of Drug Dependence conference presentations, EPPI-Centre

Database of Health Promotion Research, Google Scholar, Index to Theses in Great Britain and Ireland, International Clinical Trials Registry, NIAAA Web site, and SAMHSA Web site. Bibliographies of all screened and eligible studies and of prior narrative reviews and meta-analyses were reviewed. Hand searches of three journals were conducted: *Alcoholism: Clinical and Experimental Research*, *American Journal on Addictions*, and *Journal of Substance Abuse Treatment*.

Study Selection and Data Collection Process

All articles were double-screened for eligibility first at the title/abstract level; any potentially eligible articles were then double-screened for eligibility at the full-text level. After all eligibility decisions were completed, the research team coded information independently and in duplicate from the eligible study reports using a piloted, standardized coding protocol. At each stage of the process, disagreements were discussed until consensus was reached. If primary studies did not include the information needed to estimate effect sizes, primary study authors were contacted. We did not include studies for which data necessary to estimate effect sizes were not given or provided.

Data Items In addition to collecting data needed to estimate effect sizes, we also collected data on the following characteristics: participant demographics (age, percent male, White, fraternity/sorority membership, mandated sample); attrition; control group type; intervention name; intervention duration (minutes); time between intervention end and posttest (weeks); study location (site; country); study design (intent-to-treat [ITT] analysis [yes/no]; methods used to account for missing outcome data [type]; and monitoring of program fidelity [yes/no]).

Summary Measures

All effect sizes were transformed to the same metric, the standardized mean difference (SMD), to permit comparisons on group means measured using different continuous measurement scales (Lipsey and Wilson 2001). For binary outcomes, the Cox transformation was used to convert log odds ratio effect sizes into SMD effect sizes (Sánchez-Meca et al. 2003).

Synthesis of Results

Eligible outcomes were categorized into the three primary outcomes (frequency of heavy use, quantity, quantity during peak drinking episode) and analyzed separately, by three waves of follow-up timing (0–3 months post-

intervention; 3–6 months post-intervention; 6+ months post-intervention), resulting in nine potential networks for analysis. If a study reported multiple follow-up points within the same wave, the follow-up with the shortest duration and/or most different in duration to the other studies in that category was dropped and only the longer/similar duration was used in the analysis. Sensitivity analyses for these outcomes were conducted.

This study conducted network meta-analysis within a frequentist approach using the network package (White 2015) in StataSE (14.2; StataCorp 2015). Network plots (Chaimani et al. 2013) are a visual tool to understand the components of a particular network of treatments and display the network size and comparisons of different interventions (see Fig. 3): *nodes* (bubbles) specify each unique intervention and the number of studies providing information about that intervention (larger = more studies), whereas *edges* (lines) indicate the number of comparisons between different intervention/comparison groups (thicker = more comparisons). We used the multivariate random-effects meta-regression approach (White et al. 2012) with the use of the network package in Stata (White 2015). This approach treats the possible comparisons within a multi-arm trial as different outcomes and proceeds to fitting the model taking the correlation between treatment effects within a multi-arm trial into account. We chose the assessment-only control group (AO-CT) as the reference group and estimated the effectiveness of each intervention relative to this group. To interpret the results, emphasis is given to the results from the consistency models, which are meta-analytic models generated from the pairwise estimates using direct, indirect, and mixed evidence; however, results from the treatment rankings based on inferences about each network using the surface area under the cumulative ranking curve (SUCRA; Salanti et al. 2011) are also presented. SUCRA values represent the percentage of effectiveness of a treatment in relation to an (imaginary) treatment that always performs the best with no uncertainty: Thus, the highest SUCRA value indicates the most likely optimal treatment.¹

A network meta-analysis includes additional analyses to assess whether transitivity and consistency assumptions are met. Namely, inconsistency statistics and plots assessed global inconsistency and (local) loop-specific heterogeneity (Chaimani et al. 2013; White 2015). A global assessment of all inconsistency parameters, i.e.,

to reject the hypothesis of consistency across the network, is conducted using the Wald test statistic, which follows a χ^2 distribution. In the inconsistency plots, an inconsistency factor (IF) is calculated as the absolute difference between direct and indirect evidence. Inconsistency factors closer to zero indicate that direct and indirect evidence are in agreement. Contribution plots, which demonstrate the influence of the direct, mixed, and indirect evidence sources in the network, were used to assess whether single comparisons were unduly influencing the results (Chaimani et al. 2013). Finally, node-splitting was conducted to assess the effect of leaving one study out of the network at a time: good model fit is indicated by non-significant χ^2 test results (Dias et al. 2010).

Study Quality and Risk of Bias Across Studies Two independent raters used the Cochrane Risk of Bias tool to assess risk of bias (Higgins et al. 2011), modified to separately examine selective reporting for each of the three outcomes in the review and incomplete outcome data for each of the three waves selected. The tool was also modified to separately assess participant and personnel blinding.

Moderator and Sensitivity Analysis Given the size of the networks and number of treatment comparisons included, multivariate meta-regression for moderators was not conducted. However, seven studies enrolled students who were mandated to the BAI,² so sensitivity analyses were conducted with these studies removed. Sensitivity analyses also examined networks where one follow-up outcome was dropped in studies with multiple effect sizes. The protocol specified additional sensitivity analyses, but we were unable to quantitatively explore these factors given the small size of the networks. Finally, we explored for small study effects as a proxy for assessing potential publication bias, using the contour enhanced funnel plot: This plot allows us to explore if smaller studies more often show statistically significant results (Chaimani et al. 2013; Mavridis and Salanti 2014; Peters et al. 2008).

Results

Of 1132 identified studies, 52 trials met the inclusion criteria for this network meta-analysis (Fig. 1). Most studies involved pairwise comparisons, but three

¹ SUCRA values include the uncertainty of the different effect estimates and probabilities of assuming another possible treatment ranking so are more stable than the probability of being the best treatment: SUCRA values are thus focused on in the text; however, both values are presented in Table 3.

² Alfonso et al. 2012; Carey et al. 2011; Doumas et al. 2009; Homer 2010; Logan 2013; Terlecki 2011; Terlecki et al. 2011




Fig. 1 Study identification flow diagram

involved multi-arm trials (Appendix provides study characteristics and effect sizes). Most studies utilized AO-CT groups (AO-CT; $k=39$) and 14 utilized active comparison groups (ACT-CT). The BASICS ($k=34$) and e-CHUG ($k=9$) programs were the most prevalent interventions evaluated on college campuses, whereas the THRIVE program was the newest and least evaluated ($k=2$). On average, 58% of participants were male, 84% were White ($k=47$), 75% were reported as hazardous drinkers at baseline, and 20% were fraternity/sorority organization members ($k=24$).

Study Quality Assessment

Overall, the risk of bias was mixed across the studies (see Fig. 2 for overall summary: see Appendix for item ratings). None of the studies were rated as high risk of bias for random sequence generation or allocation concealment, although a majority were rated unclear (52% and 74%, respectively). Participant and personnel blinding was uncommon, with participant blinding rated as high risk of bias in 93% of the studies. Risk of bias for incomplete outcome reporting did not vary




Fig. 2 Risk of bias across included studies

substantially across the waves and ranged from 39 to 56% of studies at high risk of bias and 28–45% at low risk of bias. The majority of studies were ranked as low risk of bias in selective reporting for each of the outcomes. Other risks of bias were identified in 35% of studies and included factors such as trialists’ potential conflicts of interest.

Primary Outcomes

Figure 3 displays a network plot for each of the nine unique networks; however, because several networks were fairly sparse and/or not well connected, we discuss the more connected networks in the text: frequency of heavy alcohol use (0–3 months), quantity of alcohol use (0–3 and 3–6 months), and quantity of alcohol use during peak consumption (0–3 months). We also conducted network meta-analysis for the networks of heavy frequency, 3–6 months post-intervention ($k=12$) and quantity of use during peak consumption, 3–6 months post-intervention ($k=11$), the details of which are reported in full in Appendix 2 given the limited size of the networks. In brief, results from both networks indicated that no comparisons significantly reduced the occurrence of frequent heavy alcohol consumption or the quantity of alcohol consumption during peak drinking episodes for outcomes measured 3–6 months post-intervention. The following networks of outcomes for six or more months post-intervention were too sparse to be informative, although it is worth noting that they all primarily focused on BASICS: heavy frequency ($k=4$), quantity of use ($k=11$), and quantity of use during peak consumption ($k=7$).

Frequency of Heavy Alcohol Use: 0–3 Months Frequency of heavy alcohol use outcomes measured at 0–3 months post-intervention represents a connected network (Fig. 3a) of 21 studies comparing six interventions, and the AO-CT and ACT-CT groups. Two studies were multi-arm trials. As the size of the nodes indicate, THRIVE and CYD were compared the least, while BASICS was compared the most; indeed, the most informative direct evidence in this network was for BASICS versus ACT-CT (17.4% contribution). Results from the consistency model indicate that relative to the AO-CT, all comparisons reduced the frequency of heavy alcohol use (Table 2); yet, only three interventions significantly reduced the frequency of heavy drinking: BASICS (−0.36, 95% CI [−0.55, −0.18]), e-CHUG (−0.35, 95% CI [−0.59, −0.11]), and the ACT-CT (−0.29, 95% CI [−0.56, −0.02]). The global test for inconsistency was non-significant ($\chi^2(4)=3.88, p=0.4218$), and there was no loop-specific inconsistency, indicating that the results from the direct and indirect estimates were largely consistent. Node-splitting also indicated that the consistency model fits well. When each intervention was ranked against all others, BASICS had the largest SUCRA value indicating that it was most likely to be successful at reducing frequency of heavy alcohol use (Table 3): e-CHUG had the second largest. In sensitivity analysis, when the single study that utilized a mandated sample was dropped (Doumas et al. 2009), the CYD intervention was no longer included in the network, yet the results remained largely the same (Tables 2 and 3). Sensitivity analysis replacing the measurement duration from one study (4.3 versus 8.6 weeks: Terry 2012) did not change the results (Tables 2 and 3).




Fig. 3 Network plots for networks in main analysis, split by timing and outcome. **a** Frequency of heavy alcohol use, 0–3 months, 3–6 months, and 6+ months. **b** Quantity of alcohol use, 0–3 months, 3–6 months, and 6+ months. **c** Peak quantity, 0–3 months, 3–6 months, and 6+ months. Alcohol101 Alcohol 101/Alcohol 101 Plus, AlcEDU AlcoholEdu, AO-

CT assessment-only control, ACT-CT active control, BASICS Brief Alcohol Screening Intervention for College Students, CYD Check Your Drinking, CDCU College Drinker’s Check-up, ECHUG Electronic CHECKUP TO GO, THRIVE Tertiary Health Research Intervention Via Email

Table 2 Intervention effects compared to assessment-only control groups

	0–3 months frequency			0–3 months quantity			3–6 months quantity			0–3 months peak quantity		
	SMD	LCL	UCL	SMD	LCL	UCL	SMD	LCL	UCL	SMD	LCL	UCL
Alc101	-0.10	-0.41	0.22	-0.11	-0.26	0.05	-0.25	-0.69	0.18	0.38	-0.73	1.49
AlcEDU	-0.04	-0.15	0.06	-0.13	-0.22	-0.04	-0.07	-0.55	0.42	-0.41	-0.84	0.02
BASICS	-0.36	-0.55	-0.18	-0.26	-0.36	-0.16	-0.23	-0.44	-0.02	-0.34	-0.59	-0.08
CYD	-0.35	-0.95	0.24	-0.37	-0.93	0.20	-0.10	-1.02	0.82	0.12	-0.85	1.10
e-CHUG	-0.35	-0.59	-0.11	-0.25	-0.45	-0.05	-0.12	-0.39	0.16	-0.33	-0.86	0.21
CDCU	NA	NA	NA	-0.19	-0.43	0.05	-0.08	-0.74	0.59	-0.36	-0.76	0.05
THRIVE	-0.15	-0.44	0.14	-0.47	-0.60	-0.33	-0.47	-0.95	0.02	NA	NA	NA
ACT-CT	-0.29	-0.56	-0.02	-0.18	-0.39	0.02	0.03	-0.36	0.42	-0.17	-0.77	0.43
	ES replaced ^{a1}			ES replaced ^{a2}			ES replaced ^{a3}					
Alc101	-0.10	-0.41	0.21	-0.11	-0.26	0.05	-0.28	-0.62	0.06			
AlcEDU	-0.04	-0.15	0.06	-0.13	-0.22	-0.04	-0.07	-0.38	0.24			
BASICS	-0.36	-0.55	-0.18	-0.26	-0.36	-0.16	-0.21	-0.37	-0.04			
CYD	-0.36	-0.96	0.23	-0.36	-0.93	0.20	-0.11	-0.91	0.70			
e-CHUG	-0.35	-0.59	-0.11	-0.25	-0.45	-0.05	-0.11	-0.34	0.12			
CDCU	NA	NA	NA	-0.19	-0.43	0.05	-0.08	-0.63	0.47			
THRIVE	-0.15	-0.44	0.14	-0.47	-0.60	-0.33	-0.47	-0.78	-0.15			
ACT-CT	-0.30	-0.57	-0.03	-0.18	-0.39	0.02	0.03	-0.28	0.33			
	Mandated removed											
Alc101	-0.10	-0.41	0.22	-0.03	-0.34	0.27	-0.26	-0.69	0.16	0.40	-0.73	1.53
AlcEDU	-0.04	-0.15	0.06	-0.16	-0.29	-0.03	-0.07	-0.53	0.40	-0.41	-0.85	0.03
BASICS	-0.36	-0.55	-0.18	-0.26	-0.38	-0.15	-0.24	-0.44	-0.03	-0.32	-0.63	0.00
CYD	NA	NA	NA	NA	NA	NA	-0.01	-0.92	0.91	NA	NA	NA
e-CHUG	-0.35	-0.59	-0.11	-0.25	-0.46	-0.05	-0.1	-0.37	0.17	-0.32	-0.87	0.22
CDCU	NA	NA	NA	-0.19	-0.44	0.06	-0.08	-0.73	0.57	-0.36	-0.77	0.06
THRIVE	-0.15	-0.44	0.14	-0.46	-0.64	-0.28	-0.47	-0.93	0.00	NA	NA	NA
ACT-CT	-0.29	-0.56	-0.02	-0.17	-0.40	0.05	0.123	-0.27	0.52	-0.15	-0.79	0.49

Italicized effect sizes indicate intervention produced significant effects compared to assessment-only control groups

NA not applicable due to the intervention not being included in that network analysis, *Alcohol101* Alcohol 101/Alcohol 101 Plus, *AlcEDU* AlcoholEdu, *AO-CT* assessment-only control, *ACT-CT* active control, *BASICS* Brief Alcohol Screening Intervention for College Students, *CYD* Check Your Drinking, *CDCU* College Drinker’s Check-up, *ECHUG* Electronic CHECKUP TO GO, *THRIVE* Tertiary Health Research Intervention Via Email

^a This analysis involved replacing an effect size used in the original network analysis due to dependency between outcomes reported within the same follow-up wave category with the dropped effect size: (1) One study replaced the 8.6-week follow-up ES with the 4.3-week follow-up. (2) One study replaced the 8.6-week follow-up ES with the 4.3-week follow-up. (3) Three studies replaced the 25.8-week follow-up ES with the 12.9-week follow-up ES

Quantity of Alcohol Use: 0–3 Months Quantity of alcohol use measured at 0–3 months post-intervention is a connected network (Fig. 3b) of 37 studies comparing all seven eligible interventions and both types of control groups. This network also included three multi-arm trials. As the size of the nodes indicate, CYD was compared the least, while BASICS was compared the most; indeed, the most informative direct evidence in this network was for the BASICS intervention versus AO-CT (16.3% contribution). Results from the consistency model indicate that relative to the AO-CT, only THRIVE (-0.47, 95% CI [-0.60, -0.33]), BASICS (-0.26, 95% CI [-0.36, -0.16]), e-CHUG (-0.25, 95% CI [-0.45, -0.05]), and AlcoholEdu (-0.13, 95% CI [-0.22, -0.04]) significantly

reduced the quantity of alcohol use when compared to the AO-CT group (Table 2). The global test for inconsistency was non-significant ($\chi^2(6) = 3.68, p = 0.7194$), and there was no loop-specific inconsistency, indicating that the results from the direct and indirect estimates were largely consistent. Node-splitting also supported the consistency model. When compared against all other interventions, THRIVE had the largest SUCRA value, followed by CYD (Table 3). Removing the six studies with a mandated sample from this network³ resulted in removing the CYD intervention, but the results remained

³ Carey et al. 2011; Doumas et al. 2009; Homer 2010; Logan et al. 2015; Terlecki et al. 2011; Terlecki et al. 2015

Table 3 Treatment rankings

	0–3 months		0–3 months		3–6 months		0–3 months	
	Heavy frequency		Quantity		Quantity		Peak quantity	
	SUCRA	PrBest	SUCRA	PrBest	SUCRA	PrBest	SUCRA	PrBest
Alc101	32.2	1.20	27.2	0.00	65.1	13.80	17.5	4.30
AlcEDU	23.0	0.00	33.5	0.00	41.8	4.30	74.6	25.90
BASICS	<i>80.1</i>	19.40	66.0	0.20	66.2	2.90	69.6	12.40
CYD	68.5	41.40	70.6	36.20	46.9	17.70	28.3	5.50
e-CHUG	76.3	27.70	62.8	2.10	48.5	1.40	65	21.70
CDCU	NA	NA	47.4	1.00	42.7	10.20	67.7	21.20
THRIVE	42.8	3.60	<i>94.7</i>	60.30	85.3	49.50	NA	NA
ACT-CT	65.9	6.70	44.4	0.20	26.7	0.20	50.6	9.00
AO-CT	11.1	0.00	3.6	0.00	27.0	0.00	26.6	0.00
	ES removed ^{a1}		ES removed ^{a2}		ES removed ^{a3}			
Alc101	31.4	1.40	28.9	0.00	70.5	13.00		
AlcEDU	22.7	0.00	32.6	0.00	39.9	1.20		
BASICS	<i>80.0</i>	19.50	65.7	0.30	65.1	1.10		
CYD	70.6	41.20	69.2	35.90	46.6	17.20		
e-CHUG	76.0	25.80	61.9	3.40	46.1	0.60		
CDCU	NA	NA	47.4	1.40	43.8	7.10		
THRIVE	40.6	4.30	<i>94.1</i>	58.80	<i>91.4</i>	59.60		
ACT-CT	67.7	7.80	46.3	0.20	22.4	0.20		
AO-CT	11.0	0.00	4.0	0.00	24.3	0.00		
	Mandated removed							
Alc101	33.5	2.40	19.1	0.80	68.4	15.50	14.4	4.80
AlcEDU	24.8	0.00	43.6	0.10	43.3	4.90	73.9	29.10
BASICS	<i>85.9</i>	38.10	70.3	2.20	68.3	4.20	64.5	9.80
CYD	NA	NA	NA	NA	41.7	12.60	NA	NA
e-CHUG	80.1	36.20	64.5	7.70	46.9	1.20	62.6	21.80
CDCU	NA	NA	49.9	3.90	46.9	10.10	67.6	25.00
THRIVE	44.2	5.50	<i>96.9</i>	84.10	<i>86.9</i>	51.50	NA	NA
ACT-CT	70.1	17.80	47.0	1.20	18.1	0.00	45.5	9.50
AO-CT	11.4	0.00	8.6	0.00	29.7	0.00	21.6	0.00

Italicized values indicate highest SUCRA value of that intervention in that network

SUCRA surface under the curve, PrBest probability that the treatment will perform the best, NA not applicable due to the intervention not being included in that network analysis, *Alcohol101* Alcohol 101/Alcohol 101 Plus, *AlcEDU* AlcoholEdu, *AO-CT* assessment-only control, *ACT-CT* active control, *BASICS* Brief Alcohol Screening Intervention for College Students, *CYD* Check Your Drinking, *CDCU* College Drinker's Check-up, *ECHUG* Electronic CHECKUP TO GO, *THRIVE* Tertiary Health Research Intervention Via Email

^a Replaced an effect size used in the original network analysis with the dropped effect size: (1) One study replaced the 8.6-week follow-up ES with the 4.3 week follow-up. (2) One study replaced the 8.6-week follow-up ES with the 4.3-week follow-up. (3) Three studies replaced the 25.8-week follow-up ES with the 12.9-week follow-up ES

largely the same (Tables 2 and 3). Sensitivity analysis replacing the measurement duration in one study (4.3 versus 8.6 weeks: Terry 2012) did not change the final model results.

Quantity of Alcohol Use: 3–6 Months Quantity of alcohol use measured between 3 and 6 months post-intervention is a connected network (Fig. 3b) of 21 studies comparing all 7

interventions and both types of control groups. As the size of the nodes indicate, CDCU, CYD, and THRIVE were compared the least, while BASICS was compared the most, followed by e-CHUG. The most informative direct evidence in this network was for the e-CHUG intervention versus AO-CT with a total contribution of only 14.4% to network estimates. Results from the consistency model indicate that

relative to the AO-CT (Table 2), only the BASICS intervention significantly reduced the quantity of alcohol use (-0.23 , 95% CI $[-0.44, -0.02]$). The global test for inconsistency was non-significant ($\chi^2(3) = 1.57$, $p = 0.6657$), and there was no significant loop-specific inconsistency. However, the confidence interval for the Alcohol 101, BASICS, and AO-CT loop was large (IF = 0.55, 95% CI $[0.00, 1.70]$) because, although BASICS was compared often, only two studies with inconsistent results assessed Alcohol 101. Node-splitting, however, supported the consistency model. When compared against all other interventions, THRIVE held the largest SUCRA value (Table 3). Both sensitivity analyses for this network produced similar results to the original analyses, but when the single study that utilized a mandated sample was dropped (Logan et al. 2015), THRIVE now significantly reduced drinking compared to the AO-CT (Table 2). Sensitivity analysis replacing the measurement duration in three studies (12.9 versus 25.8 weeks⁴) demonstrated that THRIVE produced significantly reduced results compared to the AO-CT (Table 2).

Quantity of Alcohol Use During Peak Consumption: 0–3 Months Quantity of alcohol use during peak consumption measured at 0–3 months post-intervention is a connected network (Fig. 3c) of 18 studies comparing six interventions and both types of control groups. This network included one multi-arm trial. As the size of the nodes indicate, Alcohol101 and CYD were compared the least, while BASICS was compared the most; indeed, BASICS comparisons contributed a larger amount of direct evidence in this network compared to the other networks (21.3% compared to AO-CT and 21.7% compared to ACT-CT groups), suggesting that these results should be interpreted with caution. Results from the consistency model indicated that relative to the AO-CT (Table 2), only the BASICS intervention produced significantly improved results from the AO-CT (-0.34 , 95% CI $[-0.59, -0.08]$). The global test for inconsistency was non-significant ($\chi^2(2) = 0.29$, $p = 0.8639$), and there was no significant loop-specific inconsistency; however, the confidence interval for the BASICS, e-CHUG, and AO-CT loop was large (IF = 0.25, 95% CI $[0.00, 1.91]$). In this network, two trials utilizing e-CHUG had inconsistent results (in opposite directions) but null in both cases. Thus, the loop-specific findings suggest caution when comparing e-CHUG to other interventions in this network. Node-splitting supported the consistency model. When compared against all other interventions

AlcoholEDU had the largest SUCRA value (Table 3). Removing the four studies that utilized a mandated sample from this network⁵ removed the sole comparison involving CYD and three comparisons of BASICS, but did not substantively change the model results (Tables 2 and 3).

Small Study Bias

Funnel plots demonstrated potential small study bias for frequency of heavy alcohol use and peak quantity of alcohol use, with gaps in negative effect sizes among small studies. Funnel plots for quantity of alcohol use outcomes appeared symmetrical, although gaps in the bottom half of the funnel plots suggest that some small studies may be missing (Appendix Fig. 4). However, overall, most studies appear in the statistically non-significant area of the figure, and smaller studies do not appear to show statistically significant results more than larger studies.

Discussion

This network meta-analysis examined the comparative effectiveness of seven manualized BAIs for college students, in an effort to identify which BAIs may be more or less effective in reducing college students' problematic alcohol use. The results indicated that the AlcoholEDU, BASICS, e-CHUG, and THRIVE interventions all led to reduced problematic alcohol use among college students, relative to AO-CT groups for outcomes measured 0–3 months post-intervention. However, only the BASICS intervention consistently led to reductions across all alcohol outcomes, including heavy frequency, quantity, and quantity during a peak drinking episode. Relative to AO-CT conditions, the AlcoholEDU and e-CHUG interventions were effective in the short-term for reducing heavy frequency and quantity of alcohol use, respectively. AlcoholEDU, however, had the smallest reduction in use compared to the three other successful interventions. Finally, THRIVE yielded significant benefits over AO-CT groups for short-term quantity outcomes, and these effects were nearly double the magnitude of those observed for the other successful interventions. Sensitivity analyses also indicated that THRIVE may be particularly effective for non-mandated college students. Evidence from the studies with longer outcomes was sparse and did not suggest that any one intervention was most effective.

⁴ Labrie et al. 2013; Schaus et al. 2009; Terlecki et al. 2015

⁵ Doumas et al. 2009; Horner 2010; Terlecki et al. 2011; Terlecki et al. 2015

When all seven BAIs were ranked in terms of effectiveness, AlcoholEDU, BASICS, and THRIVE ranked the best, but the rankings varied by type of alcohol consumption outcome. That is, BASICS had the largest SUCRA value for frequency of heavy use, THRIVE the largest for quantity of alcohol use, and AlcoholEdu the largest for peak drinking quantity. However, e-CHUG and CYD also had higher SUCRA values and BASICS maintained high SUCRA values across all outcomes. The rankings are estimates of potential future intervention effectiveness that are based on evidence from direct and indirect evidence. Thus, given the size of these networks and availability of treatment comparisons, the rankings should be interpreted with caution.

One important finding to note from the current study is that, among these seven manualized BAIs, the programs varied in their success by the different measures of alcohol consumption, a finding that has previously been demonstrated in the BAI literature (Tanner-Smith and Risser 2016). Namely, this review purposely examined three different measures of problematic alcohol use. Although we anticipated that the BAIs may have operated similarly across all measures of problematic drinking, the results indicated that the mechanisms of change associated with BAIs likely vary depending on the measure of problematic drinking. All of the included BAIs reported using some form of personalized feedback, so different results could be due to better/worse tailoring by the particular intervention. That is, BASICS could be more effective because of the in-person interview component, which may provide better tailoring. It may also take a stronger intervention dose (e.g., BASICS) to change the frequency of heavy drinking, e.g., by working with students to change factors leading them to environments where heavy drinking may be encouraged, while briefer, online assessments (e.g., THRIVE) may be more appropriate for enabling students to reflect on their drinking habits and consequently drink less when engaged in social drinking situations.

Limitations This review focused only on seven manualized BAIs, and thus, it is unclear how these BAIs compare to other manualized or non-manualized BAIs. Given that the BASICS trials included in this review tended to enroll heavier drinking/mandated students, future researchers may need to consider whether these follow-up effects generalize to other populations of college students. Of note, this review did not assess frequency of any type of alcohol use outcomes, so intervention effectiveness rankings might change if we focused on that outcome. Although we sought to identify published and unpublished sources through a comprehensive and systematic literature search, there remains a potential that we missed eligible studies due

to not exhaustively searching all available databases (e.g., Embase) or of small study bias in this research area. Despite the sole inclusion of RCT designs, the risk of bias was mixed and trials often did not report enough information to accurately assess risk of bias. Although often not feasible to blind participants in behavioral intervention research (e.g., see Grant et al. 2016), blinding of study personnel is often possible, but was uncommon in the included trials.

Conclusions

This network meta-analysis builds on previous reviews by directly comparing multiple interventions for problematic alcohol use among college students and highlighting what interventions might be best for which type of risky consumption. The results indicated that the BASICS intervention yielded the most consistent beneficial effects in terms of reducing problematic alcohol use among college students, yet AlcoholEDU, CYD, e-CHUG, and THRIVE also performed well and may be particularly useful for universal prevention efforts. Given the few studies on THRIVE and CYD that nevertheless suggest their effectiveness, future research with these interventions is warranted, especially because these interventions are less resource intensive than the BASICS intervention. Considering the effectiveness-resource tradeoff, universities may choose to implement e-CHUG as a form of universal prevention and implement BASICS for smaller groups of higher risk students (e.g., mandated students). Of note, few trials employed long-term follow-up designs. Thus, although the current study finds that some BAIs may be effective in the short-term, it is unclear how most BAIs perform beyond 3 months. Finally, one important future direction for research will be examining specific mechanisms of these interventions as well as whether these interventions are effective for reducing problematic drinking patterns among subpopulations of college students.

Funding Funding for the original review was provided by the National Institute on Alcohol Abuse and Alcoholism (NIAAA) Grant 1R01AA020286-01A1 (2011–2015).

Compliance with Ethical Standards

Conflicts of Interest S.G.'s spouse is a salaried-employee of Eli Lilly and Company, and owns stock. S.G. has accompanied his spouse on company-sponsored travel. All other authors declare no known conflicts of interest.

Ethical Approval/Informed Consent Ethical approval or informed consent is not applicable, as this is a systematic review.

Appendix 1




Fig. 4 Assessment of small study bias: Contour enhanced funnel plots for networks in main analysis, split by timing and outcome

Table 4 Risk of bias

Study	Random sequence generation	Allocation concealment	Participant blinding	Personnel blinding	Incomplete outcome			Selective outcome reporting			
					0–3 months	3–6 months	6+ months	Heavy frequency	Quantity	Quantity during peak drinking	Other risk of bias identified
Alfonso et al. 2012	Low	Unclear	High	High	Unclear	Unclear	N/A	Unclear	Low	Low	No
Baer et al. 1995, 2001; Roberts et al. 2000	Low	Unclear	Unclear	High	N/A	Low	Low	Unclear	Low	Low	Yes
Borsari and Carey 2000; Borsari et al. 2009	Low	Unclear	High	High	Low	N/A	Unclear	Low	Low	Unclear	No
Bowley et al. 2013	Unclear	Unclear	High	Unclear	Low	N/A	N/A	Unclear	Low	Unclear	No
Braitman 2012	Unclear	Unclear	High	Unclear	High	N/A	N/A	Low	Low	Low	No
Bryant 2009; Bryant et al. 2013	Unclear	Unclear	High	High	High	N/A	N/A	Low	Low	Unclear	No

Table 4 (continued)

Study	Random sequence generation	Allocation concealment	Participant blinding	Personnel blinding	Incomplete outcome			Selective outcome reporting			
					0–3 months	3–6 months	6+ months	Heavy frequency	Quantity	Quantity during peak drinking	Other risk of bias identified
Butler 2007; Butler and Correia 2009	Unclear	Unclear	High	High	High	N/A	N/A	Low	Low	Low	No
Carey et al. 2011; Carey and DeMartini 2010; Reid et al. 2015	Unclear	Unclear	High	Low	Low	High	High	Unclear	Unclear	Unclear	No
Croom et al. 2009	Unclear	Unclear	High	Low	High	N/A	N/A	Low	Unclear	Unclear	Yes
Dimeff 1997; Dimeff and McNeely 2000	Low	Low	High	Low	High	N/A	N/A	Low	Low	Low	Yes
Doumas and Andersen 2009	Unclear	Low	High	High	High	N/A	N/A	Low	Low	Unclear	Yes
Doumas and Haustveit 2008	Unclear	Unclear	High	Low	High	N/A	N/A	Unclear	Low	Low	Yes
Doumas et al. 2009	Low	Unclear	High	Low	Low	N/A	N/A	Low	Low	Low	No
Doumas et al. 2010	Unclear	Unclear	High	Low	Low	Low	N/A	Low	Low	Low	No
Doumas et al. 2011, 2014	Low	Unclear	High	Unclear	High	N/A	N/A	Low	Low	Low	Yes
Eggleston 2007	Unclear	Unclear	High	High	N/A	High	N/A	Low	Low	Unclear	No
Fernandez et al. 2011; Wood et al. 2010	Low	Unclear	High	High	N/A	N/A	Low	Low	Low	High	No
Geisner et al. 2007; Geisner 2008	Low	Unclear	High	Low	Low	N/A	N/A	Unclear	Low	Low	No
Grossbard et al. 2010; Tollison et al. 2013	Low	Low	High	High	N/A	N/A	Unclear	High	Low	Low	Yes
Hallett et al. 2009; Hustad and Borsari 2010; Kypri et al. 2009	Low	Low	Low	Low	Low	High	N/A	Low	Low	Low	Yes
Henslee et al. 2006	Unclear	Unclear	High	Low	High	N/A	N/A	Low	Low	Low	No
Henslee 2008; Henslee and Correia 2009	Unclear	Unclear	High	High	High	N/A	N/A	Low	Low	Low	No
Hester et al. 2012	Unclear	Unclear	High	High	Low	N/A	Low	Unclear	Low	Low	Yes
Hester et al. 2012	Unclear	Unclear	High	Low	Low	N/A	Low	Unclear	Low	Low	Yes
Horner 2010	Unclear	Unclear	High	High	High	N/A	N/A	Unclear	Low	Low	No
Hustad et al. 2010	Low	Low	High	Low	Low	N/A	N/A	Low	Low	Low	Yes
Juarez 2001; Juarez et al. 2006	Unclear	Unclear	High	High	High	N/A	N/A	Low	Unclear	High	Yes
Kerr-Corrêa et al. 2008; Simão et al. 2008	Unclear	Unclear	High	High	N/A	N/A	High	Low	Low	Low	No
Kulesza et al. 2010	Unclear	Unclear	High	High	Low	N/A	N/A	Unclear	Low	Unclear	No
Kulesza et al. 2013	Unclear	Unclear	High	High	Low	N/A	N/A	High	Low	Unclear	Yes

Table 4 (continued)

Study	Random sequence generation	Allocation concealment	Participant blinding	Personnel blinding	Incomplete outcome			Selective outcome reporting			
					0–3 months	3–6 months	6+ months	Heavy frequency	Quantity	Quantity during peak drinking	Other risk of bias identified
LaBrie et al. 2013	Low	Low	High	Low	Low	Unclear	Unclear	Unclear	Low	Low	No
Larimer et al. 2001; O’Leary et al. 2002	Unclear	Unclear	High	High	N/A	N/A	Unclear	Low	Low	Low	Yes
Larimer et al. 2007	Low	Low	High	Low	N/A	N/A	High	Low	Low	Low	No
Lau 2006; Lau-Barraco and Dunn 2008	Unclear	Unclear	High	High	Low	N/A	N/A	Low	Low	Unclear	No
Leffingwell et al. 2005	Unclear	Unclear	High	Low	Low	Low	Unclear	Unclear	Low	Low	No
Logan 2013; Logan et al. 2015	Unclear	Unclear	High	High	High	High	N/A	Unclear	Low	Low	Yes
Lovecchio 2009; Lovecchio et al. 2010	Low	Low	High	Low	High	N/A	N/A	Low	Low	Low	No
Martin 2014	Unclear	Unclear	Unclear	Unclear	Unclear	N/A	N/A	Low	Low	Low	No
Mastroleo 2008; Orchowski et al. 2012	Low	Unclear	High	High	Low	N/A	N/A	Low	Low	Low	No
Matteucci et al. 2016; Paschall et al. 2011a, b, 2014	Unclear	Unclear	High	Low	Low	High	High	Low	Low	Low	Yes
McNally 2003; McNally et al. 2005	Unclear	Unclear	High	High	Low	N/A	N/A	Low	Low	Unclear	No
McPherson 2012	Low	Unclear	High	High	High	N/A	N/A	Unclear	Low	Low	Yes
Miller 2000	Low	Unclear	Unclear	Unclear	Low	Low	Low	Unclear	Low	Low	No
Monahan et al. 2013; Murphy et al. 2010, 2015	Low	Low	High	High	High	High	N/A	Low	Low	Unclear	No
Murphy et al. 2001	Unclear	Unclear	High	High	Low	N/A	Low	Low	Low	Unclear	No
Schaus et al. 2009	Low	Low	High	High	High	High	High	Low	Low	Low	No
Teeters et al. 2015	Low	Low	High	High	Low	Low	N/A	Low	Low	Unclear	No
Terlecki et al. 2008; Terlecki et al. 2011	Low	Low	High	High	High	N/A	N/A	High	Low	Low	Yes
Terlecki et al. 2015	Low	Unclear	High	High	High	High	High	Unclear	Low	Low	No
Terlecki 2008; Terlecki et al. 2010, 2012	Low	Low	High	High	High	N/A	N/A	High	Low	Low	Yes
Terlecki 2011	Low	Unclear	High	High	High	High	High	Unclear	Low	Low	No
Terry 2012	Low	Low	High	High	High	N/A	N/A	Low	Low	Low	No
Walters et al. 2007	Unclear	Unclear	High	Low	Unclear	Unclear	N/A	Unclear	Low	Low	No
Whiteside 2010	Low	Unclear	High	High	High	High	N/A	Low	Low	Low	No

Table 5 Included studies characteristics

Study	Country	ITT	Group	N	Male (%)	White (%)	Greek (%)	Mean age	Baseline alcohol use	Mandated
Alfonso et al. 2012	USA	No	BASICS e-CHUG	53 48	0.66 0.46	0.83 0.75	0.00 0.00	18.79 18.71	Hazardous	Yes
Baer et al. 1995, 2001; Roberts et al. 2000	USA	Yes	BASICS AO-CT	143 156	0.47 0.45	0.85 0.85	0.16 0.16	18.50 18.50	Hazardous	No
Borsari and Carey 2000; Borsari et al. 2009	USA	No	BASICS AO-CT	29 30	0.41 0.45	0.86 0.90	0.07 0.07	18.45 18.71	Hazardous	No
Bowley et al. 2013	Australia	Yes	THRIVE ACT-CT	19 20	0.75 0.80	NR NR	NR NR	22.21 20.32	Hazardous	No
Braitman 2012	USA	No	Alcohol 101 ACT-CT	38 9	0.35 0.35	0.60 0.60	0.15 0.15	NR NR	Medium-regular	No
Bryant 2009; Bryant et al. 2013	USA	No	BASICS ACT-CT	101 90	0.24 0.24	0.82 0.82	0.37 0.37	18.70 18.70	Medium-regular	No
Butler 2007; Butler and Correia 2009	USA	No	BASICS AO-CT	28 26	0.32 0.35	0.93 0.96	0.57 0.38	19.70 20.38	Hazardous	No
Carey et al. 2011; Carey and DeMartini 2010; Reid et al. 2015	USA	No	Alcohol 101 AlcoholEdu AO-CT	164 161 170	0.64 0.64	0.85 0.85	NR 0.12 NR	19.00 19.00 19.00	Hazardous	Yes
Croom et al. 2009	USA	No	AlcoholEdu AO-CT	929 962	0.51 0.48	0.63 0.63	NR NR	18.80 18.80	Medium-regular	No
Dimeff 1997; Dimeff and McNeely 2000	USA	No	BASICS AO-CT	13 20	0.41 0.33	0.71 0.75	0.12 0.42	23.00 20.73	Hazardous	No
Doumas and Andersen 2009	USA	No	e-CHUG AO-CT	18 34	0.59 0.59	0.79 0.79	NR NR	21.99 21.99	Medium-regular	No
Doumas and Haustveit 2008	USA	No	CYD ACT-CT	15 18	0.58 0.58	0.54 0.54	NR NR	18.10 18.10	Medium-regular	No
Doumas et al. 2009	USA	Yes	CYD ACT-CT	37 24	0.72 0.72	0.86 0.86	NR NR	19.24 19.24	Hazardous	Yes
Doumas et al. 2010	USA	Yes	e-CHUG ACT-CT	59 47	0.43 0.43	0.70 0.70	NR NR	18.08 18.08	Medium-regular	No
Doumas et al. 2011, 2014	USA	No	e-CHUG AO-CT	36 46	0.35 0.35	0.90 0.90	NR NR	18.00 18.00	Low/experimenters	No
Eggleston 2007	USA	Yes	BASICS AO-CT	39 37	0.49 0.49	0.84 0.87	NR NR	19.20 19.00	Hazardous	No
Fernandez et al. 2011; Wood et al. 2010	USA	Yes	BASICS AO-CT	190 202	0.43 0.43	0.89 0.89	NR NR	18.40 18.40	Hazardous	No
Geisner et al. 2007; Geisner 2008	USA	Yes	BASICS AO-CT	80 87	0.30 0.30	0.49 0.49	0.14 0.14	19.28 19.28	Medium-regular	No
Grossbard et al. 2010; Tollison et al. 2013	USA	Yes	BASICS AO-CT	228 305	0.44 0.44	0.80 0.80	NR NR	17.92 17.92	Medium-regular	No
Hallett et al. 2009; Hustad and Borsari 2010; Kypri et al. 2009	Australia	Yes	THRIVE AO-CT	966/813 944/767	0.55 0.55	NR NR	NR NR	19.70 19.70	Hazardous	No
Henslee et al. 2006	USA	No	BASICS ACT-CT	12 9	0.43 0.43	NR NR	NR NR	20.40 20.40	Hazardous	No
Henslee 2008; Henslee and Correia 2009	USA	No	BASICS ACT-CT	80 86	0.33 0.33	0.89 0.91	0.43 0.43	18.10 18.10	Hazardous	No
Hester et al. 2012	USA	Yes	CDCU AO-CT	59 71	0.63 0.62	0.62 0.51	0.11 0.13	20.51 20.29	Hazardous	No
Hester et al. 2012	USA	Yes	CDCU AO-CT	42 39	0.55 0.58	0.50 0.43	0.07 0.10	20.02 20.28	Hazardous	No
Horner 2010	USA	No	BASICS AO-CT	27 29	0.63 0.74	0.78 0.79	NR NR	18.90 18.80	Hazardous	Yes
Hustad et al. 2010	USA	No	AlcoholEdu	26	0.39	0.89	0.00	18.10	Medium-regular	No

Table 5 (continued)

Study	Country	ITT	Group	N	Male (%)	White (%)	Greek (%)	Mean age	Baseline alcohol use	Mandated
			e-CHUG	30	0.65	0.89	0.00	18.10		
			AO-CT	24	0.49	0.89	0.00	18.10		
Juarez 2001; Juarez et al. 2006	USA	Yes	e-CHUG	20	0.52	0.57	0.10	19.43	Hazardous	No
			AO-CT	21	0.46	0.57	0.10	19.43		
Kerr-Corrêa et al. 2008; Simão et al. 2008	Brazil	No	BASICS	142	0.56	NR	NR	19.60	Hazardous	No
			AO-CT	121	0.56	NR	NR	19.60		
Kulesza et al. 2010	USA	Yes	BASICS	35	0.28	0.84	NR	20.00	Hazardous	No
			AO-CT	40	0.28	0.84	NR	20.00		
Kulesza et al. 2013	USA	Yes	BASICS	81	0.29	0.86	NR	19.80	Hazardous	No
			AO-CT	97	0.30	0.90	NR	20.30		
LaBrie et al. 2013	USA	Yes	BASICS	144/147/143	0.43	0.76	0.29	19.92	Hazardous	No
			AO-CT	143/148/142	0.43	0.76	0.29	19.92		
Larimer et al. 2001; O’Leary et al. 2002	USA	Yes	BASICS	56	1.00	0.82	1.00	18.80	Medium-regular	No
			ACT-CT	64	1.00	0.82	1.00	18.80		
Larimer et al. 2007	USA	Yes	BASICS	502	0.30	0.81	NR	20.60	Medium-regular	No
			AO-CT	498	0.30	0.81	NR	20.60		
Lau 2006; Lau-Barraco and Dunn 2008	USA	No	Alcohol	64	0.44	0.76	NR	19.88	Hazardous	No
			101							
			AO-CT	39	0.51	0.76	NR	19.88		
Leffingwell et al. 2005	USA	Yes	CDCU	41	0.79	0.90	NR	19.90	Hazardous	No
			AO-CT	32	0.79	0.90	NR	19.90		
Logan 2013; Logan et al. 2015	USA	Yes	BASICS	13	0.57	0.97	0.56	19.16	Hazardous	No
			ACT-CT	11	0.57	0.97	0.56	19.16		
Lovecchio 2009; Lovecchio et al. 2010	USA	DK	AlcoholEdu	248	0.49	0.80	0.05	18.12	Medium-regular	No
			AO-CT	225	0.44	0.82	0.05	18.10		
Martin 2014	USA	Yes	BASICS	20	0.22	0.67	NR	19.89	Hazardous	No
			AO-CT	22	0.22	0.67	NR	19.89		
Mastroleo 2008; Orchowski et al. 2012	USA	Yes	BASICS	69	0.30	0.81	NR	20.60	Hazardous	No
			AO-CT	71	0.52	0.92	NR	18.12		
Matteucci et al. 2016; Paschall et al. 2011a, b, 2014	USA	Yes	AlcoholEdu	1102	0.44	0.76	0.11	18.80	Medium-regular	No
			AO-CT	1298	0.46	0.68	0.12	18.60		
McNally 2003; McNally et al. 2005	USA	Yes	BASICS	36	0.29	0.85	NR	18.58	Hazardous	No
			AO-CT	37	0.29	0.85	NR	18.58		
McPherson 2012	Australia	No	BASICS	5	0.14	NR	NR	26.60	Hazardous	No
			e-CHUG	8	0.30	NR	NR	30.00	Hazardous	No
Miller 2000	USA	Ye	Alcohol	111	0.37	0.70	NR	19.00	Low/experimenters	No
			101							
			AO-CT	121	0.37	0.70	NR	19.00		
Monahan et al. 2013; Murphy et al. 2010, 2015	USA	Yes	BASICS	41/37	0.50	0.57	NR	18.60	Hazardous	No
			e-CHUG	38/35	0.50	0.53	NR	18.60		
			AO-CT	39	0.50	0.62	NR	18.60		
Murphy et al. 2001	USA	No	BASICS	30	0.46	0.94	NR	19.60	Hazardous	No
			ACT-CT	24	0.46	0.94	NR	19.60		
Schaus et al. 2009	USA	Yes	BASICS	95/128/111	0.48	0.78	NR	20.50	Hazardous	No
			ACT-CT	114/147/125	0.48	0.77	NR	20.60		
Teeters et al. 2015	USA	Yes	BASICS	37/35	0.41	0.74	NR	21.20	Hazardous	No
			Alcohol	32/30	0.41	0.66	NR	21.20		
			101							
Terlecki et al. 2008; Terlecki et al. 2011	USA	No	BASICS	19	0.79	0.95	0.58	20.26	Hazardous	Yes
			AO-CT	24	0.92	0.83	0.42	20.29		
Terlecki 2008; Terlecki et al. 2010, 2012	USA	No	BASICS	22	0.46	0.82	0.18	20.18	Hazardous	No
			AO-CT	19	0.26	0.74	0.26	19.84		
Terlecki et al. 2015	USA	Yes	BASICS	58	0.72	0.93	0.43	20.12	Hazardous	Yes
			AO-CT	56	0.49	0.83	0.46	20.24		

Table 5 (continued)

Study	Country	ITT	Group	N	Male (%)	White (%)	Greek (%)	Mean age	Baseline alcohol use	Mandated
Terlecki 2011	USA	Yes	BASICS	57	0.80	0.91	0.30	20.14	Hazardous	No
			AO-CT	54	0.41	0.89	0.41	20.00		
Terry 2012	USA	Yes	BASICS	13	0.40	0.80	0.00	21.20	Hazardous	No
			ACT-CT	15	0.50	0.70	0.10	21.00		
Walters et al. 2007	USA	Yes	e-CHUG	53	0.52	0.73	NR	18.00	Hazardous	No
			AO-CT	53	0.52	0.73	NR	18.00		
Whiteside 2010	USA	No	BASICS	32	0.40	0.66	NR	18.92	Hazardous	No
			AO-CT	43	0.40	0.66	NR	18.92		

AO-CT assessment-only control, ACT-CT active control, DK do not know, NR not reported

Table 6 Effect sizes for networks

Frequency of Heavy Use, 0–3 months				
Study	Group 1	Group 2	SMD	SE
Hustad et al. 2010	e-CHUG	AlcoholEdu	− 0.0131397	0.2679485
Hustad et al. 2010	AlcoholEdu	AO-CT	0.4604313	0.2867894
Hustad et al. 2010	e-CHUG	AO-CT	0.482922	0.2777758
Monahan et al. 2013; Murphy et al. 2010, 2015	e-CHUG	AO-CT	0.5586971	0.232344
Monahan et al. 2013; Murphy et al. 2010, 2015	BASICS	AO-CT	0.1067247	0.2238358
Monahan et al. 2013; Murphy et al. 2010, 2015	BASICS	e-CHUG	0.0696443	0.2252482
Matteucci et al. 2016; Paschall et al. 2011a, b, 2014	AlcoholEdu	AO-CT	0.0883723	0.0409815
Lovecchio 2009; Lovecchio et al. 2010	AlcoholEdu	AO-CT	0.1209321	0.0921529
Braitman 2012	Alcohol 101	ACT-CT	− 0.5947977	0.3757533
Whiteside 2010	BASICS	AO-CT	− 0.1468197	0.2337724
Hallett et al. 2009; Hustad and Borsari 2010; Kypri et al. 2009	THRIVE	AO-CT	0.1505234	0.132653
Juarez 2001; Juarez et al. 2006	e-CHUG	AO-CT	0.3317972	0.3145816
McNally 2003; McNally et al. 2005	BASICS	AO-CT	0.3032855	0.235446
Dimeff 1997; Dimeff and McNeely 2000	BASICS	AO-CT	0.7729614	0.3687488
Butler 2007; Butler and Correia 2009	BASICS	AO-CT	0.5056834	0.2766651
Croom et al. 2009	AlcoholEdu	AO-CT	0	0.0791437
Teeters et al. 2015	BASICS	Alcohol 101	0.2975212	0.2427313
Doumas et al. 2009	CYD	ACT-CT	0.0635385	0.2621579
Lau 2006; Lau-Barraco and Dunn 2008	Alcohol 101	AO-CT	0.2423321	0.2038409
Henslee et al. 2006	BASICS	ACT-CT	0.559514	0.4493308
Henslee 2008; Henslee and Correia 2009	BASICS	ACT-CT	0.0442421	0.1553506
Bryant 2009; Bryant et al. 2013	BASICS	ACT-CT	0.0229264	0.1449603
Borsari and Carey 2000; Borsari et al. 2009	BASICS	AO-CT	0.6103175	0.2664071
Terry 2012	BASICS	ACT-CT	0.659876	0.4039006
Martin 2014	BASICS	AO-CT	0.6151792	0.3161643
Frequency of Heavy Use, 3–6 months				
Study	Group 1	Group 2	SMD	SE
Mastroleo 2008; Orchowski et al. 2012	BASICS	AO-CT	0.0255337	0.169055
Murphy et al. 2001	BASICS	ACT-CT	0.4102412	0.2766917
Doumas and Andersen 2009	e-CHUG	AO-CT	− 0.065668	0.2915626
Matteucci et al. 2016; Paschall et al. 2011a, b, 2014	AlcoholEdu	AO-CT	0.051184	0.0409683
Doumas et al. 2011, 2014	e-CHUG	AO-CT	0	0.222524
Whiteside 2010	BASICS	AO-CT	− 0.0707541	0.2392181

Table 6 (continued)

Hallett et al. 2009; Hustad and Borsari 2010; Kypri et al. 2009	THRIVE	AO-CT	0.070593	0.168367
Schaus et al. 2009	BASICS	ACT-CT	0.492206	0.1409888
Monahan et al. 2013; Murphy et al. 2010, 2015	BASICS	Alcohol 101	0.3771957	0.2509964
Doumas and Haustveit 2008	CYD	ACT-CT	−0.065289	0.3496953
Eggleston 2007	BASICS	AO-CT	0.2106767	0.2301305
Doumas et al. 2010	e-CHUG	ACT-CT	−0.0831586	0.261467

Quantity of Alcohol Use, 0–3 months

Study	Group 1	Group 2	SMD	SE
Hustad et al. 2010	e-CHUG	AlcoholEdu	0.0465842	0.2679818
Hustad et al. 2010	e-CHUG	AO-CT	0.539308	0.2787348
Hustad et al. 2010	AlcoholEdu	AO-CT	0.4273512	0.286277
Terlecki et al. 2008, 2011	BASICS	AO-CT	0.6233613	0.3143511
Hester et al. 2012	CDCU	AO-CT	0.2004084	0.1766016
LaBrie et al. 2013	BASICS	AO-CT	0.0884182	0.1151534
Kulesza et al. 2013	BASICS	AO-CT	0.309803	0.1514086
Monahan et al. 2013; Murphy et al. 2010, 2015	e-CHUG	AO-CT	0.2672501	0.2289554
Monahan et al. 2013; Murphy et al. 2010, 2015	BASICS	AO-CT	0.4747235	0.2268034
Monahan et al. 2013; Murphy et al. 2010, 2015	BASICS	e-CHUG	0.2134529	0.2258194
Matteucci et al. 2016; Paschall et al. 2011a, b, 2014	AlcoholEdu	AO-CT	0.09587	0.040985
Carey et al. 2011; Carey and DeMartini 2010; Reid et al. 2015	AlcoholEdu	AO-CT	0.0368372	0.1099799
Carey et al. 2011; Carey and DeMartini 2010; Reid et al. 2015	Alcohol 101	AO-CT	0.0840969	0.1095011
Carey et al. 2011; Carey and DeMartini 2010; Reid et al. 2015	Alcohol 101	AlcoholEdu	−0.0471421	0.1109602
Walters et al. 2007	e-CHUG	AO-CT	0.3092513	0.1954149
Lovecchio 2009; Lovecchio et al. 2010	AlcoholEdu	AO-CT	0.2116329	0.0565116
Logan 2013; Logan et al. 2015	BASICS	ACT-CT	−0.069392	0.4097957
Braitman 2012	Alcohol 101	ACT-CT	−0.5063631	0.3743723
Whiteside 2010	BASICS	AO-CT	−0.3714197	0.2354262
Hallett et al. 2009; Hustad and Borsari 2010; Kypri et al. 2009	THRIVE	AO-CT	0.4788996	0.0464172
Juarez 2001; Juarez et al. 2006	e-CHUG	AO-CT	0.1030538	0.3126477
McNally 2003; McNally et al. 2005	BASICS	AO-CT	0.2713031	0.2351786
Dimeff 1997; Dimeff and McNeely 2000	BASICS	AO-CT	0.2934548	0.3580892
Butler 2007; Butler and Correia 2009	BASICS	AO-CT	0.3756984	0.2747413
Geisner 2008	BASICS	AO-CT	−0.008391	0.1549015
Kulesza et al. 2010	BASICS	AO-CT	0.2273163	0.232198
Teeters et al. 2015	BASICS	Alcohol101	0.2123878	0.2420824
Doumas et al. 2009	CYD	ACT-CT	0.1819359	0.2626119
Horner 2010	BASICS	AO-CT	−0.2559756	0.2685234
Lau 2006; Lau-Barraco and Dunn 2008	Alcohol 101	AO-CT	0.1370167	0.2033646
Henslee et al. 2006	BASICS	ACT-CT	0.5521266	0.4491132
Henslee 2008; Henslee and Correia 2009	BASICS	ACT-CT	0.1443547	0.1555335
Hester et al. 2012	CDCU	AO-CT	0.3739442	0.2243072
Terlecki 2011	BASICS	AO-CT	0.6591333	0.1949856
Bryant 2009; Bryant et al. 2013	BASICS	ACT-CT	0.0509202	0.1449789
Borsari and Carey 2000; Borsari et al. 2009	BASICS	AO-CT	0.566366	0.2655833
Leffingwell et al. 2005	CDCU	AO-CT	−0.023189	0.2358896
McPherson 2012	BASICS	e-CHUG	−0.2274426	0.57183
Terry 2012	BASICS	ACT-CT	0.5074358	0.3996419
Martin 2014	BASICS	AO-CT	0.4491305	0.3128194
Bowley et al. 2013	THRIVE	ACT-CT	0.0051054	0.3203622
Terlecki 2008; Terlecki et al. 2010, 2012	BASICS	AO-CT	0.2340654	0.3142519
Terlecki et al. 2015	BASICS	AO-CT	0.3364387	0.1886663

Quantity of Alcohol Use, 3–6 months

Study	Group 1	Group 2	SMD	SE
Mastroleo 2008; Orchowski et al. 2012	BASICS	AO-CT	0.1264261	0.1692169
Murphy et al. 2001	BASICS	ACT-CT	0.2147281	0.2746396
Doumas and Andersen 2009	e-CHUG	AO-CT	0.203054	0.2921708
LaBrie et al. 2013	BASICS	AO-CT	0	0.1184705
Monahan et al. 2013; Murphy et al. 2010, 2015	BASICS	e-CHUG	0.2705603	0.2368688
Monahan et al. 2013; Murphy et al. 2010, 2015	e-CHUG	AO-CT	−0.0735571	0.2329142
Monahan et al. 2013; Murphy et al. 2010, 2015	BASICS	AO-CT	0.1873487	0.2299978
Matteucci et al. 2016; Paschall et al. 2011a, b, 2014	AlcoholEdu	AO-CT	0.0668791	0.040973
Walters et al. 2007	e-CHUG	AO-CT	−0.027574	0.1942664

Table 6 (continued)

Doumas et al. 2011, 2014	e-CHUG	AO-CT	0.3220704	0.2239406
Logan 2013; Logan et al. 2015	BASICS	ACT-CT	− 0.3980921	0.42153
Whiteside 2010	BASICS	AO-CT	− 0.1710859	0.2395801
Hallett et al. 2009; Hustad and Borsari 2010; Kypri et al. 2009	THRIVE	AO-CT	0.4668047	0.0463849
Schaus et al. 2009	BASICS	ACT-CT	0.4007094	0.1402939
Teeters et al. 2015	BASICS	Alcohol 101	0.2978334	0.2501741
Doumas and Haustveit 2008	CYD	ACT-CT	0.1306662	0.3499727
Horner 2010	BASICS	AO-CT	− 0.137309	0.2677464
Eggleston 2007	BASICS	AO-CT	0.3614697	0.2313605
Terlecki 2011	BASICS	AO-CT	1.01573	0.2017665
Miller 2000	Alcohol 101	AO-CT	0.4582334	0.1331391
Doumas et al. 2010	e-CHUG	ACT-CT	0.2787872	0.1964494
Leffingwell et al. 2005	CDCU	AO-CT	0.0772968	0.2359686
McPherson 2012	BASICS	e-CHUG	− 0.152507	0.586367
Peak Quantity of Alcohol Use, 0–3 months				
Study	Group 1	Group 2	SMD	SE
Hustad et al. 2010	e-CHUG	AlcoholEdu	− 0.2078722	0.2686646
Hustad et al. 2010	e-CHUG	AO-CT	0.4893903	0.2778806
Hustad et al. 2010	AlcoholEdu	AO-CT	0.689066	0.2913354
Terlecki et al. 2008, 2011	BASICS	AO-CT	0.6434117	0.3148205
Hester et al. 2012	CDCU	AO-CT	0.3578802	0.1775564
LaBrie et al. 2013	BASICS	AO-CT	0.0447274	0.1151116
Lovecchio 2009; Lovecchio et al. 2010	AlcoholEdu	AO-CT	0.1971226	0.0618565
Braitman 2012	Alcohol 101	ACT-CT	− 0.5507706	0.3750387
Geisner 2008	BASICS	AO-CT	− 0.1464643	0.155108
Doumas et al. 2009	CYD	ACT-CT	− 0.2924118	0.2634284
Horner 2010	BASICS	AO-CT	− 0.1405973	0.2677616
Henslee et al. 2006	BASICS	ACT-CT	0.3142058	0.4436159
Henslee 2008; Henslee and Correia 2009	BASICS	ACT-CT	0.1097072	0.1554483
Hester et al. 2012	CDCU	AO-CT	0.5151408	0.226028
Terlecki 2011	BASICS	AO-CT	0.9787726	0.200942
Leffingwell et al. 2005	CDCU	AO-CT	0.1923832	0.2364186
McPherson 2012	BASICS	e-CHUG	0.1058105	0.5704653
Martin 2014	BASICS	AO-CT	0.6454946	0.316883
Terlecki 2008; Terlecki et al. 2010, 2012	BASICS	AO-CT	0.2546021	0.3144466
Terlecki et al. 2015	BASICS	AO-CT	0.6002765	0.1915174
Peak Quantity of Alcohol Use, 3–6 months				
Study	Group 1	Group 2	SMD	SE
LaBrie et al. 2013	BASICS	AO-CT	0.1391386	0.1186138
Doumas et al. 2011, 2014	e-CHUG	AO-CT	0.228673	0.2232392
Schaus et al. 2009	BASICS	ACT-CT	1.235356	0.15149
Doumas and Haustveit 2008	CYD	ACT-CT	0.100941	0.3498237
Horner 2010	BASICS	AO-CT	− 0.1820898	0.2679848
Terlecki 2011	BASICS	AO-CT	1.028616	0.2020603
Miller 2000	Alcohol 101	AO-CT	0.422202	0.1328821
Doumas et al. 2010	e-CHUG	ACT-CT	0.071086	0.195575
Alfonso et al. 2012; Alfonso and Deschenes 2013	BASICS	e-CHUG	0.2179615	0.199841
Leffingwell et al. 2005	CDCU	AO-CT	− 0.2748331	0.2369759
McPherson 2012	BASICS	e-CHUG	− 0.7404641	0.6047333

Appendix 2 – Additional Networks

Frequency of Heavy Alcohol Use: 3–6 Months Frequency of heavy alcohol use outcomes measured at 3–6 months post-intervention represents a limited network (Fig. 3a) of 12 studies comparing five interventions, and the AO-CT and ACT-CT groups. As the size of the nodes indicate, BASICS was compared the most, while the other interventions were only studied

once, with the exception of e-CHUG, which was included in two trials. The most informative direct evidence in this network was for BASICS versus AO-CT as it contributed 20.9%, which is likely overly influential given the representation of other interventions in this network. Results from the consistency model indicate that relative to the AO-CT, no comparisons significantly reduced the frequency of heavy drinking (Table 7). The global test for inconsistency was non-significant ($\chi^2(1) = 0.03$, $p = 0.8518$), and for the single loop available, there was no evidence

Table 7 Intervention effects compared to assessment-only control groups

	3–6 months			3–6 months		
	Heavy frequency			Peak quantity		
	SMD	LCL	UCL	SMD	LCL	UCL
Alc101	0.33	−0.22	0.87	−0.42	−1.48	0.64
AlcEDU	0.05	−0.03	0.13	NA	NA	NA
BASICS	−0.05	−0.28	0.18	−0.36	−0.94	0.22
CYD	0.49	−0.27	1.25	0.30	−1.26	1.87
e-CHUG	0.04	−0.24	0.33	−0.16	−0.91	0.59
CDCU	NA	NA	NA	0.27	−0.85	1.40
THRIVE	−0.07	−0.40	0.26	NA	NA	NA
ACT-CT	0.42	0.09	0.76	0.41	−0.55	1.36
	ES replaced ^{a1}			ES replaced ^{a2}		
Alc101	0.33	−0.22	0.87	−0.42	−0.99	0.14
AlcEDU	0.05	−0.03	0.13	NA	NA	NA
BASICS	−0.05	−0.28	0.18	−0.22	−0.55	0.11
CYD	0.43	−0.33	1.19	−0.13	−1.13	0.88
e-CHUG	0.04	−0.24	0.33	−0.17	−0.62	0.28
CDCU	NA	NA	NA	0.27	−0.41	0.96
THRIVE	−0.07	−0.40	0.26	NA	NA	NA
ACT-CT	0.37	0.05	0.68	−0.03	−0.56	0.51
				Mandated removed		
Alc101				−0.42	−1.66	0.82
AlcEDU				NA	NA	NA
BASICS				−0.54	−1.33	0.26
CYD				0.14	−1.67	1.96
e-CHUG				−0.30	−1.30	0.71
CDCU				0.27	−1.02	1.57
THRIVE				NA	NA	NA
ACT-CT				0.24	−0.92	1.41

NA not applicable, *Alcohol101* Alcohol 101/Alcohol 101 Plus, *AlcEDU* AlcoholEdu, *AO-CT* assessment-only control, *ACT-CT* active control, *BASICS* Brief Alcohol Screening Intervention for College Students, *CYD* Check Your Drinking, *CDCU* College Drinker’s Check-up, *ECHUG* Electronic CHECKUP TO GO, *THRIVE* Tertiary Health Research Intervention Via Email

^a This analysis involved replacing an effect size used in the original network analysis due to dependency between outcomes reported within the same follow-up wave category with the dropped effect size: (1) One study replaced the 25.8-week follow-up ES with the 12.9-week follow-up. (2) Three studies replaced the 25.8-week follow-up ES with the 12.9-week follow-up ES

of loop-specific inconsistency. Node-splitting also indicated that the consistency model fit well. When compared against all other interventions BASICS had the largest SUCRA value, followed by THRIVE (Table 8)

Sensitivity Analysis: Frequency of Heavy Alcohol Use, 3–6 Months Sensitivity analysis replacing the measurement

Table 8 Treatment rankings

	3–6 months		3–6 months	
	Heavy frequency		Peak quantity	
	SUCRA	PrBest	SUCRA	PrBest
Alc101	27.2	3.9	73.6	42.1
AlcEDU	53.5	0.6	NA	NA
BASICS	77.9	31.4	77.8	26
CYD	18.6	4.1	35	11.6
e-CHUG	60.2	14.8	61.3	11.8
CDCU	NA	NA	33.3	7.2
THRIVE	77	39.8	NA	NA
ACT-CT	14.8	0	21.8	0.4
AO-CT	70.8	5.4	47.3	0.9
	ES removed ^{a1}		ES removed ^{a2}	
Alc101	26.2	4.9	81.2	52.4
AlcEDU	52.2	1.9	NA	NA
BASICS	77.7	26.8	68.3	9.7
CYD	21.8	7.1	52.5	24.7
e-CHUG	58.6	14.2	58.6	8.7
CDCU	NA	NA	15.2	1.7
THRIVE	75.7	39.6	NA	NA
ACT-CT	17.3	0	39.7	2.8
AO-CT	70.5	5.5	34.5	0
			Mandated removed	
Alc101			66.4	31.1
AlcEDU			NA	NA
BASICS			78.5	29.6
CYD			40.1	15.1
e-CHUG			61.2	15.3
CDCU			33.6	6.5
THRIVE			NA	NA
ACT-CT			28.4	1.3
AO-CT			42	1.1

Italicized values indicate highest SUCRA value of that intervention in that network

SUCRA Surface under the curve, *PrBest* Probability that the treatment will perform the best, *NA* not applicable, *Alcohol101* Alcohol 101/Alcohol 101 Plus, *AlcEDU* AlcoholEdu, *AO-CT* Assessment-only control, *ACT-CT* active control, *BASICS* Brief Alcohol Screening Intervention for College Students, *CYD* Check Your Drinking, *CDCU* College Drinker’s Check-up, *ECHUG* Electronic CHECKUP TO GO, *THRIVE* Tertiary Health Research Intervention Via Email

^a This analysis involved replacing an effect size used in the original network analysis due to dependency between outcomes reported within the same follow-up wave category with the dropped effect size: (1) One study replaced the 25.8-week follow-up ES with the 12.9-week follow-up. (2) Three studies replaced the 25.8-week follow-up ES with the 12.9-week follow-up ES

duration from one study (12.9 versus 25.8 weeks: Schaus et al. 2009) did not change the results (Tables 7 and 8)

Quantity of Alcohol Use During Peak Consumption: 3–6 Months Quantity of alcohol use during peak consumption measured at 3–6 months post-intervention is a network (Fig. 3c) of 11 studies comparing five interventions and both types of control groups. As the size of the nodes indicate, BASICS was compared the most, while the other interventions were only studied once, with the exception of e-CHUG, which was included in four trials. The most informative direct evidence in this network was for e-CHUG versus AO-CT as it contributed 17.3%. Results from the consistency model indicate that relative to the AO-CT, no comparisons significantly reduced the quantity of alcohol consumption during peak drinking episodes (Table 7). The global test for inconsistency was non-significant ($\chi^2(2) = 1.73, p = 0.4216$); however, one loop (BASICS, e-CHUG, ACT-CT) demonstrated significant inconsistency (IF = 1.04, 95% CI [0.42, 1.65]). Node-splitting, however, supported the consistency model. When compared against all other interventions BASICS had the largest SUCRA value, followed by Alcohol101 (Table 8)

Sensitivity Analysis: Quantity of Alcohol Use during Peak Consumption, 3–6 Months Removing the two studies that utilized a mandated sample from this network (Alfonso et al. 2012; Horner 2010) resulted in removing one comparison involving e-CHUG and two comparisons of BASICS, but did not substantively change the model results (Tables 7 and 8). The global test for inconsistency remained non-significant ($\chi^2 = 2.35, df = 2, p = 0.3086$). There was, however, evidence of significant loop-specific inconsistency for two evidence loops: (1) BASICS, e-CHUG, ACT-CT, IF = 1.89, 95% CI [0.59, 3.20] and (2) BASICS, e-CHUG, AO-CT, IF = 0.87, 95% CI [0.00, 2.17]

Sensitivity analysis replacing the measurement duration from three studies (12.9 versus 25.8 weeks: Labrie et al. 2013; Schaus et al. 2009; Terlecki et al. 2015) did not change the results of the effects compared to the AO-CT, but Alcohol101 and BASICS switched their rankings such that Alcohol 101 had the largest SUCRA value (Tables 7 and 8)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Abbey, A., Wegner, R., Woerner, J., Pegram, S. E., & Pierce, J. (2014). Review of survey and experimental research that examines the relationship between alcohol consumption and men's sexual aggression perpetration. *Trauma, Violence, & Abuse, 15*, 265–282.
- Alfonso, J., & Deschenes, S. D. (2013). Do drinking games matter? An examination by game type and gender in a mandated student sample. *The American Journal of Drug and Alcohol Abuse, 39*(5), 312–319. <https://doi.org/10.3109/00952990.2013.770519>.
- Barry, A. E. (2007). Using theory-based constructs to explore the impact of Greek membership on alcohol-related beliefs and behaviors: A systematic literature review. *Journal of American College Health, 56*, 307–315.
- Buscemi, J., Murphy, J. G., Martens, M. P., McDevitt-Murphy, M. E., Dennhardt, A. A., & Skidmore, J. R. (2010). Help-seeking for alcohol-related problems in college students: Correlates and preferred resources. *Psychology of Addictive Behaviors, 24*, 571.
- Carey, K. B., Scott-Sheldon, L. A., Garey, L., Elliott, J. C., & Carey, M. P. (2016). Alcohol interventions for mandated college students: A meta-analytic review. *Journal of Consulting and Clinical Psychology, 84*, 619–632.
- Chaimani, A., Caldwell, D. M., Li, T., Higgins, J. P., & Salanti, G. (2017). Additional considerations are required when preparing a protocol for a systematic review with multiple interventions. *Journal of Clinical Epidemiology, 83*, 65–74.
- Chaimani, A., Higgins, J. P., Mavridis, D., Spyridonos, P., & Salanti, G. (2013). Graphical tools for network meta-analysis in Stata. *PLoS One, 8*, e76654.
- Cipriani, A., Higgins, J. P., Geddes, J. R., & Salanti, G. (2013). Conceptual and technical challenges in network meta-analysis. *Annals of Internal Medicine, 159*(2), 130–137.
- Dias, S., Welton, N. J., Caldwell, D. M., & Ades, A. E. (2010). Checking consistency in mixed treatment comparison meta-analysis. *Statistics in Medicine, 29*, 932–944.
- Dimeff, L. A., Baer, J. S., Kivlahan, D. R., & Marlatt, G. A. (Ed.). (1999). *Brief alcohol screening and intervention for college students (BASICS): A harm reduction approach*. Guilford Press.
- Fachini, A., Aliane, P. P., Martinez, E. Z., & Furtado, E. F. (2012). Efficacy of brief alcohol screening intervention for college students (BASICS): A meta-analysis of randomized controlled trials. *Substance Abuse Treatment, Prevention, and Policy, 7*, 1.
- Foster, C., Caravelis, C., & Kopak, A. (2014). National College Health Assessment Measuring Negative Alcohol-Related Consequences among college students. *American Journal of Public Health Research, 2*, 1–5.
- Grant, S., Pedersen, E. R., Osilla, K. C., Kulesza, M., & D'amico, E. J. (2016). Reviewing and interpreting the effects of brief alcohol interventions: Comment on a Cochrane review about motivational interviewing for young adults. *Addiction, 111*, 1521–1527.
- Ham, L. S., & Hope, D. A. (2003). College students and problematic drinking: A review of the literature. *Clinical Psychology Review, 23*, 719–759.
- Higgins, J. P., Altman, D. G., Gøtzsche, P. C., Jüni, P., Moher, D., Oxman, A. D., et al. (2011). The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ, 343*, d5928. <https://doi.org/10.1136/bmj.d5928>.
- Hingson, R., & White, A. (2014). New research findings since the 2007 Surgeon General's Call to Action to Prevent and Reduce Underage Drinking: A review. *Journal of Studies on Alcohol and Drugs, 75*(1), 158–169.
- Huh, D., Mun, E. Y., Larimer, M. E., White, H. R., Ray, A. E., Rhew, I. C., et al. (2015). Brief motivational interventions for college student drinking may not be as powerful as we think: An individual participant-level data meta-analysis. *Alcoholism: Clinical and Experimental Research, 39*, 919–931.
- Hutton, B., Salanti, G., Caldwell, D. M., Chaimani, A., Schmid, C. H., Cameron, C., ... & Mulrow, C. (2015). The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: Checklist and explanations. *Annals of Internal Medicine, 162*(11), 777–784.
- Jennison, K. M. (2004). The short-term effects and unintended long-term consequences of binge drinking in college: A 10-year follow-up study. *The American Journal of Drug and Alcohol Abuse, 30*, 659–684.
- Kingree, J. B., & Thompson, M. (2015). A comparison of risk factors for alcohol-involved and alcohol-uninvolved sexual aggression perpetration. *Journal of Interpersonal Violence, 30*, 1478–1492.

- Lipsey, M. W., & Wilson, D. B. (2001). *Practical meta-analysis*. Sage Publications, Inc.
- Mavridis, D., Giannatsi, M., Cipriani, A., & Salanti, G. (2015). A primer on network meta-analysis with emphasis on mental health. *Evidence Based Mental Health, 18*, 40–46.
- Mavridis, D., & Salanti, G. (2014). Exploring and accounting for publication bias in mental health: A brief overview of methods. *Evidence Based Mental Health, 17*, 11–15. <https://doi.org/10.1136/eb-2013-101700>.
- Meisel, M. K., & Barnett, N. P. (2017). Protective and risky social network factors for drinking during the transition from high school to college. *Journal of Studies on Alcohol and Drugs, 78*, 922–929.
- Pedersen, D. E. (2013). Gender differences in college binge drinking: Examining the role of depression and school stress. *The Social Science Journal, 50*, 521–529.
- Peters, J. L., Sutton, A. J., Jones, D. R., Abrams, K. R., & Rushton, L. (2008). Contour-enhanced meta-analysis funnel plots help distinguish publication bias from other causes of asymmetry. *Journal of Clinical Epidemiology, 61*, 991–996.
- Petropoulou, M., Nikolakopoulou, A., Veroniki, A. A., Rios, P., Vafaei, A., Zarin, W., ... & Egger, M. (2017). Bibliographic study showed improving statistical methodology of network meta-analyses published between 1999 and 2015. *Journal of Clinical Epidemiology, 82*, 20–28. <https://doi.org/10.1016/j.jclinepi.2016.11.002>
- Pildal, J., Hrobjartsson, A., Jørgensen, K. J., Hilden, J., Altman, D. G., & Gøtzsche, P. C. (2007). Impact of allocation concealment on conclusions drawn from meta-analyses of randomized trials. *International Journal of Epidemiology, 36*, 847–857.
- Salanti, G. (2012). Indirect and mixed-treatment comparison, network, or multiple-treatments meta-analysis: Many names, many benefits, many concerns for the next generation evidence synthesis tool. *Research Synthesis Methods, 3*, 80–97.
- Salanti, G., Ades, A. E., & Ioannidis, J. P. (2011). Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: An overview and tutorial. *Journal of Clinical Epidemiology, 64*, 163–171.
- SAMHSA (2016a). 2016 National Survey on Drug Use and Health (NSDUH). In *Table 6.80B—Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2015 and 2016*. Available at: <https://www.samhsa.gov/data/sites/default/files/NSDUH-DetTabs-2016/NSDUH-DetTabs-2016.htm#tab6-80B>. Accessed 23 Jan 2018.
- SAMHSA (2016b). 2016 National Survey on drug use and health (NSDUH). In *Table 6.79B*. Available at: <https://www.samhsa.gov/data/sites/default/files/NSDUH-DetTabs-2016/NSDUH-DetTabs-2016.htm#tab6-79B>. Accessed 23 Jan 2018.
- SAMHSA (2018). *Key substance use and mental health indicators in the United States: Results from the 2017 National Survey on Drug Use and Health (HHS Publication No. SMA 18-5068, NSDUH Series H-53)*. Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Retrieved from: <https://www.samhsa.gov/data/>.
- Sánchez-Meca, J., Marin-Martínez, F., & Chacón-Moscó, S. (2003). Effect-size indices for dichotomized outcomes in meta-analysis. *Psychological Methods, 8*(4), 448.
- Schulenberg, J. E., Johnston, L. D., O'Malley, P. M., Bachman, J. G., Meich, R. A., & Patrick, M. E. (2017). *Monitoring the future national survey results on drug use, 1975–2016: Volume 2, college students and adults ages 19–55*. Ann Arbor: Institute for Social Research, The University of Michigan.
- StataCorp. (2015). *Stata statistical software: Release 14*. College Station: StataCorp LP.
- Steinka-Fry, K. T., Tanner-Smith, E. E., & Grant, S. (2015). Effects of 21st birthday brief interventions on college student celebratory drinking: A systematic review and meta-analysis. *Addictive Behaviors, 50*, 13–21. <https://doi.org/10.1016/j.addbeh.2015.06.001>.
- Tanner-Smith, E. E., & Lipsey, M. W. (2015). Brief alcohol interventions for adolescents and young adults: A systematic review and meta-analysis. *Journal of Substance Abuse Treatment, 51*, 1–18.
- Tanner-Smith, E. E., & Risser, M. D. (2016). A meta-analysis of brief alcohol interventions for adolescents and young adults: Variability in effects across alcohol measures. *The American Journal of Drug and Alcohol Abuse, 42*, 140–151.
- The Century Council (1997). *Alcohol 101*. Los Angeles.
- The Century Council (2003). *Alcohol 101 Plus [Computer Software] 2003*. Retrieved from <http://www.alcohol101plus.org/home.html>. Accessed 23 Jan 2018.
- Valenstein-Mah, H., Larimer, M., Zoellner, L., & Kaysen, D. (2015). Blackout drinking predicts sexual revictimization in a college sample of binge-drinking women. *Journal of Traumatic Stress, 28*, 484–488.
- Voloshyna, D. M., Bonar, E. E., Cunningham, R. M., Ilgen, M. A., Blow, F. C., & Walton, M. A. (2016). Blackouts among male and female youth seeking emergency department care. *American Journal of Drug and Alcohol Abuse, 1–11*.
- Wechsler, H., Davenport, A., Dowdall, G., Moeykens, B., & Castillo, S. (1994). Health and behavioral consequences of binge drinking in college: A National Survey of students at 140 campuses. *Journal of the American Medical Association, 272*, 1672–1677. <https://doi.org/10.1001/jama.1994.03520210056032>.
- White, I. R. (2015). Network meta-analysis. *Stata Journal, 15*, 951–985.
- White, I. R., Barrett, J. K., Jackson, D., & Higgins, J. P. (2012). Consistency and inconsistency in network meta-analysis: model estimation using multivariate meta-regression. *Research Synthesis Methods, 3*(2), 111–125.

Further Reading (References of studies included in the review)

- Alfonso, J., Hall, T. V., & Dunn, M. E. (2012). Feedback-based alcohol interventions for mandated students: An effectiveness study of three modalities. *Clinical Psychology and Psychotherapy, 19*, 1786. <https://doi.org/10.1002/cpp.1786>.
- Baer, J. S., Kivlahan, D. R., & Marlatt, G. A. (1995). High-risk drinking across the transition from high school to college. *Alcoholism: Clinical and Experimental Research, 19*, 54–61.
- Baer, J. S., Kivlahan, D. R., Blume, A. W., McKnight, P., & Marlatt, G. A. (2001). Brief intervention for heavy-drinking college students: 4-year follow-up and natural history. *American Journal of Public Health, 91*, 1310–1316. <https://doi.org/10.2105/ajph.91.8.1310>.
- Borsari, B., & Carey, K. B. (2000). Effects of a brief motivational intervention with college student drinkers. *Journal of Consulting and Clinical Psychology, 68*, 728–733. <https://doi.org/10.1037/0022-006x.68.4.728>.
- Borsari, B., Murphy, J. G., & Carey, K. B. (2009). Readiness to change in brief motivational interventions: A requisite condition for drinking reductions? *Addictive Behaviors, 34*, 232–235. <https://doi.org/10.1016/j.addbeh.2008.10.010>.
- Bowley, C., Faricy, C., Hegarty, B., Johnstone, S. J., Smith, J. L., Kelly, P. J., & Rushby, J. A. (2013). The effects of inhibitory control training on alcohol consumption, implicit-alcohol related cognitions and brain electrical activity. *International Journal of Psychophysiology, 89*, 342–348. <https://doi.org/10.1016/j.ijpsycho.2013.04.011>.
- Braitman, A. L. (2012). *The effects of personalized boosters for a computerized intervention targeting college student drinking*. (Doctoral dissertation, Old Dominion University).

- Bryant, Z. (2009). *Testing the effectiveness of e-mailed BASICS feedback with college students* (Doctoral dissertation, Auburn University). Dissertation Abstracts International, Ph.D., 198. (University Microfilms No. 3386183).
- Bryant, Z. E., Henslee, A. M., & Correia, C. J. (2013). Testing the effects of e-mailed personalized feedback on risky alcohol use among college students. *Addictive Behaviors*, 38, 2563–2567. <https://doi.org/10.1016/j.addbeh.2013.06.007>.
- Butler, L. H. (2007). *Brief alcohol intervention with college students using BASICS: Face-to-face versus computerized feedback* (Doctoral dissertation, Auburn University). Dissertation Abstracts International, M.S., 83.
- Butler, L. H., & Correia, C. J. (2009). Brief alcohol intervention with college student drinkers: Face-to-face versus computerized feedback. *Psychology of Addictive Behaviors*, 23, 163–167. <https://doi.org/10.1037/a0014892>.
- Carey, K. B., & DeMartini, K. S. (2010). The motivational context for mandated alcohol interventions for college students by gender and family history. *Addictive Behaviors*, 35, 218–229. <https://doi.org/10.1016/j.addbeh.2009.10.011>.
- Carey, K. B., Carey, M. P., Henson, J. M., Maisto, S. A., & DeMartini, K. S. (2011). Brief alcohol interventions for mandated college students: Comparison of face-to-face counseling and computer-delivered interventions. *Addiction*, 106, 528–537. <https://doi.org/10.1111/j.1360-0443.2010.03193.x>.
- Croom, K., Lewis, D., Marchell, T., Lesser, M. L., Reyna, V. F., Kubicki-Bedford, L., Feffer, M., & Staiano-Coico, L. (2009). Impact of an online alcohol education course on behavior and harm for incoming first-year college students: Short-term evaluation of a randomized trial. *Journal of American College Health*, 57, 445–454. <https://doi.org/10.3200/jach.57.4.445-454>.
- Dimeff, L. A. (1997). *Brief intervention for heavy and hazardous college drinkers in a student primary health care setting* (Doctoral dissertation, University of Washington). Dissertation Abstracts International, Ph.D.(58, 12-B), 96. (University Microfilms No. 9819231).
- Dimeff, L. A., & McNeely, M. (2000). Computer-enhanced primary care practitioner advice for high-risk college drinkers in a student primary health-care setting. *Cognitive and Behavioral Practice*, 7, 82–100. [https://doi.org/10.1016/s1077-7229\(00\)80010-3](https://doi.org/10.1016/s1077-7229(00)80010-3).
- Doumas, D. M., & Andersen, L. L. (2009). Reducing alcohol use in first-year university students: Evaluation of a web-based personalized feedback program. *Journal of College Counseling*, 12, 18–32. <https://doi.org/10.1002/j.2161-1882.2009.tb00037.x>.
- Doumas, D. M., & Haustveit, T. (2008). Reducing heavy drinking in intercollegiate athletes: Evaluation of a web-based personalized feedback program. *The Sport Psychologist*, 22, 212–228.
- Doumas, D. M., Haustveit, T., & Coll, K. M. (2010). Reducing heavy drinking among first year intercollegiate athletes: A randomized controlled trial of web-based normative feedback. *Journal of Applied Sport Psychology*, 22, 247–261. <https://doi.org/10.1080/10413201003666454>.
- Doumas, D. M., Kane, C. M., Navarro, B., & Roman, J. (2011). Decreasing heavy drinking in first-year students: Evaluation of a web-based personalized feedback program administered during orientation. *Journal of College Counseling*, 14, 5–20. <https://doi.org/10.1002/j.2161-1882.2011.tb00060.x>.
- Doumas, D. M., McKinley, L. L., & Book, P. (2009). Evaluation of two web-based alcohol interventions for mandated college students. *Journal of Substance Abuse Treatment*, 36, 65–74. <https://doi.org/10.1016/j.jsat.2008.05.009>.
- Doumas, D. M., Nelson, K., DeYoung, A., & Renteria, C. C. (2014). Alcohol-related consequences among first-year university students: Effectiveness of a web-based personalized feedback program. *Journal of College Counseling*, 17, 150–162. <https://doi.org/10.1002/j.2161-1882.2014.00054.x>.
- Eggleston, A. M. (2007). *Components analysis of a brief intervention for college drinkers* (Doctoral dissertation, The Ohio State University). Dissertation Abstracts International, Ph.D.(68, 08), 133. (University Microfilms No. 3276684).
- Fernandez, A. C., Wood, M. D., Laforge, R., & Black, J. T. (2011). Randomized trials of alcohol-use interventions with college students and their parents: Lessons from the transitions project. *Clinical Trials*, 8, 205–213. <https://doi.org/10.1177/1740774510396387>.
- Geisner, I. (2008). *A randomized clinical trial of a brief, mailed intervention for depressed mood in a college student sample* (doctoral dissertation, University of Washington). Dissertation abstracts international, Ph.D., 139. (University Microfilms No. 3328400).
- Geisner, I. M., Neighbors, C., Lee, C. M., & Larimer, M. E. (2007). Evaluating personal alcohol feedback as a selective prevention for college students with depressed mood. *Addictive Behaviors*, 32, 2776–2787. <https://doi.org/10.1016/j.addbeh.2007.04.014>.
- Grossbard, J. R., Mastrole, N. R., Kilmer, J. R., Lee, C. M., Turrisi, R., Larimer, M. E., & Ray, A. (2010). Substance use patterns among first-year college students: Secondary effects of a combined alcohol intervention. *Journal of Substance Abuse Treatment*, 39, 384–390. <https://doi.org/10.1016/j.jsat.2010.07.001>.
- Hallett, J., Maycock, B., Kypri, K., Howat, P., & McManus, A. (2009). Development of a web-based alcohol intervention for university students: Processes and challenges. *Drug and Alcohol Review*, 28, 31–39. <https://doi.org/10.1111/j.1465-3362.2008.00008.x>.
- Henslee, A. (2008). *Providing personalized feedback regarding alcohol use in a group format to college freshmen* (Doctoral dissertation, Auburn University). Dissertation Abstracts International, Ph.D.(69, 10), 97. (University Microfilms No. 3333129).
- Henslee, A. M., & Correia, C. J. (2009). The use of freshmen seminar programs to deliver personalized feedback. *Journal of Alcohol and Drug Education*, 53, 39–52.
- Henslee, A. M., Irons, J. G., Day, J. M., Butler, L., Benson, T. A., & Correia, C. J. (2006). Using national alcohol screening day to deliver personalized feedback: A pilot study. *Journal of Drug Education*, 36, 271–278. <https://doi.org/10.2190/1u8h-41u2-5k03-0j24>.
- Hester, R. K., Delaney, H. D., & Campbell, W. (2012). The college drinker's check-up: Outcomes of two randomized clinical trials of a computer-delivered intervention (STUDY ID 3099 IS FOR EXPERIMENT 2). *Psychology of Addictive Behaviors*, 26, 1–12. <https://doi.org/10.1037/a0024753>.
- Horner, K. (2010). *Brief motivational interviewing: An intervention for alcohol abusing college students* (doctoral dissertation, State University of New York at Albany). Dissertation Abstracts International, Ph.D.(71, 05), 146. (University Microfilms No. 3402364).
- Hustad, J. T. P., & Borsari, B. (2010). Web-based screening and brief motivational intervention reduces alcohol use in heavy-drinking undergraduates at up to 6 months. *Evidence Based Medicine*, 15, 17–18. <https://doi.org/10.1136/ebm1015>.
- Hustad, J. T. P., Barnett, N. P., Borsari, B., & Jackson, K. M. (2010). Web-based alcohol prevention for incoming college students: A randomized controlled trial. *Addictive Behaviors*, 35, 183–189. <https://doi.org/10.1016/j.addbeh.2009.10.012>.
- Juarez, P. (2001). *A randomized trial of motivational interviewing and feedback on heavy drinking college students* (Doctoral dissertation, University of New Mexico, Albuquerque, New Mexico). Dissertation Abstracts International, M.S., 147.
- Juarez, P., Walters, S. T., Daugherty, M., & Radi, C. (2006). A randomized trial of motivational interviewing and feedback with heavy drinking college students. *Journal of Drug Education*, 36, 233–246. <https://doi.org/10.2190/753n-8242-727t-g63l>.
- Kerr-Corrêa, F., Simão, M. O., Sumaia, S. I., Trinca, L. A., Floripes, T. M. F., Dalben, I., Martins, R. A., Oliveira, J. B., Cavarani, M. B., & Turchi, A. M. (2008). Prevention of risk drinking among university students in a Brazilian university: A clinical trial with a two year

- follow-up. In *Paper presented at the meeting of the 5th international conference of Inebria Alcohol & Drug Problems in developing countries, Ribeirão Preto, Brazil*.
- Kulesza, M., Apperson, M., Larimer, M. E., & Copeland, A. L. (2010). Brief alcohol intervention for college drinkers: How brief is? *Addictive Behaviors*, *35*, 730–733. <https://doi.org/10.1016/j.addbeh.2010.03.011>.
- Kulesza, M., McVay, M. A., Larimer, M. E., & Copeland, A. L. (2013). A randomized clinical trial comparing the efficacy of two active conditions of a brief intervention for heavy college drinkers. *Addictive Behaviors*, *38*, 2094–2101. <https://doi.org/10.1016/j.addbeh.2013.01.008>.
- Kypri, K., Hallett, J., Howat, P., McManus, A., Maycock, B., Bowe, S., & Horton, N. J. (2009). Randomized controlled trial of proactive web-based alcohol screening and brief intervention for university students. *Archives of Internal Medicine*, *169*, 1508–1514. <https://doi.org/10.1001/archinternmed.2009.249>.
- LaBrie, J. W., Lewis, M. A., Atkins, D. C., Neighbors, C., Zheng, C., Kenney, S. R., ... Ghaidarov, T. M. (2013). RCT of web-based personalized normative feedback for college drinking prevention: Are typical student norms good enough? *Journal of Consulting and Clinical Psychology*, *81*, 1074–1086. <https://doi.org/10.1037/a0034087>.
- Larimer, M. E., Lee, C. M., Kilmer, J. R., Fabiano, P. M., Stark, C. B., Geisner, I. M., Mallett, K. A., Lostutter, T. W., Cronce, J. M., Feeney, M., & Neighbors, C. (2007). Personalized mailed feedback for college drinking prevention: A randomized clinical trial. *Journal of Consulting and Clinical Psychology*, *75*, 285–293. <https://doi.org/10.1037/0022-006x.75.2.285>.
- Larimer, M. E., Turner, A. P., Anderson, B. K., Fader, J. S., Kilmer, J. R., Palmer, R. S., & Cronce, J. M. (2001). Evaluating a brief alcohol intervention with fraternities. *Journal of Studies on Alcohol*, *62*, 370–380.
- Lau-Barraco, C., & Dunn, M. E. (2008). Evaluation of a single-session expectancy challenge intervention to reduce alcohol use among college students. *Psychology of Addictive Behaviors*, *22*, 168–175. <https://doi.org/10.1037/0893-164x.22.2.168>.
- Lau, H. C. (2006). *Development and evaluation of a single-session expectancy challenge intervention to reduce alcohol use among heavy drinking college students* (doctoral dissertation, University of Central Florida). Dissertation abstracts international, Ph.D.(67, 9-B). (University Microfilms No. 3233658).
- Leffingwell, T. R., Leedy, M. J., & Lack, C. W. (2005). A multimedia computer-based intervention for college student drinking: Short-term outcomes of a randomized trial. In *Paper presented at the meeting of the association for behavioral and cognitive therapies, Washington, DC*.
- Logan, D. E. (2013). *Alcohol interventions for mandated students* (project AIMS) (doctoral dissertation, University of Washington).
- Logan, D. E., Kilmer, J. R., King, K. M., & Larimer, M. E. (2015). Alcohol interventions for mandated students: Behavioral outcomes from a randomized controlled pilot study. *Journal of Studies on Alcohol and Drugs*, *76*, 31–37. <https://doi.org/10.15288/jsad.76.1.31>.
- Lovecchio, C. P. (2009). *On-line alcohol education: Impact on knowledge, attitudes and behaviors of first-year college students* (Doctoral Dissertation, Villanova University).
- Lovecchio, C. P., Wyatt, T. M., & DeJong, W. (2010). Reductions in drinking and alcohol-related harms reported by first-year college students taking an online alcohol education course: A randomized trial. *Journal of Health Communication*, *15*, 805–819. <https://doi.org/10.1080/10810730.2010.514032>.
- Martin, J. L. (2014). Does personalized feedback enhance outcomes for students who drink to cope? A pilot study. *Alcoholism: Clinical and Experimental Research*, *38*, 195–199. <https://doi.org/10.1111/acer.12288>.
- Mastroleto, N. (2008). *Comparison of supervision training techniques in a motivational enhancement intervention on college student drinking* (Doctoral dissertation, The Pennsylvania State University). Dissertation Abstracts International, Ph.D.(69, 11), 180. (University Microfilms No. 3336087).
- Matteucci, A., Reif, S., & Paschall, M. J. (2016). Detecting variation in college student drinking patterns: An argument for creating alcohol-free environments and implementing campus policies. *Alcoholism: Clinical and Experimental Research*, *40*, 122A–122A.
- McNally, A. M. (2003). *Motivational interventions for problematic alcohol use: In search of the mechanisms of action* (Doctoral dissertation, Boston University). Dissertation Abstracts International, Ph.D.(63, 10-B), 144. (University Microfilms No. 3067197).
- McNally, A. M., Palfai, T. P., & Kahler, C. W. (2005). Motivational interventions for heavy drinking college students: Examining the role of discrepancy-related psychological processes. *Psychology of Addictive Behaviors*, *19*, 79–87. <https://doi.org/10.1037/0893-164x.19.1.79>.
- McPherson, P. (2012). *Efficacy of brief alcohol interventions in an Australian tertiary education setting* (Doctoral dissertation, Royal Melbourne Institute of Technology University, Melbourne, Australia). Dissertation Abstracts International, Ph.D., 274.
- Miller, E. T. (2000). *Preventing alcohol abuse and alcohol-related negative consequences among freshmen college students: Using emerging computer technology to deliver and evaluate the effectiveness of brief intervention efforts* (Doctoral dissertation, University of Washington). Dissertation Abstracts International, Ph.D.(61, 8-B), 271. (University Microfilms No. 9983525).
- Monahan, C. J., McDevitt-Murphy, M. E., Dennhardt, A. A., Skidmore, J. R., Martens, M. P., & Murphy, J. G. (2013). The impact of elevated posttraumatic stress on the efficacy of brief alcohol interventions for heavy drinking college students (STUDY 3035 is for BASICS-ECHUG STUDY). *Addictive Behaviors*, *38*, 1719–1725. <https://doi.org/10.1016/j.addbeh.2012.09.004>.
- Murphy, J. G., Dennhardt, A. A., Skidmore, J. R., Martens, M. P., & McDevitt-Murphy, M. E. (2010). Computerized versus motivational interviewing alcohol interventions: Impact on discrepancy, motivation, and drinking (STUDY 3035 is for BASICS-ECHUG STUDY). *Psychology of Addictive Behaviors*, *24*, 628–639. <https://doi.org/10.1037/a0021347>.
- Murphy, J. G., Dennhardt, A. A., Yurasek, A. M., Skidmore, J. R., Martens, M. P., MacKillop, J., & McDevitt-Murphy, M. E. (2015). Behavioral economic predictors of brief alcohol intervention outcomes (STUDY 3035 is for BASICS-ECHUG STUDY). *Journal of Consulting and Clinical Psychology*, *83*, 1033–1043. <https://doi.org/10.1037/ccp0000032>.
- Murphy, J. G., Duchnick, J. J., Vuchinich, R. E., Davison, J. W., Karg, R. S., Olson, A. M., ... Coffey, T. T. (2001). Relative efficacy of a brief motivational intervention for college student drinkers. *Psychology of Addictive Behaviors*, *15*, 373–379. <https://doi.org/10.1037/0893-164x.15.4.373>.
- O'Leary, T. A., Brown, S. A., Colby, S. M., Cronce, J. M., D'Amico, E. J., Fader, J. S., ... Monti, P. M. (2002). Treating adolescents together or individually? Issues in adolescent substance abuse interventions. *Alcoholism: Clinical and Experimental Research*, *26*, 890–899. <https://doi.org/10.1111/j.1530-0277.2002.tb02619.x>.
- Orchowski, L. M., Mastroleto, N. R., & Borsari, B. (2012). Correlates of alcohol-related regretted sex among college students. *Psychology of Addictive Behaviors*, *26*, 782–790. <https://doi.org/10.1037/a0027840>.
- Paschall, M. J., Antin, T., Ringwalt, C. L., & Saltz, R. F. (2011a). Effects of AlcoholEdu for college on alcohol-related problems among freshmen. A randomized multicampus trial. *Journal of Studies on Alcohol and Drugs*, *72*, 642–650.
- Paschall, M. J., Antin, T., Ringwalt, C. L., & Saltz, R. F. (2011b). Evaluation of an internet-based alcohol misuse prevention course for college freshmen: Findings of a randomized multi-campus trial.

- American Journal of Preventive Medicine*, 41, 300–308. <https://doi.org/10.1016/j.amepre.2011.03.021>.
- Paschall, M. J., Ringwalt, C., Wyatt, T., & DeJong, W. (2014). Effects of an online alcohol education course among college freshmen: An investigation of potential mediators. *Journal of Health Communication*, 19, 392–412. <https://doi.org/10.1080/10810730.2013.811328>.
- Reid, A. E., Carey, K. B., Merrill, J. E., & Carey, M. P. (2015). Social network influences on initiation and maintenance of reduced drinking among college students. *Journal of Consulting and Clinical Psychology*, 83, 36–44. <https://doi.org/10.1037/a0037634>.
- Roberts, L. J., Neal, D. J., Kivlahan, D. R., Baer, J. S., & Marlatt, G. A. (2000). Individual drinking changes following a brief intervention among college students: Clinical significance in an indicated preventive context. *Journal of Consulting and Clinical Psychology*, 68, 500–505. <https://doi.org/10.1037/0022-006x.68.3.500>.
- Schaus, J. F., Sole, M. L., McCoy, T. P., Mullett, N., & O'Brien, M. C. (2009). Alcohol screening and brief intervention in a college student health center: A randomized controlled trial. *Journal of Studies on Alcohol and Drugs*, (Suppl. 16), 131–141.
- Simão, M. O., Kerr-Corrêa, F., Smaira, S. I., Trinca, L. A., Floripes, T. M. F., Dalben, I., . . . Tucci, A. M. (2008). Prevention of “risky” drinking among students at a Brazilian university. *Alcohol and Alcoholism*, 43, 470–476. doi:<https://doi.org/10.1093/alcalc/agn019>.
- Teeters, J. B., Borsari, B., Martens, M. P., & Murphy, J. G. (2015). Brief motivational interventions are associated with reductions in alcohol-impaired driving among college drinkers (STUDY 3090 is for BASICS-ALC101 STUDY). *Journal of Studies on Alcohol and Drugs*, 76, 700–709.
- Terlecki, M. (2011). The long-term effect of a brief motivational alcohol intervention for heavy drinking mandated college students (doctoral dissertation, Louisiana State University, Baton Rouge, Louisiana). Dissertation Abstracts International, 73.
- Terlecki, M. A. (2008). *Alcohol use, negative consequences, and readiness to change in mandated and volunteer college student heavy drinkers before and after a brief alcohol intervention* (Doctoral dissertation, Louisiana State University). Dissertation Abstracts International, M.A., 79.
- Terlecki, M. A., Buckner, J. D., Larimer, M. E., & Copeland, A. L. (2011). The role of social anxiety in a brief alcohol intervention for heavy-drinking college students. *Journal of Cognitive Psychotherapy*, 25, 7–21. <https://doi.org/10.1891/0889-8391.25.1.7>.
- Terlecki, M. A., Buckner, J. D., Larimer, M. E., & Copeland, A. L. (2012). Brief motivational intervention for college drinking: The synergistic impact of social anxiety and perceived drinking norms. *Psychology of Addictive Behaviors*, 26, 917–923. <https://doi.org/10.1037/a0027982>.
- Terlecki, M. A., Buckner, J. D., Larimer, M. E., & Copeland, A. L. (2015). Randomized controlled trial of brief alcohol screening and intervention for college students for heavy-drinking mandated and volunteer undergraduates: 12-month outcomes. *Psychology of Addictive Behaviors*, 29, 2–16. <https://doi.org/10.1037/adb0000056>.
- Terlecki, M. A., Copeland, A. L., Larimer, M. E., & Taylor, H. A. (2008). A brief alcohol intervention for heavy drinking mandated and volunteer college students. *Alcoholism: Clinical & Experimental Research*, 32, 242A–242A.
- Terlecki, M. A., Larimer, M. E., & Copeland, A. L. (2010). Clinical outcomes of a brief motivational intervention for heavy drinking mandated college students: A pilot study. *Journal of Studies on Alcohol and Drugs*, 71, 54–60.
- Terry, D. L. (2012). *Screening and brief intervention for hazardous alcohol use: A pilot study in a college counseling center* (Doctoral dissertation, Syracuse University). Dissertation Abstracts International, Ph.D.(74, 02-E), 137. (University Microfilms No. 3527869).
- Tollison, S. J., Mastroleo, N. R., Mallett, K. A., Witkiewitz, K., Lee, C. M., Ray, A. E., & Larimer, M. E. (2013). The relationship between baseline drinking status, peer motivational interviewing microskills, and drinking outcomes in a brief alcohol intervention for matriculating college students: A replication. *Behavior Therapy*, 44, 137–151.
- Walters, S. T., Vader, A. M., & Harris, T. R. (2007). A controlled trial of web-based feedback for heavy drinking college students. *Prevention Science*, 8, 83–88. <https://doi.org/10.1007/s11121-006-0059-9>.
- Whiteside, U. (2010). *A brief personalized feedback intervention integrating a motivational interviewing therapeutic style and dialectical behavioral therapy skills for depressed or anxious heavy drinking young adults* (Zotero’s generic thesis/dissertation reference - override with correct option). University of Washington, Ann Arbor.
- Wood, M. D., Fairlie, A. M., Fernandez, A. C., Borsari, B., Capone, C., Laforge, R., & Carmona-Barros, R. (2010). Brief motivational and parent interventions for college students: A randomized factorial study. *Journal of Consulting and Clinical Psychology*, 78, 349–361. <https://doi.org/10.1037/a0019166>.