



## Correction to: Intranasal sufentanil given in the emergency department triage zone for severe acute traumatic pain—a randomized double-blind controlled trial: comment

Simon-Pierre Corcostegui<sup>1</sup> · Damien Commeau<sup>1</sup> · Julien Galant<sup>1</sup> · Fabien Ramon<sup>1</sup> · Cédric Boutillier du Retail<sup>1</sup>

Published online: 29 March 2019

© Società Italiana di Medicina Interna (SIMI) 2019

**Correction to: Internal and Emergency Medicine**  
<https://doi.org/10.1007/s11739-019-02069-5>

In the original publication, the doses of sufentanil were incorrectly published.

Instead of a dose of 0.6 mg/kg of sufentanil, the dosage is 0.6 µg/kg.

Instead of the dosage of 0.4 mg/kg, the dosage is 0.4 µg/kg.

---

The original article can be found online at <https://doi.org/10.1007/s11739-019-02069-5>.

---

✉ Simon-Pierre Corcostegui  
spcorcostegui@gmail.com

Damien Commeau  
damien.commeau@gendarmerie.interieur.gouv.fr

Julien Galant  
julien.galant@gendarmerie.interieur.gouv.fr

Fabien Ramon  
fabien.ramon@gendarmerie.interieur.gouv.fr

Cédric Boutillier du Retail  
cedric.boutillier-du-retail@gendarmerie.interieur.gouv.fr

<sup>1</sup> French Military Medical Service, 1ère Antenne Médicale Spécialisée, 34 rue de la Martinière, 78000 Versailles, France