



# Preoperative 5-aminolevulinic acid administration for brain tumor surgery is associated with an increase in postoperative liver enzymes: a retrospective cohort study

Jun-Hyun Kim<sup>1</sup> · Hyun-Kyu Yoon<sup>2</sup> · Hyung-Chul Lee<sup>2</sup> · Hee-Pyoung Park<sup>2</sup> · Chul-Kee Park<sup>3</sup> · Yun-Sik Dho<sup>4</sup> · Jung-Won Hwang<sup>5</sup>

Received: 3 April 2019 / Accepted: 22 August 2019 / Published online: 31 August 2019  
© Springer-Verlag GmbH Austria, part of Springer Nature 2019

## Abstract

**Background** Besides 5-aminolevulinic acid (5-ALA), liver enzyme elevation after brain tumor surgery can be caused by anesthesia and medications. In this retrospective study, we determined whether preoperative 5-ALA administration is associated with postoperative elevation of liver enzymes (PELE) in brain tumor patients and identified predictive factors for PELE in patients treated with 5-ALA.

**Methods** In 179 patients undergoing brain tumor surgery with preoperative normal values of liver enzymes, laboratory data on serum alanine transaminase (ALT), aspartate transaminase (AST), alkaline phosphatase (ALP), and total bilirubin (T.bil) levels were collected preoperatively and through postoperative day (POD) 45.

**Results** Ninety-nine PELEs (ALT, 56; AST, 34; ALP, 5; and TB, 4) were observed in 62 (34.6%) patients. Four (4.2%) patients treated with 5-ALA showed grade 3 elevation of transaminases based on the Common Terminology Criteria for Adverse Effects. Preoperative 5-ALA treatment was predictive of PELE (odds ratio [95% confidence interval], 2.30 [1.14–4.67];  $P=0.021$ ). In patients treated with 5-ALA ( $n=95$ ), 70 PELEs (ALT, 39; AST, 22; ALP, 5; and TB, 4) were observed in 41 (43.2%) patients and significant predictive factors for PELE were preoperative ALT level (1.10 [1.04–1.17];  $P=0.001$ ) and body mass index (BMI, 1.29 [1.08–1.56];  $P=0.006$ ). In patients treated with 5-ALA, 13 and 36 patients, of 39 patients whose maximum postoperative ALT levels  $>40$  U/L, showed the normal value of serum ALT on PODs 14 and 45, respectively. Only three patients showed ALT elevation  $>40$  U/L on PODs 15–45, with a downward trend.

**Conclusions** The use of 5-ALA for brain tumor surgery in patients with preoperative normal values of liver enzymes was associated with increased transient PELE, but a low incidence of severely elevated liver transaminases levels. When 5-ALA is administered to patients with the upper normal value of preoperative serum ALT and overweight, attention is paid to PELE.

**Keywords** 5-aminolevulinic acid · Brain tumor surgery · Liver enzymes

---

**Presentation at a conference** None

---

This article is part of the Topical Collection on *Brain Tumors*

---

✉ Jung-Won Hwang  
jungwon@snuh.org

<sup>1</sup> Department of Anesthesiology and Pain Medicine, Ilsan Paik Hospital, Inje University College of Medicine, Goyang, South Korea

<sup>2</sup> Department of Anesthesiology and Pain Medicine, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, South Korea

<sup>3</sup> Department of Neurosurgery, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, South Korea

<sup>4</sup> Department of Neurosurgery, Chungbuk National University Hospital, Chungbuk National University College of Medicine, Cheongju, South Korea

<sup>5</sup> Department of Anesthesiology and Pain Medicine, Seoul National University Bundang Hospital, Seoul National University College of Medicine, 173-2 Gumi-ro, Bundang-gu, Seongnam 13620, South Korea

## Introduction

High-grade glioma, a malignant neoplasm of the central nervous system, responds poorly to radiotherapy and chemotherapy [22, 25]. Surgical removal of the tumor is therefore important in disease treatment. However, complete tumor removal is difficult in clinical practice because the tumor's low degree of differentiation obscures the surgical margins. 5-Aminolevulinic acid (5-ALA) is a precursor of protoporphyrin IX (PpIX). Systemic administration of 5-ALA results in selective PpIX accumulation in high-grade glioma cells. PpIX emits red light under surgical microscope fluorescence, enabling the differentiation of a malignant tumor from normal, functional brain tissue [19]. For this reason, the use of 5-ALA is known to improve the surgical resection of high-grade gliomas [21].

One side effect of 5-ALA is hepatocellular toxicity, which is indicated by increased postoperative activities of liver enzymes, such as alanine transaminase (ALT), aspartate transaminase (AST), alkaline phosphatase (ALP), and total bilirubin (T.bil) [4, 12, 21, 23]. However, all previous studies investigating 5-ALA-induced hepatobiliary dysfunction have at least one weakness. Stummer et al. [21] did not evaluate the incidence and extent of hepatobiliary dysfunction in their comparative study. Otherwise, all other clinical investigations have been conducted only in patients who were administered with 5-ALA [4, 12, 23].

Postoperative elevation of liver enzymes (PELE) after brain tumor surgery can be caused by 5-ALA as well as other multiple factors, such as anesthetics, antibiotics, antiepileptic drugs, anemia, and intraoperative hypotension [5, 9, 13]. To confirm the detrimental effects of 5-ALA on the hepatobiliary system, multivariate analysis including data on postoperative liver enzymes in patients with brain tumors who have not received 5-ALA treatment is necessary. Also, various confounding factors mentioned the above need to be adjusted in multivariate analysis. However, no study has determined in multivariate analysis whether 5-ALA is a significant predictor of PELE in patients undergoing craniotomy for the removal of brain tumors. Moreover, no clinical investigation to identify predictive factors of PELE in patients with preoperative 5-ALA administration has yet been done.

In this retrospective study, we determined whether preoperative 5-ALA administration is associated with the development of PELE in patients undergoing craniotomy for brain tumor removal. In addition, we determined which factors were predictive of PELE in patients treated with 5-ALA.

## Materials and methods

### Patients

This retrospective study was conducted after approval from the Institutional Review Board of Seoul National University

Hospital (IRB no. 1811-078-985). We reviewed the electronic medical records of patients who underwent brain tumor surgery between January and December 2017 and who had preoperative liver enzyme levels within normal limits. Adult patients aged 18–80 years were included. Exclusion criteria were anesthesia time < 2 h (i.e., brain tumor biopsy), history of a postoperative medical condition that could cause liver enzyme elevation (i.e., gall bladder stones, cardiopulmonary resuscitation, drug toxicity), failure to collect liver enzyme data through postoperative day (POD) 45, and porphyria. The requirement for written informed content was waived because of the retrospective design of the study.

### Data collection

Enrolled patients were divided into two groups (PELE and no PELE) according to the presence or absence of PELE. Data on demographic characteristics, American Society of Anesthesiologists physical status class, comorbidities, preoperative hemoglobin (Hb) level, preoperative systolic blood pressure (SBP), and mean blood pressure (MBP) in the ward were collected. Intra- and postoperative parameters, including anesthesia duration, surgery duration, doses of propofol and remifentanyl used, fluid balance, average SBP and MBP, maximal levels of lactate and glucose, intraoperative transfusion, incidence of intraoperative vasopressor use and continuous phenylephrine infusion, immediate postoperative Hb level, pathologic diagnosis for brain tumor, and postoperative transfusion were also recorded. Serum ALT, AST, ALP, and T.bil levels were measured preoperatively (T0), immediate postoperatively (T1), and on PODs 1 (T2), 2 (T3), 3–6 (T4), 7–14 (T5), and 15–45 (T6). When liver enzymes were measured repeatedly during the study period (T4–6), the highest value for each enzyme was selected and used for data analysis. In our hospital, the upper limit of normal (ULN) for ALT, AST, ALP, and T.bil is 40 U/L, 40 U/L, 118 U/L, and 1.3 mg/dL, respectively. PELE was defined as postoperative elevation above the reference value of one or multiple liver enzymes. The maximal postoperative liver enzyme values were recorded and the incidence of patients with PELE was determined. Maximal postoperative liver enzyme values were classified as grade 1 (ALT and AST, >ULN– 3.0 × ULN; ALP, >ULN– 2.5\*ULN; T.bil, >ULN– 1.5 × ULN), grade 2 (ALT and AST, > 3.0–5.0 × ULN; ALP, > 2.5–5.0 × ULN; T.bil, > 1.5–3.0 × ULN), grade 3 (ALT, AST, and ALP, > 5.0–20.0 × ULN; T.bil, > 3.0–10.0 × ULN), and grade 4 (ALT, AST, and ALP, > 20.0 × ULN; T.bil, > 10.0 × ULN) elevation of liver enzymes based on the Common Terminology Criteria for Adverse Events (CTCAE, version 5.0) of investigations [18]. In patients who received chemotherapy and/or radiotherapy at about 1 month postoperatively, only the liver enzymes obtained immediately before the therapy were entered into the data analysis.

**Table 1** Comparison of postoperative laboratory data between patients with postoperative elevation of liver enzymes (PELE) and those without

	PELE (n = 62)	No PELE (n = 117)	P value
Postoperative ALT			
Maximal postoperative ALT (U/L)	58.0 (48.0–83.8)	22.0 (17.0–30.5)	< 0.001
Maximal postoperative ALT > 40 U/L (n, %)	56 (90.3%)	0 (0.0%)	< 0.001
Maximal postoperative ALT (U/L)	63.5 (50.0–88.3)	No value	NA
CTCAE of investigations			
Grade 1	48 (77.4%)	0 (0.0%)	< 0.001
Grade 2	4 (6.5%)	0 (0.0%)	0.013
Grade 3	4 (6.5%)	0 (0.0%)	0.013
Postoperative AST			
Maximal postoperative AST (U/L)	42.0 (28.5–69.8)	20.0 (16.0–24.0)	< 0.001
Maximal postoperative AST > 40 U/L (n, %)	34 (54.8%)	0 (0.0%)	< 0.001
Maximal postoperative AST (U/L)	61.5 (44.0–96.3)	No value	NA
CTCAE of investigations			
Grade 1	29 (46.8%)	0 (0.0%)	< 0.001
Grade 2	4 (6.5%)	0 (0.0%)	0.013
Grade 3	1 (1.6%)	0 (0.0%)	0.346
Postoperative ALP			
Maximal postoperative ALP (U/L)	68.0 (57.0–82.3)	58.0 (47.0–70.5)	< 0.001
Maximal postoperative ALP > 118 U/L (n, %)	5 (8.1%)	0 (0.0%)	0.004
Maximal postoperative ALP (U/L)	157.0 (138.5–194.5)	No value	NA
CTCAE of investigations			
Grade 1	5 (8.1%)	0 (0.0%)	0.004
Grade 2	0 (0.0%)	0 (0.0%)	NA
Grade 3	0 (0.0%)	0 (0.0%)	NA
Postoperative T.bil			
Maximal postoperative T.bil (mg/dL)	0.7 (0.6–1.0)	0.7 (0.5–0.9)	0.075
Maximal postoperative T.bil > 1.3 mg/dL (n, %)	4 (6.5%)	0 (0.0%)	0.013
Maximal postoperative T.bil (mg/dL)	1.5 (1.4–1.9)	No value	NA
CTCAE of investigations			
Grade 1	3 (4.8%)	0 (0.0%)	0.040
Grade 2	1 (1.6%)	0 (0.0%)	0.346
Grade 3	0 (0.0%)	0 (0.0%)	NA
Postoperative ASL/ALT ratio			
Immediate postoperative	0.89 (0.67–1.29)	1.00 (0.80–1.36)	0.069
POD 1	0.85 (0.71–1.37)	1.13 (0.84–1.50)	0.010
POD 2	0.66 (0.48–1.06)	0.91 (0.69–1.13)	< 0.001
PODs 3–6	0.61 (0.44–0.77)	0.81 (0.65–1.00)	< 0.001
PODs 7–14	0.63 (0.45–0.88)	0.81 (0.65–1.00)	< 0.001
PODs 14–45	0.81 (0.62–1.17)	1.00 (0.70–1.30)	0.052

Data are expressed as number (%) or median (IQR). *ALT*, alanine transaminase; *AST*, aspartate transaminase; *ALP*, alkaline phosphatase; *T.bil*, total bilirubin; *NA*, non-applicable; *POD*, postoperative day

## Patient management

All patients were treated with dexamethasone and levetiracetam 1 day before brain surgery and with cefazolin on the day of surgery. In patients treated with 5-ALA, 5-ALA (20 mg/kg; Gliolan; Medac, Wedel, Germany) was administered orally

3–8 h before anesthetic induction. Patients were instructed to avoid exposure to ultraviolet light for 24 h after 5-ALA ingestion. In all patients, anesthesia was induced and maintained with total intravenous anesthesia by target-controlled infusion of propofol and remifentanyl. Intraoperative target MBP was within 20% of the preoperative values, and a bolus or

**Table 2** Comparisons of demographic, preoperative laboratory, and pre- and intraoperative hemodynamic data in all patients with versus without postoperative elevated liver enzymes (PELE) and in 5-ALA-treated patients with versus without PELE

	PELE in all patients		P value	PELE in 5-ALA-treated patients		P value
	Yes (n = 62)	No (n = 117)		Yes (n = 41)	No (n = 54)	
Age (year)	55.5 (43.5–62.0)	57.0 (43.5–67.0)	0.166	55.0 (43.0–60.5)	58.0 (40.8–67.3)	0.121
Male gender (n, %)	38 (61.3%)	63 (53.8%)	0.425	28 (68.3%)	31 (57.4%)	0.384
Body mass index (m/kg <sup>2</sup> )	24.5 (23.3–26.7)	23.4 (21.2–25.0)	0.001	24.5 (23.3–27.2)	23.2 (21.3–25.0)	0.001
Body mass index > 25 m/kg <sup>2</sup> (n, %)	29 (50.0%)	29 (24.8%)	0.005	20 (48.8%)	13 (24.1%)	0.022
ASA class (n, %)			0.637			0.060
I	17 (27.4%)	32 (27.4%)		12 (29.3%)	12 (22.2%)	
II	37 (59.7%)	62 (53.0%)		27 (65.9%)	30 (55.6%)	
III	8 (12.9%)	22 (18.8%)		2 (4.9%)	12 (22.2%)	
IV	0 (0.0%)	1 (0.9%)		0 (0.0%)	0 (0.0%)	
Comorbidity (n, %)						
Previous craniotomy	11 (17.7%)	13 (11.1%)	0.313	8 (19.5%)	9 (16.7%)	0.930
Cerebrovascular	4 (6.5%)	4 (3.4%)	0.451	1 (2.4%)	2 (3.7%)	1.000
Cardiac	5 (8.1%)	6 (5.1%)	0.517	1 (2.4%)	3 (5.6%)	0.631
Pulmonary	10 (16.1%)	22 (18.8%)	0.811	3 (7.3%)	8 (14.8%)	0.340
Diabetes	7 (11.3%)	16 (13.7%)	0.827	1 (2.4%)	7 (13.0%)	0.132
Hypertension	21 (33.9%)	26 (22.2%)	0.132	12 (29.3%)	10 (18.5%)	0.325
Hepatic	0 (0.0%)	1 (0.9%)	1.000	0 (0.0%)	0 (0.0%)	NA
Renal	1 (1.6%)	1 (0.9%)	1.000	1 (2.4%)	0 (0.0%)	0.432
Extracranial malignancy	9 (14.5%)	25 (21.4%)	0.362	2 (4.9%)	5 (9.3%)	0.695
Others	5 (8.1%)	9 (7.7%)	1.000	3 (7.3%)	3 (5.6%)	1.000
Pathologic diagnosis (n, %)						
Astrocytoma	6 (9.7%)	12 (10.3%)	1.000	5 (12.2%)	11 (20.4%)	0.408
Ependymoma	3 (4.8%)	2 (1.7%)	0.343	1 (2.4%)	0 (0.0%)	0.432
Glioma	4 (6.5%)	5 (4.3%)	0.500	4 (9.8%)	4 (7.4%)	0.723
Glioblastoma	23 (37.1%)	33 (28.2%)	0.293	22 (53.7%)	33 (61.1%)	0.604
Hemangiocyoma	1 (1.6%)	2 (1.7%)	1.000	0 (0.0%)	0 (0.0%)	NA
Hemangioblastoma	0 (0.0%)	2 (1.7%)	0.545	0 (0.0%)	0 (0.0%)	NA
Meningioma	8 (12.9%)	28 (23.9%)	0.120	1 (2.4%)	0 (0.0%)	0.432
Metastatic tumor	8 (12.9%)	20 (17.1%)	0.604	2 (4.9%)	0 (0.0%)	0.184
Neurocytoma	2 (3.2%)	0 (0.0%)	0.119	1 (2.4%)	0 (0.0%)	0.432
Oligodendrocytoma	5 (8.1%)	5 (4.3%)	0.318	5 (12.2%)	5 (9.3%)	0.741
Schwannoma	2 (3.2%)	5 (4.3%)	1.000	0 (0.0%)	0 (0.0%)	NA
Inflammation	0 (0.0%)	3 (2.6%)	0.552	0 (0.0%)	1 (1.9%)	1.000
Preoperative Hb (g/dL)	14.1 (13.2–15.0)	13.6 (12.8–14.6)	0.017	14.6 (13.6–15.2)	13.9 (12.9–15.0)	0.128
Intraoperative transfusion (n, %)	0 (0.0%)	1 (0.9%)	1.000	0 (0.0%)	1 (1.9%)	1.000
Immediate postoperative Hb (g/dL)	11.7 ± 1.6	11.0 ± 1.6	0.014	11.8 ± 1.7	11.1 ± 1.6	0.065
Postoperative transfusion (n, %)	6 (9.7%)	13 (11.1%)	0.967	3 (7.3%)	5 (9.3%)	1.000
Preoperative ALT (U/L)	24.5 (17.8–33.3)	17.0 (13.0–23.5)	< 0.001	28.0 (19.0–35.0)	17.0 (13.8–24.3)	< 0.001
Preoperative AST (U/L)	20.0 (16.8–24.0)	18.0 (15.0–20.5)	0.002	20.0 (16.5–24.0)	17.0 (15.0–21.0)	0.046
Preoperative AST/ALT ratio	0.87 (0.70–1.11)	1.00 (0.75–1.29)	0.011	0.78 (0.62–1.05)	1.00 (0.77–1.19)	0.005
Preoperative ALP (U/L)	65.2 ± 16.6	60.7 ± 14.2	0.061	65.1 ± 16.4	61.2 ± 13.8	0.212
Preoperative T.bil (mg/dL)	0.6 (0.5–0.7)	0.6 (0.4–0.8)	0.984	0.6 (0.5–0.7)	0.6 (0.4–0.8)	0.814
Average SBP at ward (mmHg)	122.5 ± 11.3	118.0 ± 11.3	0.014	122.1 ± 11.8	116.2 ± 10.3	0.011
Average MBP at ward (mmHg)	90.5 (85.3–98.4)	86.0 (81.7–92.5)	0.001	89.0 (84.7–98.8)	85.8 (80.8–90.5)	0.006
Anesthesia duration (min)	312.5 (248.8–368.8)	300.0 (252.5–345.0)	0.439	315.0 (272.5–365.0)	302.5 (258.8–370.0)	0.644
Operation duration (min)	245.0 (183.8–305.0)	235.0 (190.0–277.5)	0.640	245.0 (202.5–305.0)	245.0 (197.5–315.0)	0.937
5-ALA (n, %)	41 (66.1%)	54 (46.2%)	0.017	41 (100.0%)	54 (100.0%)	NA
Propofol (mg/kg/h)	8.1 (7.5–9.2)	8.7 (7.8–9.9)	0.022	8.1 (7.6–9.3)	8.9 (8.1–10.2)	0.069
Remifentanyl (µg/kg/h)	8.1 (7.5–9.1)	8.6 (7.7–10.3)	0.027	8.4 (7.5–9.3)	8.7 (7.7–10.1)	0.400
Fluid balance (ml)	376.4 ± 766.5	213.3 ± 644.2	0.151	288.5 ± 744.1	408.3 ± 545.7	0.411
Average SBP during operation (mmHg)	109.0 (106.0–116.3)	109.0 (105.0–116.5)	0.654	109.0 (106.0–116.0)	109.0 (105.0–115.3)	0.857
Average MBP during operation (mmHg)	75.5 (71.0–79.0)	76.0 (71.0–80.0)	0.491	75.0 (70.5–79.0)	72.0 (70.0–78.3)	0.389
Maximal lactate during operation (mmol/L)	2.0 (1.5–2.7)	2.0 (1.2–2.5)	0.211	2.0 (1.7–3.0)	2.0 (1.2–2.8)	0.640
Maximal glucose during operation (mg/dL)	139.0 (128.8–155.3)	138.0 (125.5–153.0)	0.738	139.0 (129.0–157.0)	141.5 (130.8–153.0)	0.730
Vasopressor use (n, %)	49 (79.0%)	95 (81.2%)	0.881	33 (80.5%)	45 (83.3%)	0.930
Continuous phenylephrine infusion (n, %)	36 (58.1%)	62 (53.0%)	0.623	25 (61.0%)	33 (61.1%)	1.000

Data are expressed as number (%), mean ± SD, or median (IQR). 5-ALA, 5-aminolevulinic acid; ASA, American Society of Anesthesiologists; NA, not applicable; Hb, hemoglobin; ALT, alanine transaminase; AST, aspartate transaminase; ALP, alkaline phosphatase; T.bil, total bilirubin; SBP, systolic blood pressure; MBP, mean blood pressure

continuous infusion of phenylephrine was used as a vasopressor. After the surgery, all patients were transferred to the intensive care unit without emergence after brain computed tomography examination. In patients with glioblastoma, the Stupp chemo-radiotherapy regimen was initiated about 1 month after the surgery.

### Statistical analysis and sample size

Continuous variables were compared using Student's *t* test or the Mann–Whitney *U* test according to the results of the Kolmogorov–Smirnov test. For categorical variables, the chi-squared test or Fisher's exact test was used, according

to the frequency. *P* values < 0.05 were considered to be significant. Repeated-measures analysis of variance was performed to determine a significant difference in changes in liver enzyme levels over time within group. Multivariate regression analysis was used to identify predictive factors of PELE in patients treated with 5-ALA, and variables with *P* values ≤ 0.10 in the univariate analysis were entered into a multivariate logistic regression analysis with the forward stepwise conditional method. Receiver operating characteristic (ROC) curve analysis was performed to evaluate the diagnostic value of the identified risk factors. The optimal cut-off point was determined by maximizing the sum of sensitivity and specificity. All statistical analyses were con-

**Table 3** Univariate and multivariate analyses for predictive factors associated with postoperative elevated liver enzymes

	Univariate analysis			Multivariate analysis		
	Odds ratio	95% CI	<i>P</i> value	Odds ratio	95% CI	<i>P</i> value
Postoperative elevated liver enzymes in all patients*						
Body mass index (kg/m <sup>2</sup> )	1.21	1.08–1.35	0.001	1.13	1.00–1.28	0.045
Meningioma	0.47	0.20–1.11	0.084			
Hypertension	1.79	0.91–3.55	0.094			
Propofol (mg/kg/h)	0.81	0.66–0.99	0.038			
Remifentanyl (μg/kg/h)	0.86	0.74–1.01	0.058			
Average MBP at ward (mmHg)	1.07	1.03–1.11	0.001	1.05	1.00–1.10	0.033
5-ALA	2.28	1.20–4.32	0.012	2.30	1.14–4.67	0.021
Preoperative Hb (g/dL)	1.36	1.06–1.74	0.014			
Preoperative ALT (U/L)	1.09	1.05–1.13	< 0.001	1.07	1.02–1.11	0.002
Preoperative AST (U/L)	1.09	1.03–1.15	0.003			
Preoperative AST/ALT ratio	0.32	0.13–0.77	0.011			
Preoperative ALP (U/L)	1.02	1.00–1.04	0.064			
Postoperative elevated liver enzymes in 5-ALA-treated patients†						
Body mass index (kg/m <sup>2</sup> )	1.23	1.10–1.53	0.002	1.29	1.08–1.56	0.006
ASA class ( <i>n</i> , %)			0.095			
I	Reference					
II	0.90	0.35–2.34	0.829			
III	0.17	0.03–0.91	0.039			
Preoperative Hb (g/dL)	1.30	0.95–1.78	0.102			
Average MBP at ward (mmHg)	1.09	1.03–1.15	0.003			
Preoperative ALT (U/L)	1.11	1.05–1.17	< 0.001	1.10	1.04–1.17	0.001
Preoperative AST (U/L)	1.07	1.00–1.16	0.062			
Preoperative AST/ALT ratio	0.22	0.06–0.75	0.016			

ASA, American Society of Anesthesiologists; *Hb*, hemoglobin; *MBP*, mean blood pressure; *5-ALA*, 5-aminolevulinic acid; *ALT*, alanine transaminase; *AST*, aspartate transaminase

\*In multivariate analysis with the forward stepwise conditional method, meningioma, hypertension, propofol and remifentanyl infusion rates, preoperative hemoglobin concentration, preoperative AST and ALP levels, and preoperative AST/ALT ratio were adjusted. Nagelkerke *R*<sup>2</sup> statistic in step 4 is 0.254. Hosmer and Lemeshow goodness of fit test in step 4 is not significant at 5% (*P* = 0.607). †In multivariate analysis with the forward stepwise conditional method, ASA class, preoperative and immediate postoperative hemoglobin concentrations, average MBP at ward, preoperative AST level, and preoperative AST/ALT ratio were adjusted. Nagelkerke *R*<sup>2</sup> statistic in step 3 is 0.375. Hosmer and Lemeshow goodness of fit test in step 3 is not significant at 5% (*P* = 0.453)

ducted using SPSS statistical software for Windows, version 25.0 (IBM, Armonk, NY, USA) and MedCalc software, version 18.6 (MedCalc Software, Ostend, Belgium).

In a previous study, 42% of patients with brain tumors who received 5-ALA preoperatively showed PELE [12]. Assuming the incidence of PELE in this study conducted in mixed patients (patients with 5-ALA treatment and the rest without 5-ALA treatment) would be reduced by 8% compared with that reported in the previous study conducted in only patients with 5-ALA treatment, a minimum of 176 subjects was needed to acquire statistical significance with a confidence level of 95% and a total width of the confidence interval of 0.14.

## Results


We screened 215 patients. Thirty-six patients (20 with anesthetic times < 2 h, 13 with incomplete laboratory data, and 3 with abrupt PELE due to cardiac arrest or drug toxicity) were excluded from the study. Finally, a total of 179 patients were included in the analyses.

Ninety-nine PELEs were observed in 62 (34.6%) patients (Table 1). Postoperative elevation of ALT, AST, ALP, and TB was shown in 56, 34, 5, and 4 patients respectively. In patients

with 5-ALA treatment, three (3.2%), zero (0.0%), and one (1.1%) patient met the ALT, AST, and both criteria for grade 3 on CTCAE, respectively. However, in all four of these patients, the ALT and AST levels had returned to preoperative baseline levels (<ULN) at T6.

Patients with PELE had a higher body mass index (BMI), preoperative and immediate postoperative Hb concentrations, SBP and MBP in the ward, and preoperative ALT and AST levels and had a lower preoperative AST/ALT ratio and remifentanyl and propofol doses than those without PELE (Table 2). 5-ALA was more frequently used in patients with PELE. Multivariate analyses showed that significant predictive factors for PELE were 5-ALA treatment (odds ratio [95% confidence interval], 2.30 [1.14–4.67];  $P=0.021$ ), preoperative ALT level (1.07 [1.02–1.11];  $P=0.002$ ), BMI (1.13 [1.00–1.28];  $P=0.045$ ), and MBP in the ward (1.05 [1.00–1.10];  $P=0.033$ ; Table 3). Among 95 patients treated with 5-ALA, 41 (43.2%) patients showed 70 PELEs. Postoperative elevation of ALT, AST, ALP, and TB was shown in 39, 22, 5, and 4 patients, respectively. Patients with PELE had a higher BMI, SBP, and MBP in the ward and preoperative ALT level than those without PELE (Table 2). Multivariate analysis showed that significant predictive factors for PELE were preoperative ALT level (odds ratio [95% confidence interval], 1.10 [1.04–1.17];  $P=0.001$ ) and BMI (1.29 [1.08–1.56];

**Fig. 1** Serial changes in alanine transaminase (ALT, **a**), aspartate transaminase (AST, **b**), alkaline phosphatase (ALP, **c**), and total bilirubin (T.bil, **d**) during the study period in patients with (●, closed circle, solid line) and without (○, open circle, broken line) 5-aminolevulinic acid administration. \* $P<0.01$  vs. T0 within group, † $P<0.05$  vs. T0 within group. T0, preoperative; T1, immediate postoperative; T2, postoperative day (POD) 1; T3, POD 2; T4, PODs 3–6; T5, PODs 7–14; T6, POD 15–45



$P = 0.006$ ; Table 3). In the ROC analysis, the preoperative ALT value and BMI showed the area under the curve of 0.74 and 0.70, respectively. The optimal cut-off value for preoperative ALT was 27.5 U/L, and PELE developed more frequently in cases with preoperative ALT levels  $\geq 28$  U/L (9.26 [3.25–26.40];  $P < 0.001$ ). The optimal cut-off value for BMI was 23.8 m/kg<sup>2</sup>, and PELE developed more frequently in cases with BMI  $\geq 23.8$  m/kg<sup>2</sup> (3.97 [1.67–9.40];  $P = 0.002$ ). When the cut-off value for BMI was 25 m/kg<sup>2</sup>, PELE developed more frequently in cases with BMI  $> 25$  m/kg<sup>2</sup> (3.00 [1.25–7.20];  $P = 0.014$ ). BMI was positively correlated with preoperative serum ALT levels (correlation coefficient, 0.289;  $P = 0.005$ ).

Serial changes in liver enzymes during the study period are shown in Fig. 1. In patients treated with 5-ALA, the ALT value at T2, T3, T4, T5, and T6 was higher than at preoperative baseline ( $P < 0.01$ , respectively). In patients treated with 5-ALA, 39 patients had maximum postoperative ALT levels  $> 40$  U/L, and their ALT value returned to the normal value ( $< 40$  U/L) in one (2.6%), four (10.3%), eight (20.5%), and 23 (59.0%) patients at T3, T4, T5, and T6, respectively. Three patients had ALT levels  $> 40$  U/L at T6, with a downward trend.

## Discussion

In this clinical report, 62 (34.6%) patients showed 99 episodes of PELE. Postoperative ALT elevation was observed the most frequently and followed by AST, ALP, and T.bil elevation. Four (4.2%) patients treated with 5-ALA showed grade 3 elevation of transaminases based upon CTCAE of investigations. 5-ALA treatment was predictive of PELE. In all patients undergoing brain tumor surgery as well as those treated with 5-ALA, preoperative ALT level and BMI were independent risk factors for PELE.

5-ALA is used to facilitate high-grade glioma resection, but it can increase postoperative liver enzymes due to its hepatotoxic effect. In this study, the use of 5-ALA was associated with PELE, with an odds ratio of 2.3. Similarly, previous studies have documented transient and self-limiting liver enzyme abnormalities after the use of 5-ALA [4, 7, 11, 12, 23]. Exogenous loading of 5-ALA bypasses the rate-limiting step of heme synthesis, leading to PpIX accumulation in lesions in the skin and other tissues. Because of its high molecular weight and hydrophilic characteristics, PpIX cannot be excreted by the kidney; its excretion is solely dependent on the hepatobiliary system. Large amounts of PpIX can be hepatotoxic, resulting in PELE [17].

The CTCAE is used to categorize liver enzyme levels into four grades based on the extent of adverse effects [18]. In this study, the overall incidence of PELE was

34.6%. Specifically, the incidence of PELE was 43.2% in patients treated with 5-ALA and 25.0% in those without 5-ALA administration. Similarly, in a previous report, the incidence of 5-ALA-induced temporary PELE was 42% [12]. Another previous study showed that preoperative 5-ALA administration resulted in significant increases in the incidence of postoperative elevation of ALT (36.5%), GGT (37.7%), AST (19.2%), and T.bil (5.4%) levels [23]. More importantly, four (4.2%) patients treated with 5-ALA showed grade 3 elevation of liver enzymes (ALT,  $n = 3$ ; ALT and AST,  $n = 1$ ) during the early postoperative period. However, their ALT and AST levels returned to preoperative baseline levels ( $< \text{ULN}$ ) on PODs 15–45. Similarly, Stummer et al. [20] observed grade 3 elevation of liver enzymes in 3 (5.8%) of 52 patients on POD 7, which resolved to grade 1 or 2 on POD 14.

Notably, our study showed PELE in up to 25% of patients without 5-ALA administration. This finding suggests that PELE is associated with factors other than 5-ALA use. Previous clinical studies have shown that PELE can result from various factors, such as anesthesia, surgery, and use of medications such as antibiotics and antiepileptics [5, 9, 13, 14]. In this study, all patients were anesthetized with propofol. A previous report showed that propofol resulted in concentration-dependent inhibition of cytochrome P450 enzyme in the liver and may therefore alter the metabolism of drugs that depend on this enzyme [3]. Moreover, because it reduces blood flow to the liver, other drugs metabolized in the liver may be eliminated slowly, thereby causing more hepatotoxicity [10]. In addition to propofol, dexamethasone, antiepileptic drugs, and cefazolin are known to be associated with liver enzyme elevation [1, 2, 26]. In this study, all three drugs were administered preoperatively, which may in part explain the occurrence of PELE.

The time profile of the ALT level after brain tumor surgery is likely to be different between patients treated with 5-ALA and those without. In patients with 5-ALA administration, ALT levels were significantly higher than preoperative baseline from POD 1 to POD 45 with a peak on PODs 3–6. On PODs 15–45, these levels returned to the normal value ( $< 40$  U/L) in most patients. Specifically, of 39 patients treated with 5-ALA whose maximum postoperative ALT levels  $> 40$  U/L, 13 and 36 patients showed the normal value of serum ALT on PODs 14 and 45, respectively. Only three patients showed ALT elevation  $> 40$  U/L on PODs 15–45, with a downward trend. Consistent with our results, previous studies investigating 5-ALA-induced hepatobiliary dysfunction showed similar changes in liver enzymes during the postoperative period [12, 20, 21, 23, 24].

In this study, the preoperative ALT level was a significant predictive factor for PELE after brain tumor surgery, regardless of 5-ALA treatment. In the subgroup of patients treated with 5-ALA, ALT levels  $\geq 28$  U/L increased the risk of

PELE approximately 9.3-fold. In addition, preoperative AST level was higher and preoperative AST/ALT ratio was lower in patients with PELE, although these variables were not significant predictive factors in multivariate analysis. Although these findings are not surprising, they suggest the need to pay particular attention to postoperative liver enzyme levels in patients with high preoperative transaminase levels, who are vulnerable to the development of PELE after brain tumor surgery.

This study showed that BMI was associated with PELE in patients undergoing brain tumor surgery, irrespective of 5-ALA treatment. In the subgroup of patients treated with 5-ALA, patients with BMI  $\geq 23.8$  and  $> 25$  m/kg<sup>2</sup> had a 4.0- and 3.0-fold higher risk of PELE development, respectively. A possible explanation for such findings was that BMI was strongly associated with serum level of liver enzymes, especially ALT. High BMI is known to be a significant independent risk factor for elevated serum ALT level [8]. In addition, a previous study showed a significant correlation between BMI and the extent of hepatic steatosis [15]. Therefore, patients with high BMI have a high chance of preoperative upper normal value of serum ALT due to hepatic steatosis, which may be in part responsible for PELE in such patients.

This study showed that the MBP measured in the ward was associated with PELE. Patients with preoperative MBP  $> 87.5$  mmHg had a 2.8-fold higher risk of PELE development. A possible explanation for this finding is that patients with preoperative MBP  $> 87.5$  mmHg were more susceptible to relative intraoperative hypotension, which may have caused hepatocellular injury by ischemia. Intraoperative MBP decreased more significantly from preoperative MBP in patients with preoperative MBP  $> 87.5$  mmHg (by 19 mmHg) than in those with preoperative MBP  $< 87.5$  mmHg (by 7 mmHg). Although intraoperative MBP did not differ between patients with and without PELE, such relative changes in MBP may cause hepatocellular injury. Hepatic blood flow is significantly reduced during hypotension, especially during general anesthesia, due to the unique blood supply and regulatory mechanism [13]. In addition, ample evidence shows that hypertensive patients tend to have right-shifted autoregulation curves (higher blood pressure) and are therefore less tolerant of low blood pressure [16]. Taken together, these findings suggest that maintaining appropriate intraoperative MBP will help to decrease the occurrence of PELE in patients undergoing brain tumor surgery, especially those with high preoperative MBPs.

Our result showed that 5-ALA was more frequently used in patients with PELE than those without (66.1 vs. 46.2%, a mean difference of 19.9%). In the post hoc sample size calculation based upon our result, 74 patients for the PELE group

and 140 for the no PELE group were needed to achieve statistical significance with  $\alpha = 0.05$  and  $\beta = 0.2$ . Therefore, the post hoc power analysis showed that the number of patients enrolled in each group was inadequate in this study.

This study has some limitations. First, because of its retrospective design, unavoidable biases could have affected the results. Second, we measured only four liver enzymes (ALT, AST, ALP, and T.bil). GGT is known to be a surrogate marker of cholestatic liver injury, and previous studies have documented temporary postoperative elevation of GGT in patients with 5-ALA administration [12, 23]. Although the GGT level is indicative of liver function, it is not routinely measured in patients with brain tumors during postoperative periods in our clinical practice. Thus, we did not examine the effect of 5-ALA on postoperative GGT levels in this study. In addition, we did not measure plasma PpIX level, which results in damage to cholangiocytes and hepatocytes. Therefore, we had limitations in revealing the pathophysiology of 5-ALA-induced PELE associated with plasma PpIX level. Third, patients with preoperative elevation of liver enzymes were excluded because this condition could bias the interpretation of 5-ALA-induced PELE; thus, we did not examine the toxic hepatobiliary effect of 5-ALA in patients with preoperative liver enzyme elevation. Fourth, because a fixed dose (20 mg/kg) of 5-ALA was used in this study, we did not evaluate the dose-dependent effects of 5-ALA on PELE. Fifth, caution is necessary when interpreting predictive factors associated with PELE because the predictive model has relatively weak explanatory power, which indicates that some clinically significant risk factors may have been missed. Also, some preoperative and intraoperative variables, which are able to act as confounders, showed a significant difference between PELE and no PELE group, although they were adjusted in multivariate analysis. A propensity score matching of these variables between the two groups may strengthen our main findings. More importantly, a sample size was relatively small in this study. A further large-scaled investigation is needed to identify significant predictive factors of PELE especially when 5-ALA is used in brain tumor surgery. Finally, whether the extent of PELE associated with the use of 5-ALA is clinically relevant remains questionable. Although the use of 5-ALA was associated with increased PELE development, the incidence of grade 3 elevation of liver enzymes was very low and the maximum values of ALT and AST were within a range of  $5 \times$  ULN to  $8 \times$  ULN in this study. A previous study showed that liver damage by ischemia or other forms of liver damage were characterized by much higher concentrations of liver enzymes (e.g., aminotransferase,  $> 10 \times$  ULN to  $> 50 \times$  ULN in ischemia;  $> 10 \times$  ULN in toxic injury;  $5\text{--}10 \times$  ULN to  $> 10 \times$  ULN in acute viral hepatitis) [6].

## Conclusions

The use of 5-ALA for brain tumor surgery was associated with an increased incidence of PELE, especially ALT elevation, but a low incidence of severely elevated liver transaminases levels in patients with preoperative normal values of liver enzymes. Although 5-ALA caused a substantial increase in serum ALT level during the early postoperative period, this elevation was mostly transient and self-limited. Overall, these findings suggested that the use of 5-ALA is relatively safe for brain tumor surgery. In subgroup analysis, preoperative ALT level and BMI were associated with PELE in patients treated with 5-ALA. Therefore, when 5-ALA is administered to patients with the upper normal value of serum ALT and overweight, attention is paid to PELE.

**Funding information** Seoul National University Bundang Hospital provided financial support in the form of academic funding (grant number: 02-2014-033).

## Compliance with ethical standards

**Disclaimer** The sponsor had no role in the design or conduct of this research.

**Conflict of interest** The authors declare that they have no conflict of interest.

**Ethical approval** This retrospective study was conducted after approval from the Institutional Review Board of Seoul National University Hospital (IRB no. 1811-078-985). All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee (name of institute/committee) and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Because this was a retrospective study, for this type of study formal consent is not required.

**Informed consent** This article does not contain any studies with human participants performed by any of the authors.

## References

- Björnsson E (2008) Hepatotoxicity associated with antiepileptic drugs. *Acta Neurol Scand* 118:281–290
- Björnsson ES (2017) Drug-induced liver injury due to antibiotics. *Scand J Gastroenterol* 52:617–623
- Chen TL, Ueng TH, Chen SH, Lee PH, Fan SZ, Liu CC (1995) Human cytochrome P450 mono-oxygenase system is suppressed by propofol. *Br J Anaesth* 74:558–562
- Chung IWH, Eljamel S (2013) Risk factors for developing oral 5-aminolevulinic acid-induced side effects in patients undergoing fluorescence guided resection. *Photodiagn Photodyn Ther* 10: 362–367
- Cowan RE, Jackson BT, Grainger SL, Thompson RP (1991) Effects of anesthetic agents and abdominal surgery on liver blood flow. *Hepatology* 14:1161–1166
- Giannini EG, Testa R, Savarino V (2005) Liver enzyme alteration: a guide for clinicians. *CMAJ* 172:367–379
- Honorato-Cia C, Martínez-Simón A, Cacho-Asenjo E, Guillén-Grima F, Tejada-Solís S, Díez-Valle R (2015) Safety profile of 5-aminolevulinic acid as a surgical adjunct in clinical practice: a review of 207 cases from 2008 to 2013. *J Neurosurg Anesthesiol* 27: 304–309
- Kim J, Jo I (2010) Relationship between body mass index and alanine aminotransferase concentration in non-diabetic Korean adults. *Eur J Clin Nutr* 64:169–175
- Lee WM (2003) Drug-induced hepatotoxicity. *N Engl J Med* 349: 474–485
- Lichtenbelt BJ, Olofsen E, Dahan A, van Kleef JW, Struys MMRF, Vuyk J (2010) Propofol reduces the distribution and clearance of midazolam. *Anesth Analg* 110:1597–1606
- Marbacher S, Klinger E, Schwyzer L, Fischer I, Nevzati E, Diepers M, Roelcke U, Fathi A-R, Coluccia D, Fandino J (2014) Use of fluorescence to guide resection or biopsy of primary brain tumors and brain metastases. *Neurosurg Focus* 36:E10
- Offersen CM, Skjoeth-Rasmussen J (2017) Evaluation of the risk of liver damage from the use of 5-aminolevulinic acid for intra-operative identification and resection in patients with malignant gliomas. *Acta Neurochir* 159:145–150
- Rahimzadeh P, Safari S, Faiz SHR, Alavian SM (2014) Anesthesia for patients with liver disease. *Hepat Mon* 14:e19881
- Rai R, Nagral S, Nagral A (2012) Surgery in a patient with liver disease. *J Clin Exp Hepatol* 2:238–246
- Rinella ME, Alonso E, Rao S, Whittington P, Fryer J, Abecassis M, Superina R, Flamm SL, Blei AT (2001) Body mass index as a predictor of hepatic steatosis in living liver donors. *Liver Transpl* 7:409–414
- Ruland S, Aiyagari V (2007) Cerebral autoregulation and blood pressure lowering. *Hypertension* 49:977–978
- Sachar M, Anderson KE, Ma X (2016) Protoporphyrin IX: the good, the bad, and the ugly. *J Pharmacol Exp Ther* 356:267–275
- Savarese D (2018) Common terminology criteria for adverse events - UpToDate. <https://www.uptodate.com/contents/common-terminology-criteria-for-adverse-events>. Accessed 09 Mar 2018
- Stepp H, Stummer W (2018) 5-ALA in the management of malignant glioma. *Lasers Surg Med* 50:399–419
- Stummer W, Goetz C (2000) Fluorescence-guided resection of glioblastoma multiforme by using 5-aminolevulinic acid-induced porphyrins: a prospective study in 52 consecutive patients. *J Neurosurg* 93:1003–1013
- Stummer W, Pichlmeier U, Meinel T, Wiestler OD, Zanella F, Reulen H-J, Group A-GS (2006) Fluorescence-guided surgery with 5-aminolevulinic acid for resection of malignant glioma: a randomised controlled multicentre phase III trial. *Lancet Oncol* 7: 392–401
- Stupp R, Hegi ME, Mason WP, van den Bent MJ, Taphoorn MJB, Janzer RC, Ludwin SK, Allgeier A, Fisher B, Belanger K, Hau P, Brandes AA, Gijtenbeek J, Marosi C, Vecht CJ, Mokhtari K, Wesseling P, Villa S, Eisenhauer E, Gorlia T, Weller M, Lacombe D, Cairncross JG, Mirimanoff R-O, Groups EOfRaToCBTaRO, Group NCIoCCT (2009) Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol* 10:459–466
- Teixidor P, Arráez MÁ, Villalba G, Garcia R, Tardáguila M, González JJ, Rimbau J, Vidal X, Montané E (2016) Safety and

- efficacy of 5-aminolevulinic acid for high grade glioma in usual clinical practice: a prospective cohort study. *PLoS One* 11: e0149244
24. Webber J, Kessel D, Fromm D (1997) Side effects and photosensitization of human tissues after aminolevulinic acid. *J Surg Res* 68: 31–37
  25. Weller M, van den Bent M, Hopkins K, Tonn JC, Stupp R, Falini A, Cohen-Jonathan-Moyal E, Frappaz D, Henriksson R, Balana C, Chinot O, Ram Z, Reifenberger G, Soffiatti R, Wick W, Glioma EAFN-OETFoM (2014) EANO guideline for the diagnosis and treatment of anaplastic gliomas and glioblastoma. *Lancet Oncol* 15:e395–e403
  26. Woods CP, Hazlehurst JM, Tomlinson JW (2015) Glucocorticoids and non-alcoholic fatty liver disease. *J Steroid Biochem Mol Biol* 154:94–103

## Comments

A study that once more demonstrated the safety of using 5-ALA as adjunct for malignant brain tumor surgery with regards to the medical influence of the substance. Still the most dangerous is to have the tumor and transient elevation of liver enzymes are of less relevance.

Jane Skjoth-Rasmussen  
Copenhagen, Denmark

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.