



Original research

A data-driven, meaningful, easy to interpret, standardised accelerometer outcome variable for global surveillance



Alex V. Rowlands^{a,b,c,*}, Lauren B. Sherar^d, Stuart J. Fairclough^e, Tom Yates^{a,b}, Charlotte L. Edwardson^{a,b}, Deirdre M. Harrington^{a,b}, Melanie J. Davies^{a,b}, Fehmidah Munir^d, Kamlesh Khunti^{a,f}, Victoria H. Stiles^g

^a Diabetes Research Centre, University of Leicester, Leicester General Hospital, UK

^b NIHR Leicester Biomedical Research Centre, UK

^c Alliance for Research in Exercise, Nutrition and Activity (ARENA), Sansom Institute for Health Research, Division of Health Sciences, University of South Australia, Australia

^d School of Sport, Exercise and Health Sciences, Loughborough University, UK

^e Movement Behaviours, Health, and Wellbeing Research Group, Department of Sport and Physical Activity, Edge Hill University, UK

^f NIHR Collaboration for Leadership in Applied Health Research and Care East Midlands, Leicester General Hospital, UK

^g Sport and Health Sciences, College of Life and Environmental Sciences, University of Exeter, UK

ARTICLE INFO

Article history:

Received 7 March 2019

Received in revised form 29 May 2019

Accepted 21 June 2019

Available online 1 July 2019

Keywords:

Physical activity

Population

Acceleration

Measurement

Research-grade accelerometer

Wrist-worn

ABSTRACT

Objectives: Our aim is to demonstrate how a data-driven accelerometer metric, the acceleration above which a person's most active minutes are accumulated, can (a) quantify the prevalence of meeting current physical activity guidelines for global surveillance and (b) moving forward, could inform accelerometer-driven physical activity guidelines. Unlike cut-point methods, the metric is population-independent (e.g. age) and potentially comparable across datasets.

Design: Cross-sectional, secondary data analysis.

Methods: Analyses were carried out on five datasets using wrist-worn accelerometers: children (N = 145), adolescent girls (N = 1669), office workers (N = 114), pre- (N = 1218) and post- (N = 1316) menopausal women, and adults with type 2 diabetes (N = 475). Open-source software (GGIR) was used to generate the magnitude of acceleration above which a person's most active 60, 30 and 2 min are accumulated: M60_{ACC}, M30_{ACC} and M2_{ACC}, respectively.

Results: The proportion of participants with M60_{ACC} (children) and M30_{ACC} (adults) values higher than accelerations representative of brisk walking (i.e., moderate-to-vigorous physical activity) ranged from 17 to 68% in children and 15 to 81% in adults, tending to decline with age. The proportion of pre- and post-menopausal women with M2_{ACC} values meeting thresholds for bone health ranged from 6 to 13%.

Conclusions: These metrics can be used for global surveillance of physical activity, including assessing prevalence of meeting current physical activity guidelines. As accelerometer and corresponding health data accumulate it will be possible to interpret the metrics relative to age- and sex- specific norms and derive evidence-based physical activity guidelines directly from accelerometer data for use in future global surveillance. This is where the potential advantages of these metrics lie.

© 2019 Sports Medicine Australia. Published by Elsevier Ltd. All rights reserved.

Practical implications

- The magnitude of acceleration above which a person's most active minutes are accumulated throughout the day could be useful for monitoring and surveillance, e.g. prevalence of meeting physical activity guidelines.

- Unlike cut-point methods, the metrics are population-independent (e.g. age), do not rely on calibration studies, and no person scores zero.
- The metrics can be translated post-hoc in terms of walking/running, providing a public health friendly interpretation of the results relevant to a range of clinical populations.
- These metrics will encourage use of accelerometer data for global surveillance of meeting activity guidelines. Further, as accelerometer and corresponding health data accumulate it will be possible to use the metrics to derive evidence-based physical activity

* Corresponding author.

E-mail address: alex.rowlands@leicester.ac.uk (A.V. Rowlands).

guidelines directly from accelerometer data that express recommendations in terms of representative activity type, e.g. 'brisk walking'.

1. Introduction

National and/or large-scale surveys of physical activity through accelerometers are now commonplace in many countries worldwide.^{1–5} The World Health Organisation's recent Global Activity Action Plan on Physical Activity 2018–2030⁶ highlights monitoring and surveillance, using robust and reliable data, as the cornerstone to the implementation and evaluation of national strategies. Accelerometers provide a valid measure of physical activity⁷; however, a lack of consensus on robust and consistent methods to reduce and analyse data to create meaningful and easy to interpret outcome variables, is hampering monitoring and evaluation activities.

For example, epidemiological studies and surveillance studies frequently create variables from accelerometer-assessed moderate-to-vigorous physical activity (MVPA) using intensity cut-points. The problems with using cut-points to quantify activity are well documented^{8–10} but, briefly, include: (1) cut-points are protocol-, and population- (e.g. age-group) specific, leading to results that are not comparable across studies^{8–10}; (2) two participants with similar levels of activity score very differently if one has activity falling just above the cut-point and one has activity falling just below the cut-point; (3) many participants fail to obtain any activity above cut-points (particularly in the vigorous range), consequently a large number of people simply score zero minutes. Recently, in an examination of how cut-points influence estimates of physical activity, Migueles et al.^{11, p1} stated that it was 'not possible (and probably will never be) to know the prevalence of meeting the physical activity guidelines based on accelerometer data'. Clearly a new approach to analysing and interpreting accelerometer data is needed.

An alternative approach is to identify the minimum acceleration value above which a person's most active minutes, for example 30 min (M30_{ACC}), is accumulated. The active minutes can be accumulated in any way across the day, with no need for the activity to be in bouts, in line with recent physical activity recommendations.¹² With this approach the metric is population-independent and derived from directly measured acceleration, thus not relying on assumptions as cut-points do,⁹ and the intensity is captured regardless of level of activity with no person scoring zero. This bears similarities to the peak 30 min walking cadence (steps/min) proposed by Tudor-Locke et al.¹³ as a practical estimate of 'best natural effort' during habitual physical activity.

Moving forward, as accelerometer and corresponding health data accumulate, these data-driven population-independent metrics could be used to inform accelerometer-driven physical activity guidelines as recommended by Troiano et al.,¹⁰ rather than inappropriately evaluating physical activity assessed by accelerometer cut-points to guidelines developed from self-report data, which are conceptually different.¹⁰ For example, the M30_{ACC} and/or M60_{ACC} that is positively associated with a given health marker, e.g. adiposity, could be determined. This M30_{ACC} and/or M60_{ACC} value could then be used for surveillance which, importantly, would facilitate surveillance using the same physical activity metric as used to garner the evidence. As with the peak cadence,¹³ the metric itself is population independent, but norms will vary, e.g. by age and sex. As data accumulate, it would be possible to interpret the M30_{ACC} and M60_{ACC} relative to age- and sex-specific norms and/or relative to values associated with health markers.

To facilitate public-health recommendations, translation of the metrics to public-health friendly representative activity types is

desirable, e.g. brisk walking, and/or MVPA. This translation is necessarily population-specific and thus bears similarities to cut-point analyses. However, crucially this is only in the translation of the data for activity recommendations because all analyses are carried out on the population-independent metrics.⁹ In contrast, when using cut-points, thresholds are imposed on the data from the outset to collapse data into categories for analysis, rendering it impossible to subsequently compare any datasets deploying different cut-points.

For example, assume that a child has an M60_{ACC} value of 225 mg. Currently, we do not have the data to compare this to accelerometer-driven physical activity guidelines; however, we can assess whether the child is meeting the current 60 min daily MVPA guideline¹² by comparing their M60_{ACC} value to one of the MVPA cut-points available. According to the 200 mg MVPA cut-point published by Hildebrand et al.,¹⁴ the child exceeds the 60 min of MVPA per day recommendations,¹² while according to a more stringent 250 mg MVPA cut-point published by Phillips et al.,¹⁵ the child does not quite reach the recommendations. If a cut-point approach had been used to analyse the data, the child's score could not be compared to any alternative cut-point or threshold.

For the purposes of a simple demonstration of how these metrics could be used for surveillance of adherence to current physical activity guidelines,¹² we looked at the daily average acceleration above which the most active 30 min (M30_{ACC}, adults) or 60 min (M60_{ACC}, children) was obtained. It would be possible to alter the number of minutes over which the minimum acceleration is considered, depending on the health outcome of interest or the guideline being assessed. For example, in a large cross-sectional observational study, Stiles et al.¹⁶ demonstrated that accumulating 1–2 min of accelerometer-assessed high intensity activity, equivalent to running, was associated with bone health in pre- and post-menopausal women.

The primary aim of this paper is to demonstrate how the acceleration above which a person's most active minutes are accumulated, can be used to quantify prevalence of meeting existing physical activity guidelines. A secondary aim is to illustrate that, as accelerometer and corresponding health data accumulate, there is potential for these population-independent metrics to be used to inform accelerometer-driven physical activity guidelines.

2. Methods

Secondary data analyses were carried out on five diverse datasets: 10 year old children,¹⁷ adolescent girls,^{18,19} adult office workers,²⁰ pre- and post-menopausal women,¹⁶ and adults with type 2 diabetes.²¹ All participants gave assent (children and adolescent girls) or informed consent (adults). Parents/guardians of the children gave written informed consent and parents/guardians of the adolescent girls returned an opt-out consent form if they did not want their child to participate. All studies received the appropriate institutional ethics approval.

In all samples, wrist worn accelerometers were worn 24 h a day for up to 7 days. The children and adult office workers wore the ActiGraph GT9X Link (ActiGraph, Pensacola, FL, USA), the adolescent girls and the adults with type 2 diabetes wore the GENEActiv (ActivInsights Ltd., Cambridgeshire, UK) and the pre- and post-menopausal women wore the Axivity AX3 (Axivity, Newcastle, UK). The pre- and post-menopausal women wore the monitor on their dominant wrist, all other samples wore monitors on the non-dominant wrist. All monitors were initialised to record accelerations at 100 Hz, except the adult office workers whose monitors were initialised at 30 Hz.

ActiGraphs were initialised and downloaded using ActiLife version 6.11.9 (ActiGraph, Pensacola, FL, USA). Data were saved in raw




Fig. 1. Percentiles for the magnitude of acceleration above which the females' most active (a) 60, (b) 30 and (c) 2 min are accumulated: $M60_{ACC}$; $M30_{ACC}$ and $M2_{ACC}$ (mg). Black dashes /dashed lines represent: (a) $M60_{ACC}$ and (b) $M30_{ACC}$ at the intensity of a brisk walk (lower dashed line) or fast walk (upper dashed line); (c) $M2_{ACC}$ at the bone health threshold: medium running for pre-menopausal women and slow running for post-menopausal women.¹⁶ The stick figures indicate representative activities associated with the acceleration for: (a) slow walk and brisk walk; (b) slow walk, brisk walk and slow run; (c) brisk walk and slow run.

format as GT3X files, before being converted to raw csv file format for signal processing. GENEActivs were initialised and data downloaded in binary format using GENEActiv PC (version 3.1). Activity data were downloaded from UK Biobank in .cwa format, auto-calibrated, resampled (100 Hz) and converted to .wav format using open-source software (Omgui Version 1.0.0.28; Open Movement, Newcastle, UK).

All accelerometer files were processed and analysed with R-package GGIR version 1.6–7 (<http://cran.r-project.org>).^{22,23} Signal processing in GGIR included auto-calibration using local gravity as a reference²² (apart from the Axiivity files which were auto-calibrated when converted to .wav files); detection of sustained abnormally high values; detection of non-wear; and calculation of the average magnitude of dynamic acceleration corrected for gravity (Euclidean Norm minus 1 g, ENMO). These were averaged over 1 or 5 s epochs (1 s: pre- and post-menopausal women (UK Biobank), 5 s: children, adolescent girls, adult office workers and adults with type 2 diabetes) and expressed in milli-gravitational units (mg). Note that as the average over an epoch is used the values are independent of epoch duration.

Participants were excluded if their accelerometer files showed: post-calibration error greater than 0.01 g (10 mg), fewer than three days of valid wear (defined as >16 h per day), or wear data was not present for each 15 min period of the 24 h cycle, i.e. data from at least one day of measurement had to be present for each 15 min period. The following metrics were generated and averaged across all valid days: average acceleration; intensity gradient (intensity distribution²¹); acceleration above which a person's most active X minutes (MX_{ACC}) are accumulated: $M60_{ACC}$ (mg); $M30_{ACC}$ (mg), $M2_{ACC}$ (mg), (within the GGIR package, these metrics are obtained in part2 using: q levels (0,24h): 1380/1440, 1410/1440 and 1438/1440). These MX_{ACC} statistics rank the acceleration for each epoch during the day in descending order to obtain the acceleration above which the person's most active X minutes are accumulated. Note this differs from the peak cadence described by Tudor-Locke et al.,¹³ which is the average of the cadence for the peak 30 min. As acceleration measured at the dominant wrist is approximately 10% higher than the non-dominant,²⁴ magnitudes of $M60_{ACC}$, $M30_{ACC}$ and $M2_{ACC}$ were reduced by 10% for dominant wrist placement (pre- and post-menopausal women).




Fig. 2. Percentiles for the magnitude of acceleration above which the males' most active (a) 60, (b) 30 and (c) 2 min are accumulated: M60_{ACC}; M30_{ACC} and M2_{ACC} (mg). Black dashes/dashed lines represent: (a) M60_{ACC} and (b) M30_{ACC} at the intensity of a brisk walk (lower dashes/dashed line) or fast walk (upper dashes/dashed line). The stick figures indicate representative activities associated with the acceleration for: (a) slow walk and brisk walk; (b) slow walk, brisk walk and slow run; (c) brisk walk and slow run.

Analyses: Descriptive statistics were calculated using mean (standard deviation (SD)) for continuous variables and percentage for categorical variables.

Percentiles (5th–95th percentile) were graphed for females (all samples) and males (where available) for the M60_{ACC}, M30_{ACC} and M2_{ACC}. Presenting percentiles for each metric illustrates the magnitude of the most active X minutes, from the least to the most active participants, within each sample. To address our primary aim, the proportion of each sample meeting the MVPA physical activity guidelines, operationalised for the purposes of this demonstration as a daily average of 30 min for adults and 60 min for children and adolescents, was calculated. For MVPA, we estimated acceleration values representative of a brisk walk (5 km/h, ≈ 3.6 METs: 170 mg adults; 200 mg children¹⁴) and of a fast walk (5.6 km/h, ≈ 4.5 METs: 250 mg adults; 300 mg children^{14,25,26}). To do this we took the MET values associated with the walking speeds from activity energy expenditure compendiums for adults²⁵ and children,²⁶ converted these to VO₂ (assuming 1 MET=3.5 ml/kg/min for adults²⁵ and 1 MET=5 ml/kg/min for children²⁶), then used the child and adult regression equations presented by Hildebrand et al.¹⁴ to obtain estimates of acceleration values representa-

tive of brisk and fast walking. In addition, the proportion of pre- and post-menopausal women meeting the recently proposed accelerometer-driven guide of 2 min high-intensity activity associated with bone health¹⁶ was calculated. The thresholds (>1000 mg (medium run) pre-menopausal, >750 mg, post-menopausal (slow run)) were generated using dominant wrist data,¹⁶ so are adjusted by -10%.²⁴

3. Results

Valid accelerometer data files were available for 64% of 10 year old children (N=145, age (mean (SD))=9.6(0.3) year, 57% female), 96% of adolescent girls (age 11–12 year: N=974, age 12.3(0.4) year; age 13–14 year, age 13.6(0.4) year, 100% female), 78% of adult office workers (N=114, age 41.2(10.9) year, 80% female) and 99% of adults with type 2 diabetes (N=475, age 64.2(8.7) year, 36% female). All accelerometer files for the pre- and post-menopausal women from UK Biobank meeting the criteria of Stiles et al.¹⁶ were included (N=1218 pre-menopausal, age 46.2(3.9) year, N=1316 post-menopausal, age 59.0(5.1) year). Further descriptive characteristics are presented in Supplementary material. As expected,

Table 1
Proportion of each sample meeting MVPA guidelines operationalised as 60 min per day (children) or 30 min per day (adults) of brisk walking or fast walking.

		Brisk walk		Fast walk	
		Female	Male	Female	Male
Children 10 y olds	60 min	>200 mg	>200 mg	>300 mg	>300 mg
	% meeting guideline	39%	68%	2%	21%
	NO: M60 _{ACC}	164.9 (21.3)	157.8 (27)	192.0 (42.2)	212.1 (53.1)
	YES: M60 _{ACC}	242.3 (28.4)	289.1 (74.4)	302.2 (1.64)	377.2 (70.2)
11–12 year olds	% meeting guideline	26%		2%	
	NO: M60 _{ACC}	161.4 (23.3)		177.6 (36.8)	
	YES: M60 _{ACC}	235.6 (39.0)		355.2 (43.30)	
13–14 year olds	% meeting guideline	17%		1%	
	NO: M60 _{ACC}	154.6 (25.1)		164.4 (25.6)	
	YES: M60 _{ACC}	228.1 (30.1)		320.1 (10.5)	
Adults Office workers	30 min	>170 mg	>170 mg	>250 mg	>250 mg
	% meeting guideline	45%	55%	11%	14%
	NO: M30 _{ACC}	141.7 (18.2)	147.7 (16.0)	161.4 (30.9)	167.5 (27.4)
	YES: M30 _{ACC}	246.8 (133.6)	212.8 (46.0)	427.9 (188.5)	282.8 (19.5)
Pre-menopausal women	% meeting guideline	81%		25%	
	NO: M30 _{ACC}	150.8 (15.1)		191.2 (31.3)	
	YES: M30 _{ACC}	244.5 (86.0)		334.5 (107.6)	
Post-menopausal women	% meeting guideline	63%		11%	
	NO: M30 _{ACC}	145.6 (17.6)		177.3 (33.5)	
	YES: M30 _{ACC}	219.0 (53.4)		303.4 (69.9)	
Adults with type 2 diabetes	% meeting guideline	15%	16%	2%	3%
	NO: M30 _{ACC}	118.7 (23.9)	123.9 (26.5)	129.2 (35.1)	132.8 (33.9)
	YES: M30 _{ACC}	203.8 (28.4)	221.2 (89.1)	268.8 (15.4)	354.5 (0)

physical activity volume (average acceleration) tended to decrease and the intensity distribution (intensity gradient) worsen with increasing sample mean age, see Supplementary material.

Figs. 1 and 2 show percentile plots for M60_{ACC}, M30_{ACC} and M2_{ACC} for females and males, respectively, in order of increasing sample mean age. Stick figures mark accelerations associated with a brisk walk (5 km/h), fast walk (5.6 km/h) and slow run (≥ 8 km/h) on the y-axes to illustrate how the data could be translated in a public-health friendly way.¹⁴ The expected age-related decline in intensity of physical activity appeared to be relatively greater the fewer minutes considered (i.e. there is a steeper drop from childhood to adulthood for the M2_{ACC} than the M30_{ACC}), but also for higher percentiles (i.e. higher intensity; the top percentiles dropped more steeply than the lower percentiles) within a given outcome (Figs. 1b–c, 2 b–c). Sex differences were most evident in 10 year old children, with the intensity of boys' activity greater than that of girls' (Figs. 1a–c compared to 2 a–c).

Table 1 shows the proportion of each sample meeting MVPA guidelines operationalised as 60 min per day (children) or 30 min per day (adults) of brisk walking or fast walking. The MX_{ACC} above which the most active time is accumulated is shown for those meeting and not meeting the guidelines. The proportions of pre- and post-menopausal women meeting the recent accelerometer-derived guide proposed for bone health (2 min > 1000 mg (medium run) pre-menopausal, >750 mg (slow run) post-menopausal)¹⁶ were 6% and 13%, respectively (Fig. 1c).

4. Discussion

Given the rising use of accelerometers, including their use in large-scale surveys e.g. Refs. 1–5, it is important to have simple to derive and easy to interpret accelerometer variables that has potential for use to compare physical activity across datasets/populations/countries. This would facilitate global surveillance and the development of evidence-based physical activity guidelines directly from accelerometer data. As data accumulate, physical activity of groups and individuals can be interpreted relative to age- and sex-specific norms and/or relative to

values associated with health markers. While the values themselves are not immediately intuitive, this is also true of many metrics that are commonly used by researchers, clinicians and the public.²⁸ For example, risk thresholds for health markers such as body mass index, blood pressure, and cholesterol are routinely used and widely understood. As outlined by Welk et al.,²⁸ a range of instruments are used to obtain measures of blood pressure, but the use of a standardised metric makes it possible for researchers, clinicians and patients to discuss a common number. This would also be possible with widespread use of standardised population-independent accelerometer measures of physical activity.

In this paper, we demonstrate how presenting percentiles for population-independent metrics such as the M60_{ACC} and M30_{ACC} can be used now to estimate adherence to current MVPA guidelines. The numerous problems associated with applying cut-points to accelerometer data^{8–10} are avoided as the data and results presented are data-driven. Comparison to any representative activity, cut-point or, more importantly, any future health-related accelerometer threshold is possible and can be carried out post-hoc with no access to the original data needed. Further, demographic-specific translations can be carried out post-hoc to facilitate public-health friendly recommendations using accelerations representative of typical activities. Crucially, population-specific translation is only for interpretation and has no bearing on analyses or results presented. This means the metrics and results retain their population-independence.⁹

Further, comparison or translation is not tied to an exact acceleration value for a representative activity or cut-point. For example, if a child accumulates 60 min of activity in the acceleration range of 185–199 mg, their M60_{ACC} will be 185 mg. Another child may accumulate 60 min with accelerations just exceeding 200 mg. With the cut-point method, these similar activity levels look very disparate; 0 min of MVPA and 60 min of MVPA, respectively. Their M60_{ACC}, on the other hand reflects the smaller discrepancy in activity level that is evident; 185 mg and 200 mg. At a group level, presenting percentiles for the MX_{ACC} values as illustrated herein (Figs. 1 and 2), displays the proportion of a sample achieving X min at any given intensity. In contrast, once cut-points have been

applied, any activity accumulated just below a given cut-point will always be disregarded, irrespective of how the data are presented.

By decreasing the number of minutes of interest the metric can be used to focus on aspects of health that benefit from short, high-intensity bursts of activity, e.g. bone health.^{16,29} Accelerometer-derived physical activity intensity guides for bone health have recently been proposed for pre- and post-menopausal women using data from a UK Biobank¹⁶; these metrics could be used to further test this recommendation and to derive guidelines from accelerometer data specific to bone health in men and children.

To aid translation, we expressed the acceleration magnitudes in relation to representative activities, e.g. brisk walk, fast walk and run. Currently there are limited data from which to draw these estimates. To enhance translation of these metrics there is a need to generate more data showing the acceleration ranges associated with representative activities across a wide range of demographics. Note, this is only for translation and is not necessary for generation of the accelerometer metrics from data, or for developing the evidence base necessary to derive physical activity guidelines directly from accelerometer data.

The acceleration magnitudes tended to be higher for the pre-menopausal women who wore the Axivity on their dominant wrist than for the slightly younger office workers who wore the ActiGraph on their non-dominant wrist. While this may be due to the sedentary nature of the office job, it could reflect the non-representative nature of the samples, indicate that the –10% reduction in acceleration for dominant wrist placement²⁴ was insufficient, and/or that there were differences between the ActiGraph and the Axivity. While raw data from the GENEActiv and Axivity accelerometers compare well,²⁴ ‘raw’ data from the ActiGraph GT9X is passed through a filter that suppresses higher intensity accelerations. As the filter suppresses higher accelerations this will be most evident in active populations and for the shorter MX_{ACC} durations and will limit the comparability of data collected with the ActiGraph to data collected with the Axivity or GENEActiv. Further, the accelerometer sampling frequency and epoch differed between some studies. As the metrics are sampling frequency independent this should not impact on the outcomes generated with GGIR, but this needs to be confirmed empirically. It is also possible that the use of 1 s and 5 s epochs may have impacted on the MX_{ACC} outcomes, however, in our previous study data summarised in 1 s and 5 s epochs were comparable.³⁰ Crucially, as with any accelerometer outcome, the metrics will differ according to where the monitor is worn, e.g. output collected at the wrist is not comparable to data collected at the hip. Any norms generated for these, or any other, metrics will have to be placement-specific. It is important to consider all of the above when attempting to compare these metrics between studies.

5. Conclusion

Cut-point approaches to analysing accelerometer data are not appropriate for assessing the prevalence of meeting guidelines globally.¹¹ Metrics reflecting the acceleration above which the most active minutes are accumulated are a standardised, easy to interpret, and population-independent method that has potential for assessing prevalence of physical activity and comparing activity between demographics and/or studies. These simple to derive variables facilitate global surveillance and dose–response studies. Furthermore, translating the metrics in terms of representative activities (e.g. brisk walking) can provide a public-health friendly interpretation of the results.⁹ Currently, guidelines are largely derived from self-report data.¹⁰ As accelerometer and corresponding health data accumulate it will be possible to

derive evidence-based physical activity guidelines directly from accelerometer data.

Conflicts of interest

Declarations of interest: none

Funding

The Active Schools: Skelmersdale (ASSK) physical activity intervention study was funded by West Lancashire Sport Partnership UK, West Lancashire Community Leisure UK, and Edge Hill University Ormskirk UK. The adolescent girls’ data are from the Girls Active evaluation, which was funded by the NIHR Public Health Research Programme (13/90/30). The adult office workers data are from the SMArT Work trial, which was sponsored by Loughborough University. The project was funded by the Department of Health Policy Research Programme (project No PR-R5-0213-25004). The pre- and post-menopausal data are from UK Biobank. The processing and analysis of these data was supported by an internal grant from the University of Exeter (UK) Project Development Fund (Science).

University of Leicester authors are supported by the NIHR Leicester Biomedical Research Centre, and the Collaboration for leadership in Applied Health Research and Care (CLAHRC) East Midlands. The views expressed are those of the authors and not necessarily those of the NHS, NIHR, or Department of Health.

Authors’ contributions

AR conceived the study. **AR, LS, SF** and **VS** designed the study. **SF, CE, DH, MD, FM** and **VS** were involved in data acquisition. **AR** analysed the data and wrote the manuscript. All authors were involved in data presentation and interpretation and all actively engaged in collaboratively drafting and revising the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors thank project staff involved with ASSK and Dr Sarah Taylor for data collection; all researchers and project staff involved in the Girls Active evaluation, SMArT Work trial, UK Biobank and CODEC (adults with type 2 diabetes) for access to the data used herein. Analysis of the pre-menopausal and post-menopausal samples was conducted using the UK Biobank Resource (Reference 10995).

We also thank: the participating schools in ASSK, children and teachers for their participation; the pupils and teachers who took part in the Girls Active evaluation study, the Youth Sport Trust (YST); the participants in the SMArT Work trial; the participants in UK Biobank; and participants in the CODEC study.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.jsams.2019.06.016>.

References

1. da Silva ICM, van Hees VT, Ramires VV et al. Physical activity levels in three Brazilian birth cohorts as assessed with raw triaxial wrist accelerometry. *Int J Epidemiol* 2014; 43(6):1959–1968.
2. Doherty A, Jackson D, Hammerla N et al. Large scale population assessment of physical activity using wrist worn accelerometers: the UK biobank study. *PLoS One* 2017; 12(2). <http://dx.doi.org/10.1371/journal.pone.0169649>, e0169649.
3. Troiano RP, Berrigan D, Dodd KW et al. Physical activity in the United States measured by accelerometer. *Med Sci Sport Exerc* 2008; 40:181–188.

4. Li X, Kearney PM, Keane E et al. Levels and sociodemographic correlates of accelerometer-based physical activity in Irish children: a cross-sectional study. *J Epidemiol Commun Health* 2017; 71(6):521–527. <http://dx.doi.org/10.1136/jech-2016-207691>.
5. Menai M, van Hees VT, Elbaz A et al. Accelerometer assessed moderate-to-vigorous physical activity and successful ageing: results from the Whitehall II study. *Sci Rep* 2017; 8:45772. <http://dx.doi.org/10.1038/srep45772>.
6. World Health Organisation. *Global action plan on physical activity 2018–2030: more active people for a healthier world*. Geneva, World Health Organization, 2018. Licence: CC BY-NC-SA 3.0 IGO. 104 p. Available from: <http://apps.who.int/iris/bitstream/handle/10665/272722/9789241514187-eng.pdf>. [Accessed 25 November 2018].
7. White T, Westgate K, Wareham NJ et al. Estimation of physical activity energy expenditure during free-living from wrist accelerometry in UK adults. *PLoS One* 2016; 11(12):e0167472. <http://dx.doi.org/10.1371/journal.pone.0167472>.
8. Brazendale K, Beets MW, Bornstein DB et al. Equating accelerometer estimates among youth: the Rosetta stone 2. *J Sci Med Sport* 2016; 19:242–249.
9. Rowlands AV. Moving forward with accelerometer-assessed physical activity: two strategies to ensure meaningful, interpretable & comparable measures. *Pediatr Exerc Sci* 2018; 30:450–456. <http://dx.doi.org/10.1123/pes.2018-0201>.
10. Troiano RP, McClain JJ, Brychta RJ et al. Evolution of accelerometer methods for physical activity research. *Br J Sports Med* 2014; 48:1019–1023.
11. Migueles JH, Cadenas-Sanchez C, Tudor-Locke C et al. Comparability of published cut-points for the assessment of physical activity: implications for data harmonization. *Scand J Med Sci Sports* 2019; 1–9. <http://dx.doi.org/10.1111/sms.13356>.
12. *Physical activity guidelines for Americans 2018*, 2nd ed. Washington, DC, US Department of Health and Human Services, 2018, 118 p. Available from: https://health.gov/paguidelines/secondedition/pdf/Physical_Activity_Guidelines_2nd_edition.pdf. [Accessed 18 November 2018].
13. Tudor-Locke C, Han H, Aguiar EJ et al. How fast is fast enough? Walking cadence (steps/min) as a practical estimate of intensity in adults: a narrative review. *Br J Sports Med* 2018; 52(12):776–788. <http://dx.doi.org/10.1136/bjsports-2017-097628>.
14. Hildebrand M, Van Hees VT, Hansen BH. Age-group comparability of raw accelerometer output from wrist- and hip-worn monitors. *Med Sci Sport Exerc* 2014; 46:1816–1824.
15. Phillips LRS, Parfitt CG, Rowlands AV. Calibration of the GENEA accelerometer for assessment of physical activity intensity in children. *J Sci Med Sport* 2013; 6:124–128.
16. Stiles VS, Metcalf BS, Knapp KM et al. A small amount of precisely measured high intensity habitual physical activity predicts bone health in pre- and post-menopausal women in UK Biobank. *Int J Epidemiol* 2017; 46:1847–1856. <http://dx.doi.org/10.1093/ije/dyx080>.
17. Taylor S, Noonan R, Knowles Z et al. Evaluation of a pilot school-based physical activity clustered randomised controlled trial—active schools: Skelmersdale. *Int J Environ Res Public Health* 2018; 15(5):1011.
18. Edwardson CL, Harrington DM, Yates T et al. A cluster randomised controlled trial to investigate the effectiveness and cost effectiveness of the ‘Girls Active’ intervention: a study protocol. *BMC Public Health* 2015; 15(1):526.
19. Harrington D, Davies MJ, Bodicoat DH et al. Effectiveness of the ‘Girls Active’ school-based physical activity programme: a cluster randomised controlled trial. *Int J Behav Nutr Phys Act* 2018; 15:40. <http://dx.doi.org/10.1186/s12966-018-0664-6>.
20. Edwardson CL, Yates T, Biddle SJH et al. Effectiveness of the Stand More AT (SMArT) work intervention: cluster randomised controlled trial. *BMJ* 2018; 363:k3870. <http://dx.doi.org/10.1136/bmj.k3870>.
21. Rowlands AV, Edwardson CL, Davies MJ et al. Beyond cut-points: accelerometer metrics that capture the physical activity profile. *Med Sci Sport Exerc* 2018; 50(6):1323–1332.
22. van Hees VT, Fang Z, Langford J et al. Auto-calibration of accelerometer data for free-living physical activity assessment using local gravity and temperature: an evaluation on four continents. *J Appl Physiol* 2014; 117(7):738–744.
23. van Hees VT, Gorzelniak L, Dean León EC et al. Separating movement and gravity components in an acceleration signal and implications for the assessment of human daily physical activity. *PLoS One* 2013; 8(4):e61691. <http://dx.doi.org/10.1371/journal.pone.0061691>.
24. Rowlands AV, Plekhanova T, Yates T et al. A basis for harmonisation of accelerometer physical activity outcomes in epidemiology. *J Meas Phys Behav* 2019 [in press].
25. Ainsworth BE, Haskall WL, Whitt MC et al. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sport Exerc* 2000; 32:S498–504.
26. Lyden K, Keadle SK, Staudenmeyer J et al. Energy cost of common activities in children and adolescents. *J Phys Act Health* 2013; 10:62–69.
28. Welk GJ, Bai Y, Lee J-M et al. Standardizing analytic methods and reporting in activity monitor validation studies: guidelines to advance research and practice. *Med Sci Sport Exerc* 2019. <http://dx.doi.org/10.1249/MSS.0000000000001966>.
29. Turner CH, Robling AG. Designing exercise regimens to increase bone strength. *Exerc Sport Sci Rev* 2003; 31:45–50.
30. Rowlands AV, Cliff DP, Fairclough SJ et al. Moving forward with backwards compatibility: translating wrist accelerometer data. *Med Sci Sport Exerc* 2016; 48:2142–2149. <http://dx.doi.org/10.1249/MSS.0000000000001015>.