



Original article

A genetic variant in the cytochrome P450 family 2 subfamily R member 1 determines response to vitamin D supplementation

Afsane Bahrami ^{a, b, 1}, Mehrane Mehramiz ^{b, 1}, Majid Ghayour-Mobarhan ^{b, 1}, Hamidreza Bahrami-Taghanaki ^c, Kiana Sadeghi Ardekani ^b, Maryam Tayefi ^d, Mahsa Sadeghzade ^b, Mohammad Rashidmayvan ^b, Maryam Safari Ghalezou ^b, Gordon A. Ferns ^e, Amir Avan ^{b, **}, Hamid Reza Sadeghnia ^{f, *}

^a Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

^b Metabolic Syndrome Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

^c Complementary and Chinese Medicine, Persian and Complementary Medicine Faculty, Mashhad University of Medical Sciences, Mashhad, Iran

^d Clinical Research Unit, Mashhad University of Medical Sciences, Mashhad, Iran

^e Brighton & Sussex Medical School, Division of Medical Education, Falmer, Brighton, Sussex, BN1 9PH, UK

^f Pharmacological Research Center of Medicinal Plants, Mashhad University of Medical Sciences, Mashhad, Iran



ARTICLE INFO

Article history:

Received 26 November 2017

Accepted 31 March 2018

Keywords:

25(OH)D

Supplementation

CYP1R2

rs10766197

SUMMARY

Background: Globally, about 1 billion people have inadequate levels of serum vitamin D and it is prevalent in all ethnicities and age groups. Few foods naturally contain sufficient vitamin D; therefore, most people get their requirements through supplementation. Hence vitamin D status is affected by genetic and environmental determinants including season of measurement, diet habitual, health status, body mass index and concurrent medication. Further studies are necessary to understand how genetic variation influences vitamin D metabolism. We aimed to explore the association between a potential vitamin D-related polymorphism (the rs10766197 polymorphism in the CYP2R1 gene) with the response to supplementation of vitamin D in 253 healthy Iranian girls.

Material and method: A total of 253 healthy subjects received 50,000 IU of vitamin D3 weekly for 9 weeks. Serum 25(OH)D concentrations and metabolic profiles were measured at baseline and after 9 weeks of supplementation. The genotypes of the CYP2R1 variant (rs10766197) were identified using TaqMan genotyping assays.

Results: Serum 25(OH)D during the supplementation, increased in all individuals. Subjects with a AA major genotype at this locus had higher vitamin D concentrations after intervention (Changes (%) 448.4% ± 425% in AA vs 382.7% ± 301% in GG). This genetic variant modulated the response to supplementation ($p < 0.001$ and p -value SNP = 0.05). Regression analysis showed that the probability of affecting serum 25(OH)D, in individuals who had homozygous major allele GG was two-fold higher than carriers of the uncommon allele A (OR = 2.1 (1–4.2); $p = 0.03$). Interestingly, the Hs-CRP was reduced in AA carriers while was elevated in individuals with GG and AG genotypes, after high-dose vitamin D supplementation.

Conclusion: Changes in serum vitamin D and metabolic profile following high dose supplementation with vitamin D were associated with CYP2R1 polymorphism. Although carriers of the common G allele showed a greater response in the serum vitamin D.

© 2018 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

1. Introduction

Diet and other environmental factors such as the intake of vitamin D supplements and exposure to sunlight are known to influence serum vitamin D concentrations [1]. The assessment of serum 25-hydroxyvitamin D (25(OH)D) is the best biomarker of

* Corresponding author. Fax: +985138002287.

** Corresponding author. Fax: +985118002298.

E-mail addresses: avana@mums.ac.ir, amir_avan@yahoo.com (A. Avan), SadeghniaHR@mums.ac.ir (H.R. Sadeghnia).

¹ Contributed equally as first authors.

vitamin D status; however, the optimal serum concentration is unclear [2,3]. A study in the United States, has suggested that a serum 25(OH)D concentration of 50 nmol/L is sufficient for normal bone health in most individuals [4] whilst other studies have suggested that 60 nmol/L is necessary for reduction in the risks of falling and fractures risk [5,6]. Vitamin D has functions other than bone health. It is involved in the regulation of more than 2000 genes. Vitamin D deficiency may be associated with several non-skeletal diseases, including cancer [7], obesity [8], asthma [9], diabetes [10], cardiovascular diseases (CVD) [11] and metabolic syndrome (MS) [12] and has been reported as a major public health concern, even in regions with high levels of sunlight [13], for example it is common in the Middle East, India, Africa, Australia and South America [14–16].

In line with this, there is increasing evidence for a high prevalence of vitamin D deficiency in Iran; with reports of deficiency in >80% of the adolescence in Tehran and Arak [17,18], about 60% of school-age girls in Yazd [19] and >70% in newborn infants in Zanjan [20]. Few foods naturally contain enough vitamin D, the most natural way to get vitamin D is cutaneous production when skin is exposed to the sunlight [3]. Public concern about the high prevalence of vitamin D deficiency has caused increasing demand for supplementation and testing. Since individual responses to supplementation is variable, a more tailored approach to supplementation may be required. The variation in serum 25(OH)D level response after supplementation has been attributed to body mass [21], baseline serum 25(OH)D level [22], supplement dose [23], and the season [22]; however, there is also convincing evidence that vitamin D status is affected by genotype [24]. Several studies have reported polymorphisms in candidate genes associated with serum vitamin D that include CYP24A1 and CYP2R1 [25,26]. Each cytochrome P450 gene is known with CYP, implied that is part of the cytochrome P450 gene family. The common SNP, rs10766197, located in the promoter region of CYP2R1 gene, were reliable predictor of serum 25(OH)D levels [27].

The current study was carried out to examine whether treatment with high dose vitamin D supplementation is influenced by a variant in the CYP2R1 gene, using data obtained from a randomized controlled trial of vitamin D supplementation in healthy Iranian school-age girls of 12–18 years old; a group in which vitamin D deficiency is common.

2. Material and method

2.1. Study population

The 253 adolescent girls were recruited between January and April 2015 in Mashhad city, using a randomized cluster sampling method. Informed consent was collected from all participants using protocols approved by the Ethics Committee of the Mashhad University of Medical Sciences.

Participants with any chronic diseases history, or who were taking any kinds of dietary supplements and anti-depressant or psychotropic drugs were excluded from study.

Individuals with history of infectious disease, diabetes mellitus, family history of stroke, and myocardial infarction were excluded from study. Subjects received 50,000 IU vitamin D/week for 9 weeks. Serum 25(OH)D and metabolic profiles were measured at baseline and after 9 weeks.

2.2. Anthropometric and biochemical measurements

Anthropometric parameters (e.g., height, body weight, waist and hip circumference) were measured. BMI levels among teens

expressed relative to other children of the same sex and age. Percentiles were calculated using CDC growth charts (US Centers for Disease Control and Prevention (CDC) growth reference), which were based on national survey data collected from 1963 to 65 to 1988–94 [28]. Biochemical factors including serum calcium (Ca), and phosphate (P), fasting blood glucose (FBG), creatinine, blood urea nitrogen (BUN) and lipid profile; total cholesterol (TC), triglyceride (TG), and high-density lipoprotein cholesterol (HDL-C), measured by using commercial kits (Pars Azmun, Karaj, Iran) and the BT-3000 auto-analyzer (Biotechnica, Rome, Italy). Low-density lipoprotein cholesterol (LDL-C) was estimated using Friedewald formula if serum TG concentrations < 4.52 mmol/L [29–31]. High sensitivity C-reactive protein (Hs-CRP) was quantified using an immunoturbidimetry method, with limit of detection (LoD) 0.06 mg/L (Biosystems, Spain). Cut of value for Hs-CRP was <1.90 mg/L in 5–18 years woman according to the manufacturer's instructions.

An electrochemiluminescence method (ECL, Roche, Basel, Switzerland) was performed to measure serum 25-OH vitamin D. The LoD for the 25-OH vitamin D assay was 10 nmol/L for the ECL (Roche) and intra- and inter-assay variation were 5.7% and 9.9%, respectively.

2.3. DNA extraction and genotyping

Genomic DNA was extracted from blood samples using QIAamp[®] DNA Mini-Kit (Qiagen, San Diego, CA) according to the manufacturer's instructions. The purity and concentration of DNA samples were determined using the NanoDrop[®]-1000-Detector (NanoDrop-Technologies, Wilmington, USA). Genotype analysis of CYP2R1-rs10766197 polymorphism was carried out using TaqMan[®]-probes-based assay; PCR reactions were performed in 12.5 ml total volume, using 20 ng of DNA in TaqMan[®]n Universal MasterMix with specific primers and probes (Applied Biosystems Foster City, CA). To assess the allelic content. The ABI PRISM-7500 instrument equipped with the SDS version-2.0 software was used.

2.4. Statistics analysis

Data was analyzed using SPSS version 20, IBM (SPSS Inc., IL, USA). Variables are reported as mean \pm standard deviation (SD). Continuous variables were analyzed for normality using the Kolmogorov–Smirnov test. Analysis of variance (ANOVA) was performed to compare changes in biomarkers after intervention in different genotype groups. Post hoc analysis was done using Tukey's test. A Chi square test with continuity correction was used to determine whether genotype frequencies followed the Hardy–Weinberg Equilibrium. Repeated measures analysis of covariance (ANCOVA) was performed to investigate the effect of the genotypes. Logistic regression was performed to study the probability of change in serum 25(OH) D in the genetic dominant model. Significance was set at $p < 0.05$.

3. Results

3.1. Influences of supplementation on circulation 25(OH)D in CYP2R1 variant

In the total population of 253 healthy school-age Iranian girls, 88.1% suffered from vitamin D deficiency at baseline and only 4% of the total had a desirable vitamin D level. However, after intervention, 59.7% of the subjects were at a desirable concentration of 25(OH)D. About 20.2% of the subjects remained vitamin D deficient (Fig. 1). To examine the influence of CYP2R1 variant on the circulation levels of vitamin D after intervention, subjects were




Fig. 1. Comparison of the vitamin D status before and after 9 weeks of vitamin D supplementation. Deficiency: Serum 25(OH)D level <50 nmol/L. Sufficiency: 50 nmol/L < Serum 25(OH)D level <75 nmol/L. Desirable \geq 75 nmol/L [54].

categorized across rs10766197 genotype. The results revealed no significant trend in distribution of vitamin D status (desirable, sufficiency and deficiency) among different genotypes at baseline (P -trend = 0.4). However, supplementation for 9 weeks led to significant trend (P -trend = 0.05) (Table 1), with a reduction in the percent of subjects with a low serum vitamin D. The serum 25 (OH) D responses was dependent on the SNP in CYP1 (Fig. 2). During the supplementation, serum (OH) D increased in all groups, but carriers who had the common G allele, had higher vitamin D concentrations after 9 weeks of intervention. The SNP rs10766197 modulated response to vitamin D supplementation (p -value of intervention effect <0.001 and p -value SNP = 0.05) (Fig. 2). Regression analysis also indicated that the probability of altering serum 25(OH)D, in individuals who had homozygous major allele GG was two-fold higher than carriers of the uncommon A allele (OR = 2.1 (1–4.2); p value = 0.03). The regression model also was significant using a dominant model (OR = 1.8 (1–3.1); p value = 0.05) (Table 3). Data was adjusted for potential confounders such as age and BMI percentile.

3.2. Influence of supplementation on metabolic profile in CYP2R1 variant

Further analysis showed that fasting blood glucose and triglyceride concentration reduced in all subjects but carriers of a GG genotype showed a greater reduction in FBG and carriers of AA genotype showed a greater reduction in serum TG (Table 2). Interestingly, Hs-CRP was also reduced in AA carriers whilst the individuals with GG and AG genotypes, inflammation increased after 9 week of vitamin D supplementation (Table 2). Change in levels of Ca, BUN, creatinine and P after supplementation was not statistically significant among different genetic models (Table 2).

Table 1
Vitamin D status before and after 9 week of vitamin D supplementation according to CYP2R1 genotypes.

Vitamin D status (N = 253)	GG (N = 72)		AG (N = 119)		AA (N = 62)	
	Baseline	Follow-up	Baseline	Follow-up	Baseline	Follow-up
Desirable	4 (5.6)	50 (69.4)	2 (1.7)	70 (58.8)	4 (6.5)	31 (50)
Sufficiency	6 (8.3)	12 (16.7)	10 (8.4)	23 (19.3)	4 (6.5)	16 (25.8)
deficiency	62 (86.1)	10 (13.9)	107 (89.9)	26 (21.8)	54 (87.1)	15 (24.2)

Note: Σ 2 test showed a P_{trend} of 0.4 at baseline; P_{trend} at 12-month follow-up is 0.05. Data is presented as frequencies (%). Deficiency: Serum 25(OH)D level < 50 nmol/L. Sufficiency: Serum 25(OH) D level between 50 and 75 nmol/L. Desirable: Serum 25(OH)D level \geq 75 nmol/L.




Fig. 2. Serum 25(OH)D stratified by a polymorphism in CYP2R1 gene. Values are means \pm SD. Two-way ANCOVA repeated measures adjusted for multiple comparisons by Bonferroni test for serum 25(OH)D levels. Covariates used: age, BMI percentile and serum 25(OH)D at baseline.

4. Discussion

4.1. Influence of supplementation on circulation 25(OH)D in CYP2R1 variant

In the present study, we explored the association of rs10766197 of the CYP2R1 vitamin D-related gene with serum 25(OH)D concentrations and found that this polymorphism was significantly associated with the serum 25(OH)D concentrations after 9 weeks of vitamin supplementation and it appeared that carriers of dominant G allele were better responder to vitamin D in respect to elevation serum vitamin D. Animal and human studies have shown that different cytochrome P450 enzymes 2(CYP) including CYP2R1, CYP2D25, CYP3A4CYP27A1 are vitamin D 3 25-hydroxylases and cause 25-hydroxylation of vitamin D 3 and related metabolites [25]. Unlike others 25-hydroxylases, CYP2R1 hydroxylates both vitamin D 2 and vitamin D 3 [32]. Therefore, genetic variations including rs10766197, in the promoter region of this gene, may influence 25(OH)D synthesis. Our data indicated although this genotypic variant was not associated with baseline 25(OH)D level, it influenced on the response to the supplementation. It is possible that the regulation of 25(OH)D synthesized by skin might be different from supplementation. In agreement with our study, Nissen et al. examined variants in some vitamin D-related genes in 201 healthy Danish population. They reported a significant association between serum 25(OH)D and rs10766197. Similarly, in a study by Engelman et al. in a female population, all individuals who had no risk alleles of rs4588 and rs2060793, consuming about 670 IU/d vitamin D, the circulation level of 25(OH)D concentrations were at sufficient level (>50 nmol/L). For carriers with 1 and more risk alleles whose intakes were at least 670 IU/d vitamin D, only more than 50% of subjects had serum 25(OH)D > 50 nmol/L [26]. Thacher et al. in the cohort study on rickets Nigerian children, reported that individuals with a defective CYP2R1 allele had a mild form of VDDR1B and produce less 25(OH)D after intervention with vitamin D₂ or vitamin D₃. While, subjects who are homozygous for CYP2R1 mutations showed a severe form of VDDR1B and had minimal rise in serum

Table 2

Comparisons of the variables before and after 9 weeks of vitamin D supplementation in different genetic modes.

		GG	AG	AA	P value in different genetic models		
					Additive	Recessive	Dominant
BMI Percentile	Baseline	62 ± 56	58.4 ± 28	55.7 ± 26	N _s	N _s	N _s
	Follow-up	56 ± 30	54.6 ± 29	49 ± 27			
	Change	-17.6 ± 19	-12.4 ± 26.3	-15.9 ± 22.9			
Blood pressure SBP (mm Hg)	Baseline	101 ± 11	100.7 ± 13	99.6 ± 12	N _s	N _s	N _s
	Follow-up	99 ± 13	98.5 ± 13.6	100 ± 12			
	Change	-1 ± 15.5	-1.2 ± 15	2.1 ± 16			
DBP (mm Hg)	Baseline	67 ± 10	67.3 ± 10	67.1 ± 10.5	N _s	N _s	N _s
	Follow-up	65 ± 11	63 ± 10	64.1 ± 11			
	Change	-0.5 ± 14	-4.1 ± 12	-2 ± 9.1			
Serum fasted lipids Total Cholesterol (mg/dL)	Baseline	164.3 ± 26	165 ± 31	158 ± 28	N _s	N _s	N _s
	Follow-up	150.5 ± 24.5	156 ± 26.5	153 ± 27			
	Change	-7.4 ± 10.7	-4.6 ± 14.1	-0.1 ± 28			
TG (mg/dL)	Baseline	87.7 ± 41	79.5 ± 29	83 ± 33	0.03	N _s	0.03
	Follow-up	80.5 ± 40	81 ± 32	70.7 ± 26			
	Change	-0.8 ± 35.1	4.9 ± 29.6	-7.3 ± 27.1			
HDL (mg/dL)	Baseline	46.4 ± 8	48.9 ± 9	46.1 ± 8	N _s	N _s	N _s
	Follow-up	44.2 ± 8	46.3 ± 10	44.8 ± 7.5			
	Change	-3.2 ± 14.1	-2.5 ± 14.5	-1.7 ± 14.3			
LDL (mg/dL)	Baseline	100.7 ± 20	101.1 ± 27	99.4 ± 21	N _s	N _s	N _s
	Follow-up	89.1 ± 20	92.4 ± 22	93 ± 22			
	Change (%)	-11.9 ± 17	-9.1 ± 19	-7.1 ± 17.1			
Other blood parameters FBG	Baseline	90 ± 13	88.6 ± 10	86.6 ± 9	N _s	N _s	0.05
	Follow-up	87 ± 12	85 ± 12	85.7 ± 10.6			
	Change	-3 ± 10.8	-4.1 ± 11	-0.9 ± 13.4			
WBC(10 ⁹ /L)	Baseline	6.3 ± 1.8	6.06 ± 1.6	6.1 ± 1.5	N _s	N _s	N _s
	Follow-up	6.1 ± 1.6	6.1 ± 1.4	5.5 ± 1.3			
	Change	-1.8 ± 24.6	0.1 ± 27	-4.4 ± 28			
Serum Hs-CRP (mg/L)	Baseline	1.3 ± 1.6	1.1 ± 1.1	1.8 ± 1.7	N _s	0.003	0.05
	Follow-up	1.6 ± 2	1.4 ± 1.3	1.1 ± 1.4			
	Change	17.7 ± 13	61.6 ± 17.6	-26.8 ± 8.4			
25-OH vitamin D (nmol/L)	Baseline	27.5 ± 25	24 ± 18	29 ± 34	(AA vs. GG) 0.03 (AA vs. AG) 0.04	0.049	N _s
	Follow-up	99.3 ± 42	86 ± 40	84.4 ± 46			
	Change	447.3 ± 414.6	423.6 ± 380.4	433.0 ± 426.9			
Ca (mg/dL)	Baseline	9.6 ± 0.5	9.4 ± 0.5	9.4 ± 0.7	N _s	N _s	N _s
	Follow-up	9.7 ± 0.5	9.7 ± 0.5	9.7 ± 0.5			
	Change	0.1 ± 0.6	0.3 ± 0.57	0.3 ± 0.8			
Phosphate (mg/dL)	Baseline	4 ± 0.5	3.9 ± 0.4	3.9 ± 0.5	N _s	N _s	N _s
	Follow-up	4 ± 0.4	4 ± 0.4	4 ± 0.4			
	Change	0 ± 0.4	0.1 ± 0.4	0.1 ± 0.4			
Creatinine (mg/dL)	Baseline	10.6 ± 3.8	6.6 ± 11.5	5.6 ± 11.8	N _s	N _s	N _s
	Follow-up	0.7 ± 0.09	0.7 ± 0.1	0.7 ± 0.1			
	Change	8.7 ± 12.9	10.3 ± 15.2	8.0 ± 13			
BUN(mg/dL)	Baseline	12.3 ± 3	12.5 ± 3	12.6 ± 3	N _s	N _s	N _s
	Follow-up	13.8 ± 4	13.6 ± 3.2	14 ± 3			
	Change	16.3 ± 34.1	15.5 ± 35.3	14.4 ± 30.6			

Note: Change = ((Follow up – Baseline)/Baseline)/100; p values presented for the changes in different variables after vitamin D supplementation according to genotypes; Additive genetic model (GG genotype vs. AG genotype vs. AA genotype); Recessive genetic model (GG genotype vs. AG + AA genotypes); Dominant genetic model (GG + AG genotypes vs. AA genotype).

BMI: body mass index; TC: total cholesterol; TG: triglycerides; HDL-C: high-density lipoprotein cholesterol; LDL-C: low-density lipoprotein cholesterol; Hs-CRP: high-sensitivity Creative protein; FBG: fasting blood glucose; SBP: systolic blood pressure; DBP: diastolic blood pressure; BUN: Blood Urea nitrogen; Ca: Calcium; FBG: Fasting blood glucose.

A significance of bold values are $p < 0.05$.

Table 3

Association of CPY2R1 variant-rs10766197 with the changes in circulation levels of 25(OH)D after 9 weeks of supplementation (under different genetic models).

Model	Reference (Risk group)	AG	GG
Additive model	AA	OR (95% CI), p value	OR (95% CI)
	1	1.6 (0.8–3.3), 0.1	2.1 (1–4.2), 0.03
Recessive model	AG/AG	GG	
	Reference (Risk group)	OR (95% CI), p value	
Dominant model	1	1.5 (0.9–2.6), 0.1	
	AA	AG/GG	
	Reference (Risk group)	OR (95% CI), p value	
	1	1.8 (1.3–0.1), 0.05	

Data was adjusted for age, BMI percentile.

A significance of bold values are $p < 0.05$.

25(OH)D after administration of vitamin D, and improvement would be only with high doses of vitamin D [25,33]. In the study of Bu et al. they found that rs10741657 and rs10766197 were significantly associated with serum 25(OH)D concentrations in 496 healthy Caucasian people [34]. Based on similar results obtained from several studies on different population [26,35,36] it appeared that variants in the CYP2R1 gene predict serum 25(OH) D concentrations.

4.2. Influence of supplementation on fasted lipid profile and fasting blood glucose in CYP2R1 variant

We found that, an intake of 50,000 IU/D vitamin D per week had beneficial effects not only on 25(OH)D concentrations in all genotype groups but also on glycemic and lipid profile. However, these effects were greater in the subjects who had GG and AG genotypes at the rs10766197 locus. Noticeable that although carriers of the uncommon allele A, showed an increase in vitamin D concentration that was less than for other genotypes, the reduction in TG was more considerable. It was suggested that vitamin D has both direct and indirect effects on modifying the lipid profile.

An underlying mechanism on improving lipid profile may be through regulatory action of vitamin D in the stimulation of lipoprotein lipase [37] and reduced intestinal absorption and synthesis [38].

Cross-sectional studies have reported a negative relationship between circulation levels of 25(OH) D and serum Triglyceride. However, the influence of 25(OH) D on TG concentrations in interventional studies after supplementation with vitamin D is inconsistent [39]. Pittas et al. illustrated that in the individuals with impaired fasting glucose, administration of vitamin D and calcium might ameliorate insulin resistance [40]. Jorde et al. in a cross sectional studies examined 8018 non-smoking individuals, found a significant positive relationship between serum 25(OH)D and serum HDL-C, TC, and LDL-C and also a significant inverse associations between serum 25(OH)D with both LDL-C/HDL-C ratio and TG [41]. In an interventional study on 438 obese Norwegian, they found no statistical association between supplementation with vitamin D and lipid profile [42]. Similarly, Sieda et al. in a meta-analysis showed no significant improvement in glucose parameters [43]. These controversial illustrations might be partly attributed to the inherent limitations and heterogeneity of the studied cohorts. Some common factors may be influenced on both the high serum 25(OH)D levels and favorable lipid profile include exercise, diet habitual and genetic profile.

4.3. Influence of supplementation on inflammation in CYP2R1 variant

Our data revealed that inflammation increased in the carriers of dominant allele G after intervention by vitamin D supplementation while individuals who had AA genotype showed significant reduction in serum Hs-CRP after supplementation. It appeared that clinical outcome in response to vitamin D supplementation was genetic-related. Emerging evidence has reported a relationship between vitamin D supplementation and serum levels of proinflammatory and inflammatory markers such as cytokines and CRP. Some studies have reported a positive association with circulation levels of 25(OH)D and others showed an inverse association while some declared no relationship [44–47]. Vitamin D has been shown to suppress *in vitro* and *in vivo* the production of proinflammatory cytokines and modulate both the innate and adaptive immune systems [48,49]. It is proposed that macrophages, dendritic cells and activated lymphocytes influence on vitamin D receptor, implying a crucial role of vitamin D in the immune system [14].

Furthermore, the enzyme 25-hydroxyvitamin is produced by the immune system [50]. On the other hand, the activated vitamin D down-regulates proinflammatory mediators, such as interleukin (IL)-6, IL-8, tumor necrosis factor (TNF) α , and monocyte chemo-attractant protein (MCP)-1 [51,52]. However, it is suggested that while supplementation with vitamin D elevate the 25(OH)D, the conversion of inactive 25(OH)D to active 1,25(OH) D in the kidneys is not immediate, and may not be efficient. Both the inactive and active form of vitamin D bind to the vitamin D receptor (VDR), only the 1,25-D allow VDR to perform its functions beneficially and the 25(OH)D inhibits the VDR functions. Since VDR is the “gate-keeper” of the innate immune system and modulate by thousand genes so increased levels of 25-D might show immunosuppressive effects [53].

Generally, discrepancies in the different literature indicate the need for further studies both in healthy and disease population to find out more details about the potential association between serum levels of vitamin D and inflammation biomarkers. On the other hand, regulation of serum vitamin D in human body is a complex process that varies with individual genetic profiles and their health status. Research in genetic epidemiology of vitamin D is in its infancy and further comprehensive studies would be needed to understand how genetic variations modulate clinical outcomes of vitamin D supplementation.

5. Conclusion

We have found that although individuals with a GG genotype of CYP2R1 variant had a greater response to vitamin D supplements, the inflammation status was worsened. However, carriers of AA genotype showed less increase in 25(OH)D than others, but inflammation status only improved in this group. We conclude that personalized advice and recommendations tailored to individual's genetics seems help to determine how different individuals with various genetic background respond to the supplementation. People may need different health recommendations based on their genetic profiles, in order to elevate their serum 25(OH)D concentrations thereby avoiding adverse health outcomes.

Conflict of Interest

The authors have no conflict of interest to disclose.

Funding

This study was support by grants 941524 (Hamid R. Sadeghnia) from Mashhad University of Medical Sciences.

References

- [1] Khayyat-zadeh SS, Vatanparast H, Avan A, Bagherniya M, Bahrami A, Kiani MA, et al. Serum transaminase concentrations and the presence of irritable bowel syndrome are associated with serum 25-hydroxy vitamin D concentrations in adolescent girls who are overweight and obese. *Ann Nutr Metabol* 2017;71(3–4):234–41.
- [2] Wacker M, Holick MF. Vitamin D—effects on skeletal and extraskelatal health and the need for supplementation. *Nutrients* 2013;5(1):111–48.
- [3] Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metabol* 2011;96(7):1911–30.
- [4] Del Valle HB, Yaktine AL, Taylor CL, Ross AC. Dietary reference intakes for calcium and vitamin D. National Academies Press; 2011.
- [5] Bischoff-Ferrari HA, Willett WC, Orav EJ, Lips P, Meunier PJ, Lyons RA, et al. A pooled analysis of vitamin D dose requirements for fracture prevention. *N Engl J Med* 2012;367(1):40–9.
- [6] Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, Orav JE, Stuck A, Theiler R, et al. Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *BMJ* 2009;339:b3692.

- [7] Gandini S, Boniol M, Haukka J, Byrnes G, Cox B, Sneyd MJ, et al. Meta-analysis of observational studies of serum 25-hydroxyvitamin D levels and colorectal, breast and prostate cancer and colorectal adenoma. *Int J Cancer* 2011;128(6):1414–24.
- [8] Saliba W, Barnett-Griness O, Rennert G. The relationship between obesity and the increase in serum 25 (OH) D levels in response to vitamin D supplementation. *Osteoporos Int* 2013;24(4):1447–54.
- [9] Brown SD, Calvert HH, Fitzpatrick AM. Vitamin D and asthma. *Derm Endocrinol* 2012;4(2):137–45.
- [10] Sung C-C, Liao M-T, Lu K-C, Wu C-C. Role of vitamin D in insulin resistance. *BioMed Res Int* 2012;2012.
- [11] Kienreich K, Tomaschitz A, Verheyen N, Pieber T, Gaksch M, Gröbler MR, et al. Vitamin D and cardiovascular disease. *Nutrients* 2013;5(8):3005–21.
- [12] Weinstock-Guttman B, Mehta BK, Ramanathan M, Karmon Y, Henson LJ, Halper J, et al. Vitamin D and multiple sclerosis. *Neurol* 2012;18(4):179–83.
- [13] Muhairi SJ, Mehairi AE, Khouri AA, Naqbi MM, Maskari FA, Al Kaabi J, et al. Vitamin D deficiency among healthy adolescents in Al Ain, United Arab Emirates. *BMC Publ Health* 2013;13(1):33.
- [14] Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;2007(357):266–81.
- [15] Marwaha RK, Tandon N, Reddy DRH, Aggarwal R, Singh R, Sawhney RC, et al. Vitamin D and bone mineral density status of healthy schoolchildren in northern India. *Am J Clin Nutr* 2005;82(2):477–82.
- [16] Thacher TD, Fischer PR, Strand MA, Pettifor JM. Nutritional rickets around the world: causes and future directions. *Ann Trop Paediatr* 2006;26(1):1–16.
- [17] Neyestani TR, Hajifaraji M, Omidvar N, Eshraghian MR, Shariatzadeh N, Kalayi A, et al. High prevalence of vitamin D deficiency in school-age children in Tehran, 2008: a red alert. *Publ Health Nutr* 2012;15(2):324–30.
- [18] Talaei A, Yadegari N, Rafee M, Rezvanfar MR, Moini A. Prevalence and cut-off point of vitamin D deficiency among secondary students of Arak, Iran in 2010. *Indian J Endocrinol Metabol* 2012;16(5):786.
- [19] Shakiba M, Nafei Z, Lotfi MH, Shajari A. Prevalence of vitamin D deficiency among female students in secondary guidance school in Yazd City. *Acta Med Iran* 2009;47(3):209–14.
- [20] Kazemi A, Sharifi F, Jafari N, Mousavinasab N. High prevalence of vitamin D deficiency among pregnant women and their newborns in an Iranian population. *J Wom Health* 2009;18(6):835–9.
- [21] Didriksen A, Grimnes G, Hutchinson MS, Kjærgaard M, Svartberg J, Joakimsen RM, et al. The serum 25-hydroxyvitamin D response to vitamin D supplementation is related to genetic factors, BMI, and baseline levels. *Eur J Endocrinol* 2013;169(5):559–67.
- [22] Zhao L-J, Zhou Y, Bu F, Travers-Gustafson D, Ye A, Xu X, et al. Factors predicting vitamin D response variation in non-Hispanic white postmenopausal women. *J Clin Endocrinol Metabol* 2012;97(8):2699–705.
- [23] Chung M, Balk EM, Brendel M, Ip S, Lau J, Lee J, et al. Vitamin D and calcium: a systematic review of health outcomes. *Evid Rep Technol Assess (Full Rep)* 2009;183(183):1–420.
- [24] Bahrami A, Sadeghnia HR, Tabatabaieizadeh SA, Bahrami-Taghanaki H, Behboodi N, Esmaeili H, et al. Genetic and epigenetic factors influencing vitamin D status. *J Cell Physiol* 2017;233(5):4033–43.
- [25] Cheng JB, Levine MA, Bell NH, Mangelsdorf DJ, Russell DW. Genetic evidence that the human CYP2R1 enzyme is a key vitamin D 25-hydroxylase. *Proc Natl Acad Sci USA* 2004;101(20):7711–5.
- [26] Engelman CD, Meyers KJ, Iyengar SK, Liu Z, Karki CK, Igo RP, et al. Vitamin D intake and season modify the effects of the GC and CYP2R1 genes on 25-hydroxyvitamin D concentrations. *J Nutr* 2013;143(1):17–26.
- [27] Ahn J, Yu K, Stolzenberg-Solomon R, Simon KC, McCullough ML, Gallicchio L, et al. Genome-wide association study of circulating vitamin D levels. *Hum Mol Genet* 2010;19(13):2739–45.
- [28] Kuczmarski RJ, Ogden CL, Guo SS, Grummer-Strawn LM, Flegal KM, Mei Z, et al. 2000 CDC Growth Charts for the United States: methods and development. *Vital and health statistics. Series 11. Data Natl Health Surv* 2002;(246):1–190.
- [29] Bahrami A, Mazloum SR, Maghsoudi S, Soleimani D, Khayatizadeh SS, Arekhi S, et al. High dose vitamin D supplementation is associated with a reduction in depression score among adolescent girls: a nine-week follow-up study. *J Diet Suppl* 2017:1–10.
- [30] Tabatabaieizadeh SA, Avan A, Bahrami A, Khodashenas E, Esmaeili H, Ferns GA, et al. High-dose supplementation of vitamin D affects measures of systemic inflammation: reductions in High-Sensitivity C-Reactive Protein level and Neutrophil to lymphocyte ratio (NLR) distribution. *J Cell Biochem* 2017;118(12):4317–22.
- [31] Bahrami A, Avan A, Sadeghnia HR, Esmaeili H, Tayefi M, Ghasemi F, et al. High dose vitamin D supplementation can improve menstrual problems, dysmenorrhea, and premenstrual syndrome in adolescents. *Gynecol Endocrinol* 2018;1–5.
- [32] Cheng JB, Motola DL, Mangelsdorf DJ, Russell DW. De-orphanization of cytochrome P450 2R1 a microsomal vitamin D 25-hydroxylase. *J Biol Chem* 2003;278(39):38084–93.
- [33] Casella SJ, Reiner BJ, Chen TC, Holick MF, Harrison HE. A possible genetic defect in 25-hydroxylation as a cause of rickets. *J Pediatr* 1994;124(6):929–32.
- [34] Bu F-X, Armas L, Lappe J, Zhou Y, Gao G, Wang H-W, et al. Comprehensive association analysis of nine candidate genes with serum 25-hydroxy vitamin D levels among healthy Caucasian subjects. *Hum Genet* 2010;128(5):549–56.
- [35] Lasky-Su J, Lange N, Brehm JM, Damask A, Soto-Quiros M, Avila L, et al. Genome-wide association analysis of circulating vitamin D levels in children with asthma. *Hum Genet* 2012;131(9):1495–505.
- [36] Zhang Z, He JW, Fu WZ, Zhang CQ, Zhang ZL. An analysis of the association between the vitamin D pathway and serum 25-hydroxyvitamin D levels in a healthy Chinese population. *J Bone Miner Res* 2013;28(8):1784–92.
- [37] Wang J-H, Keisala T, Solakivi T, Minasyan A, Kalueff AV, Tuohimaa P. Serum cholesterol and expression of ApoAI, LXR β and SREBP2 in vitamin D receptor knock-out mice. *J Steroid Biochem Mol Biol* 2009;113(3):222–6.
- [38] Christensen R, Lorenzen JK, Svith CR, Bartels E, Melanson E, Saris W, et al. Effect of calcium from dairy and dietary supplements on faecal fat excretion: a meta-analysis of randomized controlled trials. *Obes Rev* 2009;10(4):475–86.
- [39] Jorde R, Grimnes G. Vitamin D and metabolic health with special reference to the effect of vitamin D on serum lipids. *Prog Lipid Res* 2011;50(4):303–12.
- [40] Pittas AG, Harris SS, Stark PC, Dawson-Hughes B. The effects of calcium and vitamin D supplementation on blood glucose and markers of inflammation in nondiabetic adults. *Diabetes Care* 2007;30(4):980–6.
- [41] Jorde R, Figenschau Y, Hutchinson M, Emaus N, Grimnes G. High serum 25-hydroxyvitamin D concentrations are associated with a favorable serum lipid profile. *Eur J Clin Nutr* 2010;64(12):1457–64.
- [42] Jorde R, Sneve M, Torjesen PA, Figenschau Y, Gøransson LG, Omdal R. No effect of supplementation with cholecalciferol on cytokines and markers of inflammation in overweight and obese subjects. *Cytokine* 2010;50(2):175–80.
- [43] Seida JC, Mitri J, Colmers IN, Majumdar SR, Davidson MB, Edwards AL, et al. Clinical review: effect of vitamin D3 supplementation on improving glucose homeostasis and preventing diabetes: a systematic review and meta-analysis. *J Clin Endocrinol Metabol* 2014;99(10):3551.
- [44] Barker T, Rogers VE, Levy M, Templeton J, Goldfine H, Schneider ED, et al. Supplemental vitamin D increases serum cytokines in those with initially low 25-hydroxyvitamin D: a randomized, double blind, placebo-controlled study. *Cytokine* 2015;71(2):132–8.
- [45] Peterson CA, Heffernan ME. Serum tumor necrosis factor- α concentrations are negatively correlated with serum 25 (OH) D concentrations in healthy women. *J Inflamm* 2008;5(1):10.
- [46] Yusupov E, Li-Ng M, Pollack S, Yeh JK, Mikhail M, Aloia JF. Vitamin D and serum cytokines in a randomized clinical trial. *Int J Endocrinol* 2010;2010.
- [47] García-Bailo B, Roke K, Mutch DM, El-Sohemy A, Badawi A. Association between circulating ascorbic acid, α -tocopherol, 25-hydroxyvitamin D, and plasma cytokine concentrations in young adults: a cross-sectional study. *Nutr Metabol* 2012;9(1):102.
- [48] McNally P, Coughlan C, Bergsson G, Doyle M, Taggart C, Adorini L, et al. Vitamin D receptor agonists inhibit pro-inflammatory cytokine production from the respiratory epithelium in cystic fibrosis. *J Cyst Fibros* 2011;10(6):428–34.
- [49] Yim S, Dhawan P, Ragunath C, Christakos S, Diamond G. Induction of cathelicidin in normal and CF bronchial epithelial cells by 1, 25-dihydroxyvitamin D 3. *J Cyst Fibros* 2007;6(6):403–10.
- [50] Lang P, Samaras N, Samaras D, Aspinall R. How important is vitamin D in preventing infections? *Osteoporos Int* 2013;24(5):1537–53.
- [51] Ding C, Wilding JP, Bing C. 1, 25-dihydroxyvitamin D3 protects against macrophage-induced activation of NF κ B and MAPK signalling and chemokine release in human adipocytes. *PLoS One* 2013;8(4):e61707.
- [52] Yin K, Agrawal DK. Vitamin D and inflammatory diseases. *J Inflamm Res* 2014;7:69.
- [53] Albert PJ, Proal AD, Marshall TG. Vitamin D: the alternative hypothesis. *Autoimmun Rev* 2009;8(8):639–44.
- [54] Wang X, Liu L, Li P, Ma J, Jiang R, Wang R, et al. Reference and influential factors of serum bone markers in Chinese adolescents. *Sci Rep* 2017;7(1):17340.