



“A National Study Demonstrating the Need for Improved Frailty Indices for Preoperative Risk Assessment of Common Urologic Procedures”

Katherine A. Amin, Una J. Lee, Chengshi Jin, John Boscardin, Andrew R. Medendorp, Jennifer T. Anger, and Anne M. Suskind

OBJECTIVE	To compare the associations between frailty indices and postoperative complications among older adults undergoing common urologic procedures. Frailty is known to be strongly associated with poor postoperative complications; however, the optimal way to measure frailty remains unknown.
METHODS	We identified the 20 most common urologic procedures from 2013-2016 in the American College of Surgeons National Surgical Quality Improvement Program (NSQIP) database. Frailty was measured using the NSQIP frailty index, simplified frailty index, and Risk Analysis Index. Multivariable logistic regression models were performed with each index and the American Society of Anesthesiologists (ASA) classification system with postoperative complications (any, major, or minor) as the outcomes. Statistical models were compared using the following fit parameters: area under the curve, Akaike information criterion, and Bayesian information criterion.
RESULTS	A total of 158,855 procedures were identified. All frailty indices (NSQIP frailty index, simplified frailty index, and Risk Analysis Index) and ASA were associated with increased odds for any, major, and minor complications (all P values $<.001$). ASA demonstrated stronger model fit parameters for any, major and minor complications compared to all other indices, with an area under the curve of 0.63, 0.64, and 0.64, respectively (all P values $<.001$). Adding ASA to each frailty index resulted in slight improvement of model fit parameters (P value $<.001$).
CONCLUSION	ASA slightly outperforms current frailty indices in predicting postoperative complications among individuals undergoing commonly performed urologic procedures. Our findings highlight the need for improved frailty measures for preoperative risk assessment. UROLOGY 132: 87–93, 2019. © 2019 Elsevier Inc.

More than two-thirds of all urologic procedures are performed in patients ≥ 65 years of age,¹ a large proportion of whom are frail.² Frailty is defined as a decline in physiologic reserve and function among older adults that is associated with an increased risk of poor health outcomes.²⁻⁶ Numerous studies using

data from the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) demonstrate that frailty is associated with increased rates of postoperative complications and decline in functional status, longer hospital stays, and higher rates of discharge to skilled or assisted nursing facilities.^{2,7-9}

Historically, the American Society of Anesthesiologists (ASA) classification system, which describes the overall health of a patient, has been used in literature to predict postoperative complications,^{10,11} however has been widely criticized for its subjectivity. Several frailty indices have been developed to objectively improve upon the ability to predict postoperative complications among older patients, however, there is a lack of consensus on which index to use. Many studies use the NSQIP frailty index (NSQIP-FI),^{2,12-16} which is a well-known 11-item cumulative deficits model developed from the 70-item questionnaire from the Canadian Study on Health and Aging. In response to recent changes in ACS-NSQIP variables and

Financial Disclosures: Pepper Center Research Career Development Core (RCDC), NIH-NIDDK K12 DK83021-09; K12 Urologic Research (KURe) Career Development Program.

Conflict of Interest: 1. Jennifer T. Anger, M.D.: Expert witness for Boston scientific; investigator for Renovia. 2. Anne Suskind:

Funding: NIH-NIA R01AG058616-01.

From the Section of Urology and Renal Transplantation, Virginia Mason Medical Center, Seattle, WA; the Department of Epidemiology and Biostatistics, University of California San Francisco (UCSF), San Francisco, CA; the Department of Surgery, Division of Urology, Tripler Army Medical Center, HI; the Department of Surgery, Division of Urology, Cedars-Sinai Medical Center, Los Angeles, CA; and the Department of Urology, University of California San Francisco (UCSF), San Francisco, CA

Address correspondence to: Katherine Amin, M.D., 1100 Ninth Ave., C7-URO, Seattle, WA 98101. E-mail: katherine.a.amin@gmail.com

Submitted: March 30, 2019, accepted (with revisions): July 5, 2019

a desire to simplify frailty measurement,^{17,18} other modified frailty indices have been developed including the simplified Frailty Index (sFI)^{8,19} and the Risk Analysis Index (RAI).⁷ The comparative predictive ability of each of these indices to determine postoperative complications, however, remains unknown.

Using data from ACS-NSQIP from 2013 to 2016, we examined and compared the associations of each tool (NSQIP-FI, sFI, RAI, and ASA) with 30-day postoperative complications among older adults undergoing common urologic procedures. Findings from this study will be helpful to determine which tools should be used to evaluate postoperative complications using ACS-NSQIP and similar data sources among individuals undergoing common urologic procedures.

MATERIALS AND METHODS

Patients and Database

This is a retrospective cohort study of men and women undergoing urologic surgery from 2013 to 2016 using the ACS-NSQIP Participant Data Use File. The ACS-NSQIP database is a prospectively maintained national database of validated data collected from trained Surgical Clinic Reviewers. Surgical Clinic Reviewers examine medical records and classify perioperative clinical data through 30-day postoperative outcomes across a sample of surgical procedures of randomly assigned patients.^{20,21} As of 2017 the Participant Use Data File contained 1,028,713 cases that were submitted by 708 participating sites nationwide.²⁰ From this database, we identified all urologic procedures using the urology surgical specialty variable and included the 20 most commonly performed procedures (Appendix A). Given the fact that female sling procedures are performed by both urologists and gynecologists, this particular procedure (Current Procedure Terminology [CPT-4] code 57288: sling operation for stress incontinence, eg fascia or synthetic) included both specialty types. Common urologic procedures were grouped together to ensure adequate power for predicting risk of complications for each frailty index. This study was determined to be exempt by our institutional review board.

Outcomes

Data on 30-day complications were obtained from the ACS-NSQIP database. Complications were classified as either minor (eg urinary tract infection [UTI], superficial wound infection, pneumonia, or blood transfusion) or major (eg hospital readmission, return to the operating room, sepsis, deep vein thrombosis [DVT], cerebral vascular accident, reintubation, renal failure, myocardial infarction, pulmonary embolus, dehiscence, cardiac arrest, deep wound infection, coma, or death) in accordance with previous studies examining complications using these data.^{2,7,8,12,15-16}

Covariates

Frailty was assessed using several indices previously designed and validated specifically for ACS-NSQIP data including (1) NSQIP-FI, (2) sFI, and (3) RAI. The NSQIP-FI is a cumulative deficits model and includes 11 items in total: history of diabetes mellitus; impaired functional status; history of chronic obstructive pulmonary disease, or pneumonia; history of congestive heart failure (CHF); history of myocardial infarction within

6 months of surgery; history of percutaneous coronary intervention, cardiac surgery, or angina; use of antihypertensive medications; peripheral vascular disease or rest pain; impaired sensorium; transient ischemic attack or cerebral vascular accident. Items are summed and divided by the total number of items (11) for the composite NSQIP-FI score, whereby a NSQIP-FI score of 1 out of 11 would be 0.09, a NSQIP-FI score of 2 out of 11 would be 0.18, etc. We categorized patients into groups based on corresponding NSQIP-FI score (0, 0.09, 0.18, and 0.27+). This index was selected because it has been shown to have excellent correlation with both mortality and morbidity in all surgical subspecialties, including urology, and distinctly measures frailty, as opposed to comorbidity.^{12,16}

The sFI was derived from the 5 most common components of NSQIP-FI and was first introduced to improve the practical usefulness of NSQIP-FI.⁸ The 5 variables included in this index are history of diabetes mellitus, functional status, history of chronic obstructive pulmonary disease, history of CHF, and hypertension requiring treatment. To calculate sFI, the presence of each variable is attributed 1 point, with a maximum score of 5 points possible.⁸ We classified patients into 3 categories, similar to a previously published study,¹⁹ denoted as a score of 0, 1, and ≥ 2 due to the nature of the compressed scale and reduced frequency of patients with a score ≥ 3 .

The RAI was adapted from the Minimum Data Set Mortality Risk Index-Revised and is based on clinically focused variables including gender, age, diagnosis of cancer, history of weight loss, renal failure, CHF, dyspnea, cognitive deterioration, residence other than living independently, and functional status.⁷ Each variable has weighted numerical value and the RAI is calculated using a scoring system with a range of 0-81.⁷ We categorized patients from our cohort into 4 groups: (1) RAI score of 0-15, (2) score of 16-25, (3) score of 26-35, and (4) score of ≥ 36 similar to distributions from previously published literature.⁷

The ASA classification system, while not a frailty index, was first introduced clinically in 1941 as a method to describe a patient's underlying systemic illnesses prior to undergoing a procedure with an anesthesia professional.^{10,11} The ASA scale assigns increasing order of risk from 1-6. In order to appropriately compare frailty indices, we grouped patients as ASA 1 (normal, healthy), 2 (mild systemic disease), 3 (severe systemic disease), or 4+ ([4] severe systemic disease, a constant threat to life,⁵ moribund, not expected to survive without the operation, or⁶ declared brain-dead for donor organ harvest).

Additional covariates extracted from the ACS-NSQIP database included patient characteristics (age, gender, race, BMI, and smoking status) and procedure characteristics (surgical setting [inpatient vs outpatient], type of anesthesia [general, monitored anesthesia care, or other], and calendar year of operation).

Statistical Analysis

Patient- and procedure-level characteristics as well as frailty index and ASA scores of patients were reported as frequencies and percentages. Unadjusted and adjusted logistic regression models were created to assess predictive ability of any, major, and minor postoperative complications. Models were adjusted for age, gender, race, BMI, and smoking status and separate models were created for each index individually (eg, NSQIP-FI, sFI, RAI, and ASA). To determine the combined effect of ASA with each frailty index, we performed separate multivariable logistic regression analyses for each frailty index in combination with ASA. Quality of statistical models were estimated and compared by performing model selection methods including area

under the curve (AUC), Akaike information criterion (AIC), and Bayesian information criterion (BIC). AUC should be interpreted such that a value of 0.5 reflects an accuracy of chance and a model with perfect accuracy has an AUC of 1.0. AIC estimates the quality of each model relative to other models, whereby a lower value suggests a better model. Similarly, BIC is used as a criterion for model selection and the model with the lowest BIC is preferred. All analyses were performed with SAS, version 9.4. All *P* values were 2-sided and 0.05 was set as the level of statistical significance.

RESULTS

Based on inclusion criteria, 158,855 relevant urologic procedures were identified in the ACS-NSQIP data set between 2013 and 2016. **Table 1** outlines patient and procedure-level characteristics, in addition to frailty and ASA scores for our cohort. The cohort mean age was 64.6 ± 13.2 years, mean BMI was 29.2 ± 6.0 kg/m², 22.9% were female, and 16.0% were smokers. Fifty-four percent of cases were performed in the inpatient setting and 91.0% were done under general anesthesia. Mean NSQIP-FI score was 0.073 ± SD 0.07, while 16.9% and 1.8% had a NSQIP-FI score of 0.18 and ≥ 0.27, respectively. Mean sFI score was 0.8 ± 0.2 and 18.8% of the cohort had a sFI score of ≥ 2. Mean RAI score was 7.4 ± SD 4.3 and 4.55% of the cohort had an RAI score of >15. Finally, mean ASA was 2.5 ± SD 0.7 and 3.9% of the cohort had an ASA of ≥ 4. **Appendix B** shows complication rates by each individual complication within the patient cohort. The rate of major complications was 8.2% and the rate of minor complications was 8.6%. The most common major complication was readmission (5.8%) followed by return to the operating room (2.0%) and the most common minor complication was blood transfusion (4.8%) followed by UTI (3.1%).

Logistic regression models for each frailty index and ASA (adjusted for age, gender, race, BMI, and smoking status) are shown in **Table 2**. These models demonstrate that increasing severity of frailty (for each frailty index) and increasing ASA score correlated with increased odds for any, major, and minor complications (all *P* values <0.001) (**Table 2**). NSQIP-FI score of 0.27+ was associated with major complications (adjusted odd ratio [aOR] 3.1, 95% confidence interval [CI] 2.8-3.4). In comparison, a sFI score of ≥ 2 was associated with an aOR of 1.9 (95% CI 1.8-2.0) and an RAI score of ≥ 36 was associated with an aOR of 6.5 (95% CI 4.1-10.5) for risk of major complications respectively. However, an ASA of 4+ had the highest association of major complications with an aOR of 7.2 (95% CI 5.9-8.8). **Table 3** summarizes the model fit parameters for each of the frailty indices and ASA. ASA demonstrated stronger model fit parameters for any, major, and minor complications (compared to all of the frailty indices) with an AUC of 0.63, 0.64, and 0.64, respectively. In comparison, NSQIP-FI and sFI both had the lowest AUC values of 0.59, 0.60, and 0.59 for any, major and minor complications, while RAI had AUC values of 0.59, 0.61, and 0.59 for any, major and minor complications. Values for AIC and BIC showed similar trends.

Because ASA demonstrated the strongest fit parameters for postoperative complications in our models, we decided to model ASA combined with each frailty index to determine whether or not this combination improved model fit. **Table 4** summarizes these findings and shows that there was slight improvement of all fit parameters (AUC, AIC, and BIC) when ASA was combined with each frailty index. For example,

Table 1. Patient characteristics, procedure characteristics, and frailty index/ASA scores of patients undergoing common urologic procedures

<i>Patient Characteristics</i>		
Age (y), frequency (%)		
Age (Mean ± SD (N))	64.6 ± 13.2 (N = 158,855)	
<65	73,157	(46.1)
65-74	49,934	(31.4)
75-84	27,127	(17.%)
≥ 85	8637	(5.4)
Gender, frequency (%)		
Male	122400	(77.1)
Female	36455	(22.9)
Race, frequency (%)		
White	111589	(86.8)
Black	11602	(9.0)
Other	5368	(4.2)
BMI (kg/m ²), frequency, (%)		
BMI (Mean ± SD (N))	29.2 ± 6.0 (N = 157,457)	
Underweight (<18.5)	1507	(1.0)
Normal (18.5-24.9)	35,777	(22.7)
Overweight (25-29.9)	60,775	(38.6)
Obese (≥ 30.0)	59,398	(37.7)
Smoking status, frequency (%)		
Smoker	25,474	(16.0)
<i>Procedure Characteristics</i>		
Surgical setting, frequency (%)		
Inpatient	85,800	(54.0)
Outpatient	73,055	(46.0)
Type of anesthesia, frequency (%)		
General	144,597	(91.0)
Monitored Anesthesia care (MAC)	4092	(2.6)
Other	10,125	(6.4)
Year of operation, frequency (%)		
2013	30,512	(19.2)
2014	34,578	(21.8)
2015	43,207	(27.2)
2016	50,537	(31.8)
<i>Frailty index and ASA scores</i>		
NSQIP-FI score, frequency (%)		
NSQIP-FI score (Mean ± SD (N))	0.07 ± 0.07 (N = 158,855)	
0	63,737	(40.1)
0.09	65,391	(41.2)
0.18	26,797	(16.9)
≥ 0.27	2930	(1.8)
sFI score, frequency (%)		
sFI score (Mean ± SD (N))	0.8 ± 0.2 (N = 158,855)	
0	63,737	(40.1)
1+	65,391	(41.2)
≥ 2	29,727	(18.8)
RAI score, frequency (%)		
RAI Score (Mean ± SD [N])	7.4 ± 4.3 (N=158,855)	
0-15	151,688	(95.5)
16-25	6183	(3.9)
26-35	898	(0.6)
≥36	86	(0.05)
ASA, frequency (%)		
ASA (Mean ± SD (N))	2.5 ± 0.7 (N = 158,492)	
1	7757	(4.9)
2	71,545	(45.1)
3	72,996	(46.1)
≥ 4	6194	(3.9)

Table 2. Summary of logistic regression models for each frailty index and ASA demonstrating odds of any complications, major and minor complications*

	Any Complication				Major Complications				Minor Complications			
	Odds Ratio	Lower 95% CI	Upper 95% CI	P Value	Odds Ratio	Lower 95% CI	Upper 95% CI	P Value	Odds Ratio	Lower 95% CI	Upper 95% CI	P Value
NSQIP-FI												
0.00	1.0				1.0				1.0			
0.09	1.4	1.3	1.4	<.001	1.4	1.3	1.4	<.001	1.4	1.3	1.4	<.001
0.18	1.7	1.6	1.8	<.001	1.8	1.7	1.9	<.001	1.7	1.6	1.8	<.001
≥ 0.27	2.8	2.5	3.1	<.001	3.1	2.8	3.4	<.001	2.4	2.2	2.7	<.001
sFI												
0	1.0				1.0				1.0			
1	1.4	1.3	1.4	<.001	1.4	1.3	1.4	<.001	1.4	1.3	1.4	<.001
≥ 2	1.8	1.7	1.9	<.001	1.9	1.8	2.0	<.001	1.8	1.7	1.9	<.001
RAI												
0-15	1.0				1.0				1.0			
16-25	2.8	2.6	3.0	<.001	2.7	2.5	2.9	<.001	2.8	2.6	3.0	<.001
26-35	4.0	3.4	4.7	<.001	4.2	3.5	5.1	<.001	3.5	2.9	4.2	<.001
≥ 36	8.7	5.4	14.2	<.001	6.5	4.1	10.5	<.001	6.2	3.9	10.0	<.001
ASA												
1	1.0				1.0				1.0			
2	2.0	1.7	2.3	<.001	1.9	1.5	2.3	<.001	2.1	1.8	2.6	<.001
3	4.0	3.5	4.6	<.001	3.7	3.1	4.5	<.001	4.6	3.8	5.6	<.001
≥ 4	7.9	6.8	9.2	<.001	7.2	5.9	8.8	<.001	9.0	7.4	11.0	<.001

Models are adjusted for age, gender, race, BMI, and smoking status.

*Each model includes age, BMI, smoking status, recent weight loss as independent variables.

NSQIP-FI alone had an AUC of 0.59, whereas, the combination of the NSQIP-FI and ASA demonstrated an AUC of 0.63, *P* value <0.001 comparing the 2 models. This trend was consistent when ASA was combined with each frailty index. The combination of RAI and ASA was the strongest predictor of odds of any, major, and minor complications with AUCs of 0.64, 0.65, and 0.64, respectively.

COMMENT

This national study highlights the strong association between increased frailty and postoperative complications among older adults undergoing common urologic procedures. However, our findings suggest that the current tools commonly used to quantify frailty (NSQIP-FI, sFI, and RAI) are relatively poor, as shown by our statistical models demonstrating model fit parameters. Interestingly, we found that ASA was marginally superior to all frailty indices. Additionally, there was modest improvement of the accuracy to predict postoperative complications when fit parameters were measured for each frailty index in combination with ASA.

Other studies that examined specific subsets of urologic patients from the ACS-NSQIP database have previously compared fit parameters between different frailty indices. One study compared NSQIP-FI, sFI, and ASA to evaluate the predictive ability of sFI as a risk assessment tool in patients who underwent radical cystectomy. This study compared AUCs between NSQIP-FI, sFI, and ASA for models predicting Clavien Grade III-V complications and demonstrated slightly higher AUCs for the frailty indices (AUCs for NSQIP-FI and sFI were both 0.56) compared to ASA (AUC of 0.544).⁸ This finding differs from our study,

which found the reverse to be true, whereby AIC (and other fit parameters) demonstrated that ASA slightly outperformed the frailty indices in predicting complications. This difference is likely due to the contrasting patient populations evaluated in each study. Cystectomy patients represent only 4.2% of our study population and are likely to be among the most frail individuals compared to patients undergoing other common urologic procedures included in our cohort. Regardless of the frailty of the patient population studied, however, differences between ASA and frailty indices in predicting postoperative complications are relatively small, highlighting the point that there may not be much difference between the each type of instrument.

Another study showed the validity of RAI predicting postoperative outcomes among surgical patients at a Veterans Affairs institution. However, the NSQIP-FI and RAI models also demonstrated weak predictability of postoperative mortality and prompted authors to call on future efforts for determining an optimal preoperative frailty assessment.⁷ This finding is similar to our study such that NSQIP-FI and RAI fitness models demonstrating the poor quality of these tools as predictors for postoperative complications among older patients.

While other reports have demonstrated the association between higher ASA class and worsened perioperative morbidity,^{22,23} the utility of ASA as a prognostic metric for postoperative complications across surgical subspecialties remains poor. Particularly in select patient populations (ie, vascular surgery patients or very elderly patients), the ASA classification system has been shown to have limited predictive utility because these patients are most often confined to ASA grade III or IV as a result of having several comorbid conditions. These studies

Table 3. Model fit parameters of frailty indices and ASA predicting odds of complications (adjusted for independent variables: age, gender, race, BMI, and smoking status)*

	Any Complication				Major Complication				Minor Complication			
	NSQIP-FI	sFI	RAI	ASA	NSQIP-FI	sFI	RAI	ASA	NSQIP-FI	sFI	RAI	ASA
AUC	0.59	0.59	0.59	0.63	0.60	0.60	0.61	0.64	0.59	0.59	0.59	0.64
AIC	102237.45	102329.89	101482.98	100316.60	71648.81	71736.22	71324.64	70628.42	74233.37	74261.16	73738.19	72670.54
BIC	102374.10	102456.77	101619.62	100453.22	71785.45	71863.10	71461.28	70765.04	74370.01	74388.04	73874.83	72807.16

*All P values were <0.001 when each index was compared to ASA for any complication, major complication, minor complication.

Table 4. Summary of regression models predicting odds of complications with ASA added to the each frailty index*

	Any Complication			Major Complications			Minor Complications		
	NSQIP-FI & ASA	sFI & ASA	RAI & ASA	NSQIP-FI & ASA	sFI & ASA	RAI & ASA	NSQIP-FI & ASA	sFI & ASA	RAI & ASA
AUC	0.63	0.63	0.64	0.64	0.64	0.65	0.64	0.64	0.64
AIC	100191.00	100233.68	99330.60	70492.18	70538.74	70034.37	72625.90	72631.98	72125.74
BIC	100356.90	100389.82	99496.50	70658.07	70694.87	70200.27	72791.79	72788.12	72291.64

*all P values were <0.001 when each index was compared in combination with ASA.

have argued to distinguish such patients with the addition of an independent tool to assess functional capacity in order to improve operative risk evaluation.²⁴⁻²⁶ Furthermore, numerous articles have explored and quantified interobserver reliability among anesthesia trained physicians^{27,28} and reduced inter-rater variability between anesthesia trained physicians compared to other clinicians, particularly in patients with higher ASA scores.^{29,30} Overall, despite the slight advantage of ASA compared to frailty indices as shown in our study, there continues to be a lack of a high-quality and accurate method to predict an older patient's ability to withstand the demands of surgery.

The findings of this study should be considered with certain limitations in mind. First, due to the nature of the ACS-NSQIP database, the variables that may be utilized to define frailty are limited and we recognize that such variables may not truly capture what it means to be frail. Second, the ACS-NSQIP database is limited by a short time-frame of 30-day postoperative follow-up and the absence of long-term complications. Due to the nature of this database, specific complications may be under-represented, particularly postoperative UTI's. While this may be true, 30-day complications have become an important metric to report for evaluating quality of care and carry significant value in the current healthcare landscape.

CONCLUSION

While several tools to measure frailty exist, they remain relatively poor predictors of any, major, and minor complications following common urologic procedures. ASA, often thought of as an historic and subjective tool describing the overall health of a patient, also showed relatively low utility as a predictive tool but had slight advantage compared to frailty indices (eg, NSQIP-FI, sFI, and RAI). Our findings highlight the need for improved frailty assessment tools for preoperative risk stratification for older adults undergoing common urologic procedures.

SUPPLEMENTARY MATERIALS

Supplementary material associated with this article can be found in the online version at <https://doi.org/10.1016/j.urology.2019.07.001>.

References

1. Drach GW, Griebing TL. Geriatric urology. *J Am Geriatr Soc.* 2003;51(7 Suppl):S355-S358.
2. Suskind AM, Walter LC, Jin C, et al. Impact of frailty on complications in patients undergoing common urological procedures: a study from the American College of Surgeons National Surgical Quality Improvement database. *BJU Int.* 2016;117:836-842. <https://doi.org/10.1111/bju.13399>.
3. Robinson TN, Wu DS, Pointer L, Dunn CL, Cleveland JC, Moss M. Simple frailty score predicts postoperative complications across surgical specialties. *Am J Surg.* 2013;206:544-550. <https://doi.org/10.1016/j.amjsurg.2013.03.012>.
4. Robinson TN, Wu DS, Stigmann GV, Moss M. Frailty predicts increased hospital and six-month healthcare cost following colorectal surgery in older adults. *Am J Surg.* 2011;202:511-514. <https://doi.org/10.1016/j.amjsurg.2011.06.017>.
5. Partridge JSL, Fuller M, Harari D, et al. Frailty and poor functional status are common in arterial vascular surgical patients and affect postoperative outcomes. *Int J Surg.* 2015;18:57-63. <https://doi.org/10.1016/j.ijso.2015.04.037>.
6. Makary MA, Segev DL, Pronovost PJ, et al. Frailty as a predictor of surgical outcomes in older patients. *J Am Coll Surg.* 2010;210:901-908. <https://doi.org/10.1016/j.jamcollsurg.2010.01.028>.
7. Hall DE, Arya S, Schmid KK, et al. Development and initial validation of the risk analysis index for measuring frailty in surgical populations. *JAMA Surg.* 2017;152:175-182. <https://doi.org/10.1001/jamasurg.2016.4202>.
8. Sathianathen NJ, Jarosek S, Lawrentschuk N, Bolton D, Konety BR. A simplified frailty index to predict outcomes after radical cystectomy. *Eur Urol Focus.* 2018. <https://doi.org/10.1016/j.euf.2017.12.011>.
9. Isharwal S, Johanning JM, Dwyer JG, Schmid KK, LaGrange CA. Preoperative frailty predicts postoperative complications and mortality in urology patients. *World J Urol.* 2017;35:21-26. <https://doi.org/10.1007/s00345-016-1845-z>.
10. Saklad M. Grading of patients for surgical procedures. *Anesthesiology.* 1941;2:281-284.
11. Dripps RD. New classification of physical status. *Anesthesiology.* 1963;24:111.
12. Suskind AM, Jin C, Walter LC, Finlayson E. Frailty and the role of obliterative versus reconstructive surgery for pelvic organ prolapse: a national study. *J Urol.* 2017;197:1502-1506. <https://doi.org/10.1016/j.juro.2016.12.001>.
13. Levy I, Finkelstein M, Bilal KH, Palese M. Modified frailty index associated with Clavien-Dindo IV complications in robot-assisted radical prostatectomies: a retrospective study. *Urol Oncol.* 2017;35:425-431. <https://doi.org/10.1016/j.urolonc.2017.01.005>.
14. Etzioni DA, Wasif N, Dueck AC, et al. Association of hospital participation in a surgical outcomes monitoring program with inpatient complications and mortality. *JAMA.* 2015;313:505-511. <https://doi.org/10.1001/jama.2015.90>.
15. Molina CS, Thakore RV, Blumer A, Obremskey WT, Sethi MK. Use of the National Surgical Quality Improvement Program in orthopaedic surgery. *Clin Orthop.* 2015;473:1574-1581. <https://doi.org/10.1007/s11999-014-3597-7>.
16. Velanovich V, Antoine H, Swartz A, Peters D, Rubinfeld I. Accumulating deficits model of frailty and postoperative mortality and morbidity: its application to a national database. *J Surg Res.* 2013;183:104-110. <https://doi.org/10.1016/j.jss.2013.01.021>.
17. Liu Y, Cohen ME, Hall BL, Ko CY, Bilimoria KY. Evaluation and enhancement of calibration in the American College of Surgeons NSQIP surgical risk calculator. *J Am Coll Surg.* 2016;223:231-239. <https://doi.org/10.1016/j.jamcollsurg.2016.03.040>.
18. Subramaniam S, Aalberg JJ, Soriano RP, Divino CM. New 5-factor modified frailty index using American College of Surgeons NSQIP Data. *J Am Coll Surg.* 2018;226:173-181.e8. <https://doi.org/10.1016/j.jamcollsurg.2017.11.005>.
19. Chimukangara M, Helm MC, Frelich MJ, et al. A 5-item frailty index based on NSQIP data correlates with outcomes following paraesophageal hernia repair. *Surg Endosc.* 2017;31:2509-2519. <https://doi.org/10.1007/s00464-016-5253-7>.
20. Am Coll Surg Natl Qual Improv Program ACS NSQIP Particip Use Data File. 2017. <https://www.facs.org/quality-programs/acs-nsqip/participant-use>.
21. Shiloach M, Frencher SK, Steeger JE, et al. Toward robust information: data quality and inter-rater reliability in the American College of Surgeons National Surgical Quality Improvement Program. *J Am Coll Surg.* 2010;210:6-16. <https://doi.org/10.1016/j.jamcollsurg.2009.09.031>.
22. Menke H, Klein A, John KD, Junginger T. Predictive value of ASA classification for the assessment of the perioperative risk. *Int Surg.* 1993;78:266-270.

23. Hackett NJ, De Oliveira GS, Jain UK, Kim JYS. ASA class is a reliable independent predictor of medical complications and mortality following surgery. *Int J Surg Lond Engl*. 2015;18:184–190. <https://doi.org/10.1016/j.ijvsu.2015.04.079>.
24. Dosluoglu HH, Wang J, Defranks-Anain L, Rainstein M, Nader ND. A simple subclassification of American Society of Anesthesiology III patients undergoing peripheral revascularization based on functional capacity. *J Vasc Surg*. 2008;47:766–772. <https://doi.org/10.1016/j.jvs.2007.11.053>. discussion 722-723.
25. Crawford RS, Cambria RP, Abularrage CJ, et al. Preoperative functional status predicts perioperative outcomes after infrainguinal bypass surgery. *J Vasc Surg*. 2010;51:351–358. <https://doi.org/10.1016/j.jvs.2009.08.065>. discussion 358-359.
26. Visnjevac O, Davari-Farid S, Lee J, et al. The effect of adding functional classification to ASA status for predicting 30-day mortality. *Anesth Analg*. 2015;121:110–116. <https://doi.org/10.1213/ANE.0000000000000740>.
27. Mak PHK, Campbell RCH, Irwin MG, American Society of Anesthesiologists. The ASA Physical Status Classification: inter-observer consistency. *American Society of Anesthesiologists. Anaesth Intensive Care*. 2002;30:633–640.
28. Sankar A, Johnson SR, Beattie WS, Tait G, Wijeyesundera DN. Reliability of the American Society of Anesthesiologists physical status scale in clinical practice. *Br J Anaesth*. 2014;113:424–432. <https://doi.org/10.1093/bja/aeu100>.
29. Curatolo C, Goldberg A, Maerz D, Lin H-M, Shah H, Trinh M. ASA physical status assignment by non-anesthesia providers: do surgeons consistently downgrade the ASA score preoperatively? *J Clin Anesth*. 2017;38:123–128. <https://doi.org/10.1016/j.jclinane.2017.02.002>.
30. Hurwitz EE, Simon M, Vinta SR, et al. Adding examples to the ASA-physical status classification improves correct assignment to patients. *Anesthesiology*. 2017;126:614–622. <https://doi.org/10.1097/ALN.0000000000001541>.