



Correction to: A retrospective analysis of weight changes in HIV-positive patients switching from a tenofovir disoproxil fumarate (TDF)- to a tenofovir alafenamide fumarate (TAF)-containing treatment regimen in one German university hospital in 2015–2017

Mario Gomez¹ · Ulrich Seybold¹ · Julia Roider¹ · Georg Härter² · Johannes R. Bogner¹

Published online: 19 November 2018
© The Author(s) 2018

Correction to: Infection
<https://doi.org/10.1007/s15010-018-1227-0>

The original version of this article unfortunately contained mistakes. The presentation of Fig. 1 was incorrect and the Acknowledgements were missing. Please find the Acknowledgement here:

Acknowledgement

This work was partly supported by DZIF project TI 02.001.

The corrected figure is given below.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

The original article can be found online at <https://doi.org/10.1007/s15010-018-1227-0>.

✉ Mario Gomez
mariogomezcamacho@gmail.com

¹ Sektion Klinische Infektiologie, Medizinische Klinik und Poliklinik IV, Klinikum der Universität, Ludwig-Maximilians-Universität München, Pettenkoferstrasse 8a, 80336 Munich, Germany

² Medicovert Ulm MVZ, Münsterplatz 6, 89073 Ulm, Germany

Fig. 1 Mean change in weight in percent \pm standard deviation through 360 days for TDF and TAF separately; **a** results of patients after switch to TAF ($n=129$) and **b** results of patients receiving TDF; this arm comprises 241 patients, since switch patients initially received TDF (pooled data from the switch group and control group)

