

Alcohol Expectancy and Cerebral Responses to Cue-Elicited Craving in Adult Nondependent Drinkers

Simon Zhornitsky, Sheng Zhang, Jaime S. Ide, Herta H. Chao, Wuyi Wang, Thang M. Le, Robert F. Leeman, Jinbo Bi, John H. Krystal, and Chiang-shan R. Li

ABSTRACT

BACKGROUND: Positive alcohol expectancy (AE) contributes to excessive drinking. Many imaging studies have examined cerebral responses to alcohol cues and how these regional processes related to problem drinking. However, it remains unclear how AE relates to cue response and whether AE mediates the relationship between cue response and problem drinking.

METHODS: A total of 61 nondependent drinkers were assessed with the Alcohol Expectancy Questionnaire and Alcohol Use Disorder Identification Test and underwent functional magnetic resonance imaging while exposed to alcohol and neutral cues. Imaging data were processed and analyzed with published routines, and mediation analyses were conducted to examine the interrelationships among global positive score of the Alcohol Expectancy Questionnaire, Alcohol Use Disorder Identification Test score, and regional responses to alcohol versus neutral cues.

RESULTS: Alcohol as compared with neutral cues engaged the occipital, retrosplenial, and medial orbitofrontal cortex as well as the left caudate head and red nucleus. The bilateral thalamus showed a significant correlation in cue response and in left superior frontal cortical connectivity with global positive score in a linear regression. Mediation analyses showed that global positive score completely mediated the relationship between thalamic cue activity as well as superior frontal cortical connectivity and Alcohol Use Disorder Identification Test score. The alternative models that AE contributed to problem drinking and, in turn, thalamic cue activity and connectivity were not supported.

CONCLUSIONS: The findings suggest an important role of the thalamic responses to alcohol cues in contributing to AE and at-risk drinking in nondependent drinkers. AEs may reflect a top-down modulation of the thalamic processing of alcohol cues, influencing the pattern of alcohol use.

Keywords: Alcohol, Craving, Cue, Expectancy, fMRI, Thalamus

<https://doi.org/10.1016/j.bpsc.2018.11.012>

Along with an impaired ability to control the urges to drink, craving is a hallmark of alcohol abuse and dependence (1). It is well known that alcohol-related cues evoke craving and expectations of positive outcomes contributed to drinking (2,3). Alcohol expectancy (AE) represents subjective beliefs about the extent to which drinking will lead to particular outcomes (e.g., positive expectancy would be associated with statements such as “drinking makes me feel good; alcohol makes me worry less”) (4). According to the outcome expectancy model of craving, expectancies can be divided into informational and motivational components (5). The former represents specific beliefs or expectancies about alcohol's effects, whereas the latter reflects the yearning for those effects. For example, seeing one's friends drink may, along with AE, generate anticipation that alcohol will produce relaxation, pleasure, or relief from withdrawal and lead to the desire to experience those feelings (6). The desire, in turn, triggers urges to drink and precipitates alcohol consumption (5). Thus, AE

may interact with environmental cues to contribute to at-risk alcohol use.

AE is an important moderator of problem drinking. For example, AE accounted for a significant proportion of the variance in drinking-related measures (4). AE discriminated between adolescent non-problem drinkers and those subsequently engaged in problem drinking (7). AE related to habitual consumption of alcohol among problem and non-problem adult drinkers (8), with higher expectancies associated with increased levels of consumption. Problem drinkers as compared with non-problem drinkers reported significantly higher AE from adolescence through middle adulthood (9). Expectancies about alcohol-enhancing social behaviors were particularly relevant to close-friend alcohol use and consequences in college students (10). In an alcohol self-administration study, high responders reported heavier drinking patterns and lower expectancies for negative consequences (11). In a treatment study, lower

expectancies of alcohol-produced relaxation were related to abstinence during a 1-year period (8).

Numerous studies have examined the effects and neural processes of environmentally triggered craving. Following administration of a nonalcoholic lager, participants reported craving in relation to how much they liked and felt stimulated by the drink (12). In an imaging study of 326 heavy drinkers, alcohol compared with neutral taste cues evoked greater activation in the dorsal striatum, insula, orbitofrontal cortex (OFC), anterior cingulate cortex (ACC), and ventral tegmental area, with activation in the dorsal striatum, insula, and precuneus in correlation with alcohol use severity (13). Olfactory alcohol cues elicited craving along with increased activation in the nucleus accumbens and ventral tegmental area among heavy drinkers, in contrast to control participants (14). In another study, alcohol-related visual cues activated the ventral striatum, OFC, and other structures in the medial prefrontal cortex (PFC) in alcohol-dependent individuals as compared with healthy subjects (15). Across cue modalities, a meta-analysis of 28 functional magnetic resonance imaging studies showed robust activation of limbic prefrontal regions, including the ACC and ventromedial PFC, in 679 cases of heavy/dependent drinkers (16). When compared with control participants, case participants exhibited more activation in the parietal and temporal regions, including the posterior cingulate, precuneus, and superior temporal gyrus. In region of interest (ROI) analyses that interrogated only limbic regions, cue-elicited activation of the ventral striatum was most frequently correlated with drinking measures reduced by treatment (16). Taken together, the studies have suggested that cue-elicited craving is associated with activation in brain regions that support reward and incentive salience and that individuals with alcohol misuse show more cue-evoked activation in these regions.

Expectation to have access to alcohol may influence craving and related psychological states. For instance, in social drinkers craving was increased both after receiving alcohol and after receiving placebo (albeit to a lesser extent) but not after receiving a nonalcoholic drink (17). Importantly, alcohol approach tendencies were more pronounced after both alcohol and placebo compared with the control beverage, with no difference between alcohol and placebo. In another study of social drinkers, alcohol urge and other subjective states were measured before and after an initial drink (alcohol, placebo, or nonalcohol) was consumed (12). Both alcohol and placebo produced increased sedated feelings, and after placebo, urge was positively related to liking and enjoying the “alcoholic” drinks and feeling more stimulated. A number of studies have specifically examined how expectation to drink or smoke modulated cue-elicited brain responses. For instance, the expectation of receiving an alcoholic drink enhanced activation in the ACC and other prefrontal regions among social drinkers performing a working memory task (18). Healthy individuals with a positive family history of alcoholism showed enhanced striatal dopamine release during expectation of alcohol (19). A study of cigarette smokers demonstrated cue-elicited activations of limbic and prefrontal structures in individuals expecting to smoke immediately after the scan but not in those not allowed to smoke despite similar levels of craving (20). A stepwise linear regression analysis revealed a correlation

between smoking cue-induced craving scores and activation in the PFC differentially modulated by the state of expectation. Another study similarly demonstrated cue-evoked ventromedial, ventrolateral, and dorsolateral prefrontal cortical activation that was modulated by the option to smoke (21). Together, the studies suggest that cue-elicited activation of the limbic prefrontal striatal circuits depends on the subjective awareness of drug accessibility.

On the other hand, the literature is limited with regard to the interactions between AE and craving. Although “expectation” and “expectancy” were used interchangeably in some studies, unlike expectation to access alcohol, AE reflects one’s belief and knowledge of positive outcomes of alcohol use and interacts with environmental primer to precipitate craving and alcohol consumption. AE was associated with increases in craving following administration of a placebo drink of chilled lemonade served in a vodka-rimmed glass (22). AE has been associated with ACC activation during a vigilance task in adolescents (23). An earlier electroencephalographic study suggested frontal but not parietal electroencephalographic power as a predictor of AE, although the prefrontal neuropsychological performance was associated with AE less than consistently across testing batteries (24). A previous structural imaging study showed that AE best predicted problem drinking in women and interacted with impulsivity to predict problem drinking in men, each in association with decreased gray matter volume of the right posterior insula and the left thalamus (25). More recently, we demonstrated how thalamic subregional functional connectivities were interrelated with AE and at-risk alcohol use in nondependent drinkers (26). However, no imaging studies have addressed the potential influence of AE on cue-induced craving or the interrelationships among AE, cue-elicited brain response, and at-risk drinking.

Here, we examined the relationship between AE and cerebral responses in a cue-elicited craving functional magnetic resonance imaging paradigm in 61 adult nondependent drinkers. We hypothesized that brain responses to alcohol cues would be modulated by AE and the extent of risky alcohol use and explored the relationships between the brain activity and connectivity, AE, and the severity of at-risk drinking.

METHODS AND MATERIALS

Subjects and Assessments

Potential candidates were recruited from the greater New Haven, Connecticut, area and were screened according to the Structured Clinical Interview for DSM-IV (27). A total of 61 nondependent adult drinkers met eligibility requirements and participated in this study (Table 1). All subjects were physically healthy with no major medical illnesses or current use of prescription medications. None reported having a history of head injury or neurological illness. Other exclusion criteria included current or past dependence on a psychoactive substance (except nicotine) and current or history of Axis I disorders according to the Structured Clinical Interview for DSM-IV (27). The Human Investigation Committee at the Yale University School of Medicine approved all study procedures, and all subjects signed an informed consent form prior to participation.

Table 1. Demographics and Drinking Measures of Male and Female Participants

Subject Characteristic	Men (<i>n</i> = 33)	Women (<i>n</i> = 28)	<i>p</i> Value ^a
Age, Years	30.8 ± 8.1	30.4 ± 8.9	.82
AUDIT Score	11.3 ± 11.3	9.6 ± 9.1	.53
Duration of Alcohol Use, Years	12.6 ± 8.0	12.9 ± 9.6	.91
Number of Drinking Days per Month, Prior Year	8.2 ± 6.0	10.9 ± 5.0	.07
Number of Drinks per Occasion	3.8 ± 2.6	3.4 ± 2.3	.50
Number of Drinks per Month, Prior Year	38 ± 45.4	41.2 ± 40.9	.78
Alcohol Expectancy GP Score	13.3 ± 6.0	14.8 ± 6.1	.33
FTND Score	0.42 ± 1.4	1.25 ± 2.6	.12
Current Smoker, Yes/No	6/27	9/19	.21

Values are mean ± SD or *n*.

AUDIT, Alcohol Use Disorder Identification Test; FTND, Fagerström Test for Nicotine Dependence; GP, global positive subscore of the Alcohol Expectancy Questionnaire.

^aTwo-tailed two-sample *t* test except for smoker status, which used chi-square test.

All participants were assessed with the Alcohol Use Disorders Identification Test (AUDIT) (28), which has been widely used to examine alcohol use behavior and alcohol-related problems. Participants were also assessed with the Alcohol Expectancy Questionnaire (29). The Alcohol Expectancy Questionnaire consists of 40 items to address both positive AE (six subscales) and negative AE (two subscales). Each subscale contains four to six statements that can be endorsed on a 6-point scale, from “disagree strongly” (1) to “agree strongly” (6). The global positive (GP) subscale contains five items and thus ranges from 5 to 30 in total score, with a greater score indicating higher GP AE. Although the expectancy sub-components are statistically discernible, the high subscale intercorrelations (ranging from $r = .42$ to $r = .92$, mean = .78) suggest that the degree of distinctiveness among the subscales is at best modest (29). Thus, in the current study, we focused on the GP subscore as a variable to quantify individual variation in AE. Participants were also assessed with the Fagerström Test for Nicotine Dependence (30) and averaged 0.8 ± 2.0 (mean ± SD) in Fagerström Test for Nicotine Dependence score, suggesting low dependence.

Behavioral Task

We employed a cue-induced alcohol craving task. In alternating blocks, participants viewed alcohol-related or neutral pictures and reported alcohol craving. Briefly, a cross appeared on the screen to engage attention at the beginning of each block. After 2 seconds, six pictures displaying alcohol-related cues (alcohol block) or neutral visual scenes (neutral block) were shown for 6 seconds each. Participants were asked to look at the pictures and think about how they may relate to the scenes. The pictures were collected from the internet and independently reviewed by two investigators. Alcohol pictures included images of alcoholic drinks, people holding or drinking alcoholic beverages, and bar scenes,

whereas neutral pictures comprised natural sceneries. At the end of each block, participants were asked to report how much they craved alcohol on a visual analog scale from 0 (no craving) to 10 (highest craving ever experienced). Each block lasted about 45 seconds (including time for craving rating), and a total of six alcohol and six neutral blocks took approximately 9 minutes to complete. Each participant completed two runs of the task.

Imaging Protocol and Data Preprocessing

The imaging protocol is described in detail in the Supplement. Data were analyzed with SPM following established routines (31,32), as in the Supplement.

Imaging Data Modeling

Alcohol and neutral cue blocks were first distinguished. A statistical analytical block design was constructed for each individual subject using a general linear model (GLM), with block onsets convolved with a canonical hemodynamic response function and with the temporal derivative of the canonical hemodynamic response function and entered as regressors in the model. Because each block was associated with a craving rating, we included a column of block onset parametrically modulated by its corresponding craving score as a regressor in the model. Realignment parameters in all six dimensions were also entered in the model. Serial autocorrelation caused by aliased cardiovascular and respiratory effects was corrected by a first-degree autoregressive model. The GLM estimated the component of variance that could be explained by each of the regressors.

In the first-level analysis, we constructed for each individual subject statistical contrasts of alcohol picture versus neutral picture. These contrasts allowed us to evaluate brain regions that responded differently to viewing of alcohol and neutral pictures. The contrast (difference in beta) images of the first-level analysis were then used for the second-level group statistics (random-effects analysis). Following current reporting standards, all imaging results were evaluated with voxel $p < .001$, uncorrected, in combination with cluster $p < .05$, familywise error corrected, on the basis of Gaussian random field theory, as implemented in SPM.

In ROI analysis, we used MarsBaR (<http://marsbar.sourceforge.net/>) to derive for each individual subject the activity (beta contrast) for the ROIs. Functional ROIs were defined based on clusters obtained from whole-brain analysis. All voxel activations were presented in Montreal Neurological Institute coordinates.

General Psychophysiological Interaction

To explore circuit activities, we examined psychophysiological interaction (PPI) of the ROIs with differential response to alcohol versus neutral cues. PPI describes functional connectivity between brain regions contingent on a psychological context. We used a generalized form of context-dependent PPI (general PPI [gPPI]; <http://brainmap.wisc.edu/PPI>) (33). Briefly, in gPPI the hemodynamic responses to alcohol picture and neutral picture formed the psychological regressors, whereas in conventional PPI only alcohol picture > neutral picture would be included in the GLM. The inclusion of task regressors

in gPPI reduces the likelihood that the functional connectivity estimates were driven by simple coactivation. The extracted mean time series of the blood oxygen level-dependent signals were temporally filtered, mean corrected, and deconvolved to generate the signal time series of the ROIs for each subject to compose the physiological variable. These time series were then multiplied by the onset times of the alcohol picture and neutral picture separately and reconvolved with the canonical hemodynamic response function to obtain the interaction term or PPI variable. Finally, the psychological regressors of alcohol picture and neutral picture, physiological variable of the ROIs, and PPI variables of alcohol picture and neutral picture were entered as regressors in a whole-brain GLM. gPPI analysis was performed for each individual subject, and the resulting contrast images were used in random-effects group analysis. Likewise, the results were evaluated at voxel $p < .001$, uncorrected, in combination with cluster $p < .05$, familywise error corrected, according to current reporting standards.

Mediation Analysis

Owing to space limitations, mediation analyses are presented in the [Supplement](#).

RESULTS

Cue-Induced Craving and Regional Activations to Alcohol Cue Exposure

Alcohol, as compared with neutral cue, elicited a higher craving rating (3.1 ± 2.5 vs. 1.9 ± 2.1 , $p < .0001$, paired t test). Alcohol- but not neutral cue-elicited craving was also correlated positively with GP score across subjects ($r = .43$, $p < .0006$ and $r = .16$, $p = .2141$, respectively).

In examining regional responses to alcohol versus neutral cues, we first conducted a two-sample t test to compare men and women. Even at a relaxed threshold at voxel $p < .005$, uncorrected, no clusters showed a significant sex difference. Thus, men and women were combined in the analysis. Exposure to alcohol as compared with neutral cues engaged higher activation of cortical and subcortical structures, including the occipital cortex, medial OFC, retrosplenial cortex/parieto-occipital sulcus, left caudate head, and a cluster in the midbrain predominantly in the area of the red nucleus. Conversely, neutral as compared with alcohol cues involved higher activation in the superior parietal gyrus/cuneus, supra-marginal gyrus, and posterior cingulate gyrus ([Figure 1](#)). These clusters are summarized in [Table 2](#). We extracted the beta contrast of alcohol versus neutral cue response for each of these clusters, and none showed a significant correlation with craving rating during the alcohol block (all $ps > .24$) or with difference in craving rating between the alcohol and neutral blocks (all $ps > .13$).

Cue Reactivity in Relation to AE and Problem Drinking

As with the analyses of regional responses to alcohol versus neutral cues, we compared men and women in voxelwise regression against GP score and observed no sex differences at voxel $p < .005$, uncorrected. With men and women combined and in a whole-brain linear regression of alcohol versus

neutral cue exposure against GP score for all subjects with age as a covariate, bilateral thalamus ($x = 12$, $y = -13$, $z = 13$, volume = 1971 mm^3 , $Z = 3.43$; $x = -15$, $y = -19$, $z = 10$, volume = 1458 mm^3 , $Z = 3.29$), in the area of the pulvinar and medial dorsal nucleus, showed activation in positive correlation with GP score ([Figure 2A](#)). The analyses with sex as an additional covariate identified essentially the same clusters: $x = 12$, $y = -13$, $z = 13$, $Z = 3.37$, volume = 1944 mm^3 ; $x = -15$, $y = -19$, $z = 10$, $Z = 3.34$, volume = 1404 mm^3 . We extracted the beta contrast of thalamic activation to alcohol versus neutral cue for individual subjects. [Figure 2B, C](#) shows the linear regression of GP and AUDIT scores against cue-elicited thalamic activity for men and women separately. In a slope test, we observed that the correlation between thalamic cue activation with GP score and that with AUDIT score did not differ between men and women ($p = .95$ and $p = .44$, respectively) ([34](#)).

Psychophysiological Interaction

We used bilateral thalamus clusters as a seed region in gPPI analysis. The results showed a number of cortical and subcortical regions with higher interaction with the thalamus during alcohol versus neutral cue blocks ([Figure 3](#) and [Table 3](#)). Of these nine clusters, we examined whether any of these regional interactions correlated with the GP and AUDIT scores. Because GP and AUDIT scores were highly correlated, we corrected for the number of clusters with a $p = .05/9 = .0056$ in examining the results. The gPPI magnitude of the cluster located at the left superior frontal gyrus (SFG)/superior frontal sulcus showed a positive correlation with both the GP score ($r = .38$, $p = .0023$) and AUDIT score ($r = .39$, $p = .0022$) ([Figure 4A](#)). A slope test showed no difference between men and women in the regression of the gPPI against GP score ($p = .79$) or AUDIT score ($p = .27$) ([Figure 4B, C](#)).

Mediation Analysis

With mediation analysis, we further examined the interrelationship among thalamic activation to alcohol (vs. neutral) cue exposure, AE (GP score), and problem drinking (AUDIT score). AUDIT score was highly correlated with GP score in men and women combined ($r = .5934$, $p = 4.66 \times 10^{-7}$), and thalamic activity during alcohol versus neutral cue exposure was correlated with GP score ($r = .4218$, $p = .0007$) and with AUDIT score ($r = .2708$, $p = .03476$). However, the interrelationships between these neural and clinical measures remained open. We performed mediation analyses to test two specific hypotheses, namely that 1) thalamic activity contributed to higher AE and, in turn, problem drinking, and 2) higher AE led to problem drinking and, in turn, altered thalamic activity during cue exposure. The results showed that GP score completely mediated the correlation between the thalamic response to alcohol versus neutral cue and AUDIT score in men and women combined ([Figure 5A](#)). The alternative models where AUDIT score mediated the correlation between GP score and thalamic activity were not supported ([Figure 5B](#)). Likewise, we conducted mediation analyses on thalamic connectivity with the SFG, GP score, and AUDIT score. The results showed that GP score completely mediated the correlation between the gPPI strength and AUDIT score in men and women combined




Figure 1. Regional activations to alcohol (A) vs. neutral (N) cues at $p < .001$, uncorrected. All clusters with cluster $p < .05$, familywise error corrected, are shown in Table 2. CN, caudate nucleus; L, left; mOFC, medial orbitofrontal cortex; OC, occipital cortex; PCG, posterior cingulate gyrus; R, right; RN, red nucleus; RSC/POS, retrosplenial cortex/parieto-occipital sulcus; SMG, supramarginal gyrus; SPG/Cu, superior parietal gyrus/cuneus.

Table 2. Regional Activations to Alcohol vs. Neutral Cue Exposure

Volume (mm ³)	Peak Voxel (Z)	MNI Coordinates (mm)			Side	Identified Brain Region
		x	y	z		
Alcohol > Neutral Cues						
19,494 ^a	6.66	21	−91	4	R	Superior/Middle occipital gyrus/Cuneus
20,412 ^a	6.48	−24	−85	−5	L	Superior/Middle occipital gyrus/Cuneus
7263 ^a	5.42	−12	−43	13	L/R	Retrosplenial cortex/Parieto-occipital sulcus
5481 ^a	4.77	−9	44	−11	L/R	Medial orbitofrontal cortex
3699	4.08	−6	59	31	L/R	Dorsomedial prefrontal cortex
3807	4.01	−3	−16	−8	L/R	Red nucleus
3402	4.00	−12	20	7	L	Caudate nucleus
Neutral > Alcohol Cues						
7857 ^a	4.69	9	−79	31	L/R	Superior parietal gyrus/Cuneus
4617	4.33	−60	−28	13	L	Superior temporal gyrus
5643	4.14	57	−40	37	R	Supramarginal gyrus
3780	4.05	15	−34	46	L/R	Posterior cingulate gyrus

Voxel $p < .001$, uncorrected, and cluster level $p < .05$, familywise error corrected.

L, left; MNI, Montreal Neurological Institute; R, right.

^aClusters with voxel peak meeting $p < .05$, familywise error corrected.

(Figure 5C). The alternative models where AUDIT score mediated the correlation between GP score and connectivity strength were not supported (Figure 5D).

DISCUSSION

We identified regional activations in response to alcohol versus neutral cues in the occipital, retrosplenial, and medial OFC as well as in the left caudate head and red nucleus, in accordance with earlier imaging studies of cue-related responses (35) and reward-related responses (36,37). Although not showing higher responses to alcohol versus neutral cues, the bilateral thalamus demonstrated a positive correlation in cue response with GP score and AUDIT score in a linear regression across participants. Psychophysiological interaction analyses showed higher thalamic connectivity with a number of cortical and subcortical structures, including the left SFG during cue exposures. Thalamic SFG connectivity was also correlated with

both GP and AUDIT scores. Furthermore, mediation analyses showed that GP score completely mediated the relationship between thalamic cue activity as well as thalamic SFG connectivity and AUDIT score. These findings suggested that AE was reflected in thalamic cue responses and a potentially unique role of cue-elicited thalamic responses in supporting the influence of the expectancy of positive alcohol effects on drinking behavior.

Comprising subnuclei with distinct anatomical connections that relay and integrate information between cortical and subcortical structures, the thalamus is instrumental in supporting multiple cognitive and affective processes (38,39). For example, the medial dorsal nucleus responds to reward anticipation (40,41) and mediates working memory and executive control, which are often compromised following excessive alcohol consumption (42,43). The anterior thalamic nucleus is part of the Papez circuit and supports episodic memory and emotional expression. Deficits in episodic




Figure 2. (A) Bilateral thalamus showing regional activations to alcohol > neutral cues in correlation with global positive alcohol expectancy (AE) score in all subjects at voxel $p < .001$, uncorrected, in combination with cluster $p < .05$, familywise error corrected. (B) Regression of thalamic activity (beta contrast: alcohol > neutral cue) against AE score separately for men (M) ($p = .05$, $r = .37$) and women (W) ($p = .007$, $r = .46$) and for all subjects ($p = .0007$, $r = .42$). A slope test showed no difference between men and women ($p = .95$). (C) Regression of thalamic activity (alcohol > neutral cue) against Alcohol Use Disorder Identification Test (AUDIT) score separately for men ($p = .05$, $r = .34$) and women ($p = .43$, $r = .15$) and for all subjects ($p = .03$, $r = .27$). A slope test showed no difference between men and women ($p = .44$).




Figure 3. Brain regions showing a higher psychophysiological interaction with the bilateral thalamus during alcohol (A) vs. neutral (N) cue blocks (warm color) and during neutral vs. alcohol cue blocks (cool color) at $p < .001$, uncorrected. Clusters meeting cluster $p < .05$, familywise error corrected, are summarized in Table 3. AG, angular gyrus; CB, cerebellum; HG, hippocampal gyrus; MTG, middle temporal gyrus; OC, occipital cortex; PCL/PMC, paracentral lobule/pre-motor cortex; PCu, precuneus; SFS, superior frontal sulcus; Th/CN/Pa, thalamus/caudate nucleus/pallidum.

Table 3. Brain Regions Showing Psychophysiological Interaction With Bilateral Thalamus During Exposure to Alcohol vs. Neutral Cue

Volume (mm ³)	Peak Voxel (Z)	MNI Coordinates (mm)			Side	Identified Brain Region
		x	y	z		
Positive						
32,184	5.79	24	−88	−11	R	Occipital cortex/cerebellum
	5.42	−24	−88	−11	L	Occipital cortex/cerebellum
2511 ^a	4.87	33	−16	−20	R	Hippocampus
1593 ^a	4.85	−54	−52	−11	L	Middle temporal gyrus
8154 ^a	4.82	−6	−31	55	L/R	Paracentral lobule/PMC
1620 ^a	4.81	−3	−61	19	L/R	Precuneus
7398 ^a	4.75	−6	−7	10	L/R	Thalamus/caudate/pallidum
1782 ^a	4.75	−27	11	55	L	Superior frontal sulcus
2457 ^a	4.68	−9	−46	−14	L/R	Cerebellum
3672 ^a	4.59	−36	−70	37	L	Occipital cortex
Negative						
None						

Voxel $p < .001$, uncorrected, and cluster level $p < .05$, familywise error corrected.

L, left; MNI, Montreal Neurological Institute; PMC, premotor cortex; R, right.

^aClusters with voxel peak $p < .05$, familywise error corrected.

and emotional memory are key manifestations of the Wernicke–Korsakoff syndrome in alcohol-addicted individuals (43,44). The pulvinar supports attention and cross-modal integration of information (45). There is a substantial literature of thalamic dysfunction in alcohol misuse, with studies reporting both increased (46,47) and decreased (40,44,48) thalamic activity and connectivity in drinkers relative to nondrinkers.

Drug cue reactivity is known to be a psychologically complex process and would likely engage the thalamus. On the other hand, imaging studies of cue reactivity did not typically implicate the thalamus (49). As shown in a meta-analytic review, alcohol cue exposure most consistently engaged the ventromedial PFC, posterior cingulate cortex, and striatum, although ventral striatal activations were reported largely in studies that interrogated only the limbic circuits with ROI

analysis (16). There was substantial intra- and interstudy variability in brain responses to drug cues, suggesting that cue reactivity is amenable to modulation by a variety of experimental variables. In an earlier study of more than 300 participants, authors reported robust thalamic response to alcohol versus litchi juice drinks (13), suggesting that gustatory stimulation may have more powerful effects on the thalamus than visual stimulation. Furthermore, of the clinical variables that influenced cue-elicited brain responses, length of use and addiction severity appeared to influence activities of the thalamus, amygdala, and dorsal ACC, among other regions of the mesolimbic circuit (49).

Here, as with the majority of imaging studies, we did not observe increased thalamic activation during exposure to pictorial alcohol versus neutral cues. However, bilateral thalamus clusters exhibited higher response to alcohol versus




Figure 4. (A) Left superior frontal sulcus/gyrus showing a higher psychophysiological interaction with the bilateral thalamus during alcohol vs. neutral cue blocks at voxel $p < .001$, uncorrected, in combination with cluster $p < .05$, familywise error corrected. (B) Regression of the general psychophysiological interaction (gPPI) magnitude against global positive alcohol expectancy (AE) score separately for men (M) ($p = .01$, $r = .42$) and women (W) ($p = .11$, $r = .30$) and for all subjects ($p = .0023$, $r = .38$). A slope test showed no difference between men and women ($p = .79$). (C) Regression of the gPPI magnitude against Alcohol Use Disorder Identification Test (AUDIT) score separately for men ($p = .0034$, $r = .50$) and women ($p = .18$, $r = .26$) and for all subjects ($p = .0022$, $r = .39$). A slope test showed no difference between men and women ($p = .27$).




Figure 5. Mediation analysis. **(A)** The global positive (GP) alcohol expectancy score completely mediated the correlation between the thalamic response to alcohol vs. neutral cue and Alcohol Use Disorder Identification Test (AUDIT) score in men and women combined. **(B)** The alternative model where AUDIT score mediated the correlation between the GP score and thalamic response was not supported. **(C)** Likewise, the GP score completely mediated the correlation between the strength of thalamic superior frontal gyrus (SFG) connectivity and AUDIT score. **(D)** The alternative model where AUDIT score mediated the correlation between the GP score and connectivity strength was not supported. The *p* values associated with mediation are for the *a***b* path (see [Supplemental Methods](#)). **p* < .05. gPPI, general psychophysiological interaction; thal., thalamic.

neutral cues in association with AE. These clusters comprised primarily the dorsomedial nucleus and pulvinar in the area of the frontal and parietal association thalamus, according to a tractography study (38), integrating multiple modalities of sensory inputs to support cognition. Interestingly, although the thalamus did not show higher response to alcohol versus neutral cues, thalamic responses appeared to play an important role in distinguishing relapsers from nonrelapsers in treatment studies and in predicting individual vulnerability to relapse (50–53), as recently reviewed (54). Because AE is conducive to alcohol use, along with these earlier studies, the current results support thalamic cue response as a useful biomarker of problem drinking and alcohol addiction.

The thalamus interacted with the left SFG during cue exposure, and the magnitude of psychophysiological connectivity was also positively correlated with both the GP and AUDIT scores. As part of the prefrontal task network, the SFG has been widely implicated in inhibitory control and other executive functions. However, it is important to distinguish the roles of right- and left-hemispheric PFCs and the exact locale of cortical regions in these executive processes. Whereas the right-hemispheric PFC is involved in action control (55), the roles of the left prefrontal cortical regions appear to be more complex and in many instances antithetical to those of their right-hemispheric counterparts. For instance, in studies of the stop signal task, we showed that response speeding as compared with slowing, as a behavioral index of risk taking,

engaged predominantly left prefrontal and subcortical structures, whereas post-error slowing involved the right-hemispheric ventrolateral PFC (56,57).

Studies of electrical stimulation provided additional evidence in support of hemispheric differences in prefrontal cortical control of impulsive behavior. High-definition transcranial anodal direct current stimulation, which increased neuronal activity as compared with cathodal stimulation, of the left dorsolateral PFC at a location near the SFG (F3) increased risky choices in the Balloon Analog Risk Task (58). This finding was consistent with other reports that left anodal/right cathodal and right anodal/left cathodal transcranial anodal direct current stimulation of the dorsolateral PFC each increased (59,60) and diminished (60–62) risk-taking behavior. In a recent work, reward expectancy was associated with higher left ventrolateral PFC activity in a decision-making task (63). Together, these studies spoke to distinct roles of the left- and right-hemispheric PFC in facilitating approach and avoidance behavior. The current findings of increased thalamic–left SFG connectivity in association with higher AE may provide a new circuit marker of at-risk drinking.

In relation to alcohol misuse, young adults who used alcohol on a regular basis showed significantly higher activation than those who did not use alcohol regularly in the left SFG, despite similar behavioral performance, in an imaging study of the go/no-go task (64). Although interpreted as a compensatory process by the authors, the latter findings may reflect a distinct

role of the left SFG in impulsive response, as discussed earlier. In addicted individuals, alcohol dependence severity was negatively associated with activation in the right SFG during impulsive relative to delayed decisions in a delayed discounting task (65), again suggesting contrasting roles of the left and right SFG in cognitive control. It would be of interest to further explore the role of the thalamus, left SFG, and thalamic–prefrontal cortical connectivity in cue-elicited responses and whether these responses translate into alcohol-seeking behavior in a laboratory or real-life setting.

More broadly, the effects of expectancy on subjective experience of environmental stimuli have been most thoroughly investigated for placebo analgesia—expectations that a treatment will produce pain relief cause pain reduction even when the treatment itself is inert (66). In behavioral terms, individuals learn to expect a certain outcome and harness physiological resources to support such expectations (67). Indeed, as the current results showed, AE was highly correlated with cue-elicited craving during the alcohol blocks but not the neutral blocks. Notably, imaging studies showed that placebo effects involved reduced activation of the ACC and thalamus to pain stimulation (66). In contrast, the nocebo effect—negative expectation of the manipulation or treatment—was associated with increased activation of the thalamus, amygdala, and hippocampus (68). These results suggested flexible thalamic response to expectancy that associated individual variation in expectancy, acquired via conditioning or instrumental learning, with physiological effects.

A number of limitations need to be considered. First, it is worth noting that although a substantial number of participants reported an AUDIT score greater than 8, it remains to be seen whether the current findings would generalize to heavier drinking populations, including those with an alcohol use disorder. Second, cue-related regional activations did not appear to relate to acute craving rating. This may have reflected the nature of the experimental design; the alternating presentation of alcohol and neutral cue blocks may have masked the differences in craving elicited by alcohol and neutral cues. Furthermore, the alcohol cue pictures have not been validated independently, and some of the alcohol cue stimuli involved human faces known to elicit emotions that may complicate the measures of alcohol cue response. In particular, human faces elicited activation of occipital and temporal areas, including the fusiform gyrus (69), which may be conflated with cue-elicited activation. Third, it is important to note that although the thalamic effect size of cue response, GP score, and AUDIT score were correlated pairwise, their interrelationship remained to be clarified. Mediation analyses delineated how AE mediated the contribution of thalamic effect size to at-risk drinking, but the causal link among thalamic cue response, AE, and problem alcohol use needs to be confirmed in a longitudinal setting.

In conclusion, the current study demonstrated the cue-elicited thalamic activity and connectivity in association with AE. To our knowledge, these findings are the first to relate AE to cue-elicited brain responses and provide new markers of alcohol misuse. The etiologies of alcohol misuse are multifaceted. The current findings of altered thalamic activation and connectivity in relation to AE may complement work of other psychological processes and neural circuits to more fully capture the biological markers of at-risk alcohol consumption (70). Identifying these regional markers of at-risk alcohol use

may also facilitate research of effective treatment, such as noninvasive brain stimulation, of individuals with alcohol use disorders (71,72).

ACKNOWLEDGMENTS AND DISCLOSURES

This study was supported by National Institutes of Health Grant Nos. AA021449 (to CRL) and P50AA12870 (to JHK) and by the Department of Veterans Affairs National Center for Posttraumatic Stress Disorder (to JHK).

JHK has individual consultant agreements at less than \$10,000 per year with AstraZeneca Pharmaceuticals, Biogen, Idec, MA, Biomedisyn Corp., Bionomics Ltd. (Australia), Boehringer Ingelheim International, Concert Pharmaceuticals Inc., Heptares Therapeutics Ltd. (UK), Janssen Research & Development, L.E.K. Consulting, Otsuka America Pharmaceutical Inc., Spring Care Inc., Sunovion Pharmaceuticals Inc., Takeda Industries, and Taisho Pharmaceutical Co. Ltd. JHK is on the scientific advisory boards of Bioasis Technologies Inc., Biohaven Pharmaceuticals, Blackthorn Therapeutics Inc., Broad Institute of MIT and Harvard, Cadent Therapeutics, Lohocla Research Corp., Pfizer Pharmaceuticals, and Stanley Center for Psychiatric Research at the Broad Institute. JHK has stocks or stock options on ArRETT Neuroscience Inc., Blackthorn Therapeutics Inc., Biohaven Pharmaceuticals Medical Sciences, and Spring Care Inc. JHK receives income greater than \$10,000 as the editor of *Biological Psychiatry*. JHK has the following patents or patent applications: Dopamine and noradrenergic reuptake inhibitors in treatment of schizophrenia (U.S. Patent No. 5,447,948); Glutamate modulating agents in the treatment of mental disorders (U.S. Patent No. 8,778,979); Intranasal administration of ketamine to treat depression (No. 14/197,767 filed on March 5, 2014); Methods for treating suicidal ideation (No. 14/197,767 filed on March 5, 2014); Composition and methods to treat addiction (No. 61/973/961 filed on April 2, 2014); Treatment selection for major depressive disorder (USPTO Docket No. Y0087.70116US00 filed on June 3, 2016); Compounds, compositions and methods for treating or preventing depression and other diseases (No. 62/444,552 filed on January 10, 2017); Combination therapy for treating or preventing depression or other mood diseases [No. 047162-7177P1 (00754) filed on August 20, 2018]. JHK receives the following nonfederal research support: AstraZeneca Pharmaceuticals provides the drug, Saracatinib, for research related to the National Institute on Alcohol Abuse and Alcoholism grant “Center for Translational Neuroscience of Alcoholism; Pfizer Pharmaceuticals provides an investigational drug, PF-03463275, for research related to the National Institutes of Health grant “Translational Neuroscience Optimization of GlyT1 Inhibitor.” The other authors report no biomedical financial interests or potential conflicts of interest.

ARTICLE INFORMATION

From the Department of Psychiatry (SZho, SZha, JSI, WW, TML, RFL, JHK, CRL), Department of Medicine (HHC), and Department of Neuroscience (JHK, CRL), Yale University School of Medicine, and Interdepartmental Neuroscience Program (JHK, CRL), Yale University, New Haven; VA Connecticut Healthcare System (HHC), West Haven; Department of Computer Science & Engineering (JB), School of Engineering, University of Connecticut, Storrs; and Department of Community Medicine and Health Care (JB), School of Medicine, University of Connecticut Health Center, Farmington, Connecticut; and Department of Health Education & Behavior (RFL), University of Florida, Gainesville, Florida.

SZho and SZha contributed equally to this work.

Address correspondence to Chiang-shan R. Li, M.D., Ph.D., Connecticut Mental Health Center, S112, 34 Park Street, New Haven, CT 06519; E-mail: chiang-shan.li@yale.edu.

Received Sep 11, 2018; revised Nov 21, 2018; accepted Nov 28, 2018.

Supplementary material cited in this article is available online at <https://doi.org/10.1016/j.bpsc.2018.11.012>.

REFERENCES

1. American Psychiatric Association (1994): *Diagnostic and Statistical Manual of Mental Disorders IV*. Washington, DC: American Psychiatric Press.

2. Volkow ND, Li TK (2004): Drug addiction: The neurobiology of behaviour gone awry. *Nat Rev Neurosci* 5:963–970.
3. Noori HR, Cosa Linan A, Spanagel R (2016): Largely overlapping neuronal substrates of reactivity to drug, gambling, food and sexual cues: A comprehensive meta-analysis. *Eur Neuropsychopharmacol* 26:1419–1430.
4. Hittner JB (1997): Alcohol-related outcome expectancies: Construct overview and implications for primary and secondary prevention. *J Prim Prev* 17:297–314.
5. Tiffany ST (1999): Cognitive concepts of craving. *Alcohol Res Health* 23:215–224.
6. Marlatt GA (1985): Cognitive factors in the relapse process. In: Marlatt GGJ, editor. *Relapse Prevention*. New York: Guilford, 128–200.
7. Christiansen BA, Smith GT, Roehling PV, Goldman MS (1989): Using alcohol expectancies to predict adolescent drinking behavior after one year. *J Consult Clin Psychol* 57:93–99.
8. Brown SA (1985): Do alcohol expectancies mediate drinking patterns of adults? *J Consult Clin Psychol* 53:512–519.
9. Vilenne A, Quertemont E (2015): Explicit and implicit positive alcohol expectancies in problem and non-problem drinkers: Differences across age groups from young adolescence to adulthood. *Front Psychol* 6:1773.
10. Walther CAP, Pedersen SL, Cheong J, Molina BSG (2017): The role of alcohol expectancies in the associations between close friend, typical college student, and personal alcohol use. *Subst Use Misuse* 52:1656–1666.
11. Stangl BL, Vatsalya V, Zametkin MR, Cooke ME, Plawecki MH, O'Connor S, et al. (2017): Exposure-response relationships during free-access intravenous alcohol self-administration in nondependent drinkers: Influence of alcohol expectancies and impulsivity. *Int J Neuropsychopharmacol* 20:31–39.
12. Rose AK, Hobbs M, Drummond C (2013): Differentiating the contribution of pharmacological from alcohol expectancy effects to changes in subjective response and priming over successive drinks. *Alcohol Clin Exp Res* 37:687–695.
13. Claus ED, Ewing SW, Filbey FM, Sabbineni A, Hutchison KE (2011): Identifying neurobiological phenotypes associated with alcohol use disorder severity. *Neuropsychopharmacology* 36:2086–2096.
14. Kareken DA, Claus ED, Sabri M, Dziedzic M, Kosobud AE, Radnovich AJ, et al. (2004): Alcohol-related olfactory cues activate the nucleus accumbens and ventral tegmental area in high-risk drinkers: Preliminary findings. *Alcohol Clin Exp Res* 28:550–557.
15. Sjoerds Z, van den Brink W, Beekman AT, Penninx BW, Veltman DJ (2014): Cue reactivity is associated with duration and severity of alcohol dependence: An fMRI study. *PLoS One* 9:e84560.
16. Schacht JP, Anton RF, Myrick H (2013): Functional neuroimaging studies of alcohol cue reactivity: A quantitative meta-analysis and systematic review. *Addict Biol* 18:121–133.
17. Christiansen P, Rose AK, Cole JC, Field M (2013): A comparison of the anticipated and pharmacological effects of alcohol on cognitive bias, executive function, craving and ad-lib drinking. *J Psychopharmacol* 27:84–92.
18. Gundersen H, Specht K, Gruner R, Ersland L, Hugdahl K (2008): Separating the effects of alcohol and expectancy on brain activation: An fMRI working memory study. *NeuroImage* 42:1587–1596.
19. Kegeles LS, Horga G, Ghazzaoui R, Rosengard R, Ojeil N, Xu X, et al. (2018): Enhanced striatal dopamine release to expectation of alcohol: A potential risk factor for alcohol use disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging* 3:591–598.
20. McBride D, Barrett SP, Kelly JT, Aw A, Dagher A (2006): Effects of expectancy and abstinence on the neural response to smoking cues in cigarette smokers: An fMRI study. *Neuropsychopharmacology* 31:2728–2738.
21. Wilson SJ, Sayette MA, Delgado MR, Fiez JA (2005): Instructed smoking expectancy modulates cue-elicited neural activity: A preliminary study. *Nicotine Tob Res* 7:637–645.
22. Christiansen P, Jennings E, Rose AK (2016): Anticipated effects of alcohol stimulate craving and impair inhibitory control. *Psychol Addict Behav* 30:383–388.
23. Pulido C, Anderson KG, Armstead AG, Brown SA, Tapert SF (2009): Family history of alcohol-use disorders and spatial working memory: Effects on adolescent alcohol expectancies. *J Stud Alcohol Drugs* 70:87–91.
24. Deckel AW, Hesselbrock V, Bauer L (1995): Relationship between alcohol-related expectancies and anterior brain functioning in young men at risk for developing alcoholism. *Alcohol Clin Exp Res* 19:476–481.
25. Ide JS, Zhornitsky S, Hu S, Zhang S, Krystal JH, Li CR (2017): Sex differences in the interacting roles of impulsivity and positive alcohol expectancy in problem drinking: A structural brain imaging study. *Neuroimage Clin* 14:750–759.
26. Zhornitsky S, Ide SJ, Wang W, Choa H, Zhang S, Hu S, et al. (2018): Problem drinking, alcohol expectancy and thalamic resting-state functional connectivity in non-dependent adult drinkers. *Brain Connect* 8:487–502.
27. First MS R, Williams J, Gibbon M (1995): *Structured Clinical Interview for DSM-IV (SCID)*. Washington, DC: American Psychiatric Press.
28. Babor TF, Higgins-Biddle JC, Saunders JB, Monteiro MG (2001): *The Alcohol Use Disorders Identification Test*. Geneva: World Health Organization, Department of Mental Health and Substance Dependence.
29. George WH, Frone MR, Cooper ML, Russell M, Skinner JB, Windle M (1995): A revised Alcohol Expectancy Questionnaire: Factor structure confirmation, and invariance in a general population sample. *J Stud Alcohol* 56:177–185.
30. Heatherton TF, Kozlowski LT, Frecker RC, Fagerström KO (1991): The Fagerström Test for Nicotine Dependence: A revision of the Fagerström Tolerance Questionnaire. *Br J Addict* 86:1119–1127.
31. Friston K, Ashburner J, Frith C, Polone J, Heather J, Frackowiak R (1995): *Spatial registration and normalization of images*. Hum Brain Mapp 2:165–189.
32. Ashburner J, Friston KJ (1999): Nonlinear spatial normalization using basis functions. *Hum Brain Mapp* 7:254–266.
33. McLaren DG, Ries ML, Xu G, Johnson SC (2012): A generalized form of context-dependent psychophysiological interactions (gPPI): A comparison to standard approaches. *NeuroImage* 61:1277–1286.
34. Zar JH (1999): *Biostatistical Analysis*, 4th ed. Englewood Cliffs, NJ: Prentice Hall.
35. Schacht JP, Anton RF, Randall PK, Li X, Henderson S, Myrick H (2014): Varenicline effects on drinking, craving and neural reward processing among non-treatment-seeking alcohol-dependent individuals. *Psychopharmacology* 231:3799–3807.
36. van Holst RJ, Clark L, Veltman DJ, van den Brink W, Goudriaan AE (2014): Enhanced striatal responses during expectancy coding in alcohol dependence. *Drug Alcohol Depend* 142:204–208.
37. Baker TE, Castellanos-Ryan N, Schumann G, Cattel A, Flor H, Nees F, et al. (2019): Modulation of orbitofrontal-striatal reward activity by dopaminergic functional polymorphisms contributes to a predisposition to alcohol misuse in early adolescence. *Psychol Med* 49:801–810.
38. Behrens TE, Johansen-Berg H, Woolrich MW, Smith SM, Wheeler-Kingshott CA, Boulby PA, et al. (2003): Non-invasive mapping of connections between human thalamus and cortex using diffusion imaging. *Nat Neurosci* 6:750–757.
39. Mink JW (1996): The basal ganglia: Focused selection and inhibition of competing motor programs. *Prog Neurobiol* 50:381–425.
40. Camchong J, Stenger VA, Fein G (2013): Resting-state synchrony in short-term versus long-term abstinent alcoholics. *Alcohol Clin Exp Res* 37:794–803.
41. Alexander GE, DeLong MR, Strick PL (1986): Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu Rev Neurosci* 9:357–381.
42. Chanraud S, Martelli C, Delain F, Kostogianni N, Douaud G, Aubin HJ, et al. (2007): Brain morphometry and cognitive performance in detoxified alcohol-dependents with preserved psychosocial functioning. *Neuropsychopharmacology* 32:429–438.
43. Pitel AL, Segabin SH, Ritz L, Eustache F, Beaunieux H (2015): Thalamic abnormalities are a cardinal feature of alcohol-related brain dysfunction. *Neurosci Biobehav Rev* 54:38–45.

44. Kim E, Ku J, Namkoong K, Lee W, Lee KS, Park JY, *et al.* (2009): Mammillothalamic functional connectivity and memory function in Wernicke's encephalopathy. *Brain* 132:369–376.
45. Sherman SM (2016): Thalamus plays a central role in ongoing cortical functioning. *Nat Neurosci* 19:533–541.
46. Muller-Oehring EM, Jung YC, Pfefferbaum A, Sullivan EV, Schulte T (2015): The resting brain of alcoholics. *Cereb Cortex* 25:4155–4168.
47. Shokri-Kojori E, Tomasi D, Wiers CE, Wang GJ, Volkow ND (2017): Alcohol affects brain functional connectivity and its coupling with behavior: Greater effects in male heavy drinkers. *Mol Psychiatry* 22:1185–1195.
48. Wang J, Fan Y, Dong Y, Ma M, Ma Y, Dong Y, *et al.* (2016): Alterations in brain structure and functional connectivity in alcohol dependent patients and possible association with impulsivity. *PLoS One* 11:e161956.
49. Jasinska AJ, Stein EA, Kaiser J, Naumer MJ, Yalachkov Y (2014): Factors modulating neural reactivity to drug cues in addiction: A survey of human neuroimaging studies. *Neurosci Biobehav Rev* 38:1–16.
50. Grusser SM, Wrase J, Klein S, Hermann D, Smolka MN, Ruf M, *et al.* (2004): Cue-induced activation of the striatum and medial prefrontal cortex is associated with subsequent relapse in abstinent alcoholics. *Psychopharmacology* 175:296–302.
51. McClemon FJ, Hiott FB, Liu J, Salley AN, Behm FM, Rose JE (2007): Selectively reduced responses to smoking cues in amygdala following extinction-based smoking cessation: Results of a preliminary functional magnetic resonance imaging study. *Addict Biol* 12:503–512.
52. Janes AC, Pizzagalli DA, Richardt S, Frederick BD, Chuzy S, Pachas G, *et al.* (2010): Brain reactivity to smoking cues prior to smoking cessation predicts ability to maintain tobacco abstinence. *Biol Psychiatry* 67:722–729.
53. Versace F, Engelmann JM, Robinson JD, Jackson EF, Green CE, Lam CY, *et al.* (2014): Prequit fMRI responses to pleasant cues and cigarette-related cues predict smoking cessation outcome. *Nicotine Tob Res* 16:697–708.
54. Courtney KE, Schacht JP, Hutchison K, Roche DJ, Ray LA (2016): Neural substrates of cue reactivity: Association with treatment outcomes and relapse. *Addict Biol* 21:3–22.
55. Hu S, Ide JS, Zhang S, Li CR (2016): The right superior frontal gyrus and individual variation in proactive control of impulsive response. *J Neurosci* 36:12688–12696.
56. Li CS, Huang C, Yan P, Paliwal P, Constable RT, Sinha R (2008): Neural correlates of post-error slowing during a stop signal task: A functional magnetic resonance imaging study. *J Cogn Neurosci* 20:1021–1029.
57. Li CS, Chao HH, Lee TW (2009): Neural correlates of speeded as compared with delayed responses in a stop signal task: An indirect analog of risk taking and association with an anxiety trait. *Cereb Cortex* 19:839–848.
58. Guo H, Zhang Z, Da S, Sheng X, Zhang X (2018): High-definition transcranial direct current stimulation (HD-tDCS) of left dorsolateral prefrontal cortex affects performance in Balloon Analogue Risk Task (BART). *Brain Behav* 8:e884.
59. Boggio PS, Campanha C, Valasek CA, Fecteau S, Pascual-Leone A, Fregni F (2010): Modulation of decision-making in a gambling task in older adults with transcranial direct current stimulation. *Eur J Neurosci* 31:593–597.
60. Luo J, Ye H, Zheng H, Chen S, Huang D (2017): Modulating the activity of the dorsolateral prefrontal cortex by tDCS alters distributive decisions behind the veil of ignorance via risk preference. *Behav Brain Res* 328:70–80.
61. Gilmore CS, Dickmann PJ, Nelson BG, Lamberty GJ, Lim KO (2018): Transcranial direct current stimulation (tDCS) paired with a decision-making task reduces risk-taking in a clinically impulsive sample. *Brain Stimul* 11:302–309.
62. Fecteau S, Knoch D, Fregni F, Sultani N, Boggio P, Pascual-Leone A (2007): Diminishing risk-taking behavior by modulating activity in the prefrontal cortex: A direct current stimulation study. *J Neurosci* 27:12500–12505.
63. Chase HW, Fournier JC, Bertocci MA, Greenberg T, Aslam H, Stiffler R, *et al.* (2017): A pathway linking reward circuitry, impulsive sensation-seeking and risky decision-making in young adults: Identifying neural markers for new interventions. *Transl Psychiatry* 7:e1096.
64. Hatchard T, Mioduszewski O, Fall C, Byron-Alhassan A, Fried P, Smith AM (2017): Neural impact of low-level alcohol use on response inhibition: An fMRI investigation in young adults. *Behav Brain Res* 329:12–19.
65. Lim AC, Cservenka A, Ray LA (2017): Effects of alcohol dependence severity on neural correlates of delay discounting. *Alcohol Alcoholism* 52:506–515.
66. Atlas LY, Wager TD (2012): How expectations shape pain. *Neurosci Lett* 520:140–148.
67. Beecher HK (1955): The powerful placebo. *JAMA* 159:1602–1606.
68. Jensen KB, Kaptchuk TJ, Chen X, Kirsch I, Ingvar M, Gollub RL, *et al.* (2015): A neural mechanism for nonconscious activation of conditioned placebo and nocebo responses. *Cereb Cortex* 25:3903–3910.
69. Furl N, Garrido L, Dolan RJ, Driver J, Duchaine B (2011): Fusiform gyrus face selectivity relates to individual differences in facial recognition ability. *J Cogn Neurosci* 23:1723–1740.
70. Grodin EN, Sussman L, Sundby K, Brennan GM, Diazgranados N, Heilig M, *et al.* (2018): Neural correlates of compulsive alcohol seeking in heavy drinkers. *Biol Psychiatry Cogn Neurosci Neuroimaging* 3:1022–1031.
71. Loheswaran G, Barr MS, Rajji TK, Zomorodi R, Le Foll B, Daskalakis ZJ (2016): Brain stimulation in alcohol use disorders: Investigational and therapeutic tools. *Biol Psychiatry Cogn Neurosci Neuroimaging* 1:5–13.
72. Hanlon CA, Dowdle LT, Correia B, Mithoefer O, Kearney-Ramos T, Lench D, *et al.* (2017): Left frontal pole theta burst stimulation decreases orbitofrontal and insula activity in cocaine users and alcohol users. *Drug Alcohol Depend* 178:310–317.